RESEARCH

Open Access

On recursions for coefficients of mock theta functions

Song Heng Chan¹, Renrong Mao² and Robert Osburn^{3,4*}

*Correspondence:

robert.osburn@ucd.ie;

osburn@ihes.fr

³School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland

⁴IHÉS, Le Bois-Marie, 35, route de Chartres, F-91440 Bures-sur-Yvette, France

Full list of author information is available at the end of the article

Abstract

We use a generalized Lambert series identity due to the first author to present q -series proofs of recent results of Imamoğlu, Raum and Richter concerning recursive formulas for the coefficients of two 3rd order mock theta functions. Additionally, we discuss an application of this identity to other mock theta functions.

2010 Mathematics Subject Classification: Primary: 33D15; Secondary: 11F30

Keywords: Lambert series, Mock theta functions, q -series identities

1 Introduction

In [8], the first author proved the following generalized Lambert series identity:

$$\frac{[a_1, \dots, a_r]_\infty (q)_\infty^2}{[b_1, \dots, b_s]_\infty} = \frac{[a_1/b_1, \dots, a_r/b_1]_\infty}{[b_2/b_1, \dots, b_s/b_1]_\infty} \sum_{k=-\infty}^{\infty} \frac{(-1)^{(s-r)k} q^{(s-r)k(k+1)/2}}{1 - b_1 q^k} \left(\frac{a_1 \cdots a_r b_1^{s-r-1}}{b_2 \cdots b_s} \right)^k + \text{idem}(b_1; b_2, \dots, b_s), \quad (1.1)$$

valid for nonnegative integers $r < s$. Here and throughout, we use the following standard q -hypergeometric notation

$$(a)_n := (a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k)$$

$$(a_1, \dots, a_m)_n := (a_1, \dots, a_m; q)_n := (a_1)_n \cdots (a_m)_n$$

$$[a_1, \dots, a_m]_n := [a_1, \dots, a_m; q]_n = (a_1, q/a_1, \dots, a_m, q/a_m)_n$$

valid for $n \in \mathbb{N} \cup \{\infty\}$ and $F(a_1, a_2, \dots, a_m) + \text{idem}(a_1; a_2, \dots, a_n)$ to denote the sum

$$\sum_{i=1}^n F(a_i, a_2, \dots, a_{i-1}, a_1, a_{i+1}, \dots, a_m)$$

where the i th term of the sum is obtained from the first by interchanging a_1 and a_i .

Identity (1.1) is of interest for several reasons. For example, it generalizes a key identity used by Atkin and Swinnerton-Dyer [4] in their proof of Dyson's conjectures on the rank of a partition. Also, (1.1) played a crucial role in the construction of quasimock theta functions [7] and rank-crank PDE's [10], in proving congruences for the mock theta function $\varphi(q)$ [9], Appell-Lerch sums [12], spt-type functions [17] and partition pairs [18] and

in obtaining identities for generating functions of other types of partition pairs [11] and various rank differences [19–22].

Recently, Imamoğlu, Raum and Richter [16] proved some intriguing results concerning recursive formulas for the coefficients of the 3rd order mock theta functions

$$f(q) := \sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q)_n^2} = \sum_{n=0}^{\infty} c(f; n) q^n$$

and

$$\omega(q) := \sum_{n=0}^{\infty} \frac{q^{2n(n+1)}}{(q; q^2)_{n+1}^2} = \sum_{n=0}^{\infty} c(\omega; n) q^n.$$

Namely, if $\sigma(n) := \sum_{0 < d|n} d$, $\text{sgn}^+(n) := \text{sgn}(n)$ for $n \neq 0$ and $\text{sgn}^+(0) := 1$ and

$$d(N, \tilde{N}, t, \tilde{t}) := \text{sgn}^+(N) \text{sgn}^+(\tilde{N}) (|N + t| - |\tilde{N} + \tilde{t}|),$$

then we have the following (see Theorems 1 and 9 in [16], slightly rewritten).

Theorem 1.1. *For a fixed $n \in \mathbb{Z}^+$ and for any $a, b \in \mathbb{Z}$, set $N := \frac{1}{6}(-3a + b - 1)$ and $\tilde{N} := \frac{1}{6}(3a + b - 1)$. Then*

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + m \leq 2n}} \left(m + \frac{1}{6}\right) c\left(f; n - \frac{3}{2}m^2 - \frac{1}{2}m\right) = \frac{4}{3}\sigma(n) - \frac{16}{3}\sigma\left(\frac{n}{2}\right) - 2 \sum_{\substack{a, b \in \mathbb{Z} \\ ab=2n \\ 6|3a+b-1}} d\left(N, \tilde{N}, \frac{1}{6}, \frac{1}{6}\right). \quad (1.2)$$

For a fixed $n \in \mathbb{Z}^+$ and for any $a, b \in \mathbb{Z}$, set $N := \frac{1}{12}(3a - b - 2)$ and $\tilde{N} := \frac{1}{12}(3a + b - 4)$. Then

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + 2m \leq n}} \left(m + \frac{1}{3}\right) c\left(f; \frac{n}{2} - \frac{3}{2}m^2 - m\right) = -2 \sum_{\substack{a, b \in \mathbb{Z} \\ ab=4n+1 \\ 12|3a-b-2}} d\left(N, \tilde{N}, \frac{1}{6}, \frac{1}{3}\right). \quad (1.3)$$

Theorem 1.2. *For a fixed $n \in \mathbb{Z}^+$ and for any $a, b \in \mathbb{Z}$, set $N := \frac{1}{12}(3a - b - 4)$ and $\tilde{N} := \frac{1}{12}(3a + b - 2)$. Then*

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + 2m \leq n}} \left(m + \frac{1}{6}\right) c\left(\omega; n - 3m^2 - m\right) = (-1)^{n+1} \sum_{\substack{a, b \in \mathbb{Z} \\ ab=4n+3 \\ 12|3a-b-4}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{6}\right). \quad (1.4)$$

For a fixed $n \in \mathbb{Z}^+$ and for any $a, b \in \mathbb{Z}$, set $N := \frac{1}{6}(a - 3b - 2)$ and $\tilde{N} := \frac{1}{6}(a + 3b - 2)$, and let

$$R_n := \begin{cases} \frac{2}{3} \left(\sigma\left(\frac{n}{4}\right) - \sigma\left(\frac{n}{2}\right) \right), & \text{if } n \text{ is even;} \\ \frac{1}{3} \sigma(n), & \text{if } n \text{ is odd.} \end{cases}$$

Then

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + 2m + 1 \leq n}} \left(m + \frac{1}{3}\right) c\left(\frac{\omega(q) + \omega(-q)}{2}; n - 3m^2 - 2m - 1\right) = R_n - \sum_{\substack{a, b \in \mathbb{Z} \\ ab=n \\ 12|a-3b-8}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right), \quad (1.5)$$

and

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + 2m + 1 \leq n}} \left(m + \frac{1}{3}\right) c\left(\frac{\omega(q) - \omega(-q)}{2}; n - 3m^2 - 2m - 1\right) = -R_n + \sum_{\substack{a, b \in \mathbb{Z} \\ ab = n \\ 12|a - 3b - 2}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right). \quad (1.6)$$

The identities (1.2)–(1.6) were proven in [16] by applying holomorphic projection to the tensor product of a vector-valued harmonic weak Maass form of weight $1/2$ and vector-valued modular form of weight $3/2$ [13, 23]. In Remark 1, ii) of [16], it was stated that these identities “can sometimes also be furnished by Appell sums because these are typically expressible in terms of divisors. However, it is not clear whether Theorem 1 and 9 could be obtained using this idea”.

Motivated by this remark, the main purpose of this paper is explain how (1.1) can be used to give a q -series proof of these identities. The idea is to compare the coefficients of q^n in identities which express a modular form times either $f(q)$ or $\omega(q)$ (Appell-Lerch sums) in terms of Lambert series (divisor sums). Specifically, (1.2) and (1.3) will basically follow from (2.8) and (3.11), (1.4) from (3.17), (1.5) and (1.6) from (3.24) and (2.9), respectively.

The paper is organized as follows. In Section 2, we discuss some preliminary q -series identities. In Section 3, we prove Theorems 1.1 and 1.2. In Section 4, we discuss another application of (1.1) to other mock theta functions.

2 Preliminaries

To prove (1.2), (1.5) and (1.6), we need the following two results.

Lemma 2.1. *We have*

$$\begin{aligned} \frac{(q)_\infty^2}{(-q)_\infty^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)/2}}{1 - xq^n} &= 4x \frac{(q)_\infty^2 (q^2; q^2)_\infty^2}{[-x]_\infty [x^2; q^2]_\infty} + 2 \sum_{n=-\infty}^{\infty} \frac{q^{n(3n+1)/2}}{(1 + xq^n)^2} \\ &\quad + \sum_{n=-\infty}^{\infty} \frac{(6n-1)q^{n(3n+1)/2}}{1 + xq^n}, \end{aligned} \quad (2.1)$$

$$\begin{aligned} q \frac{(q^2; q^2)_\infty^2}{(-q; q^2)_\infty^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)}}{1 - xq^{2n}} &= \frac{q}{x} \frac{(q^2; q^2)_\infty^4 (-q; q^2)_\infty^2}{[x, -xq, -xq; q^2]_\infty} + \sum_{n=-\infty}^{\infty} \frac{q^{3n^2}}{(1 + xq^{2n-1})^2} \\ &\quad + \sum_{n=-\infty}^{\infty} \frac{(3n-1)q^{3n^2}}{1 + xq^{2n-1}}. \end{aligned} \quad (2.2)$$

Proof. Beginning with the case $r = 0$, $s = 3$ in (1.1), setting $(b_1, b_2, b_3) = (x, -xq/a, -ax)$ and multiplying by $a[-a, 1/a^2]_\infty$, we find

$$\begin{aligned} a \frac{(q)_\infty^2 [-a, 1/a^2]_\infty}{[x, -xq/a, -ax]_\infty} &= \frac{[1/a^2]_\infty}{[-1/a]_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)/2}}{1 - xq^n} + \sum_{n=-\infty}^{\infty} \frac{q^{n(3n+1)/2} a^{-3n-1}}{1 + xq^n/a} \\ &\quad - \sum_{n=-\infty}^{\infty} \frac{q^{n(3n+1)/2} a^{3n}}{1 + axq^n}. \end{aligned} \quad (2.3)$$

Differentiating (2.3) with respect to a and letting $a \rightarrow 1$, we find that

$$4 \frac{(q)_\infty^4 (-q)_\infty^2}{[x, -xq, -x]_\infty} = \frac{(q)_\infty^2}{(-q)_\infty^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)/2}}{1 - xq^n} - \sum_{n=-\infty}^{\infty} \frac{q^{n(3n+1)/2} [1 + 3n(1 + xq^n)]}{(1 + xq^n)^2} \\ - \sum_{n=-\infty}^{\infty} \frac{q^{n(3n+1)/2} [1 + (3n-1)(1 + xq^n)]}{(1 + xq^n)^2}. \quad (2.4)$$

Rearranging and simplifying, we find that (2.4) is equivalent to (2.1).

Again, for the case $r = 0, s = 3$ in (1.1), setting $(b_1, b_2, b_3) = (x, -x/a, -ax)$, and multiplying by $[-a, 1/a^2]_\infty$, we find

$$\frac{(q)_\infty^2 [-a, 1/a^2]_\infty}{[x, -x/a, -ax]_\infty} = \frac{[1/a^2]_\infty}{[-1/a]_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)/2}}{1 - xq^n} - \sum_{n=-\infty}^{\infty} \frac{q^{3n(n+1)/2} a^{-3n-2}}{1 + xq^n/a} \\ + \sum_{n=-\infty}^{\infty} \frac{q^{3n(n+1)/2} a^{3n+1}}{1 + axq^n}. \quad (2.5)$$

Replacing q by q^2 and setting $a = 1/bq$ in (2.5), we find that

$$\frac{(q^2; q^2)_\infty^2 [-1/(bq), b^2 q^2; q^2]_\infty}{[x, -xbq, -x/(bq); q^2]_\infty} = \frac{[b^2 q^2; q^2]_\infty}{[-bq; q^2]_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)}}{1 - xq^{2n}} - \sum_{n=-\infty}^{\infty} \frac{q^{3n^2-1} b^{3n-1}}{1 + x b q^{2n-1}} \\ + \sum_{n=-\infty}^{\infty} \frac{q^{3n^2-1} b^{-3n-1}}{1 + x q^{2n-1}/b}. \quad (2.6)$$

Differentiating (2.6) with respect to b and letting $b \rightarrow 1$, we arrive at

$$2 \frac{(q^2; q^2)_\infty^4 [-1/q; q^2]_\infty}{[x, -xq, -x/q; q^2]_\infty} = 2 \frac{(q^2; q^2)_\infty^2}{[-q; q^2]_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)}}{1 - xq^{2n}} \\ - \sum_{n=-\infty}^{\infty} \frac{q^{3n^2-1} [1 + (3n-2)(1 + xq^{2n-1})]}{(1 + xq^{2n-1})^2} \\ - \sum_{n=-\infty}^{\infty} \frac{q^{3n^2-1} [1 + 3n(1 + xq^{2n-1})]}{(1 + xq^{2n-1})^2}. \quad (2.7)$$

Rearranging and simplifying, we find that (2.7) is equivalent to (2.2). $\square$

Corollary 2.2. *We have*

$$\frac{(q)_\infty^3}{(-q)_\infty^2} f(q) = -4 \sum_{n=1}^{\infty} \frac{q^n}{(1 - q^n)^2} - 16 \sum_{n=1}^{\infty} \frac{q^{2n}}{(1 - q^{2n})^2} + 1 + 4 \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{q^{n(3n+1)/2}}{(1 - q^n)^2} \\ + 2 \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(6n-1)q^{n(3n+1)/2}}{1 - q^n}, \quad (2.8)$$

$$q \frac{(q^2; q^2)_\infty^3}{(-q; q^2)_\infty^2} \omega(-q) = \sum_{n=1}^{\infty} \frac{q^{2n-1}}{(1 + q^{2n-1})^2} - 2 \sum_{n=1}^{\infty} \frac{q^{2n}}{(1 - q^{2n})^2} + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{q^{3n^2}}{(1 - q^{2n})^2} \\ + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(3n-1)q^{3n^2}}{1 - q^{2n}}. \quad (2.9)$$

Proof. For (2.8), multiply both sides of (2.1) by $(1+x)(1+1/x)$, differentiate twice with respect to x , set $x = -1$ and use

$$f(q) = \frac{2}{(q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)/2}}{1+q^n}. \quad (2.10)$$

Here, we have set $x = 1$ in the identity (see (12.2.5) in [2])

$$\sum_{n=0}^{\infty} \frac{q^{2n^2+2n}}{(xq; q^2)_{n+1}(q/x; q^2)_{n+1}} = \frac{1}{(q^2; q^2)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+3n}}{1-xq^{2n+1}}.$$

For (2.9), multiply both sides of (2.2) by $(1+q/x)(1+x/q)$, differentiate twice with respect to x , set $x = -q$ and use

$$\omega(q) = \frac{1}{(q^2; q^2)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+3n}}{1-q^{2n+1}}. \quad (2.11)$$

The latter follows from taking $x = -1$ in the identity (see (12.2.3) in [2])

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(xq; q)_n(q/x; q)_n} = \frac{(1-x)}{(q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)/2}}{1-xq^n}.$$

□

To prove (1.3), (1.5) and (1.6), we also need the following.

Theorem 2.3. *For integers l and j , we have*

$$\begin{aligned} & \frac{[-q^{l+j}, q^{2l}; q^{6l}]_\infty (q^{6l}; q^{6l})_\infty^2}{[-q^j, q^l, -q^{2l+j}; q^{6l}]_\infty} \left\{ j + 2 \sum_{m=0}^{\infty} \left(\frac{q^{6lm+l}}{1-q^{6lm+l}} - \frac{q^{6lm+5l}}{1-q^{6lm+5l}} + \frac{q^{6lm+l+j}}{1+q^{6lm+l+j}} \right. \right. \\ & \quad \left. \left. - \frac{q^{6lm+5l-j}}{1+q^{6lm+5l-j}} - 2 \frac{q^{6lm+3l-j}}{1+q^{6lm+3l-j}} + 2 \frac{q^{6lm+3l+j}}{1+q^{6lm+3l+j}} \right) \right\} \\ & = \sum_{n=-\infty}^{\infty} \frac{(6n+2+j)(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} \\ & \quad + \sum_{n=-\infty}^{\infty} \frac{(6n+j)(-1)^n q^{3ln(n+1)-ln}}{1+q^{6ln+j}} \\ & \quad + \frac{4(q^{2l}; q^{2l})_\infty^2}{(q^l; q^{2l})_\infty^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+2ln+2l}}{1+q^{6ln+j+3l}}. \end{aligned} \quad (2.12)$$

Proof. First, we need to prove the following:

$$\begin{aligned} & \frac{[-q^{l+j}, q^{2l}; q^{6l}]_\infty (q^{6l}; q^{6l})_\infty^2}{[-q^j, q^l, -q^{2l+j}; q^{6l}]_\infty} \sum_{m=0}^{\infty} \left(\frac{q^{6lm+l}}{1-q^{6lm+l}} - \frac{q^{6lm+5l}}{1-q^{6lm+5l}} + \frac{q^{6lm+l+j}}{1+q^{6lm+l+j}} - \frac{q^{6lm+5l-j}}{1+q^{6lm+5l-j}} \right) \\ & = \left(2 \sum_{m=0}^{\infty} \left(\frac{q^{6lm+l}}{1-q^{6lm+l}} - \frac{q^{6lm+5l}}{1-q^{6lm+5l}} \right) \times \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} \right) \\ & \quad + \sum_{n=-\infty}^{\infty} \frac{(n+1)(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} + \sum_{n=-\infty}^{\infty} \frac{n(-1)^n q^{3ln(n+1)-ln}}{1+q^{6ln+j}} \end{aligned} \quad (2.13)$$

and

$$\begin{aligned} & \frac{[-q^{l+j}, q^{2l}; q^{6l}]_{\infty} (q^{6l}; q^{6l})_{\infty}^2}{[-q^j, q^l, -q^{2l+j}; q^{6l}]_{\infty}} \sum_{m=0}^{\infty} \left(\frac{q^{6lm+3l-j}}{1+q^{6lm+3l-j}} - \frac{q^{6lm+3l+j}}{1+q^{6lm+3l+j}} \right) \\ &= \left(\sum_{m=0}^{\infty} \left(\frac{q^{6lm+l}}{1-q^{6lm+l}} - \frac{q^{6lm+5l}}{1-q^{6lm+5l}} \right) \times \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} \right) \\ & \quad - \sum_{n=-\infty}^{\infty} \frac{n(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} - \sum_{n=-\infty}^{\infty} \frac{n(-1)^n q^{3ln(n+1)-ln}}{1+q^{6ln+j}} \\ & \quad - \frac{[q^{-2l}, q^{2l}; q^{6l}]_{\infty} (q^{6l}; q^{6l})_{\infty}^2}{[q^{3l}, q^l, q^{-l}; q^{6l}]_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+2ln+3l}}{1+q^{6ln+j+3l}}. \end{aligned} \quad (2.14)$$

As the proofs of (2.13) and (2.14) are similar, we only give details for (2.14). Replacing q, a_1, a_2, b_1, b_2 and b_3 by $q^{6l}, -q^{j+3l}, -q^{j+l}, -bq^{j+3l}, -q^{j+2l}$ and $-q^j$, respectively, in (1.1) with $r = 2, s = 3$ and after rearranging, we obtain

$$\begin{aligned} & \frac{[-q^{3l+j}, -q^{j+l}, q^{2l}, bq^{3l}; q^{6l}]_{\infty} (q^{6l}; q^{6l})_{\infty}^2}{[-q^{j+2l}, -q^j, q^{3l}, q^l, -bq^{j+3l}; q^{6l}]_{\infty}} = \frac{b[1/b, q^{-2l}/b, q^{2l}; q^{6l}]_{\infty}}{[q^{3l}, q^l, q^{-l}/b; q^{6l}]_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1)^{n+1} q^{3ln(n+1)+2ln+3l}}{1+bq^{6ln+j+3l}} \\ & \quad + \frac{[q^l, bq^{3l}; q^{6l}]_{\infty}}{[q^{3l}, bq^l; q^{6l}]_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1/b)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} \\ & \quad + \sum_{n=-\infty}^{\infty} \frac{(-1/b)^n q^{3ln(n+1)-ln}}{1+q^{6ln+j}}. \end{aligned} \quad (2.15)$$

Applying the operator $\left. \frac{d}{db} \right|_{b=1}$ on both sides of (2.15) and simplifying, we obtain (2.14). If we multiply four times (2.14), then subtract from twice (2.13), we find

$$\begin{aligned} & \frac{[-q^{l+j}, q^{2l}; q^{6l}]_{\infty} (q^{6l}; q^{6l})_{\infty}^2}{[-q^j, q^l, -q^{2l+j}; q^{6l}]_{\infty}} \sum_{m=0}^{\infty} \left\{ 2 \left(\frac{q^{6lm+l}}{1-q^{6lm+l}} - \frac{q^{6lm+5l}}{1-q^{6lm+5l}} + \frac{q^{6lm+l+j}}{1+q^{6lm+l+j}} - \frac{q^{6lm+5l-j}}{1+q^{6lm+5l-j}} \right) \right. \\ & \quad \left. - 4 \left(\frac{q^{6lm+3l-j}}{1+q^{6lm+3l-j}} - \frac{q^{6lm+3l+j}}{1+q^{6lm+3l+j}} \right) \right\} \\ &= \sum_{n=-\infty}^{\infty} \frac{(6n+2)(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} + \sum_{n=-\infty}^{\infty} \frac{6n(-1)^n q^{3ln(n+1)-ln}}{1+q^{6ln+j}} \\ & \quad + \frac{4(q^{2l}; q^{2l})_{\infty}^2}{(q^l; q^{2l})_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+2ln+2l}}{1+q^{6ln+j+3l}}. \end{aligned} \quad (2.16)$$

Replacing q, a_1, b_1, b_2 by $q^{6l}, -q^{l+j}, -q^j, -q^{j+2l}$ in (1.1) with $r = 1, s = 2$, we obtain

$$\frac{[-q^{l+j}, q^{2l}; q^{6l}]_{\infty} (q^{6l}; q^{6l})_{\infty}^2}{[-q^j, q^l, -q^{2l+j}; q^{6l}]_{\infty}} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+ln+l}}{1+q^{6ln+j+2l}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)-ln}}{1+q^{6ln+j}} \quad (2.17)$$

Finally, (2.16) plus j times (2.17) yields (2.12). $\square$

To prove (1.4), we need the following result whose proof is analogous to that of Theorem 2.3 and thus is omitted.

Theorem 2.4. *We have*

$$\begin{aligned} & \frac{[q^{2l+j}, q^{4l}, q^{6l}]_{\infty} (q^{6l}; q^{6l})_{\infty}^2}{[-q^j, -q^{2l}, -q^{4l+j}; q^{6l}]_{\infty}} \left\{ j + \sum_{m=0}^{\infty} 2 \left(\frac{q^{6lm+2l}}{1+q^{6lm+2l}} - \frac{q^{6lm+4l}}{1+q^{6lm+4l}} - \frac{q^{6lm+2l+j}}{1-q^{6lm+2l+j}} \right. \right. \\ & \quad \left. \left. + \frac{q^{6lm+4l-j}}{1-q^{6lm+4l-j}} \right) - 4 \left(\frac{q^{6lm+4l+j}}{1-q^{6lm+4l+j}} - \frac{q^{6lm+2l-j}}{1-q^{6lm+2l-j}} \right) \right\} \\ &= - \sum_{n=-\infty}^{\infty} \frac{(6n+4+j)q^{3ln(n+1)+2ln+2l}}{1+q^{6ln+j+4l}} \\ & \quad + \sum_{n=-\infty}^{\infty} \frac{(6n+j)q^{3ln(n+1)-2ln}}{1+q^{6ln+j}} \\ & \quad + \frac{2(q^{2l}; q^{2l})_{\infty}^2}{(-q^{2l}; q^{2l})_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3ln(n+1)+2ln+2l}}{1-q^{6ln+j+4l}}. \end{aligned}$$

3 Proofs of Theorems 1.1 and 1.2

Proof of Theorem 1.1. We begin with the proof of (1.2). First, note that

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{q^n}{(1-q^n)^2} + 4 \sum_{n=1}^{\infty} \frac{q^{2n}}{(1-q^{2n})^2} &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} m q^{mn} + 4 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} m q^{2mn}, \\ \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{q^{n(3n+1)/2}}{(1-q^n)^2} &= \sum_{n=1}^{\infty} \frac{q^{n(3n+1)/2}}{(1-q^n)^2} + \sum_{n=1}^{\infty} \frac{q^{n(3n-1)/2+2n}}{(1-q^n)^2} \\ &= \sum_{n=1}^{\infty} q^{n(3n+1)/2} \sum_{m=1}^{\infty} m q^{nm-n} + \sum_{n=1}^{\infty} q^{n(3n+1)/2} \sum_{m=2}^{\infty} (m-1) q^{nm-n} \\ &= \sum_{m=1}^{\infty} (2m-1) \sum_{n=1}^{\infty} q^{n(3n+2m-1)/2}, \end{aligned}$$

and

$$\begin{aligned} \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(6n-1)q^{n(3n+1)/2}}{1-q^n} &= \sum_{n=1}^{\infty} \frac{(6n-1)q^{n(3n+1)/2}}{1-q^n} + \sum_{n=1}^{\infty} \frac{(6n+1)q^{n(3n+1)/2}}{1-q^n} \\ &= 12 \sum_{n=1}^{\infty} n q^{n(3n+1)/2} \sum_{m=0}^{\infty} q^{mn} \\ &= 12 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} n q^{n(3n+2m-1)/2}. \end{aligned}$$

Therefore, (2.8) is equivalent to

$$\frac{(q)_{\infty}^3}{(-q)_{\infty}^2} f(q) = 1 - 4 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} n q^{mn} - 16 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} n q^{2mn} + 4 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} (6n+2m-1) q^{n(3n-1+2m)/2}. \quad (3.1)$$

By applying (see [5], page 114, Entry 8 (ix))

$$\sum_{n=-\infty}^{\infty} (6n+1) q^{n(3n+1)/2} = \frac{(q)_{\infty}^3}{(-q)_{\infty}^2}, \quad (3.2)$$

and extracting the coefficient of q^n on both sides of (3.1), we obtain

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + m \leq 2n}} (6m+1) c\left(f; n - \frac{3}{2}m^2 - \frac{1}{2}m\right) = -4\sigma(n) - 16\sigma\left(\frac{n}{2}\right) + 4 \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ a \not\equiv b \pmod{2}}} (3a+b). \quad (3.3)$$

We now examine the right-hand side of (1.2). Observe that both a and b have the same sign, and when $a = 3b$, N is not an integer. Hence, by splitting the sum according to the values of a and b in the range $1 \leq b < 3a$, $1 < 3a < b$, $3a < b \leq -1$, and $b < 3a < -1$, we find that

$$\begin{aligned} \sum_{\substack{a, b \in \mathbb{Z} \\ ab=2n \\ 6|3a+b-1}} d\left(N, \tilde{N}, \frac{1}{6}, \frac{1}{6}\right) &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ 1 \leq b < 3a \\ 6|3a+b-1}} \frac{b}{3} - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ 6|3a+b-1}} a - \sum_{\substack{a, b \in \mathbb{Z}^- \\ ab=2n \\ 3a < b \leq -1 \\ 6|3a+b-1}} \frac{b}{3} + \sum_{\substack{a, b \in \mathbb{Z}^- \\ ab=2n \\ b < 3a < -1 \\ 6|3a+b-1}} a \\ &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ 1 \leq b < 3a \\ 6|3a+b-1 \text{ or } 6|3a+b+1}} \frac{b}{3} - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ 6|3a+b-1 \text{ or } 6|3a+b+1}} a \\ &= \frac{1}{3} \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ 1 \leq b < 3a \\ b \not\equiv a \pmod{2}}} b - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ b \not\equiv a \pmod{2}}} a, \end{aligned} \quad (3.4)$$

where in the third equality, we note that

$$\begin{aligned} \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ 1 \leq b < 3a \\ 6|b+3a+3}} b &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ 1 \leq b < 3a \\ 3|b \\ a \not\equiv b \pmod{2}}} b = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ 3ab=2n \\ 1 \leq 3b < 3a \\ a \not\equiv b \pmod{2}}} 3b = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ 3ba=2n \\ 1 \leq 3a < 3b \\ b \not\equiv a \pmod{2}}} 3a = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ 3|b \\ a \not\equiv b \pmod{2}}} 3a = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ 6|b+3a+3}} 3a. \end{aligned}$$

Thus, by (3.3) and (3.4), it suffices to show that

$$\sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ b > 3a \\ a \not\equiv b \pmod{2}}} b = 3\sigma(n) - 4\sigma\left(\frac{n}{2}\right) - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ 1 \leq b < 3a \\ a \not\equiv b \pmod{2}}} b. \quad (3.5)$$

By elementary manipulations, we see that

$$\begin{aligned} 3\sigma(n) - 4\sigma\left(\frac{n}{2}\right) &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=n}} 3b - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ a \equiv b \equiv 0 \pmod{2}}} 2b \\ &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=n \\ b \equiv 1 \pmod{2}}} b + \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=n \\ a \equiv 1 \pmod{2}}} 2b \\ &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ a \equiv 0 \pmod{2}, b \equiv 1 \pmod{2}}} b + \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ a \equiv 1 \pmod{2}, b \equiv 0 \pmod{2}}} b \\ &= \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=2n \\ a \not\equiv b \pmod{2}}} b \end{aligned}$$

and thus (3.5) is true. This proves (1.2).

Splitting the sum on the right-hand side of (1.3) in a way similar to (3.4), we find that

$$\begin{aligned} -6 \sum_{\substack{a,b \in \mathbb{Z} \\ ab=4n+1 \\ 12|3a-b-2}} d\left(N, \tilde{N}, \frac{1}{6}, \frac{1}{3}\right) &= \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=4n+1 \\ 1 \leq b \leq 3a-2 \\ 12|3a-b-2 \text{ or } 12|3a-b+2}} b - \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=4n+1 \\ b \geq 3a+2 \\ 12|3a-b-2 \text{ or } 12|3a-b+2}} 3a \\ &=: S_1(n) - 3S_2(n). \end{aligned} \quad (3.6)$$

The generating function for $S_1(n)$ is given by

$$\sum_{n=0}^{\infty} S_1(n) q^n = \sum_{a=1}^{\infty} \sum_{\substack{b=1 \\ 12|3a-b-2 \\ 4|ab-1}}^{3a-2} b q^{(ab-1)/4} + \sum_{a=1}^{\infty} \sum_{\substack{b=1 \\ 12|3a-b+2 \\ 4|ab-1}}^{3a-2} b q^{(ab-1)/4}. \quad (3.7)$$

Note to have $12|3a-b-2$ and $4|ab-1$, we must have a odd and $b \equiv 1 \pmod{6}$. Hence we replace $a = 2k+1$ and $b = 6l+1$ in the first sum on the right-hand side of (3.7). Similarly, we replace $a = 2k+1$ and $b = 6l+5$ in the second sum. This leads us to

$$\begin{aligned} \sum_{n=0}^{\infty} S_1(n) q^n &= \sum_{k=0}^{\infty} \sum_{\substack{l=0 \\ 2|k-l}}^k (6l+1) q^{3kl+(k+3l)/2} + \sum_{k=0}^{\infty} \sum_{\substack{l=0 \\ 2|k-l}}^k (6l+5) q^{3kl+(5k+3l)/2+1} \\ &= \sum_{l=0}^{\infty} \sum_{\substack{k=l \\ 2|k-l}}^{\infty} (6l+1) q^{3kl+(k+3l)/2} + \sum_{l=0}^{\infty} \sum_{\substack{k=l \\ 2|k-l}}^{\infty} (6l+5) q^{3kl+(5k+3l)/2+1} \\ &= \sum_{l=0}^{\infty} (6l+1) q^{3l^2+2l} \sum_{\substack{k=0 \\ 2|k}}^{\infty} q^{k(6l+1)/2} + \sum_{l=0}^{\infty} (6l+5) q^{3l^2+4l+1} \sum_{\substack{k=0 \\ 2|k}}^{\infty} q^{k(6l+5)/2} \\ &= \sum_{l=0}^{\infty} (6l+1) \frac{q^{3l^2+2l}}{1-q^{6l+1}} + \sum_{l=0}^{\infty} (6l+5) \frac{q^{3l^2+4l+1}}{1-q^{6l+5}} \\ &= \sum_{l=-\infty}^{\infty} \frac{(6l+1) q^{3l^2+2l}}{1-q^{6l+1}}, \end{aligned} \quad (3.8)$$

where in the last equality, we replaced l by $-l-1$ in the second sum. Similarly, the generating function for $S_2(n)$ is

$$\sum_{n=0}^{\infty} S_2(n) q^n = \sum_{n=-\infty}^{\infty} \frac{(2n-1) q^{3n^2+2n-2}}{1-q^{6n-3}}. \quad (3.9)$$

From (2.10), applying (see [5], page 115, Entry 8 (x))

$$\sum_{n=-\infty}^{\infty} (3n+1) q^{n(3n+2)} = \frac{(q^2; q^2)_{\infty}^3}{(-q; q^2)_{\infty}^2}, \quad (3.10)$$

(3.6), (3.8) and (3.9), we see that (1.3) follows upon extracting the coefficient of q^n from both sides of

$$2 \frac{(q^2; q^2)_{\infty}^2}{(-q; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)}}{1+q^{2n}} = \sum_{n=-\infty}^{\infty} \frac{(6n+1) q^{3n^2+2n}}{1-q^{6n+1}} - 3 \sum_{n=-\infty}^{\infty} \frac{(2n-1) q^{3n^2+2n-2}}{1-q^{6n-3}}. \quad (3.11)$$

Now, to prove (3.11), we first set $l = 1$ and $j = -3$ in Theorem 2.3 to obtain

$$\begin{aligned} & \frac{[-q^{-2}, q^2; q^6]_{\infty} (q^6; q^6)_{\infty}^2}{[-q^{-3}, q, -1/q; q^6]_{\infty}} \left\{ -1 + 2 \sum_{m=0}^{\infty} \left(\frac{q^{6m+1}}{1 - q^{6m+1}} - \frac{q^{6(m+1)-1}}{1 - q^{6(m+1)-1}} + \frac{q^{6m-2}}{1 + q^{6m-2}} \right. \right. \\ & \quad \left. \left. - \frac{q^{6(m+1)+2}}{1 + q^{6(m+1)+2}} \right) \right\} \\ &= \sum_{n=-\infty}^{\infty} \frac{(6n-1)(-1)^n q^{3n^2+4n+1}}{1 + q^{6n-1}} \\ & \quad + \sum_{n=-\infty}^{\infty} \frac{(6n-3)(-1)^n q^{3n^2+2n}}{1 + q^{6n-3}} \\ & \quad + \frac{4[q^{-2}, q^2; q^6]_{\infty} (q^6; q^6)_{\infty}^2}{[q^3, q, 1/q; q^6]_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+2n+3}}{1 + q^{6n}}. \end{aligned} \quad (3.12)$$

Replacing q, a, b and c by q^6, q^5, q^5 and $-1/q^2$, respectively, in ([8], Corollary 3.2), we obtain

$$\begin{aligned} & -1 + 2 \sum_{m=0}^{\infty} \left(\frac{q^{6m+1}}{1 - q^{6m+1}} - \frac{q^{6(m+1)-1}}{1 - q^{6(m+1)-1}} + \frac{q^{6m-2}}{1 + q^{6m-2}} - \frac{q^{6(m+1)+2}}{1 + q^{6(m+1)+2}} \right) \\ &= -\frac{[q^{10}, -q^3, -q^3; q^6]_{\infty} (q^6; q^6)_{\infty}^2}{[q, q, -1/q^2, -q^8; q^6]_{\infty}}, \end{aligned}$$

which together with (3.12) gives

$$\begin{aligned} & -\sum_{n=-\infty}^{\infty} \frac{(6n-1)(-1)^n q^{3n^2+4n+1}}{1 + q^{6n-1}} - \sum_{n=-\infty}^{\infty} \frac{(6n-3)(-1)^n q^{3n^2+2n}}{1 + q^{6n-3}} \\ &= -\frac{[q, q^2, q^2, -q^2, q^3; q^6]_{\infty} (q^6; q^6)_{\infty}^4}{[-q; q^6]_{\infty}} + 4 \frac{(q^2; q^2)_{\infty}^2}{(q; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+5)+2}}{1 + q^{6n}}. \end{aligned} \quad (3.13)$$

By ([6], Eq. (2.1)), we have

$$-\frac{q^2 [q, q^2, q^2, -q^2, q^3; q^6]_{\infty} (q^6; q^6)_{\infty}^4}{[-q; q^6]_{\infty}} = -2 \frac{(q^2; q^2)_{\infty}^2}{(q; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+3)+2}}{1 + q^{6n}}. \quad (3.14)$$

Substituting (3.14) into (3.13), we obtain

$$\begin{aligned} & -\sum_{n=-\infty}^{\infty} \frac{(6n-1)(-1)^n q^{3n^2+4n+1}}{1 + q^{6n-1}} - \sum_{n=-\infty}^{\infty} \frac{(6n-3)(-1)^n q^{3n^2+2n}}{1 + q^{6n-3}} \\ &= 4 \frac{(q^2; q^2)_{\infty}^2}{(q; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+5)+2}}{1 + q^{6n}} - 2 \frac{(q^2; q^2)_{\infty}^2}{(q; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+3)+2}}{1 + q^{6n}} \\ &= 2 \frac{(q^2; q^2)_{\infty}^2}{(q; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)+2}}{1 + q^{6n}}, \end{aligned} \quad (3.15)$$

where the last step follows from the fact that

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)}}{1 + q^{6n}} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+5)}}{1 + q^{6n}}$$

and

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)}}{1+q^{2n}} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+1)}(1-q^{2n}+q^{4n})}{1+q^{6n}}.$$

As (3.15) is equivalent to (3.11), this proves (1.3). $\square$

Proof of Theorem 1.2. For (1.4), we first split the sum on the right-hand side in a way similar to (3.4) to obtain

$$6(-1)^{n+1} \sum_{\substack{a,b \in \mathbb{Z} \\ ab=4n+3 \\ 12|3a-b-4}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{6}\right) = \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=4n+3 \\ b \geq 3a+2 \\ 12|3a-b-4 \text{ or } 12|3a-b+4}} 3a - \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=4n+3 \\ 1 \leq b \leq 3a-1 \\ 12|3a-b-4 \text{ or } 12|3a-b+4}} b. \quad (3.16)$$

By (2.11), (3.2) and a calculation similar to (3.8) for the generating function of the right-hand side of (3.16), we see that (1.4) follows extracting the coefficient of q^n from both sides of

$$\frac{(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+3n}}{1-q^{2n+1}} = -3 \sum_{n=-\infty}^{\infty} \frac{(2n-1)q^{n(3n+1)-2}}{1+q^{6n-3}} - \sum_{n=-\infty}^{\infty} \frac{(6n-1)q^{n(3n+1)-1}}{1+q^{6n-1}}. \quad (3.17)$$

To prove (3.17), we set $j = -1, l = 1$ in Theorem 2.4 to get

$$\begin{aligned} & \frac{[q, q^4; q^6]_{\infty} (q^6; q^6)_{\infty}^2}{[-1/q, -q^2, -q^3; q^6]_{\infty}} \left\{ -1 + \sum_{m=0}^{\infty} 2 \left(\frac{q^{6m+2}}{1+q^{6m+2}} - \frac{q^{6m+4}}{1+q^{6m+4}} - \frac{q^{6m+1}}{1-q^{6m+1}} + \frac{q^{6m+5}}{1-q^{6m+5}} \right) \right\} \\ &= - \sum_{n=-\infty}^{\infty} \frac{(6n+3)q^{3n(n+1)+2n+2}}{1+q^{6n+3}} + \sum_{n=-\infty}^{\infty} \frac{(6n-1)q^{3n(n+1)-2n}}{1+q^{6n-1}} \\ &+ \frac{2(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+2n+2}}{1-q^{6n+3}}. \end{aligned} \quad (3.18)$$

Replacing q, a, b and c by q^6, q, q and $-q^2$, respectively, in ([8], Corollary 3.2), we obtain

$$\begin{aligned} & -1 + \sum_{m=0}^{\infty} 2 \left(\frac{q^{6m+2}}{1+q^{6m+2}} - \frac{q^{6m+4}}{1+q^{6m+4}} - \frac{q^{6m+1}}{1-q^{6m+1}} + \frac{q^{6m+5}}{1-q^{6m+5}} \right) \\ &= - \frac{[q^2, -q^3, -q^4; q^6]_{\infty} (q^6; q^6)_{\infty}^2}{[q, q, -q^2, -q^4; q^6]_{\infty}}, \end{aligned}$$

which together with (3.18) gives

$$\begin{aligned} & \sum_{n=-\infty}^{\infty} \frac{(6n+3)q^{3n(n+1)+2n+2}}{1+q^{6n+3}} - \sum_{n=-\infty}^{\infty} \frac{(6n-1)q^{3n(n+1)-2n}}{1+q^{6n-1}} \\ &= \frac{2(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+2n+2}}{1-q^{6n+3}} + \frac{q[q^2, -q^3, q^4; q^6]_{\infty} (q^6; q^6)_{\infty}^4}{[q, -q, -q^2, -q^2, -q^4; q^6]_{\infty}}. \end{aligned} \quad (3.19)$$

We note that (see [6, Eq. (2.1)])

$$\frac{q[q^2, -q^3, q^4; q^6]_{\infty} (q^6; q^6)_{\infty}^4}{[q, -q, -q^2, -q^2, -q^4; q^6]_{\infty}} = \frac{(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+3)+1}}{1-q^{6n+3}}. \quad (3.20)$$

Substituting (3.20) into (3.19), we find that

$$\begin{aligned} & \sum_{n=-\infty}^{\infty} \frac{(6n+3)q^{3n(n+1)+2n+2}}{1+q^{6n+3}} - \sum_{n=-\infty}^{\infty} \frac{(6n-1)q^{3n(n+1)-2n}}{1+q^{6n-1}} \\ &= \frac{2(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+2n+2}}{1-q^{6n+3}} + \frac{(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+1}}{1-q^{6n+3}} \\ &= \frac{(q^2; q^2)_{\infty}^2}{(-q^2; q^2)_{\infty}^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+1}}{1-q^{2n+1}}, \end{aligned} \quad (3.21)$$

where the last step follows from the fact that

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+2n+1}}{1-q^{6n+3}} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)+4n+2}}{1-q^{6n+3}}$$

and

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)}}{1-q^{2n+1}} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n(n+1)}(1+q^{2n+1}+q^{4n+2})}{1-q^{6n+3}}.$$

By (3.21), we find that, to prove (3.17), it suffices to show that

$$\sum_{n=-\infty}^{\infty} \frac{(6n+3)q^{3n(n+1)+2n+2}}{1+q^{6n+3}} = - \sum_{n=-\infty}^{\infty} \frac{(6n-3)q^{3n^2+n-1}}{1+q^{6n-3}}$$

which is easily checked to be true by replacing n by $-n$ in the sum on the left-hand side. This completes the proof of (3.17) and thus (1.4).

Now, by taking the sum and difference of (1.5) and (1.6), respectively, we obtain the two equivalent formulas

$$\begin{aligned} \sum_{\substack{m \in \mathbb{Z} \\ 3m^2+2m+1 \leq n}} \left(m + \frac{1}{3}\right) c(\omega(q); n - 3m^2 - 2m - 1) &= \sum_{\substack{a, b \in \mathbb{Z} \\ ab=n \\ 12|a-3b-2}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right) \\ &- \sum_{\substack{a, b \in \mathbb{Z} \\ ab=n \\ 12|a-3b-8}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right), \end{aligned} \quad (3.22)$$

and

$$\begin{aligned} \sum_{\substack{m \in \mathbb{Z} \\ 3m^2+2m+1 \leq n}} \left(m + \frac{1}{3}\right) c(\omega(-q); n - 3m^2 - 2m - 1) &= 2R_n \\ &- \sum_{\substack{a, b \in \mathbb{Z} \\ ab=n \\ 12|a-3b-8 \text{ or } 12|a-3b-2}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right). \end{aligned} \quad (3.23)$$

From (3.10) and noting that

$$3 \sum_{\substack{a, b \in \mathbb{Z} \\ ab=n \\ 12|a-3b-8}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right) = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=n \\ 1 \leq a \leq 3b-1 \\ 12|a-3b-8 \text{ or } 12|a-3b+8}} a - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab=n \\ a \geq 3b+1 \\ 12|a-3b-8 \text{ or } 12|a-3b+8}} 3b$$

and

$$3 \sum_{\substack{a,b \in \mathbb{Z} \\ ab=n \\ 12|a-3b-2}} d\left(N, \tilde{N}, \frac{1}{3}, \frac{1}{3}\right) = \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ 1 \leq a \leq 3b-1 \\ 12|a-3b-2 \text{ or } 12|a-3b+2}} a - \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a \geq 3b+1 \\ 12|a-3b-2 \text{ or } 12|a-b+2}} 3b,$$

we see that (3.22) follows from

$$\begin{aligned} \frac{q(q^2; q^2)_\infty^3}{(-q; q^2)_\infty^2} \omega(q) &= \sum_{n=-\infty}^{\infty} \frac{(3n+2)q^{3n^2+14n+8}}{1-q^{12n+8}} - 3 \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{nq^{3n^2+2n}}{1-q^{12n}} \\ &\quad - \sum_{n=-\infty}^{\infty} \frac{(3n+2)q^{3n^2+8n+4}}{1-q^{12n+8}} + 3 \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{nq^{3n^2+8n}}{1-q^{12n}} \end{aligned} \quad (3.24)$$

or equivalently

$$\begin{aligned} \frac{q(q^2; q^2)_\infty^3}{(-q; q^2)_\infty^2} \omega(q) &= -3 \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{nq^{3n^2+2n}}{1+q^{6n}} - \sum_{n=-\infty}^{\infty} \frac{(3n+2)q^{3n^2+8n+4}}{1+q^{6n+4}} \\ &= -3 \sum_{n=-\infty}^{\infty} \frac{nq^{3n^2+2n}}{1+q^{6n}} + \sum_{n=-\infty}^{\infty} \frac{(3n+1)q^{3n^2+4n+1}}{1+q^{6n+2}}. \end{aligned} \quad (3.25)$$

We now set $l = 1, j = 0$ and replace q by $-q$ in Theorem 2.3 to obtain

$$\begin{aligned} &2 \frac{[q, q^2; q^6]_\infty (q^6; q^6)_\infty^2}{[-1, -q, -q^2; q^6]_\infty} \sum_{m=0}^{\infty} \left(-\frac{q^{6m+1}}{1+q^{6m+1}} + \frac{q^{6m+5}}{1+q^{6m+5}} - \frac{q^{6m+1}}{1-q^{6m+1}} + \frac{q^{6m+5}}{1-q^{6m+5}} \right) \\ &= - \sum_{n=-\infty}^{\infty} \frac{(6n+2)q^{3n^2+4n+1}}{1+q^{6n+2}} + \sum_{n=-\infty}^{\infty} \frac{6nq^{3n^2+2n}}{1+q^{6n}} + 4 \frac{(q^2; q^2)_\infty^2}{(-q; q^2)_\infty^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+5n+2}}{1-q^{6n+3}}. \end{aligned} \quad (3.26)$$

By ([8], Corollary 3.1), we have

$$\begin{aligned} \sum_{m=0}^{\infty} \left(-\frac{q^{6m+1}}{1+q^{6m+1}} + \frac{q^{6m+5}}{1+q^{6m+5}} - \frac{q^{6m+1}}{1-q^{6m+1}} + \frac{q^{6m+5}}{1-q^{6m+5}} \right) &= -4 \sum_{j=-\infty}^{\infty} \frac{q^{6j+1}}{1-q^{12j+2}} \\ &= -\frac{4q [q^8; q^{12}]_\infty (q^{12}; q^{12})_\infty^2}{[q^2, q^6; q^{12}]_\infty} \end{aligned}$$

which together with (3.26) gives

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{(6n+2)q^{3n^2+4n}}{1+q^{6n+2}} - \sum_{n=-\infty}^{\infty} \frac{6nq^{3n^2+2n-1}}{1+q^{6n}} &= 2 \frac{(q^2; q^2)_\infty^3}{(-q; q^2)_\infty^2} \times \frac{[q^6; q^{18}]_\infty^3 (q^{18}; q^{18})_\infty^3}{[q^2, q^3; q^6]_\infty (q^6; q^6)_\infty^2} \\ &\quad + 4 \frac{(q^2; q^2)_\infty^2}{(-q; q^2)_\infty^2} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(3n+5)+1}}{1-q^{6n+3}}. \end{aligned}$$

This together with ([15], Eq. (5.8)) implies

$$2 \frac{(q^2; q^2)_\infty^3}{(-q; q^2)_\infty^2} \omega(q) = \sum_{n=-\infty}^{\infty} \frac{(6n+2)q^{3n^2+4n}}{1+q^{6n+2}} - \sum_{n=-\infty}^{\infty} \frac{6nq^{3n^2+2n-1}}{1+q^{6n}}$$

which is equivalent to (3.25). Thus, (3.22) is proven.

For (3.23), it suffices to prove

$$\begin{aligned}
 & \sum_{\substack{m \in \mathbb{Z} \\ 3m^2 + 2m + 1 \leq n}} (3m + 1) c(\omega(-q); n - 3m^2 - 2m - 1) \\
 &= 6R_n - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ 1 \leq a \leq 3b-1 \\ a-3b \equiv 2, 4, 8, 10 \pmod{12}}} a + \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ a \geq 3b+1 \\ a-3b \equiv 2, 4, 8, 10 \pmod{12}}} 3b \\
 &= 6R_n - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ 1 \leq a \leq 3b-1 \\ a \equiv b \pmod{2}}} a + \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ a \geq 3b+1 \\ a \equiv b \pmod{2}}} 3b \\
 &= 6R_n - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ 1 \leq a \leq 3b \\ a \equiv b \pmod{2}}} a + \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ a \geq 3b \\ a \equiv b \pmod{2}}} 3b, \tag{3.27}
 \end{aligned}$$

where in the second equality, we used the fact that

$$\sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ 1 \leq a \leq 3b-1 \\ a-3b \equiv 0 \pmod{6}}} a = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ 1 \leq 3a \leq 3b-1 \\ a \equiv b \pmod{2}}} 3a = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ba = n \\ 1 \leq 3b \leq 3a-1 \\ a \equiv b \pmod{2}}} 3b = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ a \geq 3b+1 \\ 3|a \\ a \equiv b \pmod{2}}} 3b = \sum_{\substack{a, b \in \mathbb{Z}^+ \\ ab = n \\ a \geq 3b+1 \\ a-3b \equiv 0 \pmod{6}}} 3b.$$

Next, we examine the right-hand side of (2.9). Note that

$$\begin{aligned}
 \sum_{n=1}^{\infty} \frac{q^{2n-1}}{(1+q^{2n-1})^2} - 2 \sum_{n=1}^{\infty} \frac{q^{2n}}{(1-q^{2n})^2} &= \sum_{n=1}^{\infty} \left(\frac{(2n-1)q^{2n-1}}{1-q^{4n-2}} - \frac{4nq^{2n}}{1-q^{2n}} + \frac{2nq^{4n}}{1-q^{4n}} \right) \\
 &= \sum_{n=1}^{\infty} 6\tilde{R}_n q^n,
 \end{aligned}$$

where

$$\tilde{R}_n := \begin{cases} \frac{1}{3}(\sigma(\frac{n}{4}) - 2\sigma(\frac{n}{2})), & \text{if } n \text{ is even;} \\ \frac{1}{6}\sigma(n), & \text{if } n \text{ is odd,} \end{cases}$$

and

$$\begin{aligned}
 \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{q^{3n^2}}{(1-q^{2n})^2} &= \sum_{n=1}^{\infty} \frac{q^{3n^2}}{(1-q^{2n})^2} + \sum_{n=1}^{\infty} \frac{q^{3n^2+4n}}{(1-q^{2n})^2} \\
 &= \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} q^{3n^2+2mn} (m+1) + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} q^{3n^2+2mn} (m-1) \\
 &= \sum_{n=1}^{\infty} q^{3n^2} + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} 2mq^{n(3n+2m)}.
 \end{aligned}$$

Similarly,

$$\sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(3n-1)q^{3n^2}}{1-q^{2n}} = \sum_{n=1}^{\infty} (3n-1)q^{3n^2} + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} 6nq^{n(3n+2m)}.$$

Hence, identity (2.9) is equivalent to

$$\begin{aligned}
 q \frac{(q^2; q^2)_{\infty}^3}{(-q; q^2)_{\infty}^2} \omega(-q) &= \sum_{n=1}^{\infty} 6\tilde{R}_n q^n + \sum_{n=1}^{\infty} 3nq^{3n^2} + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} (6n+2m)q^{n(3n+2m)} \\
 &= \sum_{n=1}^{\infty} 6\tilde{R}_n q^n + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} (3n+2m)q^{n(3n+2m)} + \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} 3nq^{n(3n+2m)} \\
 &= \sum_{n=1}^{\infty} 6\tilde{R}_n q^n + \sum_{n=1}^{\infty} q^n \sum_{\substack{b,m \in \mathbb{Z}^+ \\ b(3b+2m)=n}} (3b+2m) + \sum_{n=1}^{\infty} q^n \sum_{\substack{b,m \in \mathbb{Z}^+ \\ b(3b+2m-2)=n}} 3b \\
 &= \sum_{n=1}^{\infty} 6\tilde{R}_n q^n + \sum_{n=1}^{\infty} q^n \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a>3b \\ a \equiv b \pmod{2}}} a + \sum_{n=1}^{\infty} q^n \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a>3b-2 \\ a \equiv b \pmod{2}}} 3b. \quad (3.28)
 \end{aligned}$$

Applying (3.10) while extracting the coefficient of q^n from both sides of (3.28), we obtain

$$\sum_{\substack{m \in \mathbb{Z} \\ 3m^2+2m+1 \leq n}} (3m+1) c(\omega(-q); n-3m^2-2m-1) = 6\tilde{R}_n + \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a>3b \\ a \equiv b \pmod{2}}} a + \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a>3b-2 \\ a \equiv b \pmod{2}}} 3b.$$

Note that

$$\begin{aligned}
 \sum_{n=1}^{\infty} 6(R_n - \tilde{R}_n)q^n &= \sum_{n=1}^{\infty} 2\sigma(n/2)q^{2n} + \sum_{n=1}^{\infty} \sigma(2n-1)q^{2n-1} \\
 &= \sum_{n=1}^{\infty} \frac{2nq^{4n}}{1-q^{4n}} + \sum_{n=1}^{\infty} \frac{(2n-1)q^{2n-1}}{1-q^{4n-2}} \\
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} 2nq^{4mn} + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} (2n-1)q^{(2n-1)(2m-1)} \\
 &= \sum_{n=1}^{\infty} q^n \sum_{\substack{a,b \in \mathbb{Z}^+ \\ 4ab=n}} 2a + \sum_{n=1}^{\infty} q^n \sum_{\substack{a,b \in \mathbb{Z}^+ \\ (2a-1)(2b-1)=n}} (2a-1) \\
 &= \sum_{n=1}^{\infty} q^n \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a \equiv b \pmod{2}}} a.
 \end{aligned}$$

Thus, (3.27) follows upon observing that

$$\sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a \equiv b \pmod{2}}} a - \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ 1 \leq a \leq 3b \\ a \equiv b \pmod{2}}} a + \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a \geq 3b \\ a \equiv b \pmod{2}}} 3b = \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a > 3b \\ a \equiv b \pmod{2}}} a + \sum_{\substack{a,b \in \mathbb{Z}^+ \\ ab=n \\ a > 3b-2 \\ a \equiv b \pmod{2}}} 3b.$$

This proves (3.23). Adding (3.22) and (3.23), then dividing by two yields (1.5) while subtracting (3.23) from (3.22), then dividing by two implies (1.6). $\square$

4 Other applications of (1.1)

It is worth noting that (1.1) can also be used to obtain other identities involving mock theta functions. For example, in [3], Andrews, Rhoades and Zwegers consider the automorphic properties of the q -series

$$\nu_2(q) := \frac{1}{(q)_\infty^3} \left(\sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(-1)^{n+1} n q^{\frac{n(n+1)}{2}}}{1 - q^n} - \frac{1}{4} - 2 \sum_{n=1}^{\infty} \frac{q^n}{(1 + q^n)^2} \right),$$

which is related to the generating function for the number of concave compositions of n [1]. In particular, to show that $\nu_2(q)$ is a mock theta function, they require the following key identity (see Theorem 1.3 in [3])

$$\tilde{F}(q) := \frac{1}{(q)_\infty (-q)_\infty^2} \sum_{n=-\infty}^{\infty} \frac{q^{\frac{n(n+1)}{2}}}{1 + q^n} = -2\nu_2(q). \quad (4.1)$$

We now prove a generalization of (4.1).

Theorem 4.1. *We have*

$$\frac{(q)_\infty^2}{[1/b]_\infty} \sum_{k=-\infty}^{\infty} \frac{(-b)^k q^{\frac{k(k+1)}{2}}}{1 - bq^k} = - \sum_{n=0}^{\infty} \left(\frac{q^n b}{(1 - bq^n)^2} + \frac{q^{n+1}/b}{(1 - q^{n+1}/b)^2} \right) + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(-1)^n n q^{n(n+1)/2}}{1 - q^n}. \quad (4.2)$$

Proof of Theorem 4.1. Setting $r = 1$, $s = 2$ and replacing a_1 by bb_1 in (1.1), after rearranging, we obtain

$$\frac{[bb_1, b_1/b]_\infty (q)_\infty^2}{[b, b_1, b_1]_\infty} = \sum_{k=-\infty}^{\infty} \frac{(-b)^k q^{\frac{k(k+1)}{2}}}{1 - bq^k} - \frac{[b]_\infty}{b[b_1]_\infty} \sum_{k=-\infty}^{\infty} \frac{(-b_1)^{k+1} q^{\frac{k(k+1)}{2}}}{1 - b_1 q^k}. \quad (4.3)$$

Multiplying by $(1 - b_1)^2$ and applying the operator $\frac{d^2}{d^2 b_1} \Big|_{b_1=1}$, we obtain

$$\begin{aligned} & \frac{(qb, q/b)_\infty}{(1 - b)(q)_\infty^2} + \frac{[1/b]_\infty}{(q)_\infty^2} \sum_{n=1}^{\infty} \left(\frac{2q^n}{(1 - q^n)^2} - \frac{q^n b}{(1 - bq^n)^2} - \frac{q^n/b}{(1 - q^n/b)^2} \right) \\ &= \sum_{k=-\infty}^{\infty} \frac{(-b)^k q^{\frac{k(k+1)}{2}}}{1 - bq^k} + \frac{[b]_\infty}{b(q)_\infty^2} \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \left\{ \frac{(-1)^n q^{n(n+1)/2} (1 + q^n)}{(1 - q^n)^2} + \frac{(-1)^n n q^{n(n+1)/2}}{1 - q^n} \right\}. \end{aligned} \quad (4.4)$$

Letting $a, c, d, e \rightarrow 1$ and $b \rightarrow q$ in a limiting case of Watson's $8\phi_7$ transformation, [5, Eq. (7.2), p. 16]

$$\sum_{n=0}^{\infty} \frac{(aq/bc, d, e; q)_n \left(\frac{aq}{de}\right)^n}{(q, aq/b, aq/c; q)_n} = \frac{(aq/d, aq/e; q)_\infty}{(aq, aq/de; q)_\infty} \sum_{n=0}^{\infty} \frac{(a, b, c, d, e; q)_n (1 - aq^{2n}) (-a^2)^n q^{n(n+3)/2}}{(q, aq/b, aq/c, aq/d, aq/e; q)_n (1 - a)(bcde)^n},$$

we find that

$$\sum_{n=1}^{\infty} \frac{q^n}{(1 - q^n)^2} = - \sum_{n=1}^{\infty} \frac{(-1)^n q^{n(n+1)/2} (1 + q^n)}{(1 - q^n)^2}. \quad (4.5)$$

Substituting (4.5) into (4.4) and rearranging, we obtain

$$\frac{-1}{b(1-b)^2} - \sum_{n=1}^{\infty} \left(\frac{q^n b}{(1-bq^n)^2} + \frac{q^n/b}{(1-q^n/b)^2} \right) = \frac{(q)_\infty^2}{[1/b]_\infty} \sum_{k=-\infty}^{\infty} \frac{(-b)^k q^{\frac{k(k+1)}{2}}}{1-bq^k} - \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(-1)^n n q^{n(n+1)/2}}{1-q^n}$$

which implies (4.2). $\square$

Multiplying by $\frac{1-1/b}{(q;q)_\infty^3}$ on both sides of (4.2) and setting $b = -1$, we obtain (4.1). Using Theorem 4.1, one can also show

$$q(q^4; q^4)_\infty^3 B(q) = \sum_{n=0}^{\infty} \left(\frac{q^{4n+1}}{(1-q^{4n+1})^2} + \frac{q^{4n+3}}{(1-q^{4n+3})^2} \right) - \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(-1)^n n q^{2n(n+1)}}{1-q^{4n}} \quad (4.6)$$

where

$$B(q) := \sum_{n=0}^{\infty} \frac{q^n (-q; q^2)_n}{(q; q^2)_{n+1}}$$

is a 2nd order mock theta function (see [14]). To see this, replace q and b by q^4 and q , respectively in (4.2) to obtain

$$\frac{(q^4; q^4)_\infty^2}{[1/q; q^4]_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(2n+3)}}{1-q^{4n+1}} = - \sum_{n=0}^{\infty} \left(\frac{q^{4n+1}}{(1-q^{4n+1})^2} + \frac{q^{4n+3}}{(1-q^{4n+3})^2} \right) + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{(-1)^n n q^{2n(n+1)}}{1-q^{4n}}.$$

By ([15], (3.2a), (3.2b), Eq. (5.2)), we have

$$B(q) = \frac{1}{(q, q^3, q^4; q^4)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n(2n+3)}}{1-q^{4n+1}}$$

and thus (4.6) follows.

Extracting the coefficient of q^n on both sides of (4.6), we obtain the following corollary.

Corollary 4.2. *For a fixed positive integer n , we have*

$$\sum_{\substack{m \in \mathbb{Z}^+ \\ 2m^2+m+1 \leq n}} (-1)^m (2m+1) c(B, n-2m^2-2m-1) = \sum_{\substack{0 < d | n \\ \frac{n}{d} \text{ odd}}} d - \sum_{\substack{a, b \in \mathbb{Z}^+ \\ 1 \leq a < b \\ 2ab=n \\ a \not\equiv b \pmod{2}}} (-1)^a a.$$

Similar results exist, for example, for the mock theta functions $\psi(q)$, $\rho(q)$ and $\lambda(q)$ of order 6 and $V_0(q)$ of order 8 as they can be written in terms of Appell-Lerch series (see Section 5 of [15]).

Acknowledgements

The first author was partially supported by the Singapore Ministry of Education Academic Research Fund, Tier 2, project number MOE2014-T2-1-051, ARC40/14. The second author was partially supported by National Natural Science Foundation of China (Grant No. 11501398), Natural Science Foundation of Jiangsu Province (Grant No. BK20150304) and Natural Science Foundation of the Jiangsu Higher Education Institutions of China (Grant No. 15KJB110020). The third author would like to thank the Institut des Hautes Études Scientifiques for their support during the preparation of this paper. This material is based upon work supported by the National Science Foundation under Grant No. 1002477. Finally, the authors would like to thank Jeremy Lovejoy for his motivating remark in [16].

Author details

¹Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang link, Singapore 637371, Republic of Singapore. ²School of Mathematical Sciences, Soochow University, Suzhou 215006, PR China. ³School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland. ⁴IHÉS, Le Bois-Marie, 35, route de Chartres, F-91440 Bures-sur-Yvette, France.

Received: 7 September 2015 Accepted: 12 November 2015

Published online: 10 December 2015

References

1. Andrews, GE: Concave and convex compositions. *Ramanujan J.* **31**(1–2), 67–82 (2013)
2. Andrews, GE, Berndt, BC: Ramanujan's lost notebook. Part I. Springer, New York (2005)
3. Andrews, GE, Rhoades, R, Zwegers, S: Modularity of the concave composition generating function. *Algebra Number Theory.* **7**(9), 2103–2139 (2013)
4. Atkin, AOL, Swinnerton-Dyer, P: Some properties of partitions. *Proc. London Math. Soc.* **4**(3), 84–106 (1954)
5. Berndt, BC: Ramanujan's Notebooks, Part III. Springer-Verlag, New York (1991)
6. Berndt, BC, Chan, HH, Chan, SH, Liaw, W-C: Cranks and dissections in Ramanujan's lost notebook. *J. Combin. Theory Ser. A.* **109**(1), 91–120 (2005)
7. Bringmann, K, Lovejoy, J, Osburn, R: Automorphic properties of generating functions for generating rank moments and Durfee symbols. *Int. Math. Res. Not. IMRN.* **2**, 238–260 (2010)
8. Chan, SH: Generalized Lambert series identities. *Proc. London Math. Soc.* **91**(3), 598–622 (2005)
9. Chan, SH: Congruences for Ramanujan's φ function. *Acta Arith.* **153**(2), 161–189 (2012)
10. Chan, SH, Dixit, A, Garvan, F: Rank-crank-type PDEs and generalized Lambert series identities. *Ramanujan J.* **31**(1–2), 163–189 (2013)
11. Chan, SH, Mao, R: Pairs of partitions without repeated odd parts. *J. Math. Anal. Appl.* **394**(1), 408–415 (2012)
12. Chan, SH, Mao, R: Two congruences for Appell-Lerch sums. *Int. J. Number Theory.* **8**(1), 111–123 (2012)
13. Duncan, J, Griffin, M, Ono, K: Proof of the Umbral Moonshine conjecture. preprint available at <http://arxiv.org/abs/1503.01472>. Accessed 4 March 2015
14. Gordon, B, McIntosh, RJ: A survey of classical mock theta functions. In: *Partitions, q -series, and Modular Forms, Developments in Mathematics*, vol. 23, pp. 95–144. Springer, New York, (2012)
15. Hickerson, D, Mortenson, E: Hecke-type double sums, Appell-Lerch sums, and mock theta functions, I. *Proc. London Math. Soc.* **109**, 382–422 (2014)
16. Imamoğlu, O, Raum, M, Richter, O: Holomorphic projections and Ramanujan's mock theta functions. *Proc. Natl. Acad. Sci. USA.* **111**(11), 3961–3967 (2014)
17. Jennings-Shaffer, C: Exotic Bailey-Slater spt-functions I: Group A. preprint available at <http://arxiv.org/abs/1408.5395>, Accessed 22 August 2014
18. Jennings-Shaffer, C: Congruences for partition pairs with conditions. *Q. J. Math.* **66**(3), 837–860 (2015)
19. Lovejoy, J, Osburn, R: Rank differences for overpartitions. *Q. J. Math.* **59**(2), 257–273 (2008)
20. Lovejoy, J, Osburn, R: M_2 -rank differences for partitions without repeated odd parts. *J. Théor. Nombres Bordeaux.* **21**(2), 313–334 (2009)
21. Lovejoy, J, Osburn, R: M_2 -rank differences for overpartitions. *Acta Arith.* **144**(2), 193–212 (2010)
22. Mao, R: Ranks of partitions modulo 10. *J. Number Theory.* **133**(11), 3678–3702 (2013)
23. Mertens, M: Eichler-Selberg type identities for mixed mock modular forms. preprint available at <http://arxiv.org/abs/1404.5491>. Accessed 22 April 2015

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com