




Gabriel–Colman rearrangement


Reaction of the enolate of a maleimidyl acetate to provide isoquinoline 1,4-diol.



Example 1⁶



Example 2⁹



References

1. (a) Gabriel, S.; Colman, J. *Ber.* **1900**, *33*, 980–995. (b) Gabriel, S.; Colman, J. *Ber.* **1900**, *33*, 2630–2634. (c) Gabriel, S.; Colman, J. *Ber.* **1902**, *35*, 1358–1368.
2. Allen, C. F. H. *Chem. Rev.* **1950**, *47*, 275–305. (Review).
3. Gensler, W. J. *Heterocyclic Compounds*, Vol. 4, R. C. Elderfield, Ed., Wiley & Sons., New York, N.Y., **1952**, 378. (Review).
4. Hill, J. H. M. *J. Org. Chem.* **1965**, *30*, 620–622. (Mechanism).
5. Lombardino, J. G.; Wiseman, E. H.; McLamore, W. M. *J. Med. Chem.* **1971**, *14*, 1171–1175.
6. Schapira, C. B.; Perillo, I. A.; Lamdan, S. *J. Heterocycl. Chem.* **1980**, *17*, 1281–1288.
7. Lazer, E. S.; Miao, C. K.; Cywin, C. L.; *et al.* *J. Med. Chem.* **1997**, *40*, 980–989.
8. Pflum, D. A. *Gabriel–Colman Rearrangement*. In *Name Reactions in Heterocyclic Chemistry*; Li, J. J., Ed.; Wiley: Hoboken, NJ, **2005**, pp 416–422. (Review).
9. Kapatsina, E.; Lordon, M.; Baro, A.; Laschat, S. *Synthesis* **2008**, 2551–2560.