

---

# Platelets and Airway Diseases

Stefania Momi, Simon Pitchford, Paolo Gresele, and Clive P. Page

---

## Abstract

The role of platelets as inflammatory cells is now well established. Given the peculiar characteristics of the lung circulation, with a broad capillary bed, platelets are especially involved with the physiology of the lungs and play a key role in a number of inflammatory lung disorders. The platelet precursors, megakaryocytes, are detected in the lung microcirculation; moreover platelets with their endothelium-protective and vascular reparative activities contribute to the lung capillary blood barrier integrity. Given the function of the lungs as first wall against pathogen invasion, platelets participate in immune defence of the normal lung. On the other hand, platelets may turn into effectors of the inflammatory reaction of the lungs to allergens, to infectious agents, to chemical agents and may contribute strongly to the perpetuation of chronic inflammatory reactions, largely by their ability to interact with other inflammatory cells and the endothelium. In this chapter we provide an overview of the role of platelets in several inflammatory lung disorders discussing the pathophysiologic bases of platelet involvement in these conditions and the experimental and clinical evidence for a role of platelets in lung diseases.

Platelets have a well-established role in haemostasis and thrombosis, but there is now considerable evidence that they also play an important role in a number of inflammatory diseases (see Ware and Post 2017), including respiratory diseases such as asthma (Pitchford and Page, *Clin Exp Allergy* 36: 399–401, 2006) and chronic obstructive pulmonary disease (COPD) (Ferroni et al. *J Investig Med* 48:21–27, 2000). There is a relevant amount of information

on the involvement of platelets in allergic inflammatory diseases, and platelet abnormalities in patients with allergy have been reported for more than 40 years following the seminal observation by Benveniste et al. in 1972 that leukocyte-dependent histamine release from platelets involved IgE-mediated activation (Benveniste et al. *J Exp Med*. 136:1356–77, 1972). This work led to the discovery of the lipid mediator platelet-activating factor (PAF) capable of inducing eosinophilia (Arnoux et al. *Am Rev Resp Dis* 137: 855–860, 1988). It is becoming increasingly apparent that platelets are central to inflammatory diseases of the airways, being a source of inflammatory mediators (see Table 1) and spasmogens per se and being critical for the recruitment of leukocytes into tissues.

---

S. Momi • P. Gresele

Section of Internal and Cardiovascular Medicine, Department of Medicine, University of Perugia, Perugia, Italy  
e-mail: [paolo.gresele@unipg.it](mailto:paolo.gresele@unipg.it)

S. Pitchford

The Sackler Institute of Pulmonary Pharmacology, Institute of Pharmaceutical Science, King's College London, London, England, UK

C.P. Page (✉)

Sackler Institute of Pulmonary Pharmacology, GKT School of Biomedical Sciences, King's College London, 5th Floor, Hodgkin Building, Guy's Campus, London Bridge, London SE19RT, UK  
e-mail: [clive.page@kcl.ac.uk](mailto:clive.page@kcl.ac.uk)

## Platelets in Asthma

### Pathophysiologic Observations: Altered Platelet Functionality

- Platelets accumulate into the lungs upon allergen exposure.
- Platelet activation by allergen via an IgE-mediated process reveals an important causal involvement of platelets in the inflammatory response.
- Studies in animals reveal platelets are necessary for pulmonary leukocyte recruitment, chronic inflammatory events leading to airway remodelling, and the release of spasmogens to cause bronchoconstriction. Emerging reports suggest platelets are involved in initial allergen processing with antigen-presenting cells.
- Many of these processes can be attributable to a direct effect of platelets, in their capacity to migrate through lung tissue and be present in the localised environment.

In 1978, Gallagher and colleagues observed that platelets isolated from the peripheral blood of allergic patients during the allergy season often showed reduced secondary wave platelet aggregation, and it was suggested that they behaved like ‘exhausted’ platelets (Gallagher et al. 1978). Since this early observation, there have been many other reports of altered platelet behaviour in patients with allergic disease, ranging from altered arachidonic acid metabolism to greater turnover of intra-cellular signalling pathways (Audera et al. 1988; Block et al. 1990). Indeed, these platelet abnormalities may well account for the observation that patients with allergic asthma have a mild haemostatic defect and delayed thrombin generation detected using the template bleeding time (Szczeplik et al. 1986). This ‘exhausted platelet’ syndrome may be the consequence of continuous platelet activation *in vivo* and is compatible with the observation that allergic patients undergoing allergen provocation have a mild thrombocytopenia (Kowal et al. 2006; Sullivan et al. 2000; Maestrelli et al. 1990) with increased numbers of platelet–leukocyte aggregates in their peripheral blood. Such observations clearly support the conclusion that platelet activation accompanies allergic exposure in such patients (Gresele et al. 1982, 1993; Pitchford et al. 2003; Sladek et al. 1990). This may explain recent observations suggesting that patients with asthma often have co-morbidities, including cardiovascular disease (Cazzola et al. 2013) and an increased risk of pulmonary embolism (Majoor et al. 2013), particularly patients with more severe asthma possibly associated with use of glucocorticosteroids (Varas-

Lorenzo et al. 2007). This hypothesis is further supported by a growing number of clinical observations showing increased markers of platelet activation in allergic asthmatics. These include RANTES (CCL5), platelet factor 4 (PF4), CXCL4 and  $\beta$ -thromboglobulin ( $\beta$ -TG, CXCL7) in the peripheral blood of patients with asthma (Maccia et al. 1977; Kowal et al. 2006; Yasuba et al. 1991; Knauer et al. 1981), as well the demonstration of increases in some of these platelet markers in bronchoalveolar lavage (BAL) fluid obtained from such patients (Jeffery et al. 1989; Metzger et al. 1987) (Table 1). Furthermore, there are other observations demonstrating extravascular platelets in the airways of both experimental animals (Pitchford et al. 2008) and in patients with asthma (Jeffery et al. 1989; Metzger et al. 1987) (see Petito et al. 2017).

In the context of respiratory diseases, it is of interest that platelets have long been recognised as a rich source of spasmogens (Arnoux et al. 1988; Yoshimi et al. 2001) that can lead to both contraction of airway smooth muscle *in vitro* (Inoue and Kannan 1989) and bronchoconstriction *in vivo* (Robertson and Page 1987; Yoshimi et al. 2001). Indeed, we have recently reported that a range of inflammatory mediators, exemplified by bradykinin, elicit airway obstruction *in vivo* that is platelet dependent, building on earlier observations of platelet involvement in airway obstruction (Keir et al. 2015; Coyle et al. 1993). However, perhaps of greater interest for respiratory diseases are the increasing numbers of observations demonstrating a crucial role for platelets in the recruitment of leukocytes into various tissues, including the lung (Pitchford et al. 2003, 2005; Kornerup et al. 2010). Platelet-derived mediators are known to be able to induce leukocyte activation and recruitment (Page and Pitchford 2013), as well as the release of substances that contribute to the remodelling and repair of tissues after injury, making them well placed to contribute to some of the important features of asthma (Fig. 1). Furthermore, it is now recognised that platelets from allergic asthma patients express both the high- and the low-affinity IgE receptors on their surface and that exposure to sensitising antigens can lead to the generation of inflammatory mediators, such as oxygen free radical species, 5-HT and RANTES (Joseph et al. 1983, 1997; Hasegawa et al. 2001; Klouche et al. 1997). Interestingly, activation of the high-affinity IgE receptor can cause platelets to undergo chemotaxis (Pitchford et al. 2008), confirming other observations that suggest that platelets behave like primitive leukocytes and demonstrate directional movement in response to certain stimuli (Czapiga et al. 2005; Kraemer et al. 2010), stimuli that tend to be distinct from those that elicit platelet aggregation (Abi-Younes et al. 2001; Kowalska et al. 2000; Clemetson et al. 2000).

Platelets have long been recognised to contain high concentrations of 5-hydroxytryptamine (5-HT or serotonin) in their dense granules, and several clinical observations

**Table 1** Principal platelet-derived inflammatory mediators and their involvement in airway diseases

Mediator	Action
$\alpha\beta3$	NF- $\kappa$ B activation and induction of inflammatory cell responses (cell adhesion and transmigration into the lung)
CD40L	TNF superfamily: antigen-presenting cell activation, B-cell responses, endothelial cell activation
P-selectin	Selectin: leukocyte adhesion, complement activation
IL-1 $\beta$	Cytokine: acute phase response, leukocyte and endothelial activation
TXA2	Eicosanoid: T-cell differentiation, monocyte activation, smooth muscle proliferation
Nitric Oxide (NO)	Reactive oxygen species: anti-inflammatory and antithrombotic
ADP	Platelet, leukocyte, endothelial cell activation
TNF $\alpha$	Modulate air wall remodelling
LTB4	Modulate leukocyte recruitment
CXCL-7/ $\beta$ -TG	Modulate leukocyte recruitment, bronchospasm
CCL5/RANTES	Monocyte adhesion on the endothelium and recruitment
CXCL4/PF-4	Chemokine: monocyte, neutrophil and T-cell recruitment, Th differentiation, bronchospasm
TGF $\beta$	Cytokine: cell proliferation, T-cell differentiation, B-cell and macrophage phenotype regulation
PDGF	Growth factor: cell growth and differentiation, monocyte/macrophage differentiation
VWF	Platelet adhesion, PMN extravasation
SDF-1	Chemokine: T cell, monocyte and PMN chemotaxis
VEGF	Growth factor: angiogenesis, adhesion molecule expression
5-HT	Dendritic cell activation, bronchospasm


have shown altered levels of this vasoactive amine in patients with asthma (Maccia et al. 1977; Malmgren et al. 1982; Ring et al. 1980). Furthermore, some studies have shown a beneficial effect of the 5-HT antagonist ketanserin in asthma (Cazzola et al. 1990, 1992). Tryptophan hydroxylase (THP)-1 is a critical enzyme for the biosynthesis of 5-HT outside of the central nervous system (CNS), and it is of interest that Durk et al. have utilised mice genetically deficient in THP-1, as well as mast cell-deficient mice, to demonstrate that platelets, rather than mast cells, are the main source of 5-HT released during an allergic inflammatory response (Dürk et al. 2013). Unlike mice, human mast cells are not a major source of 5-HT and so it is of even more interest that Idzko and colleagues have found elevated 5-HT levels in BAL fluid following segmental bronchial allergen challenge in allergic asthmatics, as their results suggest that the 5-HT found in the lung likely comes from activated platelets (Dürk et al. 2013). These results provide further support to other clinical observations reporting platelet activation in patients with allergic asthma (Maccia et al. 1977; Kowal et al. 2006; Pitchford et al. 2003; Johansson et al. 2012). It is of particular interest that platelets have now been found extravascularly in a number of inflammatory situations (Feng et al. 1998), including in the lungs of experimental animals (Pitchford et al. 2008) and patients with asthma (Jeffery et al. 1989). It becomes intriguing to know whether the elevated 5-HT found in BAL of allergic asthmatics is derived from circulating platelets and/or from platelets that have undergone diapedesis into lung tissue. Either way the results presented by Durk et al. are of further interest as they have reported that allergic mice deficient in THP-1 or treated with an inhibitor of THP-1, PCPA, exhibit

reduced leukocyte infiltration into the lung and inhibition of the bronchial hyperresponsiveness (BHR) that normally accompanies allergen challenge, suggesting that the platelet-derived 5-HT is playing a central role in allergic inflammation in the lung. Given that there are some encouraging early clinical observations in patients with asthma administered drugs either affecting 5-HT uptake (Lechin et al. 1998) or antagonising 5-HT<sub>2</sub> receptors (Lima et al. 2007), it would seem timely to consider large trials of such agents in patients with allergic airways disease.

### Studies in Experimental Animal Models: Leukocyte Recruitment

There is now a growing body of literature that suggests that platelets play a central role in both allergic (Pitchford et al. 2003, 2005) and non-allergic (Kornerup et al. 2010) leukocyte recruitment into the lung, as well as a critical role in other manifestations of allergic asthma such as airway remodelling (Pitchford et al. 2004). The mechanisms by which platelets influence these processes are discussed below and open up the possibility of identifying a number of potential new targets for the treatment of inflammatory diseases such as asthma and COPD.

Investigation of the mechanics of blood flow (haemoreology) has uncovered interesting phenomena that have fundamental implications related to how platelets are required to initiate the 'adhesion cascade' that results in leukocyte recruitment into tissues associated with inflammatory airway diseases. In laminar flow in blood vessels, a shearing motion occurs due to the influence of wall friction that results in a



**Fig. 1** There is evidence for platelet participation in both innate immune surveillance and the adaptive response to allergen and inflammatory events that occur after subsequent secondary exposure. This

picture was originally published in Idzko, Pitchford, and Page *J Allergy Clin Immunol* (2015) 135: 1416-1423

parabolic velocity profile of the fluid (Hagan–Poiseuille Law); because blood is a non-Newtonian fluid, the viscosity of blood decreases with increasing shear rate and thus red blood cells actually aggregate to form reversible rouleaux under conditions of low shear and move inwardly (due to their relative deformability compared to platelets and leukocytes) into the inner part of the vessel (Goldsmith and Spain 1984a, b; Goldsmith et al. 1981, 1999). This phenomenon is known as the Fahraeus effect, the consequence of which is a redistribution of the blood elements, with the density of platelets and leukocytes increasing around the vessel periphery. The region of the vessel where the velocity gradient is highest is around the vessel wall; thus cells travelling at different velocities along adjacent streamlines have increased probability of cellular collisions. This peripheral zone therefore ‘traps’ leukocytes into an environment rich in platelets, greatly enhancing collisions between platelets and leukocytes (Goldsmith and Spain 1984a, b; Goldsmith et al. 1981, 1999) and leading to the tethering of platelets to leukocytes to form rosettes through P-selectin recognition steps, to the subsequent upregulation of integrin expression and to the firm adhesion to endothelium as platelet-bound leukocytes enter the capillary network. High-resolution video microscopy has revealed the existence of membrane tethers involving P-selectin/PSGL-1 bonds that regulate leukocyte rolling on platelets and P-selectin with changes in tether length and lifetime dependent on increasing shear force (Schmidtke and Diamond 2000). The biomechanics of the complex formation of activated platelets with leukocytes reveals that the high tensile strength of P-selectin/PSGL-1 interactions enables P-selectin-dependent tethering at high shear rates, whereas integrin activation may mediate platelet–leukocyte complex formation at low shear rates. Nevertheless, unstimulated leukocytes (which constitutively express PSGL-1/L-selectin) may also bind to activated platelets in an integrin-independent manner, suggesting that cell adhesion that is not purely selectin dependent may create platelet–leukocyte complexes (Xiao et al. 2006).

These events occur in the circulation of patients with asthma upon allergen challenge. Thus, circulating platelet–leukocyte complexes are found in allergic asthmatic patients after both spontaneous asthma attacks and after allergen challenge, in a biphasic manner (Gresele et al. 1993; Pitchford et al. 2003). The possible significance of these platelet–leukocyte complexes is to act as a ‘priming’ step for further leukocyte adhesion, since leukocytes attached to platelets display enhanced expression of the  $\beta$ 1 integrin MAC-1 (Pitchford et al. 2003, 2005; Johansson et al. 2012). It is therefore of considerable interest that recent research has correlated platelet activation with eosinophil inflammation in patients with asthma (Benton et al. 2010). Furthermore, in patients with asthma,  $\beta$ 1 integrin expression on eosinophils correlated with eosinophil-bound platelets expressing P-selectin and antigen challenge leads to the

disappearance from the circulation of eosinophils bearing platelet P-selectin, presumably because these complexes are sequestered in the lungs (Johansson et al. 2012). The mechanistic significance of these platelet/eosinophil interactions is that this leads to increased eosinophil adhesion to the vascular endothelium via a platelet-P-selectin-dependent mechanism (Jawień et al. 2002; Ulfman et al. 2003).

The importance of platelets in pulmonary leukocyte recruitment after allergen challenge has now been extensively proven using *in vivo* models of platelet depletion. In fact platelet depletion, either via immunological or non-immunological methods, strongly reduces pulmonary eosinophil and lymphocyte recruitment in rabbits, guinea pigs and mice (Pitchford et al. 2003, 2005; Coyle et al. 1990; Lellouch-Tubiana et al. 1988) and the recruitment of effector T cells in murine models of contact hypersensitivity (Ludwig et al. 2004; Tamagawa-Mineoka et al. 2007). This process required intact platelets, since the reinfusion of lysed platelet products was insufficient to restore leukocyte recruitment, whereas the reinfusion of intact activated platelets expressing selectins on the cell surface restored leukocyte recruitment (Pitchford et al. 2005; Tamagawa-Mineoka et al. 2007). With similarities to allergic asthmatic patients undergoing allergen challenge, mice sensitised to allergen have circulating leukocytes attached to platelets displaying significantly increased CD11b (integrin  $\alpha$ M) and VLA-4 (very late antigen-4) in comparison with leukocytes not attached to platelets (Pitchford et al. 2003, 2005). Contact with platelets therefore induces activation of leukocytes, thus enhancing the expression of integrins, presumably for firm adhesion. This has been confirmed by the experimental use of antibodies blocking P-selectin which results in the suppression of platelet–leukocyte complexes, integrin expression and subsequent tissue recruitment in models of asthma and chronic contact hypersensitivity (Pitchford et al. 2005; Ludwig et al. 2004; Tamagawa-Mineoka et al. 2007; Symon et al. 1999; Mayadas et al. 1993; Lukacs et al. 2002; De Sanctis et al. 1997; Broide et al. 1998). Indeed, the targeting of P-selectin or PSGL-1 as a novel therapeutic option is currently progressing in phase II clinical trials in patients with asthma (Bedard and Kaila 2010).


Nevertheless, many other studies also reveal that other mediators released, or expressed, by platelets can modulate leukocyte recruitment. Examples are the platelet-specific chemokines PF-4 (CXCL4),  $\beta$ -TG (CXCL-7) and RANTES (CCL5) and pleiotropic mediators such as leukotrienes, 5-HT and sphingolipids (Page and Pitchford 2013; Brandt et al. 2000; Duerschmied et al. 2013; Florey and Haskard 2009). Thus, whilst there is a requirement for platelets expressing adhesion molecules on their surface for leukocyte recruitment, this must also be sequential to the release or expression of other platelet-derived factors. However, it is not yet understood from a physiological perspective why

the process of leukocyte activation and adhesion to post-capillary venules is so inefficient and actually requires platelet activation. Perhaps the rapidity of platelet activation to danger signals combined with a higher surface area/volume ratio, allowing a critical density of adhesion molecules to be expressed compared to larger leukocytes, has resulted in an evolutionary requirement for platelets in the rheological processes that trigger the leukocyte adhesion cascade.

### Pathophysiologic Observations: The Extravascular Presence of Platelets

Current perceptions concerning platelet function in inflammation are heavily influenced by the knowledge of the intravascular role of platelets in haemostasis and thrombosis. Thus, dogma suggests that the participation of platelets in inflammation is also confined to intravascular events and therefore that it is indirect and totally dependent on the ability of platelets to influence leukocyte recruitment. However, accumulating evidence details a novel function of platelets in being able to respond to chemotactic signals and migrate extravascularly through inflamed tissue (see Petito et al. 2017) (Fig. 2).

Platelets have been observed to undergo diapedesis in sections of the lungs obtained from patients with asthma and the lungs (and BAL fluid) from allergen-sensitised and allergen-exposed mice, rabbits and guinea pigs (Jeffery et al. 1989; Metzger et al. 1987; Pitchford et al. 2008; Lellouch-Tubiana et al. 1988; Beasley et al. 1989). Using quantitative histology, we have recently reported that the migration of platelets into lung tissue and the localisation of platelets around the airway wall in allergen-sensitised and allergen-exposed mice was an IgE-mediated process and that platelets from such mice could also undergo chemotaxis to the sensitising allergen *in vitro* (Pitchford et al. 2008). Interestingly, the migratory response of platelets in the lungs *in vivo* commenced before significant leukocyte recruitment, whilst at later time points, when leukocytes had also entered lung tissue, around 50 % of platelets were still not complexed to leukocytes (Pitchford et al. 2008). The rapidity of this response highlights that platelet activation by allergen is direct and independent of activation of other cells, like mast cells (Yoshida et al. 2002). It is interesting to note that evidence is accumulating of platelet migration into tissues in other inflammatory diseases, such as into the synovial fluid in patients with rheumatoid arthritis, and transmigration across the vascular wall after long periods of ischaemia (see Ware and Post 2017) (Boilard et al. 2010; Kraemer et al. 2010). The mechanisms controlling this platelet function have not been elucidated, but platelets express a number of different chemokine receptors (CCR1, CCR3, CCR4 and CXCR4) which are functional,



**Fig. 2** Histological analysis of the lungs taken from ovalbumin (OVA)-sensitised mice after allergen inhalation. Section was stained for the platelet-specific antigen CD41 (integrin  $\alpha$ IIb). Platelets (*brown*) can be observed throughout the airway wall. Image originally published in Page & Pitchford (2014) *Clin Exp Allergy* 44: 901–913

since the ligands SDF-1  $\alpha$ , MDC and TARC can activate platelets (Abi-Younes et al. 2001; Kowalska et al. 2000; Clemetson et al. 2000). Platelets have also been shown to undergo chemotaxis to f-MLP and SDF-1 $\alpha$  and can therefore be considered as motile cells (Czapiga et al. 2005; Kraemer et al. 2010). Therefore, platelets can migrate through lung tissue and localise to specific resident cells/structures in a highly regulated process, as it is for leukocytes. Whilst the significance of platelet migration into lung tissue has not yet been fully characterised, platelets may directly influence the development of sensitisation towards allergen, BHR, bronchospasm, tissue damage and chronic inflammation leading to airway wall remodelling.

### Studies in Experimental Animal Models: Involvement with Antigen-Presenting Cells and Contact with Allergen

Platelets express both the high-affinity (Fc $\epsilon$ RI) and low-affinity (CD23) receptors for IgE, as well as functional receptors for other immunoglobulins. Platelets can be activated by specific allergen via IgE-dependent processes, as revealed when platelets are taken from patients allergic to *Dermatophagoides pteronyssinus* (Der p1) and exposed to synthetic peptides derived from the allergen Der p1 *ex vivo* (Cardot et al. 1992). We have shown that platelets from

patients with asthma will undergo chemotaxis specifically towards the known allergen (rather than allergens to which individual patients are not allergic too) (Pitchford et al. 2008). The implications of direct platelet activation and motility by allergen are not yet known, but it is interesting to note that platelets activate dendritic cells (DCs) in the airways (Dürk et al. 2013). Both CD40 and CD40L have been identified on activated platelets and might be responsible for this cellular interaction (Semple et al. 2011; Henn et al. 1998). Platelet CD40L has been described in other situations as an important link between the innate and adaptive immune response to induce DC maturation, for example, the expression of CD80 and CD83, and immunoglobulin class switching (Elzey et al. 2003; Czapiga et al. 2004; Sprague et al. 2008). Platelet CD40L has recently been reported to be involved in the promotion of allergic airway inflammation by polarising Th2 responses after allergen exposure (Tian et al. 2015). Despite this evidence, it remains to be understood whether platelets are involved in the initial allergen sensitisation process *per se*.

### Studies in Experimental Animal Models: Lung Function

The observation that platelets migrate into the lung tissue of patients with asthma, and into the lungs of allergen-challenged sensitised animals, opens the possibility that platelets may contribute directly to alterations in lung function in patients with asthma. For example, platelet depletion in allergen-sensitised rabbits and guinea pigs abolishes bronchoconstriction and anaphylaxis induced by inhaled spasmogens or allergens, respectively (Coyle et al. 1990; Lellouch-Tubiana et al. 1988). There is now some understanding of the pathways and platelet mediators involved in these processes from the observations on the effects of intravenous platelet agonists on bronchospasm and platelet accumulation in the lung (Arnoux et al. 1988; Yoshimi et al. 2001; Lellouch-Tubiana et al. 1988; Robertson and Page 1987). We have recently observed that platelet depletion inhibits bronchospasm induced by ‘indirect spasmogens’, such as capsaicin and bradykinin, whilst it does not inhibit direct-acting spasmogens, such as histamine and methacholine (Keir et al. 2015), suggesting that platelet-derived mediators contribute to airway obstruction under certain circumstances. Furthermore, the inhibition of the release of bronchoactive agents from platelets abrogated the resulting changes in airway obstruction, confirming that platelet-derived mediators might also contribute to airway tone (Arnoux et al. 1988; Yoshimi et al. 2001). Indeed, a direct participation of platelets in allergy, independent of leukocyte responses, was highlighted by the intradermal injection of supernatants from activated human platelets (but not leukocytes) inducing delayed, sustained inflammatory

responses in the skin of patients with atopic dermatitis (Matsuda et al. 1997). These effects on tissue suggest that platelets are very capable of directly inducing sustained inflammation. Human platelets synthesise and release a number of bronchoactive mediators, for example, histamine, 5-HT, TXA<sub>2</sub>, adenosine and 12-hydroxyeicosatetraenoic acid (12-HETE), contain cytotoxic compounds within their granules and generate substances capable of inducing tissue damage, such as reactive oxygen species (ROS), cationic proteins (PCPs), platelet basic proteins (PBPs) and matrix metalloproteinases (Saxena et al. 1989; Knauer et al. 1984; Busti et al. 2010). Yet it is not understood how these mediators interact, and with what (e.g. sensory nerves, airway smooth muscle, epithelium), to elicit bronchospasm and BHR.

### Studies in Experimental Animal Models: Chronic Inflammation and Lung Remodelling

One consequence of persistent, chronic inflammation is the alteration of tissue structure and function. In bronchial asthma, chronic inflammation contributes to changes in airway architecture referred to as ‘airway remodelling’. The observation that platelets migrate into lung tissue of patients with asthma and in experimental animals (Jeffery et al. 1989; Metzger et al. 1987; Pitchford et al. 2008; Lellouch-Tubiana et al. 1988; Beasley et al. 1989) suggests that platelets may contribute directly to changes in the airway architecture by releasing factors that control the synthetic phenotype of the airway epithelium and of fibroblasts and airway smooth muscle cells. Indeed, in murine models of chronic allergic inflammation, the depletion of platelets led to a comprehensive suppression of lung remodelling (smooth muscle hyperplasia, subepithelial fibrosis, collagen deposition, epithelial hyperplasia), an effect that chronic treatment with glucocorticosteroids did not attain, suggesting that platelet involvement in tissue remodelling may in some instances be independent of leukocyte-associated inflammation (Pitchford et al. 2004). In patients with asthma, platelet activation was shown to persist for some time after the late asthmatic response had occurred (Kowal et al. 2006), even though the increases in circulating platelet–leukocyte aggregates returned to basal levels 24 h after allergen exposure (Pitchford et al. 2003), thus implicating platelets in chronic inflammatory events and airway remodelling.

Platelets are a rich reservoir of mitogens and enzymes and therefore may contribute to generate an environment that induces synthetic responses in structural airway cells. Platelet mitogens include PDGF, EGF (epidermal growth factor), TGF- $\beta$  (transforming growth factor- $\beta$ ), VEGF (vascular endothelial growth factor) and the major product of arachidonic acid metabolism in platelets, TXA<sub>2</sub>, which all have proliferative activity on structural cells of the airways (Pitchford and Page 2002; Rendu and Bohard-Bohn 2002).

Platelet-derived enzymes include matrix metalloproteinases,  $\beta$ -hexosaminidases and heparanases that are released following allergen challenge in asthmatic patients and following ozone challenge in guinea pigs and may alter the composition of the extracellular matrix (Falcinelli et al. 2005; Kelly et al. 2000). Disruption of the composition and integrity of cell membranes by degradation of glycoproteins, glycolipids and glycosaminoglycans may also release membrane-bound growth factors for wound repair (McKenzie 2007). Recent evidence shows that platelet membranes are required to induce synthetic responses in airway smooth muscle cells (Svensson Holm et al. 2011), and this function was controlled by platelet 5-lipoxygenase, 12-lipoxygenase and the production of reactive oxygen species (Svensson Holm et al. 2008, 2014). These effects may be supplemented by the ability of platelets to influence the survival, recruitment, proliferation and differentiation of circulating stem cells involved in (inappropriate) tissue regeneration (Stellos et al. 2010). It can be surmised therefore that platelets may influence lung regeneration as well as inappropriate remodelling of the airways after injury, although the interplay between platelets and different structural cells is likely to be extremely complex and involve a plethora of mediators.


### Platelets in Infectious Pulmonary Disease

- Platelets have a broad repertoire of receptor molecules that enable them to sense invading pathogens and infection-induced inflammation.
- Consequently, platelets exert direct antimicrobial mechanisms and initiate an intense crosstalk with other cells of the innate and adaptive immunity, including neutrophils, monocyte/macrophages, dendritic cells and B and T lymphocytes, favouring antimicrobial activity.

In mammals, platelets have retained key structures and functions of immune effector cells which are integral to antimicrobial host defence (Morrell et al. 2014). In the bloodstream they act as quiescent ‘sentinels’ of tissue injury and microbial threat. Platelets exert activities that span from acute inflammation to adaptive immune responses that confer to them an important role as immune regulators during sepsis (Semple et al. 2011). Platelets store antimicrobial peptides, such as  $\beta$ -defensins, thrombocidins, PF-4 (CXCL4), RANTES (CCL5), connective tissue-activating peptide-3, platelet basic protein, thymosin  $\beta$ -4, fibrinopeptide B and fibrinopeptide A, and contribute to the innate immune system (Tang et al. 2002; Kraemer et al. 2011). Inflammatory lung diseases and other pulmonary disorders cause accumulation of platelets in the lungs, and tissue injury and systemic conditions such as sepsis can activate platelets in the circulation and sequester them in the pulmonary vascular bed (Fig. 3).

### Pathophysiologic Observations

Platelets participate in inflammatory and immune responses using a variety of molecular mechanisms, including the release of chemokines ( $\beta$ -TG, PF4, ENA-78, Gro- $\alpha$ , RANTES, SDF1 $\alpha$  and others), cytokines (HMBG1) and growth factors (PDGF, VEGF, EGF, FGF, TGF $\beta$ , and so on), and the synthesis of lipid mediators, like thromboxane A2 and PAF, but contribute also to the maintenance of the endothelial barrier. Key differences exist between the characteristics of endothelium in the pulmonary circulation and the systemic vasculature. Notably, ICAM-1 and integrin  $\alpha_v\beta_3$  are highly expressed by the pulmonary circulation as compared to other vascular beds (Panes et al. 1995; Singh et al. 2000). It is thus possible that platelet retention in the lungs is mediated by the constitutive expression of these two adhesion molecules by pulmonary endothelial cells. Activation of platelets, which is a requirement for their firm adhesion to endothelial cells, results in a conformational change



**Fig. 3** Direct interaction of bacteria with platelets can lead to aggregation, release of antimicrobial factors and pro-inflammatory molecules and formation of platelet–leukocyte aggregates. Under some conditions, bacterial toxins of several classes can trigger these


responses of human and murine platelets. Evidence suggests that these activities mediate immobilisation and killing of bacteria and other pathogens. In addition, however, each can contribute to local thrombosis and inflammation



in platelet  $\alpha_{IIb}\beta_3$  that increases its affinity for fibronectin, which can then act as a bridge to endothelial cell  $\alpha_v\beta_3$  in the low shear stress circulation of the pulmonary vascular bed (Goldenberg and Kuebler 2015). Moreover, platelets can induce inflammatory responses by expressing CD40L which is recognised by surface CD40 on endothelial cells which will then secrete adhesion molecules and TF (Freedman 2003). In addition, activated platelets display P-selectin which mediates their adhesion to PMN by binding to PSGL-1, favouring tethering and rolling on the endothelium, thereby guiding neutrophils to transmigrate (Weyrich et al. 2009) (Fig. 4).

Platelets express a ligand for triggering receptor expressed on myeloid cells (TREM)-1, a surface receptor that plays important roles in innate and adaptive immunity expressed by neutrophils and monocytes. Platelets bound to endothelium support the engagement of neutrophil TREM-1 and result in increased expression of pro-inflammatory chemokines and cytokines, thus amplifying the inflammatory response (Haselmayer et al. 2007). Platelets additionally function as a threshold switch for the release of neutrophil extracellular traps (NETs), web-like structures of DNA that can trap and kill microbes in the vasculature, by neutrophils (Brinkmann et al. 2004) (Rondina et al. 2017; Pryzdial et al. 2017; Foote et al. 2017). This occurs primarily in small vessels, like the liver sinusoids and pulmonary capillaries, where NETs, and particularly the histone components, are responsible for death of lung epithelial and endothelial cells, often with consequent tissue damage and organ dysfunction (Saffarzadeh et al. 2012).

Platelets also express Toll-like receptors (TLRs), a highly conserved family of pattern recognition receptors expressed from flies to mammals. TLRs bind pathogen-associated molecules that are broadly expressed by many infectious organisms. LPS is the most studied TLR ligand, and others include unmethylated cytosine guanine dinucleotide, double-stranded RNA and lipoproteins. Platelets express low levels of TLR2 (Aslam et al. 2006), TLR4 and TLR9 (Cognasse et al. 2005). In addition, additional TLRs and other secondary signalling molecules can become expressed following platelet activation. For example, high concentrations of thrombin induce the expression of TLR9 by platelets. TLR2 and TLR4 are functional in inflammatory responses such as sepsis, e.g. TLR4 triggers the expression of IL-6 and cyclooxygenase 2 (COX-2) in response to LPS (Scott and Owens 2008). LPS has been shown to induce thrombocytopenia and platelet accumulation in the lungs of wild-type but not TLR4-deficient mice. Moreover, LPS stimulation of platelet TLR4 induces platelet binding to and activation of adherent neutrophils. Indeed, stimulation of platelet TLR2 by synthetic bacterial lipopeptide directly activates the platelet's thrombotic and inflammatory response through the phosphoinositide 3-kinase (PI3-K) signalling pathway (Blair et al. 2009). In sepsis, neutrophils and



**Fig. 4** Interactions between platelets and pulmonary ECs and monocytes in pulmonary infectious disease

platelets co-localise in several organs. Upon stimulation via TLR4, platelets induce neutrophil activation and the formation of NETs that display proteolytic activity, leading to the killing of microbes (Clark et al. 2007). Furthermore, platelets directly kill bacteria by releasing microbicidal peptides or by phagocytosing bacteria (Semple et al. 2011; Kraemer et al. 2011; Krijgsveld et al. 2000). Exposure of platelets to Gram-negative bacteria induces platelet aggregation, activation, P-selectin expression and formation of platelet/neutrophil aggregates via TLR2 (Beaulieu et al. 2011). Moreover, incubation of megakaryocytes with TLR2 leads to the generation of platelets with an enhanced pro-inflammatory gene and protein expression profile (Aslam et al. 2006).

In addition to their roles in innate immunity, platelets act as mediators between innate and adaptive immune response cells (Aslam et al. 2006; Elzey et al. 2003, 2005; Semple et al. 2011). Thus, platelets serve as early effectors of dendritic cells (DCs) activation in tissue injury through the release of CD40L and IL-1 $\beta$  and DCs are essential to link innate and adaptive immunity (Medzhitov and Janeway 2000; Banchereau and Steinman 1998). Platelet-mediated leukocyte adhesion to endothelium may also enhance lymphocyte trafficking during adaptive immunity and host

defence (Diacovo et al. 1996; von Hundelshausen and Weber 2007).

### Studies in Experimental Animal Models

Platelets have a key role in immune surveillance by monitoring the surface of active macrophages (Wong et al. 2013). Platelets survey liver Kupffer cells through transient interactions via their GPIb with VWF constitutively expressed on Kupffer cells. When Kupffer cells capture blood-borne *S. aureus* or *Bacillus cereus*, they upregulate their VWF and the normally transient interaction with platelets becomes a sustained adhesion that actively encases the bacteria for eradication within the first minute of encounter. GPIb-deficient mice are significantly more susceptible to Kupffer cell damage and mortality in infections generated by the above-mentioned bacteria, ostensibly due to inefficient detection of Kupffer cell infection by platelets (Wong et al. 2013). Therefore, the targeting of infected macrophages by platelets seems to be an early and key host defence step to facilitate bacterial clearance and protection against disseminated infection (see Slaba and Kubes 2017).

The depletion of platelets in mice submitted to Klebsiella-induced pneumosepsis enhanced mortality in a way proportional to the extent of platelet depletion, by increasing the release of pro-inflammatory cytokines. Although Klebsiella sepsis did not directly enhance the expression of P-selectin on circulating platelets, the infection did increase the responsiveness of platelets to thrombin receptor stimulation (De Stoppelaar et al. 2014). In accordance, lipopolysaccharide (LPS) has been described to act as a platelet ‘primer’ (Gresele et al. 2008); in fact LPS induces platelet hypersensitivity to subthreshold concentrations of platelet agonists (Zhang et al. 2009).

During severe influenza A virus infection in mice, platelets massively infiltrated bronchoalveolar lavage fluid and were detected in ultrathin cryosections of the lungs staining positive for virus antihaemagglutinin, indicating that platelets actively incorporate influenza virus (Mazur et al. 2007).

### Platelets in Cystic Fibrosis

- Platelets from cystic fibrosis (CF) patients have increased reactivity.

- A positive feedback loop of inflammation and platelet activation, which in turn lead to more inflammation and platelet activation, may exacerbate lung disease in patients with CF.

Cystic fibrosis (CF) is a multi-organ disease which affects the pancreas, the gastrointestinal tract, the male reproductive system and especially the respiratory tract. It is a lethal genetic disorder caused by a mutation in the gene, located in the long arm of chromosome 7, coding for the cystic fibrosis transmembrane conductance regulator (CFTR), a chloride ion channel belonging to the superfamily of the ATP-binding cassette (ABC) proteins (Riordan et al. 1989). Chronic lung inflammation, due to an impairment of mucociliary clearance and to the consequent bacterial colonisation, represents the main cause of mortality and morbidity in CF patients.

### Pathophysiologic Observations

CF patients have an increase in circulating activated platelets and platelet reactivity, as determined by enhanced monocyte–platelet aggregates, neutrophil/platelet aggregates and platelet surface P-selectin (Sturm et al. 2010; O’Sullivan et al. 2005). Moreover, elevated concentrations of sCD40L and TXA2 metabolites, which correlate with decreased lung function, further implicate platelets in the pathophysiology of this lethal disorder (Ciabattini et al. 2000; Falco et al. 2004).

CFTR was found to be expressed by neutrophils and platelets, where it is localised in the cytosol, cell membranes and  $\alpha$ -granules, and when blocked a markedly decreased thrombin-induced AKT phosphorylation and an upregulation of TRAP-stimulated p38MAPK phosphorylation were observed (Mattoscoio et al. 2010). p38 MAPK activation has been associated with platelet secretion of pro-inflammatory mediators; thus the CFTR-dependent increase in p38 MAPK phosphorylation is consistent with a pro-inflammatory profile of CF platelets.

Another abnormality reported in CF patients is the reduction of plasma levels of vitamin E, which may increase fatty acid oxidation and thus the production of isoprostanes, which in turn activate platelets (Ciabattini et al. 2000). Moreover, reduced nitric oxide (NO) generation was also observed in patients with CF and found to be a negative prognostic index associated with bacterial colonisation of the lung and sustained inflammation (Balfour-Lynn et al. 1996; Grasemann and Ratjen 2012). Endothelial dysfunction may also contribute to reduce NO generation in CF

(Poore et al. 2013), and the reduction of NO may contribute to platelet hyperreactivity. Platelets also produce NO upon eNOS activation that may act as a negative feedback mechanism to limit platelet activation and platelet recruitment to a growing thrombus (Cozzi et al. 2015; Momi et al. 2014; Freedman et al. 1997). Furthermore, the blockade of CFTR blunted NO generation by platelets suggesting this as another mechanism of the platelet hyperfunction in CF (Mattoscio et al. 2010).

Platelet 12-lipoxygenase is a key enzyme in the biosynthesis of lipoxins (Romano et al. 1993), arachidonic acid (AA) metabolites with potent anti-inflammatory properties. Lipoxin formation occurs in vivo during platelet–PMN interactions (Gangemi et al. 2003) and represents a main stop signal of inflammation (Serhan et al. 2008). Of interest, CF patients display a defect in the biosynthesis of lipoxin A4 (Karp et al. 2004), whereby mutated CFTR downregulates lipoxin formation by selective inhibition of platelet 12-lipoxygenase activity.

Patients with CF have increased plasma levels of ATP, due to the dysfunction of CFTR which acts as an ATP transporter, as well as a chloride channel, that can contribute to platelet activation (Lader et al. 2000). ATP may in fact act as a platelet agonist in vivo under special circumstances, like under high shear stress conditions, and may thus also contribute to the hyperreactivity of CF platelets (Birk et al. 2002).

Altogether, these data suggest an involvement of platelets in the sustained inflammatory response of CF originating from increased pro-inflammatory platelet-activating mediators and reduced counter-regulatory signals (Mattoscio et al. 2010; O’Sullivan et al. 2005).

Patients with CF show thrombocytosis, increased expression of P-selectin on circulating platelets (O’Sullivan et al. 2005), increased platelet reactivity to ADP and TRAP (Stead et al. 1987) and increased release of platelet-derived mediators (e.g. TNF $\alpha$ , CD40L, LTB4 and interleukins) (Shoki et al. 2013). Moreover, patients with CF display increased circulating platelet–leukocyte and platelet–monocyte complexes (O’Sullivan et al. 2005) and enhanced NET formation in airway fluids correlating with impaired obstructive lung function. These phenomena are blocked by the intra-airway delivery of small-molecule antagonists of CXCR2, with inhibition of NET formation and improved lung function (Marcos et al. 2010). In CF, increased platelet number correlated with decreased PaO<sub>2</sub> (Gross and Luckey 1969), increased urinary thromboxane metabolite excretion with decreased FEV1 (Davì et al. 1995; Ciabattini et al. 2000) and increased plasma sCD40L with decreased pulmonary function (Falco et al. 2004). Taken together these studies suggest a link between platelet activation and progressive impairment of lung function in CF. Whether platelet abnormalities are the cause or consequence of lung

inflammation in CF is unclear. However, platelet dysfunction in CF could be the result of platelet abnormalities intrinsic to CF, of the accumulation of plasma factors unique to CF and/or of generalised inflammation.

## Studies in Experimental Animal Models

Mice carrying the F508del mutation of CFTR developed more severe thrombocytopenia, higher levels of plasma TxB2 and PAF levels and enhanced neutrophilic lung infiltration when challenged with LPS relative to wild-type controls (Zhao et al. 2013). Blockade of PSGL-1 (P-selectin glycoprotein ligand-1) or of P-selectin, PAF-receptor antagonism or correction of mutated CFTR trafficking by KM11060 (an analogue of sildenafil, which restores a function of the F508del-mutated CFTR chloride channel) all significantly increased plasma lipoxin A4 levels, whilst the depletion of platelets significantly decreased plasma lipoxin A4 and enhanced LPS-induced lung inflammation (Wu et al. 2014).

Therefore, a growing body of evidence suggests that platelets play a very important role in CF and along with evidence of platelet involvement in other inflammatory lung diseases suggests that they may represent a potential therapeutic target in CF. However, much work remains to be done before anti-platelet therapy can be recommended for patients with CF.

---

## Platelets in Chronic Obstructive Pulmonary Disease (COPD)

Platelet activation represents a mechanism that may contribute to the increased cardiovascular risk associated with stable COPD and with exacerbations.

Chronic obstructive pulmonary disease (COPD) is a respiratory disorder characterised by progressive lung destruction with airflow limitation associated with an airway and systemic inflammatory response (Rabe et al. 2007). Low-grade, chronic systemic inflammation in COPD is frequently complicated by acute atherothrombotic events, independent of the smoking history or other cardiovascular risk factors (Malerba et al. 2013), and coronary artery disease is one of the leading causes of death in COPD (Hansell et al. 2003). Moreover, COPD patients have an increased risk of ischaemic stroke (Truelsen et al. 2001) and venous thromboembolism, particularly during acute exacerbations (Tillie-Leblond et al. 2006). Although the mechanisms responsible for the association between COPD and atherothrombosis are still largely unknown, platelet activation is one of the proven links.

## Pathophysiologic Observations

A link between hypoxia and platelet activation has been suggested since hypoxia facilitates the activation of cyclooxygenase-1 and thus thromboxane formation (Ponick et al. 1987). More recently, a significantly enhanced urinary 11-dehydro TxB2 excretion, the urinary metabolite of platelet-released TxA2, was reported in patients with COPD, irrespective of the smoking status, that inversely correlated with arterial oxygen tension (Davì et al. 1997). Increased *in vivo* platelet activation in hypoxaemic COPD patients with secondary pulmonary hypertension was also detected by a significant increase in circulating platelet aggregates and plasma  $\beta$ -TG (Cordova et al. 1985), suggesting that activation of platelets in the pulmonary blood vessels may be associated with the induction of pulmonary hypertension.

Moreover, an elevation in arginase activity in platelets, which is associated with a reduction of NO production, has been described in COPD patients, and this may also contribute to enhanced platelet reactivity (Guzman-Grenfell et al. 2011). Increased levels of circulating soluble P-selectin, most likely of platelet origin, have also been observed in COPD patients (Ferroni et al. 2000), and platelet P-selectin is crucial for the formation of platelet–leukocyte aggregates (Totani et al. 2016). It is therefore conceivable that neutrophil recruitment to the lungs, a feature of COPD patients, is favoured by interactions with activated platelets. Platelet–leukocyte interactions mediate the process of NETosis (see Evangelista et al. 2017) and NETs are a major contributor to chronic inflammation and lung tissue damage in COPD (Grabcanovic-Musija et al. 2015). NETs were found in COPD patients and are associated with other markers of activated innate immune response, including the expression of the pro-inflammatory cytokines IL-1 $\beta$  and CXCL8 and of the inflammasome component NLRP3. Consequently NETs may contribute, through a positive feedback and the stimulation of neutrophilic chemokines and cytokines, to the persistent airway neutrophilia observed in COPD (Wright et al. 2016). Further research is required to elucidate the participation of platelets in NETosis and the exact role that NETs are playing in COPD. Furthermore, the use of anti-platelet therapy in patients has been shown to improve survival in patients with oxygen-dependent COPD (Ekstrom et al. 2013).

### Platelets in Acute Lung Injury and Acute Respiratory Distress Syndrome


- Platelet/neutrophil interactions may be major pathogenic contributors in ALI.

- The interaction between platelets and neutrophils is mediated by platelet P-selectin and neutrophil PSGL-1.
- Platelets contribute to NET formation that involves the expulsion of nuclear material embedded with histones, by neutrophils in ALI/ARDS. Histones were found in the bronchoalveolar lavage fluid in ARDS; in a transfusion-related acute lung injury model in mice, depletion of platelets was protective.

Acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) are life-threatening diseases occurring in critically ill patients. They are manifestations of an inflammatory response of the lung to direct and indirect insults and are characterised by severe hypoxaemia, hypercapnia, diffuse inflammatory lung infiltration and impairment of pulmonary compliance (Ragaller and Richter 2010). In particular, ALI is defined as an acute lung disease with bilateral pulmonary infiltrates in a chest radiogram consistent with the presence of oedema, and no evidence of pulmonary hypertension (Pepe et al. 1982). ALI is an important complication of sepsis.

## Pathophysiologic Observations

Growing evidence suggests that platelets play a role in the pathogenesis of ALI (Zarbock et al. 2006). Intrapulmonary triggers, like acid aspiration and pneumonia, and systemic inflammatory stimuli activate the lung microvascular endothelium with upregulation of adhesion molecules, including ICAM-1 (Basit et al. 2006), release and presentation of chemokines (Belperio et al. 2006) and increase of lipid mediators (Zarbock et al. 2006; Kieffmann et al. 2004). During endotoxaemia platelets roll on and adhere to pulmonary capillary endothelial cells, as shown by intravital and electron microscopy (Kieffmann et al. 2006; Zarbock and Ley 2009), and release chemokines and lipid mediators that activate leukocytes and endothelial cells (Zarbock et al. 2006) (Fig. 5). Platelet microparticles (PMPs), small (50 nm to 1  $\mu$ m) circulating cell-derived vesicles that break off from intact platelets upon activation and contain a variety of enzymes and proteins as well as mRNAs (McVey et al. 2012) (see Cointe et al. 2017), may play a role in ALI by promoting the inflammatory response of the lungs through the activation of neutrophil cell surface proteins (e.g. IL-8, MAC-1, PECAM-1, etc.) (Reutershan and Ley 2004; Bastarache et al. 2009; Shang and Yao 2014) and the production of cytokines and inflammatory mediators by endothelial cells (Eickmeier et al. 2013). A recent study showed that PMPs carry membrane-bound sCD40L, promote PMN-mediated HUVEC damage and may affect the



**Fig. 5** Platelet–platelet, platelet–endothelial, platelet–PMN and platelet–endothelial cell–PMN interactions occur in pulmonary arterioles, capillaries and venules. These events can mediate acute lung injury. Different molecular interactions mediate adhesion of

activated platelets to PMNs including P-selectin/PSGL-1, integrin MAC-1/integrin  $\alpha$ IIb $\beta$ 3 or ICAM-2, integrin MAC-1/GPIIb $\alpha$  or JAM-3; ICAM-2/ $\alpha$ L $\beta$ 2 (LFA-1). Platelets also interact with monocytes under these conditions

development of ALI (Xie et al. 2015). Moreover, PMPs from LPS-stimulated platelets induce VCAM-1 production by cultured human endothelial cells (Levy and Serhan 2014).

Neutrophils and platelets are both key players in the pathophysiology of ALI. Thus, in a transfusion-related acute lung injury model in mice, depletion of either neutrophils or platelets was protective (Looney et al. 2009).


NETs are the product of a neutrophil death process that involves the expulsion of nuclear material embedded with histones, proteases, antimicrobial proteins and peptides. Histones have been found in the BAL fluid in acute respiratory distress syndrome (ARDS) (Bosmann et al. 2013). Platelets contribute to NET formation by neutrophils (see Slaba and Kubes 2017). Using an *in vitro* flow chamber system, it was shown that platelet-bound TLR4 detected TLR4 ligands in blood, activated neutrophils and facilitated the formation of NETs in the bloodstream. Moreover, using *in vivo* imaging in mice, it appeared that the lung capillaries where platelets bound neutrophils were hot spots for bacterial trapping and that the removal of platelets impaired bacterial clearance (Clark et al. 2007). Interestingly LPS, even at high concentrations, is not able to induce the formation of NETs in the absence of platelets, suggesting that platelets are not just facilitators in this process but essential mediators. On the other hand, the histone/DNA complexes of NETs activate platelets that in turn further promote NETosis, thrombosis and coagulation (Semeraro et al. 2011). Finally, extracellular histones, known to act as strong drivers of inflammation and tissue damage, activate TLRs resulting in pro-inflammatory cytokine production contributing to ALI development (Ward and Grailer 2014). All these observations point to an involvement of platelets in ARDS. One SNP within the LRR16A gene (rs7766874), associated with variations in the platelet count, was involved in the pathophysiology of

ARDS in a large series of 629 ARDS cases and 1,026 at-risk control subjects. However, it remained unclear whether the reduced platelet counts in patients with the LRR16A SNP affected ARDS risk by a reduced platelet production, increased platelet activation/consumption or platelet sequestration in lung tissues (Wei et al. 2015).

Finally, another evidence has demonstrated that activated coagulation and impaired fibrinolysis are associated with ALI. During ALI, the coagulation system is activated, producing fibrin deposition in the lung (Idell et al. 1989) (Fig. 6). The newly produced thrombin, on the one hand, activates protease-activated receptors (PARs) on endothelial cells, inducing an inflammatory response with upregulation of cytokines as well as thrombin formation, whilst, on the other hand, it also activates platelets by binding to platelet PAR1 and 4 (Coughlin 2005). Thrombin also induces the conversion of fibrinogen to fibrin that, together with activated platelets, can induce the formation of microvascular thrombosis.

## Studies in Animal Models

Inhibition of platelet/neutrophil aggregate formation resulted in reduced neutrophil recruitment, increased survival and lower levels of hypoxia in LPS-induced ALI in mice (Zarbock et al. 2006). In a ventilator-induced lung injury model in rats, increased levels of VWF were found on freshly isolated lung endothelial cells. The increased expression of VWF was inhibited by platelet removal from the lung perfusion and by a P-selectin-blocking antibody, and it was also drastically reduced in the lungs of mice perfused with blood from P-selectin knockout animals. These findings indicate that in ventilation-induced stress, platelets transfer VWF to endothelial cells and that platelet P-selectin plays a critical role in this



**Fig. 6** Platelets and PMNs accumulate in macro- and microvessels in the lungs of patients with ALI/ARDS. Sequestration of platelets and PMNs is frequently associated with fibrin deposition and endothelial injury

transfer (Yiming et al. 2008). These data imply that during mechanical ventilation-induced lung injury, platelets deliver leukocyte-binding proteins to endothelial cells promoting leukocyte recruitment, thus playing a key role in generating a pro-inflammatory milieu.

Platelet depletion markedly reduced lung neutrophil infiltration in a murine model of lung injury induced by aerosolised LPS (Grommes et al. 2012). In the same model, antagonism of the CCL5-CXCL4 interaction reduced lung oedema, neutrophil infiltration and tissue damage, suggesting an important and causative role for platelet-derived chemokines in the development of lung injury (Grommes et al. 2012).

Moreover, in a mouse model of transfusion-associated lung injury (TRALI), platelet inhibition by aspirin or by a glycoprotein IIb/IIIa inhibitor reduced the degree of lung injury and the formation of NETs (Caudrillier et al. 2012). Therefore, platelet inhibition could reduce lung injury in ARDS by reducing the formation of NETs.

### Take-Home Messages

A role for platelets in lung disorders has been revealed by studies in experimental animal models and in patients with different forms of acute and chronic lung injury with an inflammatory component, with platelets being critical for leukocyte recruitment, tissue damage and lung dysfunction. The complex web of cellular interactions that contribute to lung tissue damage, increasingly unravelled, includes now the humble platelet, and the improved understanding of the central pathophysiologic role of this cell may hold the key to improved treatments for these common conditions. It is becoming clear that the ability of

platelets to act as inflammatory cells involves activation and signalling pathways distinct from those involved in haemostasis and thrombosis. This may provide new exciting opportunities for identifying new drug targets for the treatment of inflammatory lung disorders.

### References

- Abi-Younes S, Si-Tahar M, Luster AD (2001) The CC chemokines MDC and TARC induce platelet activation via CCR4. *Thromb Res* 101:279–289
- Arnoux B, Denjean A, Page CP, Nolibé D, Morley J, Benveniste J (1988) Accumulation of platelets and eosinophils in baboon lung after paf-acether challenge: inhibition by ketotifen. *Am Rev Resp Dis* 137:855–860
- Aslam R, Speck ER, Kim M, Crow AR, Bang KW, Nestel FP, Ni H, Lazarus AH, Freedman J, Semple JW (2006) Platelet Toll-like receptor expression modulates lipopolysaccharide-induced thrombocytopenia and tumor necrosis factor- $\alpha$  production in vivo. *Blood* 107:637–41
- Audera C, Rocklin R, Vaillancourt R, Jakubowski JA, Deykin D (1988) Altered arachidonic acid metabolism and platelet size in atopic subjects. *Clin Immunol Immunopathol* 46:352–359
- Balfour-Lynn IM, Laverty A, Dinwiddie R (1996) Reduced upper airway nitric oxide in cystic fibrosis. *Arch Dis Child* 75:319–322
- Banchereau J, Steinman RM (1998) Dendritic cells and the control of immunity. *Nature* 392:245–25
- Basit A, Reutershan J, Morris MA, Solga M, Rose CE Jr, Ley K (2006) ICAM-1 and LFA-1 play critical roles in LPS-induced neutrophil recruitment into the alveolar space. *Am J Physiol Lung Cell Mol Physiol* 291:L200–207
- Bastarache JA, Fremont RD, Kropski JA, Bossert FR, Ware LB (2009) Procoagulant alveolar microparticles in the lungs of patients with acute respiratory distress syndrome. *Am J Physiol Lung Cell Mol Physiol* 297:L1035–L1041

- Beasley R, Roche WR, Roberts JA, Holgate ST (1989) Cellular events in the bronchi in mild asthma and after bronchial provocation. *Am Rev Respir Dis* 139:806–817
- Beaulieu LM, Lin E, Morin KM, Tanriverdi K, Freedman JE (2011) Regulatory effects of TLR2 on megakaryocytic cell function. *Blood* 117:5963–5974
- Bedard PW, Kaila N (2010) Selectin inhibitors: a patent review. *Expert Opin Ther Pat* 20:781–793
- Belperio JA, Keane MP, Lynch JP 3rd, Strieter RM (2006) The role of cytokines during the pathogenesis of ventilator-associated and ventilator-induced lung injury. *Semin Respir Crit Care Med* 27:350–364
- Benton AS, Kumar N, Lerner J, Wiles AA, Foerster M, Teach SJ, Freishtat RJ (2010) Airway platelet activation is associated with airway eosinophilic inflammation in asthma. *J Investig Med* 58:987–990
- Benveniste J, Henson PM, Cochran CG (1972) Leukocyte-dependent histamine release from rabbit platelets. The role of IgE, basophils, and a platelet-activating factor. *J Exp Med* 136:1356–1377
- Birk AV, Broekman MJ, Gladek EM, Robertson HD, Drosopoulos JH, Marcus AJ, Szeto HH (2002) Role of extracellular ATP metabolism in the regulation of platelet reactivity. *J Lab Clin Med* 140:166–175
- Blair P, Rex S, Vitseva O, Beaulieu L, Tanriverdi K, Chakrabarti S, Hayashi C, Genco CA, Iafrazi M, Freedman JE (2009) Stimulation of Toll-like receptor 2 in human platelets induces a thromboinflammatory response through activation of phosphoinositide 3-kinase. *Circ Res* 104:346–354
- Block LH, Imhof E, Emmons LR, Roth M, Perruchoud AP (1990) PAF-dependent phosphatidylinositol turnover in platelets: differences between asthmatics and normal individuals. *Respiration* 57:372–378
- Boilard E, Nigrovic PA, Larabee K, Watts GF, Coblyn JS, Weinblatt ME, Massarotti EM, Remold-O'Donnell E, Farndale RW, Ware J, Lee DM (2010) Platelets amplify inflammation in arthritis via collagen-dependent microparticle production. *Science* 327:580–583
- Bosmann M, Grailer JJ, Russkamp NF, Ruemmler R, Monestier M, Zetoune FS, Sarma JV, Ward PA (2013) Extracellular histones are essential effectors of C5aR and C5L2-dependent tissue damage and inflammation in acute lung injury. *FASEB J* 27:5010–21
- Brandt E, Petersen F, Ludwig A, Ehlert JE, Bock L, Flad HD (2000) The beta-thromboglobulins and platelet factor 4: blood platelet-derived CXC chemokines with divergent roles in early neutrophil regulation. *J Leukoc Biol* 67:471–478
- Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y, Zychlinsky A (2004) Neutrophil extracellular traps kill bacteria. *Science* 303:1532–1535
- Broide DH, Sullivan S, Gifford T, Sriramarao P (1998) Inhibition of pulmonary eosinophilia in P-selectin- and ICAM-1-deficient mice. *Am J Respir Cell Mol Biol* 18:218–225
- Busti C, Falcinelli E, Momi S, Gresele P (2010) Matrix metalloproteinases and peripheral arterial disease. *Intern Emerg Med* 5:13–25
- Cardot E, Pestel J, Callebaut I, Lassalle P, Tscopoulos A, Gras-Masse H, Capron A, Joseph M (1992) Specific activation of platelets from patients allergic to Dermatophagoides pteronyssinus by synthetic peptides derived from the allergen Der p I. *Int Arch Allergy Immunol* 98:127–134
- Caudrillier A, Kessenbrock K, Gilliss BM, Nguyen JX, Marques MB, Monestier M, Toy P, Werb Z, Looney MR (2012) Platelets induce neutrophil extracellular traps in transfusion-related acute lung injury. *J Clin Invest* 122:2661–2671
- Cazzola M, Assogna G, Lucchetti G, Cicchitto G, D'Amato G (1990) Effect of ketanserin, a new blocking agent of the 5-HT<sub>2</sub> receptor, on airway responsiveness in asthma. *Allergy* 45:151–153
- Cazzola M, Matera MG, Santangelo G, Assogna G, D'Amato G, Rossi F, Girbino G (1992) Effect of the selective 5-HT<sub>2</sub> antagonist ketanserin on adenosine-induced bronchoconstriction in asthmatic subjects. *Immunopharmacology* 23:21–28
- Cazzola M, Segreti A, Calzetta L, Rogliani P (2013) Comorbidities of asthma: current knowledge and future research needs. *Curr Opin Pulm Med* 19:36–41
- Ciabattoni G, Davi G, Collura M, Iapichino L, Pardo F, Ganci A, Romagnoli R, Maclouf J, Patrono C (2000) In vivo lipid peroxidation and platelet activation in cystic fibrosis. *Am J Respir Crit Care Med* 162:1195–1201
- Clark SR, Ma AC, Tavener SA, McDonald B, Goodarzi Z, Kelly MM, Patel KD, Chakrabarti S, McAvoy E, Sinclair GD, Keys EM, Allen-Vercoe E, Devinney R, Doig CJ, Green FH, Kubes P (2007) Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood. *Nat Med* 13:463–469
- Clemetson KJ, Clemetson JM, Proudfoot AE, Power CA, Baggiolini M, Wells TN (2000) Functional expression of CCR1, CCR3, CCR4, and CXCR4 chemokine receptors on human platelets. *Blood* 96:4046–4054
- Cognasse F, Hamzeh H, Chavarin P, Acquart S, Genin C, Garraud O (2005) Evidence of toll-like receptor molecules on human platelets. *Immunol Cell Biol* 83:196–198
- Cointe S, Lacroix R, Dignat-George F (2017) Platelet-derived microparticles. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 379–392
- Cordova C, Musca A, Violi F, Alessandri C, Perrone A, Balsano F (1985) Platelet hyperfunction in patients with chronic airways obstruction. *Eur J Respir Dis* 66:9–12
- Coughlin SR (2005) Protease-activated receptors in hemostasis, thrombosis and vascular biology. *J Thromb Haemost* 3:1800–1814
- Coyle AJ, Page CP, Atkinson L, Flanagan R, Metzger WJ (1990) The requirement for platelets in allergen-induced late asthmatic airway obstruction. Eosinophil infiltration and heightened airway responsiveness in allergic rabbits. *Am Rev Respir Dis* 142:587–593
- Coyle AJ, Ackerman SJ, Irvin CG (1993) Cationic proteins induce airway hyperresponsiveness dependent on charge interactions. *Am Rev Respir Dis* 147:896–900
- Cozzi MR, Guglielmini G, Battiston M, Momi S, Lombardi E, Miller EC, De Zanet D, Mazzucato M, Gresele P, De Marco L (2015) Visualization of nitric oxide production by individual platelets during adhesion in flowing blood. *Blood* 125:697–705
- Czapiga M, Kirk AD, Lekstrom-Himes J (2004) Platelets deliver costimulatory signals to antigen-presenting cells: A potential bridge between injury and immune activation. *Exp Hematol* 32:135–139
- Czapiga M, Gao JL, Kirk A, Lekstrom-Himes J (2005) Human platelets exhibit chemotaxis using functional N-formyl peptide receptors. *Exp Hematol* 33:73–84
- Davi G, Iapichino L, Balsano V, Ganci A, Giammarresi C, Patrignani P, Patrono C (1995) Thromboxane biosynthesis and pulmonary function in cystic fibrosis. *Adv Prostaglandin Thromboxane Leukot Res* 23:369–371
- Davi G, Basili S, Vieri M, Cipollone F, Santarone S, Alessandri C, Gazzaniga P, Cordova C, Violi F (1997) Enhanced thromboxane biosynthesis in patients with chronic obstructive pulmonary disease. The Chronic Obstructive Bronchitis and Haemostasis Study Group. *Am J Respir Crit Care Med* 156:1794–9
- de Sanctis GT, Wolyniec WW, Green FH, Qin S, Jiao A, Finn PW, Noonan T, Joetham AA, Gelfand E, Doerschuk CM, Drazen JM (1985) [Reduction of allergic airway responses in P-selectin-deficient mice](#). *J Appl Physiol* 83:681–687
- de Stoppelaar SF, van't Veer C, Claushuis TA, Albersen BJ, Roelofs JJ, van der Poll T (2014) Thrombocytopenia impairs host defense in gram-negative pneumonia-derived sepsis in mice. *Blood* 124:3781–3790
- Diacovo TG, Puri KD, Warnock RA, Springer TA, von Andrian UH (1996) Platelet-mediated lymphocyte delivery to high endothelial venules. *Science* 273:252–255

- Duerschmied D, Suidan GL, Demers M, Herr N, Carbo C, Brill A, Cifuni SM, Mauler M, Cicko S, Bader M, Idzko M, Bode C, Wagner DD (2013) Platelet serotonin promotes the recruitment of neutrophils to sites of acute inflammation in mice. *Blood* 121:1008–1015
- Dürk T, Duerschmied D, Müller T, Grimm M, Reuter S, Vieira RP, Ayata K, Cicko S, Soricther S, Walther DJ, Virchow JC, Taube C, Idzko M (2013) Production of serotonin by tryptophan hydroxylase 1 and release via platelets contribute to allergic airway inflammation. *Am J Respir Crit Care Med* 187:476–85
- Eickmeier O, Seki H, Haworth O, Hilberath JN, Gao F, Uddin M, Croze RH, Carlo T, Pfeffer MA, Levy BD (2013) Aspirin-triggered resolvin D1 reduces mucosal inflammation and promotes resolution in a murine model of acute lung injury. *Mucosal Immunol* 6:256–266
- Eckstrom MP, Hermansson AB, Strom KE (2013) Effects of cardiovascular drugs on mortality in severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 187:715–720
- Elzey BD, Tian J, Jensen RJ, Swanson AK, Lees JR, Lentz SR, Stein CS, Nieswandt B, Wang Y, Davidson BL, Ratliff TL (2003) Platelet-mediated modulation of adaptive immunity. A communication link between innate and adaptive immune compartments. *Immunity* 19:9–19
- Elzey BD, Grant JF, Sinn HW, Nieswandt B, Waldschmidt TJ, Ratliff TL (2005) Cooperation between platelet-derived CD154 and CD4+ T cells for enhanced germinal center formation. *J Leukoc Biol* 78:80–84
- Evangelista V, Totani L, Manfredi AA, Maugeri N (2017) Platelet-leukocyte interactions. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 407–433
- Falcinelli E, Guglielmini G, Torti M, Gresele P (2005) Intraplatelet signaling mechanisms of the priming effect of matrix metalloproteinase-2 on platelet aggregation. *J Thromb Haemost* 3:2526–2535
- Falco A, Romano M, Lapichino L, Collura M, Davi G (2004) Increased soluble CD40 ligand levels in cystic fibrosis. *J Thromb Haemost* 2:557–560
- Feng D, Nagy JA, Pyne K, Dvorak HF, Dvorak AM (1998) Platelets exit venules by a transcellular pathway at sites of F-met peptide-induced acute inflammation in guinea pigs. *Int Arch Allergy Immunol* 116:188–195
- Ferroni P, Basili S, Martini F, Vieri M, Labbadia G, Cordova C, Alessandri C, Gazzaniga PP (2000) Soluble P-selectin as a marker of platelet hyperactivity in patients with chronic obstructive pulmonary disease. *J Investig Med* 48:21–27
- Florey O, Haskard DO (2009) Sphingosine 1-phosphate enhances Fc gamma receptor-mediated neutrophil activation and recruitment under flow conditions. *J Immunol* 183:2330–2336
- Foote S, Burgio G, McMorran B (2017) Platelets in a malarial infection: protective or pathological? In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 1107–1113
- Freedman JE (2003) CD40-CD40L and platelet function: beyond hemostasis. *Circ Res* 92:944–946
- Freedman JE, Loscalzo J, Barnard MR, Alpert C, Keaney JF, Michelson AD (1997) Nitric oxide released from activated platelets inhibits platelet recruitment. *J Clin Invest* 100:350–356
- Gallagher JS, Bernstein IL, Maccia CA, Splansky GL, Glueck HI (1978) Cyclic platelet dysfunction in IgE-mediated allergy. *J Allergy Clin Immunol* 62:229–235
- Gangemi S, Lucioti G, D'Urbano E, Mallamace A, Santoro D, Bellinghieri G, Davi G, Romano M (2003) Physical exercise increases urinary excretion of lipoxin A4 and related compounds. *J Appl Physiol* 94:2237–2240
- Goldenberg NM, Kuebler WM (2015) Endothelial cell regulation of pulmonary vascular tone, inflammation and coagulation. *Compr Physiol* 5:531–59
- Goldsmith HL, Spain S (1984a) Margination of leukocytes in blood flow through small tubes. *Microvasc Res* 27:204–222
- Goldsmith HL, Spain S (1984b) Radial distribution of white cells in tube flow. *Kroc Found Ser* 16:131–146
- Goldsmith HL, Lichtarge O, Tessier-Lavigne M, Spain S (1981) Some model experiments in hemodynamics: VI. Two body collisions between blood cells. *Biorheology* 18:531–555
- Goldsmith HL, Bell DN, Spain S, McIntosh FA (1999) Effect of red blood cells and their aggregates on platelets and white cells in flowing blood. *Biorheology* 36:461–468
- Grabcanovic-Musija F, Obermayer A, Stoiber W, Krautgartner WD, Steinbacher P, Winterberg N, Bathke AC, Klappacher M, Studnicka M (2015) Neutrophil extracellular trap (NET) formation characterises stable and exacerbated COPD and correlates with airflow limitation. *Respir Res* 16:59
- Grasemann H, Ratjen F (2012) Nitric oxide and L-arginine deficiency in cystic fibrosis. *Curr Pharm Des* 18:726–36
- Gresele P, Todisco T, Merante F, Nenci GG (1982) Platelet activation and allergic asthma. *N Engl J Med* 306:549
- Gresele P, Dottorini M, Selli ML, Iannacci L, Canino S, Todisco T, Romano S, Crook P, Page CP, Nenci GG (1993) Altered platelet function associated with the bronchial hyperresponsiveness accompanying nocturnal asthma. *J Allergy Clin Immunol* 91:894–902
- Gresele P, Falcinelli E, Momi S (2008) Potentiation and priming of platelet activation: a potential target for antiplatelet therapy. *Trends Pharmacol Sci* 29:352–60
- Grommes J, Alard JE, Drechsler M, Wantha S, Morgelin M, Kuebler WM, Jacobs M, von Hundelshausen P, Markart P, Wygrecka M, Preissner KT, Hackeng TM, Koenen RR, Weber C, Soehnlein O (2012) Disruption of platelet-derived chemokine heteromers prevents neutrophil extravasation in acute lung injury. *Am J Respir Crit Care Med* 185:628–636
- Gross S, Luckey C (1969) The oxygen tension-platelet relationship in cystic fibrosis. *Am Rev Respir Dis* 100:513–517
- Guzman-Grenfell A, Nieto-Velazquez N, Torres-Ramos Y, Montoya-Estrada A, Ramírez-Venegas A, Ochoa-Cautiño L, Flores-Trujillo F, Hicks JJ (2011) Increased platelet and erythrocyte arginase activity in chronic obstructive pulmonary disease associated with tobacco or wood smoke exposure. *J Investig Med* 59:587–92
- Hansell AL, Walk JA, Soriano JB (2003) What do chronic obstructive pulmonary disease patients die from? A multiple cause coding analysis. *Eur Respir J* 22:809–14
- Hasegawa S, Tashiro N, Matsubara T, Furukawa S, Ra C (2001) A comparison of Fc epsilon RI-mediated RANTES release from human platelets between allergic patients and healthy individuals. *Int Arch Allergy Immunol* 125(Suppl 1):42–47
- Haselmayer P, Grosse-Hovest L, von Landenberg P, Schild H, Radsak MP (2007) TREM-1 ligand expression on platelets enhances neutrophil activation. *Blood* 110:1029–1035
- Henn V, Slupsky JR, Grafe M et al (1998) CD40L on activated platelets triggers an inflammatory reaction on endothelial cells. *Nature* 391:591–594
- Idell S, James KK, Levin EG, Schwartz BS, Manchanda N, Maunder RJ, Martin TR, McLarty J, Fair DS (1989) Local abnormalities in coagulation and fibrinolytic pathways predispose to alveolar fibrin deposition in the adult respiratory distress syndrome. *J Clin Invest* 84:695–705
- Inoue T, Kannan MS (1989) Platelet-activating factor-induced functional changes in the guinea pig trachea in vitro. *Respir Physiol* 77:157–171



- Jawień J, Chłopicki S, Gryglewski RJ (2002) Interactions between human platelets and eosinophils are mediated by selectin-P. *Pol J Pharmacol* 54:157–160
- Jeffery PK, Wardlaw AJ, Nelson FC, Collins JV, Kay AB (1989) Bronchial biopsies in asthma. An ultrastructural, quantitative study and correlation with hyperreactivity. *Am Rev Respir Dis* 140:1745–1753
- Johansson MW, Han ST, Gunderson KA, Busse WW, Jarjour NN, Mosher DF (2012) Platelet activation, P-selectin, and eosinophil  $\beta 1$ -integrin activation in asthma. *Am J Respir Crit Care Med* 185:498–507
- Joseph M, Auriault C, Capron A, Vormg H, Viens P (1983) New function for platelets: IgE-dependent killing of schistosomes. *Nature* 303:810–812
- Joseph M, Gounni AS, Kusnierz JP, Vormg H, Sarfati M, Kinet JP, Tonnel AB, Capron A, Capron M (1997) Expression and functions of the high-affinity IgE receptor on human platelets and megakaryocyte precursors. *Eur J Immunol* 27:2212–2218
- Karp CL, Flick LM, Park KW, Softic S, Greer TM, Keledjian R, Yang R, Uddin J, Guggino WB, Atabani SF, Belkaid Y, Xu Y, Whitsett JA, Accurso FJ, Wills-Karp M, Petasis N (2004) Defective lipoxin-mediated anti-inflammatory activity in the cystic fibrosis airway. *Nat Immunol* 5:388–392
- Keir SD, Spina D, Page CP (2015) Bradykinin and capsaicin induced airways obstruction in the guinea pig are platelet dependent. *Pulm Pharmacol Ther* 33:25–31
- Kelly EA, Busse WW, Jarjour NN (2000) Increased matrix metalloproteinase-9 in the airway after allergen challenge. *Am J Respir Crit Care Med* 162:1157–1161
- Kiefmann R, Heckel K, Schenkat S, Dorger M, Wesierska-Gadek J, Goetz AE (2004) Platelet-endothelial cell interaction in pulmonary micro-circulation: the role of PARS. *Thromb Haemost* 91:761–770
- Kiefmann R, Heckel K, Schenkat S, Dorger M, Goetz AE (2006) Role of p-selectin in platelet sequestration in pulmonary capillaries during endotoxemia. *J Vasc Res* 43:473–481
- Klouche M, Klinger MH, Kühnel W, Wilhelm D (1997) Endocytosis, storage, and release of IgE by human platelets: differences in patients with type I allergy and nonatopic subjects. *J Allergy Clin Immunol* 100:235–241
- Knauer KA, Lichtenstein LM, Adkinson NF, Fish JE (1981) Platelet activation during antigen-induced airway reactions in asthmatic subjects. *N Engl J Med* 304:1404–1406
- Knauer KA, Kagey-Sobotka A, Adkinson NF Jr, Lichtenstein LM (1984) Platelet augmentation of IgE-dependent histamine release from human basophils and mast cells. *Int Arch Allergy Appl Immunol* 74:29–35
- Kornerup KN, Salmon GP, Pitchford SC, Liu WL, Page CP (2010) Circulating platelet-neutrophil complexes are important for subsequent neutrophil activation and migration. *J Appl Physiol* 109:758–767
- Kowal K, Pampuch A, Kowal-Bielecka O, DuBuske LM, Bodzenta-Lukaszyk A (2006) Platelet activation in allergic asthma patients during allergen challenge with *Dermatophagoides pteronyssinus*. *Clin Exp Allergy* 36:426–432
- Kowalska MA, Ratajczak MZ, Majka M, Jin J, Kunapuli S, Brass L, Poncz M (2000) Stromal cell-derived factor-1 and macrophage-derived chemokine: 2 chemokines that activate platelets. *Blood* 96:50–57
- Kraemer BF, Borst O, Gehring EM, Schoenberger T, Urban B, Ninci E, Seizer P, Schmidt C, Bigalke B, Koch M, Martinovic I, Daub K, Merz T, Schwanitz L, Stellos K, Fiesel F, Schaller M, Lang F, Gawaz M, Lindemann S (2010) PI3 kinase-dependent stimulation of platelet migration by stromal cell-derived factor 1 (SDF-1). *J Mol Med (Berl)* 88:1277–1288
- Kraemer BF, Campbell RA, Schwertz H, Cody MJ, Franks Z, Tolley ND, Kahr WH, Lindemann S, Seizer P, Yost CC, Zimmerman GA, Weyrich AS (2011) Novel anti-bacterial activities of  $\beta$ -defensin 1 in human platelets: suppression of pathogen growth and signaling of neutrophil extracellular trap formation. *PLoS Pathog* 7, e1002355
- Krijgsveld J, Zaat SAJ, Meeldijk J, van Veelen PA, Fang G, Poolman B, Brandt E, Ehlert JE, Kuijpers AJ, Engbers GH, Feijen J, Dankert J (2000) Thrombocidins, microbicidal proteins from human blood platelets, are C-terminal deletion products of CXC chemokines. *J Biol Chem* 275:20374–81
- Lader AS, Prat AG, Jackson GR, Chervinsky KL, Lapey A, Kinane TB, Cantiello HF (2000) Increased circulating levels of plasma ATP in cystic fibrosis. *Clin Pharmacol* 20:348–353
- Lechin F, van der Dijs B, Orozco B, Jara H, Rada I, Lechin ME, Lechin AE (1998) Neuropharmacologic treatment of bronchial asthma with the antidepressant tianeptine: a double-blind, crossover placebo-controlled study. *Clin Pharmacol Ther* 64:223–232
- Lellouch-Tubiana A, Lefort J, Simon MT, Pfister A, Vargaftig BB (1988) Eosinophil recruitment into guinea pig lungs after PAF-acether and allergen administration. Modulation by prostacyclin, platelet depletion, and selective antagonists. *Am Rev Respir Dis* 137:948–954
- Levy BD, Serhan CN (2014) Resolution of acute inflammation in the lung. *Annu Rev Physiol* 76:467–492
- Lima C, Souza VM, Soares AL, Macedo MS, Tavares-de-Lima W, Vargaftig BB (2007) Interference of methysergide, a specific 5-hydroxytryptamine receptor antagonist, with airway chronic allergic inflammation and remodelling in a murine model of asthma. *Clin Exp Allergy* 37:723–734
- Looney MR, Nguyen JX, Hu Y, Van Ziffle JA, Lowell CA, Matthay MA (2009) Platelet depletion and aspirin treatment protect mice in a two-event model of transfusion-related acute lung injury. *J Clin Invest* 119:3450–3461
- Ludwig RJ, Schultz JE, Boehncke WH, Podda M, Tandi C, Krombach F, Baatz H, Kaufmann R, von Andrian UH, Zollner TM (2004) [Activated, not resting, platelets increase leukocyte rolling in murine skin utilizing a distinct set of adhesion molecules.](#) *J Invest Dermatol* 122:830–836
- Lukacs NW, John A, Berlin A, Bullard DC, Knibbs R, Stoolman LM (2002) E- and P-selectins are essential for the development of cockroach allergen-induced airway responses. *J Immunol* 169:2120–2125
- Maccia CA, Gallagher JS, Ataman GMS, Glueck HI, Brooks SM, Bernstein IL (1977) Platelet thrombopathy in asthmatic patients with elevated immunoglobulin E. *J Allergy Clin Immunol* 59:101–108
- Maestrelli P, Boschetto P, Zocca E et al (1990) Venous blood platelets decrease during allergen-induced asthmatic reactions. *Clin Exp Allergy* 20:367–372
- Majoor CJ, Kamphuisen PW, Zwinderman AH et al (2013) Risk of deep vein thrombosis and pulmonary embolism in asthma. *Eur Respir J* 42:655–661
- Malerba M, Clini E, Malagola M, Avanzi GC (2013) Platelet activation as a novel mechanism of atherothrombotic risk in chronic obstructive pulmonary disease. *Expert Rev Hematol* 6(4):475–483
- Malmgren R, Grubbstrom J, Olsson P, Theorell H, Tornling G, Unge G (1982) Defective serotonin (5-HT) transport mechanism in platelets from patients with endogenous and allergic asthma. *Allergy* 37:29–39
- Marcos V, Zhou Z, Yildirim AO, Bohla A, Hector A, Vitkov L, Wiedenbauer EM, Krautgartner WD, Stoiber W, Belohradsky BH, Rieber N, Kormann M, Koller B, Roscher A, Roos D, Griese M, Eickelberg O, Döring G, Mall MA, Hartl D (2010) CXCR2 mediates NADPH oxidase-independent neutrophil extracellular

- trap formation in cystic fibrosis airway inflammation. *Nat Med* 16:1018–23
- Matsuda H, Ushio H, Geba GP, Askenase PW (1997) Human platelets can initiate T cell-dependent contact sensitivity through local serotonin release mediated by IgE antibodies. *J Immunol* 158:2891–2897
- Mattosio D, Evangelista V, De Cristofaro R, Recchiuti A, Pandolfi A, Di Silvestre S, Manarini S, Martelli N, Rocca B, Petrucci G, Angelini DF, Battistini L, Robuffo I, Pensabene T, Pieroni L, Furnari ML, Pardo F, Quattrucci S, Lancellotti S, Davì G, Romano M (2010) Cystic fibrosis transmembrane conductance regulator (CFTR) expression in human platelets: impact on mediators and mechanisms of the inflammatory response. *FASEB J* 24:3970–80
- Mayadas TN, Johnson RC, Rayburn H, Hynes RO, Wagner DD (1993) Leukocyte rolling and extravasation are severely compromised in P selectin-deficient mice. *Cell* 74:541–554
- Mazur I, Wurzer WJ, Ehrhardt C, Pleschka S, Puthavathana P, Silberzahn T, Wolff T, Planz O, Ludwig S (2007) Acetylsalicylic acid (ASA) blocks influenza virus propagation via its NF-kappaB-inhibiting activity. *Cell Microbiol* 9:1683–1694
- McKenzie EA (2007) Heparanase: a target for drug discovery in cancer and inflammation. *Br J Pharmacol* 151:1–14
- McVey M, Tabuchi A, Kuebler WM (2012) Microparticles and acute lung injury. *Am J Physiol Lung Cell Mol Physiol* 303:L364–L381
- Medzhitov R, Janeway C Jr (2000) Innate immunity. *N Engl J Med* 343:338–344
- Metzger WJ, Sjoerdsma K, Richerson HB et al (1987) Platelets in bronchoalveolar lavage from asthmatic patients and allergic rabbits with allergen-induced late phase responses. *Agents Actions Suppl* 21:151–159
- Momi S, Caracchini R, Falcinelli E, Evangelista S, Gresele P (2014) Stimulation of platelet nitric oxide production by nebulivolol prevents thrombosis. *Arterioscler Thromb Vasc Biol* 34:820–829
- Morrell CN, Aggrey AA, Chapman LM, Modjeski KL (2014) Emerging roles for platelets as immune and inflammatory cells. *Blood* 123:2759–67
- O'Sullivan BP, Linden MD, Frelinger LA, Barnard MR, Spencer-Manzon M, Morris JE, Salem RO, Laposata M, Michelson AD (2005) Platelet activation in cystic fibrosis. *Blood* 105:4635–4641
- Page C, Pitchford S (2013) Neutrophil and platelet complexes and their relevance to neutrophil recruitment and activation. *Int Immunopharmacol* 17:1176–1184
- Panes J, Perry MA, Anderson DC, Manning A, Leone B, Cepinskas G, Rosenbloom CL, Miyasaka M, Kviety PR, Granger DN (1995) Regional differences in constitutive and induced ICAM-1 expression in vivo. *Am J Physiol* 269:H1955–H1964
- Pepe PE, Potkin RT, Reus DH, Hudson LD, Carrico CJ (1982) Clinical predictors of the adult respiratory distress syndrome. *Am J Surg* 144:124–130
- Petito E, Momi S, Gresele P (2017) The migration of platelets and their interaction with other migrating cells. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 337–351
- Pitchford SC, Page CP (2002) Platelets and allergic disorders. In: Gresele P, Page C, Fuster V, Vermynen J (eds) *Platelets in thrombotic and non-thrombotic disorders*. Cambridge University Press, Cambridge, pp 852–868
- Pitchford SC, Page CP (2006) Platelet activation in asthma: integral to the inflammatory response. *Clin Exp Allergy* 36:399–401
- Pitchford SC, Yano H, Lever R, Riffo-Vasquez Y, Ciferri S, Rose MJ, Giannini S, Momi S, Spina D, O'Connor B, Gresele P, Page CP (2003) Platelets are essential for leukocyte recruitment in allergic inflammation. *J Allergy Clin Immunol* 112:109–118
- Pitchford SC, Riffo-Vasquez Y, Sousa A, Momi S, Gresele P, Spina D, Page CP (2004) Platelets are necessary for airway wall remodeling in a murine model of chronic allergic inflammation. *Blood* 103:639–647
- Pitchford SC, Momi S, Giannini S, Casali L, Spina D, Page CP, Gresele P (2005) Platelet P-selectin is required for pulmonary eosinophil and lymphocyte recruitment in a murine model of allergic inflammation. *Blood* 105:2074–2081
- Pitchford SC, Momi S, Baglioni S, Casali L, Giannini S, Rossi R, Page CP, Gresele P (2008) Allergen induces the migration of platelets to lung tissue in allergic asthma. *Am J Respir Crit Care Med* 177:604–612
- Ponick K, Sternitzky R, Mest HJ (1987) Stimulation of aggregation and thromboxane A2 formation of human platelets by hypoxia. *Prostaglandins Leukot Med* 29:49–59
- Poore S, Berry B, Eidson D, McKie KT, Harris RA (2013) Evidence of vascular endothelial dysfunction in young patients with cystic fibrosis. *Chest* 143:939–45
- Pryzdial ELG, Lin BH, Sutherland MR (2017) Virus–platelet associations. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 1089–1106
- Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, Fukuchi Y, Jenkins C, Rodríguez-Roisin R, van Weel C, Zielinski J, [Global Initiative for Chronic Obstructive Lung Disease](#) (2007) Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 176:532–555
- Ragaller M, Richter T (2010) Acute lung injury and acute respiratory distress syndrome. *J Emerg Trauma Shock* 3:43–51
- Rendu R, Bohard-Bohn B (2002) Platelet Organelles. In: Gresele P, Page C, Fuster V, Vermynen J (eds) *Platelets in thrombotic and non-thrombotic disorders*. Cambridge University Press, Cambridge, pp 104–112
- Reutershan J, Ley K (2004) Bench-to bedside review: acute respiratory distress syndrome—how neutrophils migrate into the lung. *Crit Care* 8:453–61
- Ring J, Allen DH, Mathison DA, Spiegelberg HL (1980) In vitro releasability of histamine and serotonin: studies of atopic patients. *J Clin Lab Immunol* 3:85–91
- Riordan JR, Rommens JM, Kerem B, Alon N, Rozmahel R, Grzelczak Z, Zielinski J, Lok S, Plavsic N, Chou JL et al (1989) Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* 245:1066–73
- Robertson DN, Page CP (1987) Effect of platelet agonists on airway reactivity and intrathoracic platelet accumulation. *Br J Pharmacol* 92:105–111
- Rondina MT, Garraud O, Schwertz H (2017) Platelets and bacterial infections. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 1073–1087
- Romano M, Chen XS, Takahashi Y, Yamamoto S, Funk CD, Serhan CN (1993) Lipoxin synthase activity of human platelet 12-lipoxygenase. *Biochem J* 296:127–133
- Saffarzadeh M, Juenemann C, Queisser MA, Lochnit G, Barreto G, Galuska SP, Lohmeyer J, Preissner KT (2012) Neutrophil extracellular traps directly induce epithelial and endothelial cell death: a predominant role of histones. *PLoS One* 7, e32366
- Saxena SP, McNicol A, Brandes LJ, Becker AB, Gerrard JM (1989) Histamine formed in stimulated human platelets is cytoplasmic. *Biochem Biophys Res Commun* 164:164–168
- Schmidtke DW, Diamond SL (2000) Direct observation of membrane tethers formed during neutrophil attachment to platelets or P-selectin under physiological flow. *J Cell Biol* 149:719–730

- Scott T, Owens MD (2008) Thrombocytes respond to lipopolysaccharide through Toll-like receptor-4, and MAP kinase and NF- $\kappa$ B pathways leading to expression of interleukin-6 and cyclooxygenase-2 with production of prostaglandin E2. *Mol Immunol* 45:1001–1008
- Semeraro F, Ammollo CT, Morrissey JH, Dale GL, Friese P, Esmon NL et al (2011) Extracellular histones promote thrombin generation through platelet-dependent mechanisms: involvement of platelet TLR2 and TLR4. *Blood* 118:1952–1961
- Semple JW, Italiano JE Jr, Freedman J (2011) Platelets and the immune continuum. *Nat Rev Immunol* 11:264–74
- Serhan CN, Chiang N, Van Dyke TE (2008) Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol* 8:349–361
- Shang Y, Yao S (2014) Pro-resolution of inflammation: a potential strategy for treatment of acute lung injury/acute respiratory distress syndrome. *Chin Med J* 127:801–802
- Shoki AH, Mayer-Hamblett N, Wilcox PG, Sin DD, Quon BS (2013) Systematic review of blood biomarkers in cystic fibrosis pulmonary exacerbations. *Chest* 144:1659–70
- Singh B, Fu C, Bhattacharya J (2000) Vascular expression of the alpha (v)beta(3)-integrin in lung and other organs. *Am J Physiol Lung Cell Mol Physiol* 278:L217–L226
- Slaba I, Kubes P (2017) Platelets and immunity. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 489–512
- Sladek K, Dworski R, Fitzgerald GA, Buitkus KL, Block FJ, Marney SR Jr, Sheller JR (1990) Allergen-stimulated release of thromboxane A2 and leukotriene E4 in humans. Effect of indomethacin. *Am Rev Respir Dis* 141:1441–1445
- Sprague DL, Elzey BD, Crist SA, Waldschmidt TJ, Jensen RJ, Ratliff TL (2008) Platelet-mediated modulation of adaptive immunity: unique delivery of CD154 signal by platelet-derived membrane vesicles. *Blood* 111:5028–36
- Stead RJ, Barradas MA, Mikhailidis DP, Jeremy JY, Hodson ME, Batten JC, Dandona P (1987) Platelet hyperaggregability in cystic fibrosis. *Prostaglandins Leukot Med* 26:91–103
- Stellos K, Kopf S, Paul A, Marquardt JU, Gawaz M, Huard J, Langer HF (2010) Platelets in regeneration. *Semin Thromb Hemost* 36:175–184
- Sturm A, Hebestreit H, Koenig C, Walter U, Grossmann R (2010) Platelet proinflammatory activity in clinically stable patients with CF starts in early childhood. *J Cyst Fibros* 9:179–86
- Sullivan PJ, Jafar ZH, Harbinson PL, Restrict LJ, Costello JF, Page CP (2000) Platelet dynamics following allergen challenge in allergic asthmatics. *Respiration* 67:514–517
- Svensson Holm AC, Bengtsson T, Grenegård M, Lindström EG (2008) Platelets stimulate airway smooth muscle cell proliferation through mechanisms involving 5-lipoxygenase and reactive oxygen species. *Platelets* 19:528–536
- Svensson Holm AC, Bengtsson T, Grenegård M, Lindström EG (2011) Platelet membranes induce airway smooth muscle cell proliferation. *Platelets* 22:45–55
- Svensson Holm AC, Grenegård M, Ollinger K, Lindström EG (2014) Inhibition of 12-lipoxygenase reduces platelet activation and prevents their mitogenic function. *Platelets* 25:111–117
- Symon FA, McNulty CA, Wardlaw AJ (1999) P- and L-selectin mediate binding of T cells to chronically inflamed human airway endothelium. *Eur J Immunol* 29:1324–1333
- Szczeklik A, Milner PC, Birch J, Watkins J, Martin JF (1986) Prolonged bleeding time, reduced platelet aggregation, altered PAF-acether sensitivity and increased platelet mass are a trait of asthma and hay fever. *Thromb Haemost* 56:283–287
- Tamagawa-Mineoka R, Katoh N, Ueda E, Takenaka H, Kita M, Kishimoto S (2007) The role of platelets in leukocyte recruitment in chronic contact hypersensitivity induced by repeated elicitation. *Am J Pathol* 170:2019–2029
- Tang YQ, Yaeman MR, Selsted ME (2002) Antimicrobial peptides from human platelets. *Infect Immun* 70:6524–33
- Tian J, Zhu T, Liu J, Guo Z, Cao X (2015) Platelets promote allergic asthma through the expression of CD154. *Cell Mol Immunol* 12:700–707
- Tillie-Leblond I, Marquette CH, Perez T, Scherpereel A, Zanetti C, Tonnel AB, Remy-Jardin M (2006) Pulmonary embolism in patients with unexplained exacerbation of chronic obstructive pulmonary disease: prevalence and risk factors. *Ann Intern Med* 144:390–6
- Totani L, Amore C, Di Santo A, Dell’Elba G, Piccoli A, Martelli N, Tenor H, Beume R, Evangelista V (2016) Roflumilast inhibits leukocyte-platelet interactions and prevents the prothrombotic functions of polymorphonuclear leukocytes and monocytes. *J Thromb Haemost* 14:191–204
- Truelsen T, Prescott E, Lange P, Schnohr P, Boysen G (2001) Lung function and risk of fatal and non-fatal stroke. The Copenhagen City Heart Study. *Int J Epidemiol* 30:145–51
- Ulfman LH, Joosten DP, van Aalst CW, Lammers JW, van de Graaf EA, Koenderman L, Zwaginga JJ (2003) Platelets promote eosinophil adhesion of patients with asthma to endothelium under flow conditions. *Am J Respir Cell Mol Biol* 28:512–519
- Varas-Lorenzo C, Rodriguez LAG, Maguire A et al (2007) Use of oral corticosteroids and the risk of acute myocardial infarction. *Atherosclerosis* 192:376–383
- von Hundelshausen P, Weber C (2007) Platelets as immune cells: bridging inflammation and cardiovascular disease. *Circ Res* 100:27–40
- Ward PA, Grailer JJ (2014) Acute lung injury and the role of histones. *Transl Respir Med* 2:1
- Ware J, Post SR (2017) Platelets and inflammatory disorders of connective tissue. In: Gresele P et al (eds) *Platelets in thrombotic and non-thrombotic disorders*. Springer, Cham, pp 1137–1141
- Wei Y, Wang Z, Su L, Chen F, Tejera P, Bajwa EK, Wurfel MM, Lin X, Christiani DC (2015) Platelet count mediates the contribution of a genetic variant in LRRRC16A to ARDS risk. *Chest* 147:607–617
- Weyrich AS, Schwertz H, Kraiss LW, Zimmerman GA (2009) Protein synthesis by platelets: historical and new perspectives. *J Thromb Haemost* 7:241–246
- Wong CHY, Jenne CN, Petri B, Chrobok N, Kubes P (2013) Nucleation of platelets with blood-borne pathogens on Kupffer cells precedes other innate immunity and contributes to bacterial clearance. *Nat Immunol* 14:785–792
- Wright TK, Gibson PG, Simpson JL, McDonald VM, Wood LG, Baines KJ (2016) Neutrophil extracellular traps are associated with inflammation in chronic airway disease. *Respirology* 21:467–75
- Wu H, Yang J, Su EM, Li L, Zhao C, Yang X, Gao Z, Pan M, Sun P, Sun W, Jiang Y, Su X (2014) Lipoxin A4 and platelet activating factor are involved in E. coli or LPS-induced lung inflammation in CFTR-deficient mice. *PLoS One* 9:e93003
- Xiao Z, Goldsmith HL, McIntosh FA, Shankaran H, Neelamegham S (2006) Biomechanics of P-selectin PSGL-1 bonds: shear threshold and integrin-independent cell adhesion. *Biophys J* 90:2221–2234
- Xie RF, Hu P, Wang ZC, Yang J, Yang YM, Gao L, Fan HH, Zhu YM (2015) Platelet-derived microparticles induce polymorphonuclear leukocyte-mediated damage of human pulmonary microvascular endothelial cells. *Transfusion* 55:1051–7
- Yasuba H, Chihara J, Kino T, Satake N, Oshima S (1991) Increased releasability of platelet products and reduced heparin-induced platelet factor 4 release from endothelial cells in bronchial asthma. *J Lipid Mediat* 4:5–21
- Yiming MT, Lederer DJ, Sun L, Huertas A, Issekutz AC, Bhattacharya S (2008) Platelets enhance endothelial adhesiveness in high tidal volume ventilation. *Am J Respir Cell Mol Biol* 39:569–575

- Yoshida A, Ohba M, Wu X, Sasano T, Nakamura M, Endo Y (2002) Accumulation of platelets in the lung and liver and their degranulation following antigen-challenge in sensitized mice. *Br J Pharmacol* 137:146–152
- Yoshimi Y, Fujimura M, Myou S, Tachibana H, Hirose T (2001) Effect of thromboxane A<sub>2</sub> (TXA<sub>2</sub>) synthase inhibitor and TXA<sub>2</sub> receptor antagonist alone and in combination on antigen-induced bronchoconstriction in guinea pigs. *Prostaglandins Other Lipid Mediat* 65:1–9
- Zarbock A, Ley K (2009) The role of platelet in acute lung injury. *Front Biosci* 14:150–158
- Zarbock A, Singbartl K, Ley K (2006) Complete reversal of acid-induced acute lung injury by blocking of platelet-neutrophil aggregation. *J Clin Invest* 116:3211–3219
- Zhang G, Han J, Welch EJ, Ye RD, Voyno-Yasenetskaya TA, Malik AB, Du X, Li Z (2009) Lipopolysaccharide stimulates platelet secretion and potentiates platelet aggregation via TLR4/MyD88 and the cGMP-dependent protein kinase pathway. *J Immunol* 182:7997–8004
- Zhao C, Su EM, Yang X, Gao Z, Li L, Wu H, Jiang Y, Su X (2013) Important role of platelets in modulating endotoxin-induced lung inflammation in CFTR-deficient mice. *PLoS One* 8, e82683