

# On the Page Number of Upward Planar Directed Acyclic Graphs\*

Fabrizio Frati<sup>1,2,3</sup>, Radoslav Fulek<sup>1</sup>, and Andres J. Ruiz-Vargas<sup>1</sup>

<sup>1</sup> School of Basic Sciences - École Polytechnique Fédérale de Lausanne, Switzerland  
{fabrizio.frati, radoslav.fulek, andres.ruizvargas}@epfl.ch

<sup>2</sup> Dipartimento di Informatica e Automazione, Università Roma Tre

<sup>3</sup> School of Information Technologies, University of Sydney

**Abstract.** In this paper we study the page number of upward planar directed acyclic graphs. We prove that: (1) the page number of any  $n$ -vertex upward planar triangulation  $G$  whose every maximal 4-connected component has page number  $k$  is at most  $\min\{O(k \log n), O(2^k)\}$ ; (2) every upward planar triangulation  $G$  with  $o(\frac{n}{\log n})$  diameter has  $o(n)$  page number; and (3) every upward planar triangulation has a vertex ordering with  $o(n)$  page number if and only if every upward planar triangulation whose maximum degree is  $O(\sqrt{n})$  does.

## 1 Introduction

A  $k$ -page book embedding of a graph  $G=(V, E)$  is a total ordering  $\sigma$  of  $V$  and a partition of  $E$  into subsets  $E_1, E_2, \dots, E_k$ , called *pages*, such that no two edges  $(u, v)$  and  $(w, z)$  with  $u <_{\sigma} w <_{\sigma} v <_{\sigma} z$  belong to the same set  $E_i$ . The *page number* of  $G$  is the minimum  $k$  such that  $G$  admits a  $k$ -page book embedding.

Book embeddings (first introduced by Kainen [15] and by Ollmann [19]) find applications in several contexts, such as VLSI design, fault-tolerant processing, sorting networks, and parallel matrix multiplication (see, e.g., [4,11,20,21]). Henceforth, they have been widely studied from a theoretical point of view; namely, the literature is rich of combinatorial and algorithmic contributions on the page number of various classes of graphs (see, e.g., [2,7,8,9,10,17,18]). We remark here a famous result of Yannakakis [22] stating that any planar graph has page number at most four.

Heath *et al.* [13,14] extended the notions of book embedding and page number to directed acyclic graphs (DAGs for short) in a very natural way: Given a DAG  $G=(V, E)$ , book embedding and page number of  $G$  are defined as for undirected graphs, except that the total ordering of  $V$  is now required to be a *linear extension* of the partial order of  $V$  induced by  $E$ . That is, if  $G$  contains an edge from a vertex  $u$  to a vertex  $v$ , then  $u <_{\sigma} v$  in any feasible total ordering  $\sigma$  of  $V$ . The authors of [13,14] showed that DAGs with page number equal to one can be characterized and recognized efficiently; however, they proved that, in general, determining the page number of a DAG is NP-complete.

---

\* Work partially supported by the Italian Ministry of Research, grant RBIP06BZW8, FIRB project “Advanced tracking system in intermodal freight transportation”, by the Swiss National Science Foundation 200021-125287/1, by the ESF project 10-EuroGIGA-OP-003 “Graph Drawings and Representations”, and by the MIUR of Italy, project AlgoDEEP 2008TFBWL4.

The main problem raised by Heath *et al.* and studied in, e.g., [1,6,12,13,14], is whether every *upward planar DAG* admits a book embedding in few pages. An upward planar DAG is a DAG that admits a drawing which is simultaneously *upward*, *i.e.*, each edge is represented by a curve monotonically increasing in the  $y$ -direction, and *planar*, *i.e.*, no two edges cross. Upward planar DAGs are the natural counterpart of planar graphs in the context of directed graphs. Notice that there exist DAGs which admit a planar non-upward embedding and that require  $\Omega(|V|)$  pages in any book embedding [12,14]. No upper bound better than the trivial  $O(|V|)$  and no lower bound better than the trivial  $\Omega(1)$  are known for the page number of upward planar DAGs. It is however known that *directed trees* have page number one [14], that *unicyclic DAGs* have page number two [14], and that *series-parallel DAGs* have page number two [1,6].

In this paper we study the page number of upward planar DAGs. Before stating our results we need some background.

First, it is known that every upward planar DAG  $G$  can be augmented to an *upward planar triangulation*  $G'$  [5]. That is, edges can be added to  $G$  so that the resulting graph  $G'$  is still an upward planar DAG and every face of  $G'$  is delimited by a 3-cycle. Thus, in order to establish tight bounds on the page number of upward planar DAGs, it suffices to look at upward planar triangulations, as the page number of a subgraph  $G$  of a graph  $G'$  is at most the page number of  $G'$ . In the following, unless otherwise specified, all the considered graphs are upward planar triangulations.

Second, consider a total ordering  $\sigma$  of  $V$ . A *twist* is a set of pairwise crossing edges, *i.e.*, a set  $\{(u_1, v_1), (u_2, v_2), \dots, (u_k, v_k)\}$  of edges such that  $u_1 <_\sigma u_2 <_\sigma \dots <_\sigma u_k <_\sigma v_1 <_\sigma v_2 <_\sigma \dots <_\sigma v_k$ . It is straightforward that the page number of a graph  $G$  is lower bounded by the minimum over all vertex orderings  $\sigma$  of the maximum size of a twist in  $\sigma$ . Moreover, a function of the maximum size of a twist in a vertex ordering upper bounds the page number of an  $n$ -vertex graph  $G$ , as stated in the following two lemmata.

**Lemma 1.** [3] *Let  $\sigma$  be a vertex ordering of an  $n$ -vertex graph  $G$ . Suppose that the maximum twist of  $\sigma$  has size  $k$ . Then  $G$  admits a book embedding with vertex ordering  $\sigma$  and with  $O(k \log n)$  pages.*

**Lemma 2.** [16] *Let  $\sigma$  be a vertex ordering of an  $n$ -vertex graph  $G$ . Suppose that the maximum twist of  $\sigma$  has size  $k$ . Then  $G$  admits a book embedding with vertex ordering  $\sigma$  and with  $O(2^k)$  pages.*

Thus, in order to get upper bounds for the page number of a graph, it often suffices to construct vertex orderings with small maximum twist size.

In this paper we consider the relationship between the page number of an  $n$ -vertex upward planar triangulation  $G$  and three important graph parameters of  $G$ : The connectivity, the diameter, and the degree. We show the following results. (i) In Sect. 3, we prove that an upward planar triangulation  $G$  admits a vertex ordering with maximum twist size  $O(f(n))$  if and only if every maximal 4-connected component of  $G$  does. As a corollary, upward planar 3-trees have constant page number. (ii) In Sect. 4, we prove that every upward planar triangulation  $G$  has a vertex ordering whose maximum twist size is a function of the *diameter* of  $G$ , that is, of the length of the longest directed path in  $G$ . As a corollary, every upward planar triangulation whose diameter is  $o(n/\log n)$

admits a book embedding in  $o(n)$  pages. (iii) In Sect. 5, we show that every upward planar triangulation has a vertex ordering with  $o(n)$  page number if and only if every upward planar triangulation whose maximum degree is  $O(\sqrt{n})$  does.

## 2 Definitions

A *directed graph* is a graph with direction on the edges. The *underlying graph* of a directed graph  $G$  is the undirected graph obtained from  $G$  by removing the directions on its edges. We denote by  $(u, v)$  an edge directed from a vertex  $u$ , which is called the *origin* of  $(u, v)$ , to a vertex  $v$ , which is called the *destination* of  $(u, v)$ ; edge  $(u, v)$  is *incoming*  $v$  and *outgoing*  $u$ . A *source* (resp. *sink*) is a vertex with no incoming edge (resp. with no outgoing edge). A *directed cycle* is a directed graph whose underlying graph is a cycle and containing no source and no sink. A *directed acyclic graph* (DAG for short) is a directed graph containing no directed cycle. A *directed path* is a directed graph whose underlying graph is a path and containing exactly one source and one sink. The *diameter* of a directed graph is the number of vertices in its longest directed path.

A *drawing* of a directed graph is a mapping of each vertex to a point in the plane and of each edge to a Jordan curve between its end-points. A drawing is *upward* if each edge  $(u, v)$  is a curve monotonically increasing in the  $y$ -direction and it is *planar* if no two edges intersect except, possibly, at common end-points. A drawing is *upward planar* if it is both upward and planar. An *upward planar graph* is a graph that admits an upward planar drawing. A planar drawing of a graph partitions the plane into connected regions, called *faces*. The unbounded face is the *outer face*, all the other faces are *internal faces*. Two upward planar drawings of an upward planar DAG are *equivalent* if they determine the same clockwise ordering of the edges around each vertex. An *embedding* of an upward planar DAG is an equivalence class of upward planar drawings. An *embedded upward planar graph* is an upward planar DAG together with an embedding.

An *upward planar triangulation* is an upward planar graph whose underlying graph is a maximal planar graph. Consider any two upward planar drawings  $\Gamma_1$  and  $\Gamma_2$  of an upward planar triangulation  $G$ . Then, either  $\Gamma_1$  and  $\Gamma_2$  are equivalent, or the clockwise ordering of the edges around each vertex in  $\Gamma_1$  is exactly the opposite of the one in  $\Gamma_2$ . The outer face of an upward planar drawing  $\Gamma$  of an upward planar triangulation  $G$  is delimited by a cycle composed of three edges  $(u, v)$ ,  $(u, z)$ , and  $(v, z)$ . Then,  $u$ ,  $v$ , and  $z$  are called *bottom vertex*, *middle vertex*, and *top vertex* of  $\Gamma$ , respectively. Consider the two embeddings  $\mathcal{E}_1$  and  $\mathcal{E}_2$  of an upward planar triangulation  $G$ . Then, the bottom, middle, and top vertex of  $\mathcal{E}_1$  coincide with the bottom, middle, and top vertex of  $\mathcal{E}_2$ , respectively. Hence such vertices are simply called the *bottom vertex of  $G$* , the *middle vertex of  $G$* , and the *top vertex of  $G$* , respectively.

A total vertex ordering  $\sigma$  of a DAG  $G$  is *upward* if  $G$  has no edge  $(u, v)$  such that  $v <_{\sigma} u$ . The upward vertex orderings are all and only the vertex orderings that are feasible for a book embedding of a DAG. We say that an upward vertex ordering  $\sigma$  *induces* a twist of size  $k$  if  $G$  contains edges  $(u_1, v_1), \dots, (u_k, v_k)$  such that  $u_1 <_{\sigma} \dots <_{\sigma} u_k <_{\sigma} v_1 <_{\sigma} \dots, v_k$ . The *maximum twist size* of an upward vertex ordering  $\sigma$  is the maximum number of edges in a twist induced by  $\sigma$ . Two edges  $(u_1, v_1)$  and  $(u_2, v_2)$  are *nested* in  $\sigma$  if  $u_1 <_{\sigma} u_2 <_{\sigma} v_2 <_{\sigma} v_1$ . Two edges  $(u_1, v_1)$  and  $(u_2, v_2)$  *cross* in  $\sigma$  if  $u_1 <_{\sigma} u_2 <_{\sigma} v_1 <_{\sigma} v_2$ .

An undirected graph is *k-connected* if the removal of any  $k - 1$  vertices leaves the graph connected. A directed graph is *k-connected* if its underlying graph is. A *maximal k-connected component* of a graph  $G$  is a subgraph  $G'$  of  $G$  such that  $G'$  is *k-connected* and no subgraph  $G''$  of  $G$  with  $G' \subset G''$  is *k-connected*. A *separating triangle*  $C$  in a graph  $G$  is a 3-cycle such that the removal of the vertices of  $C$  from  $G$  disconnects  $G$ . A separating triangle  $C$  in a graph  $G$  is *maximal* if  $G$  has no separating triangle  $C'$  such that  $C$  is internal to  $C'$ .

The *degree of a vertex* is the number of edges incident to it. The *degree of a graph* is the maximum among the degrees of its vertices. A DAG is *Hamiltonian* if it contains a directed path passing through all its vertices. An Hamiltonian DAG  $G$  has exactly one upward total vertex ordering. Moreover, if  $G$  is upward planar, then it has page number at most 2. A *plane 3-tree* is a maximal plane graph that can be constructed as follows. Let  $G_3$  be a 3-cycle embedded in the plane. A plane 3-tree with  $n$  vertices is a plane graph that can be constructed from a plane graph  $G_{n-1}$  with  $n - 1$  vertices by inserting a vertex inside an internal face of  $G_{n-1}$  and by connecting such a vertex to the three vertices incident to the face. A *planar 3-tree* is a planar graph that can be embedded as a plane 3-tree. An *upward plane 3-tree* is an upward planar DAG whose underlying graph is a plane 3-tree.


### 3 Page Number and Connectivity

In this section we study the relationship between the page number of an upward planar DAG and the page number of its maximal 4-connected components. We prove the following:

**Theorem 1.** *Let  $f(n)$  be any function such that  $f(n) \in \Omega(1)$  and  $f(n) \in O(n)$ . Consider any  $n$ -vertex upward planar triangulation  $G$  and suppose that every maximal 4-connected component of  $G$  has an upward vertex ordering with maximum twist size at most  $f(n)$ . Then  $G$  has an upward vertex ordering with maximum twist size  $O(f(n))$ .*

First, we define a rooted tree  $T = (V', E')$ , whose nodes correspond to subgraphs of  $G=(V, E)$ , which reflects the structure of separating triangles in  $G$ . Tree  $T$  is recursively defined as follows (see Fig. 1(a)). The root  $r$  of  $T$  corresponds to  $G'(r) = G$ . Suppose that a node  $a$  of  $T$  corresponds to a subgraph  $G'(a)$  of  $G$ . If  $G'(a)$  contains no separating triangle, then  $a$  is a leaf of  $T$ . Otherwise, consider every maximal separating triangle  $(u, v, z)$  of  $G'(a)$ ; then, insert a node  $b$  in  $T$  as a child of  $a$ , such that  $G'(b)$  is the subgraph of  $G'(a)$  induced by the vertices internal to or on the border of cycle  $(u, v, z)$ . For each node  $a \in T$ , denote as  $V'(a)$  and  $E'(a)$  the vertex set and the edge set of  $G'(a)$ . Further, for each node  $a \in T$ , let  $G(a) = (V(a), E(a))$  denote the subgraph of  $G'(a)$  induced by all the vertices which are not internal to any separating triangle of  $G'(a)$ . Note that  $G(a)$  is 4-connected for every  $a \in V'$ .

We now define a total ordering  $o(V)$  of  $V$  and we later prove that the maximum twist size of  $o(V)$  is  $O(f(n))$ . Ordering  $o(V)$  is constructed by induction on  $T$ . In the base case  $a$  is a leaf; then let  $o(V'(a))$  be any total ordering of  $V'(a)$  such that the maximum twist size of  $o(V'(a))$  is  $f(n)$ . Such an ordering exists by hypothesis, since  $G'(a)$  is 4-connected. In the inductive case, let  $a_1, \dots, a_m$  be the children of  $a$  in  $T$ ,



**Fig. 1.** (a) Tree  $T$  capturing the structure of the separating triangles in  $G$ . (b) Graph  $G'(a)$ ; the thick edges belong to  $M_0$ .

where total orderings  $o(V'(a_1)), \dots, o(V'(a_m))$  of  $V'(a_1), \dots, V'(a_m)$ , respectively, have already been computed. Compute a total ordering  $o(V(a))$  of  $V(a)$  such that the maximum twist size of  $o(V(a))$  is  $f(n)$ . Again, such an ordering exists by hypothesis, since  $G(a)$  is 4-connected. Next, we merge  $o(V'(a_1)), \dots, o(V'(a_m))$  with  $o(V(a))$ . In order to do this, we define the operation of *merging an ordering  $V_2$  into an ordering  $V_1$* , that takes as input two total vertex orderings  $o(V_1)$  and  $o(V_2)$  such that  $V_1$  and  $V_2$  share a single vertex  $v$ , and outputs a single total vertex ordering  $o(V_1 \cup V_2)$  of  $V_1 \cup V_2$  such that  $o(V_1 \cup V_2)$  coincides with  $o(V_i)$  when restricted to the vertices in  $V_i$ , for  $i = 1, 2$ , and such that every vertex of  $V_1$  that precedes  $v$  in  $o(V_1)$  (resp. follows  $v$  in  $o(V_1)$ ) precedes all the vertices of  $V_2$  in  $o(V)$  (resp. follows all the vertices of  $V_2$  in  $o(V)$ ). Denote by  $b(H)$ , by  $m(H)$ , and by  $t(H)$  the bottom vertex, the middle vertex, and the top vertex of an upward triangulation  $H$ , respectively. Then, ordering  $o(V'(a))$  is defined as follows: Let  $o_1 = o(V(a))$  and let  $o_{i+1}$  be the ordering obtained by merging  $o(V'(a_i)) \setminus \{b(G'(a_i)), t(G'(a_i))\}$  into  $o_i$ , for  $i = 1, \dots, m$ ; then  $o(V'(a)) = o_{m+1}$ . Observe that  $o(V'(a))$  is an upward vertex ordering because  $o(V(a)), o(V'(a_1)), \dots, o(V'(a_m))$  are and because of the definition of the merging operation.

We now prove that the size of the maximum twist induced by  $o(V)$  is  $O(f(n))$ . Let  $M = \{e_1=(u_1, v_1), \dots, e_k=(u_k, v_k)\}$  denote any maximal twist induced by  $o(V)$ . We have the following:

**Claim 1.** *Let  $a$  be a node of  $T$ . Let  $a_1$  and  $a_2$  be two distinct children of  $a$ . There is no pair of distinct edges  $(u_i, v_i), (u_j, v_j)$  in  $M$  such that  $(u_i, v_i) \in E'(a_1), (u_j, v_j) \in E'(a_2)$ , and  $\{u_i, v_i, u_j, v_j\} \cap V(a) = \emptyset$ .*

**Proof:** Let  $(u^1, v^1, z^1)$  and  $(u^2, v^2, z^2)$  be the separating triangles of  $G'(a)$  that delimit the outer faces of  $G'(a_1)$  and  $G'(a_2)$ , where  $v^i$  is the middle vertex of  $G'(a_i)$ , for  $i = 1, 2$ . If  $v^1 \neq v^2$ , then, by the construction of  $o(V)$ , all internal vertices of  $G'(a_1)$  precede all internal vertices of  $G'(a_2)$  or vice versa, thus  $e_i$  and  $e_j$  do not both belong to  $M$ . Otherwise,  $v^1 = v^2$ . Then, again by the construction of  $o(V)$ ,  $e_i$  and  $e_j$  are nested, thus they do not both belong to  $M$ .  $\square$

Let  $r$  be the root of  $T$ . We assume that  $G$  is “minimal”, that is, we assume that there exists no child  $a$  of  $r$  such that all the edges in  $M$  belong to  $G'(a)$ . Indeed, if such a child exists, graph  $G=G'(r)$  can be replaced by  $G'(a)$ , and the bound on the size of  $M$  can be achieved by arguing on  $G'(a)$  rather than on  $G'(r)$ . Denote by  $M_i$ , with  $i = 0, 1, 2$ , the subset of  $M$  that contains all the edges having  $i$  endpoints in  $V(r)$ . Observe that  $|M| = |M_0| + |M_1| + |M_2|$ , hence it suffices to prove that  $|M_i| \in O(f(n))$ , for

$i = 0, 1, 2$ , in order to prove the theorem. By hypothesis and since  $G(r)$  is 4-connected, we have  $|M_2| \leq f(n)$ . We now deal with the edges in  $M_1$ .

**Claim 2.**  $|M_1| \in O(f(n))$ .

**Proof:** First, we argue that  $M_1$  contains at most one edge  $e$  such that an end-vertex of  $e$  is the middle vertex of an upward planar triangulation  $G'(a)$ , for some child  $a$  of  $r$ . Indeed, by the vertex ordering's construction, any two such edges, say  $e_a$  and  $e_b$ , are either incident to the same vertex or are such that both end-vertices of  $e_a$  come before both end-vertices of  $e_b$  in  $o(V'(a))$ . Thus, it is enough to bound the number of edges in  $M_1$  whose end-vertex in  $V(r)$  is the bottom vertex or the top vertex of an upward planar triangulation  $G'(a)$ , where  $a$  is a child of  $r$ .

Let  $M_1^b$  (resp.  $M_1^t$ ) be the subset of the edges in  $M_1$  whose end-vertex in  $V(r)$  is the bottom vertex (resp. the top vertex) of an upward planar triangulation  $G'(a)$ , where  $a$  is a child of  $r$ . Observe, that by the above observation,  $|M| \leq |M_1^b| + |M_1^t| + 1$ . In the following we bound  $|M_1^b|$  (the bound for  $|M_1^t|$  can be obtained analogously).

Consider any edge  $(u, v) \in M_1^b$ , where  $u \in V(r)$ . We define a *corresponding edge* of  $(u, v)$  in  $G(r)$  as follows. Let  $a_{u,v}$  be the child of  $r$  such that  $G'(a_{u,v})$  contains edge  $(u, v)$ . Further, denote by  $m_{u,v}$  the middle vertex of  $G'(a_{u,v})$ . Then,  $(u, m_{u,v})$  is the corresponding edge of  $(u, v)$  in  $G(r)$ . Observe that edge  $(u, m_{u,v})$  exists and belongs to  $E(r)$ . Now consider the multi-set  $E_1^b$  of the corresponding edges, that is  $E_1^b = \{(u, m_{u,v}) | (u, v) \in M_1^b\}$ . First, we have that, for each vertex  $w$  in  $V(r)$ , there exist at most two edges  $(z, w)$  in  $E_1^b$ , since each vertex in  $V(r)$  is the middle vertex of at most two upward planar triangulations  $G'(a_i)$ , where  $a_i$  is a child of  $r$ , and since  $G'(a_i)$  has at most one edge in  $M_1^b$ . If there exist two edges  $(z_1, w)$  and  $(z_2, w)$  in  $E_1^b$ , then remove one of them. Then, after such deletions,  $|E_1^b| \geq |M_1^b|/2$ .

Next, we prove that each vertex in  $V(r)$  is an end-vertex of at most two edges in  $E_1^b$ . Namely, consider any two edges  $(u_1, v_1)$  and  $(u_2, v_2)$  in  $E_1^b$ . Then,  $v_1 \neq v_2$  because of the deletions performed on  $E_1^b$ , and  $u_1 \neq u_2$  as otherwise the corresponding edges in  $M_1^b$  would share a vertex, contradicting the assumption that  $M$  is a twist; thus, each vertex in  $V(r)$  is the source of at most one edge in  $E_1^b$  and the sink of at most one edge in  $E_1^b$ . Since the degree of graph  $(V(r), E_1^b)$  is two, there exists a subset  $E^*$  of  $E_1^b$  such that the degree of graph  $(V(r), E^*)$  is one and  $|E^*| \geq |E_1^b|/3$ .

Finally, we have that every two edges in  $E^*$  cross. Namely, if they do not, then by the vertex ordering's construction the corresponding edges in  $M_1^b$  would not cross either, thus contradicting the assumption that  $M$  is a twist.

Since  $E^* \subseteq E(r)$  and the maximum size of a twist of edges in  $E(r)$  is  $f(n)$ , given that  $G(r)$  is 4-connected, it follows that  $|E^*| \leq f(n)$ . Using  $|E^*| \geq |E_1^b|/3$  and  $|E_1^b| \geq |M_1^b|/2$ , we get  $|M_1^b| \leq 6f(n)$ . Such an inequality, together with the analogous bound  $|M_1^t| \leq 6f(n)$  and with  $|M| \leq |M_1^b| + |M_1^t| + 1$ , proves the theorem.  $\square$

We now proceed by bounding the size of  $M_0$ .

**Claim 3.**  $|M_0| \in O(f(n))$ .

**Proof:** By Claim 1, all the edges in  $M_0$  belong to a graph  $G'(a)$ , for a certain descendant  $a$  of  $r$ . Let us choose  $a$  so that the length of the path from  $a$  to  $r$  is maximized. Let  $w$  be the middle vertex of the separating triangle  $(u, v, w)$  delimiting  $G'(a)$ . Let  $a'$  denote

the child of  $r$  which is an ancestor of  $a$  or that coincides with  $a$ . Let  $w'$  be the middle vertex of the separating triangle  $(u', v', w')$  delimiting  $G'(a')$ .

For any edge  $(y, z) \in M_0$ , we have that  $(y, z)$  “nests around  $w'$ ”, that is,  $y$  precedes  $w'$  and  $w'$  precedes  $z$  in  $o(V)$ . Indeed, if both  $y$  and  $z$  precede  $w'$  in  $o(V)$  (or if they both follow  $w'$  in  $o(V)$ ), then only the edges in  $G'(a')$  can possibly cross  $(y, z)$ , by the construction of  $o(V)$ , thus contradicting the minimality of  $r$ .

If  $w \neq w'$ , then  $|M_0| \leq 3$ , since only the edges incident to  $u, v$  and  $w$  can belong to  $M_0$ . Otherwise we have  $w' = w$  (see Fig. 1(b)). Consider graph  $G'(a)$ ; partition the edges in  $M_0$  into two subsets, namely  $M'_0$  contains all the edges of  $M_0$  having at least one end-vertex in  $V(a)$  and  $M''_0$  contains all the edges of  $M_0$  having no end-vertex in  $V(a)$ . By definition of  $a$  and by Claim 1,  $|M'_0| > 0$ , as otherwise there would exist a child of  $a$  containing all the edges of  $M_0$ . However, by Claim 2 applied to  $G'(a)$  and by the hypothesis of the theorem, we have  $|M'_0| \in O(f(n))$ . Moreover, every edge in  $M''_0$  is in a separating triangle of  $G'(a)$  having  $w$  as middle vertex; however, any such edge is nested inside any edge of  $M'_0$ ; thus, since  $|M'_0| > 0$ , we have  $|M''_0| = 0$  and hence  $|M_0| \in O(f(n))$ , which concludes the proof.  $\square$

Since  $|M_i| \in O(f(n))$ , for  $i = 0, 1, 2$ , it follows that  $|M| \in O(f(n))$ , thus proving Theorem 1. By Lemmata 1 and 2, we have the following:

**Corollary 1.** *If every  $n$ -vertex upward planar 4-connected triangulation has  $o(\frac{n}{\log n})$  page number, then every  $n$ -vertex upward planar triangulation has  $o(n)$  page number.*

**Corollary 2.** *Every upward planar 3-tree has  $O(1)$  page number.*


## 4 Page Number and Diameter

In this section we study the relationship between the page number of an upward planar DAG and its diameter  $D$ . We show that upward planar DAGs with small diameter have sub-linear page number. Notice that such a result pairs the observation that graphs with diameter  $n - o(n)$  have sub-linear page number as well, given that upward planar Hamiltonian DAGs have page number two. We have the following:

**Theorem 2.** *Every  $n$ -vertex upward planar triangulation whose diameter is at most  $D$  admits an upward vertex ordering whose maximum twist size  $t(n)$  is a function satisfying  $t(n) \leq aD + t(\frac{n}{2}) + b$ , for some constants  $a$  and  $b$ .*

We will prove the statement for a family of upward planar DAGs that is strictly larger than the family of upward planar triangulations. Namely, we call *upward cactus* an embedded upward planar DAG  $G$  having exactly one source  $s(G)$  and such that every internal face is delimited by a 3-cycle. See Fig. 2. Observe that an upward planar triangulation is an upward cactus.

Consider an upward cactus  $G$ . We call *monotone path* any directed path  $P = (u_1, \dots, u_k)$  from  $s(G)$  to a sink of  $G$ . Consider an upward planar drawing  $\Gamma$  of  $G$  in which  $u_k$  is the vertex with highest  $y$ -coordinate. Observe that such a drawing  $\Gamma$  always exists because  $G$  is an upward cactus. Then, we define the *left side of  $P$*  as the



**Fig. 2.** An upward cactus  $G$ . The thick edges represent a monotone path  $P$

subgraph of  $G$  induced by all the vertices which are to the left of the Jordan curve representing  $P$  in  $\Gamma$ . The *right side of  $P$*  is defined analogously. Observe that the vertices of  $P$ , the vertices of the left side of  $P$ , and the vertices of the right side of  $P$  form a partition of the vertices of  $G$ . We have the following:

**Claim 4.** *In every  $n$ -vertex upward cactus there exists a monotone path  $P$  such that both the left side of  $P$  and the right side of  $P$  have less than  $\frac{n}{2}$  vertices.*

We now prove the statement of the theorem for every  $n$ -vertex upward cactus  $G$  with diameter at most  $D$ . The proof is by induction on  $n$ . If  $n \leq 3$ , then in any upward vertex ordering of  $G$  the maximum twist size is 1, hence  $t(3) \leq b$ , for any  $b \geq 1$ , thus proving the base case.

Suppose that  $n > 3$ . By Claim 4, there exists a monotone path  $P$  in  $G$  such that both the left side of  $P$  and the right side of  $P$  have less than  $\frac{n}{2}$  vertices. We now associate each vertex in the left side of  $P$  and each vertex in the right side of  $P$  to a vertex of  $P$ . Namely, we associate a vertex  $v$  in the left side of  $P$  to the vertex  $u_i$  of  $P$  such that there exists a directed path from  $u_i$  to  $v$  and such that, for every  $j > i$ , there exists no directed path from  $u_j$  to  $v$ . Observe that, for every vertex  $v$  in the left side of  $P$ , there exists a directed path from  $s(G)$  to  $v$ , since  $G$  has a unique source, hence  $v$  is associated to exactly one vertex of  $P$ . Then, we call *left bag of  $u_i$*  the set of vertices in the left side of  $P$  which are associated to  $u_i$ , for each  $i = 1, \dots, k$ . Vertices in the right side of  $P$  are associated to vertices of  $P$  analogously, thus analogously defining the *right bag of  $u_i$* , for each  $i = 1, \dots, k$ . We have the following:

**Claim 5.** *The subgraph  $G_i^L$  of  $G$  induced by the left bag of  $u_i$  and by  $u_i$  is an upward cactus, for every  $i = 1, \dots, k$ .*

An analogous claim holds for the subgraph  $G_i^R$  of  $G$  induced by the right bag of  $u_i$  and by  $u_i$ .

Next, we construct an upward vertex ordering of  $G$ . This is done as follows. First, inductively construct an upward vertex ordering  $\sigma_i^L$  of  $G_i^L$  and an upward vertex ordering  $\sigma_i^R$  of  $G_i^R$ , for  $i = 1, \dots, k$ , such that the maximum twist size of each of  $\sigma_i^R$  and  $\sigma_i^L$  is  $t(\frac{n}{2})$ . This is possible since  $G_i^L$  and  $G_i^R$  are upward cacti, by Claim 5, and they have less than  $\frac{n}{2}$  vertices, by Claim 4. Observe that  $u_i$  is the first vertex both in  $\sigma_i^L$  and in  $\sigma_i^R$ , given that it is the only source of both  $G_i^L$  and  $G_i^R$ . Then, denote by  $\sigma_i$  the vertex ordering of  $G_i^L \cup G_i^R$  which is obtained by concatenating  $\sigma_i^L$  and  $\sigma_i^R \setminus \{u_i\}$ . Finally a vertex ordering  $\sigma$  of  $G$  is obtained by concatenating  $\sigma_1, \sigma_2, \dots, \sigma_k$ .



**Claim 6.**  $\sigma$  is an upward vertex ordering.

Next, we prove that the maximum twist size  $t(n)$  of  $\sigma$  is at most  $aD + t(\frac{n}{2}) + b$ , for some constants  $a$  and  $b$ .

First, observe that the edges that have both end-vertices in  $P$  create twists of size at most two, since the graph induced by the vertices of  $P$  is upward planar Hamiltonian.

Second, we discuss the size of a twist composed of *intra-bag* edges, which are edges whose both end-vertices are associated to the same vertex of  $P$ . Consider any edge  $e_i^L$  of  $G_i^L$  and any edge  $e_i^R$  of  $G_i^R$ . Such edges do not cross. Namely, if such edges are both incident to  $u_i$ , then they do not cross by definition. If  $e_i^R$  is not incident to  $u_i$ , then both end-vertices of  $e_i^R$  come after both end-vertices of  $e_i^L$ , by construction, hence such edges do not cross. Moreover, if  $e_i^R$  is incident to  $u_i$  and  $e_i^L$  is not, then  $e_i^L$  is nested inside  $e_i^R$ , by construction, hence such edges do not cross. It follows that the maximum size of a twist of intra-bag edges is equal to the maximum twist size of  $\sigma$  restricted to the vertices in  $G_i^a$  for some  $a \in \{L, R\}$  and some  $1 \leq i \leq k$ . By Claim 5, graph  $G_i^a$  is an upward cactus. Moreover, by Claim 4,  $G_i^a$  has at most  $\frac{n}{2}$  vertices, hence the maximum size of a twist of intra-bag edges is at most  $t(\frac{n}{2})$ .

Third, we discuss the maximum size of a twist composed of *inter-bag* edges, which are edges whose end-vertices are associated to distinct vertices of  $P$ . We show that the maximum size of a twist composed of inter-bag edges in the left side of  $P$  is  $2D$ . An analogous proof shows that the maximum size of a twist composed of inter-bag edges in the right side of  $P$  is also  $2D$ .

Consider any two inter-bag edges  $(w_1, w_2)$  and  $(w_3, w_4)$  in the left side of  $P$ . Suppose that  $(w_1, w_2)$  and  $(w_3, w_4)$  cross in  $\sigma$ . Denote by  $u_{j_1}, u_{j_2}, u_{j_3}$ , and  $u_{j_4}$ , such that  $u_{j_1} < u_{j_2}$  and  $u_{j_3} < u_{j_4}$ , the vertices of  $P$  vertices  $w_1, w_2, w_3$ , and  $w_4$  have been assigned to, respectively. The following claim asserts that any two inter-bag edges  $(w_1, w_2)$  and  $(w_3, w_4)$  that cross in  $\sigma$  either have their sources assigned to the same vertex of  $P$ , or have their destinations assigned to the same vertex of  $P$ , or the source of one of them and the destination of the other of them are assigned to the same vertex of  $P$ .

**Claim 7.** *At least one of the following holds:  $j_1 = j_3 < j_2, j_4$ , or  $j_1 < j_2 = j_3 < j_4$ , or  $j_3 < j_4 = j_1 < j_2$ , or  $j_1, j_3 < j_2 = j_4$ .*

Hence, if there are more than  $2D$  inter-bag edges pairwise crossing in the left side of  $P$ , then either there are more than  $D$  inter-bag edges pairwise crossing in the left side of  $P$  such that the origins of such edges have all been assigned to the same vertex of  $P$ , or there are more than  $D$  inter-bag edges pairwise crossing in the left side of  $P$  such that the destinations of such edges have all been assigned to the same vertex of  $P$ . In the following, we discuss such two cases.

**Claim 8.** *Suppose that  $G$  contains inter-bag edges  $(v_1, w_1), (v_2, w_2), \dots, (v_k, w_k)$  in the left side of  $P$ , where  $v_1 <_\sigma v_2 <_\sigma \dots <_\sigma v_k <_\sigma w_1 <_\sigma w_2 <_\sigma \dots <_\sigma w_k$  and where all the vertices  $w_i$  have been assigned to the same vertex  $u_1$  of  $P$ , for  $i = 1, \dots, k$ , or all the vertices  $v_i$  have been assigned to the same vertex  $u_1$  of  $P$ , for  $i = 1, \dots, k$ . Then, there exists a directed path starting at  $u_1$  and passing through  $w_1, w_2, \dots, w_k$ .*

Since by hypothesis any directed path contains at most  $D$  vertices, then, by Claim 8, the maximum size of a twist of inter-bag edges sharing their destinations in the left side of  $P$  is at most  $D$  and the maximum size of a twist of inter-bag edges sharing their origins in the left side of  $P$  is at most  $D$ . Hence, by Claim 7, the maximum size of a twist of inter-bag edges in the left side of  $P$  is at most  $2D$  and the maximum size of a twist of inter-bag edges is at most  $4D$ . Since every edge of  $G$  is either an edge having both end-vertices in  $P$ , or is an intra-bag edge, or is an inter-bag edge, it follows that the maximum size of a twist in  $\sigma$  is  $t(n) = 2 + t(\frac{n}{2}) + 4D$ , thus proving Theorem 2.

By Lemma 1, we have the following:

**Corollary 3.** *Every  $n$ -vertex upward planar triangulation whose diameter is  $o(\frac{n}{\log n})$  has  $o(n)$  page number.*

## 5 Page Number and Degree


In this section we discuss the relationship between the page number of a graph and its degree. We prove the following theorem.

**Theorem 3.** *Let  $f(n)$  be any function such that  $f(n) \in \Omega(\sqrt{n})$  and  $f(n) \in O(n)$ . Suppose that every  $n$ -vertex upward planar triangulation whose degree is  $O(f(n))$  admits a book embedding with  $O(g(n))$  pages, for some function  $g(n) \in \Omega(1)$  and  $g(n) \in O(n)$ . Then, every  $n$ -vertex upward planar triangulation admits a book embedding with  $O(g(n) + \frac{n}{f(n)})$  pages.*

Consider any  $n$ -vertex upward planar triangulation  $G$ . We transform  $G$  into an  $O(n)$ -vertex upward planar triangulation  $G'$  with degree  $O(f(n))$  as follows. Fix any constant  $c > 0$  and denote by  $u_1, \dots, u_k$  any ordering of the vertices of  $G$  whose degree is greater than  $cf(n)$ .

For  $i = 1, \dots, k$ , consider vertex  $u_i$ . Suppose that  $u_i$  is an internal vertex of  $G$ , the case in which  $u_i$  is an external vertex being analogous. Since it is an upward planar triangulation,  $G$  has exactly two faces  $(v_1, v_2, u_i)$  and  $(v_3, v_4, u_i)$  incident to  $u_i$  such that edges  $(v_1, u_i)$  and  $(v_4, u_i)$  are incoming  $u_i$  and such that edges  $(u_i, v_2)$  and  $(u_i, v_3)$  are outgoing  $u_i$ . Assume, w.l.o.g., that  $(v_1, u_i)$ ,  $(u_i, v_2)$ ,  $(u_i, v_3)$ , and  $(v_4, u_i)$  appear in this clockwise order around  $u_i$ . Denote by  $w_1 = v_2, w_2, \dots, w_{x-1}, w_x = v_3, w'_1 = v_4, w'_2, \dots, w'_{y-1}, w'_y = v_1$  the clockwise order of the neighbors of  $u_i$  (see Fig. 3(a)). Remove  $u_i$  and its incident edges from  $G$ . Let  $M = \lceil \frac{x}{f(n)-1} \rceil$  and  $N = \lceil \frac{y}{f(n)-1} \rceil$ . Insert  $M + N + 2$  vertices  $z_1, \dots, z_{M+N+2}$  in  $G$  inside the cycle of the neighbors of  $u_i$ . Insert an edge from  $z_j$  to  $z_{j+1}$ , for  $j = 1, \dots, M$ , insert an edge from  $z_{j+1}$  to  $z_j$ , for  $j = M + 1, \dots, M + N + 1$ , and insert edges from  $z_{M+2}$  to  $z_1, \dots, z_M$  and from  $z_{M+3}, \dots, z_{M+N+2}$  to  $z_1$ . Insert edges from  $v_1$  to  $z_1$ , from  $z_1$  to  $v_2$ , from  $v_4$  to  $z_{M+2}$ , and from  $z_{M+2}$  to  $v_3$ . Insert edges from  $z_j$  to  $w_{(j-2)(f(n)-1)+1}, w_{(j-2)(f(n)-1)+2}, \dots, w_{(j-1)(f(n)-1)}$ , for  $j = 2, \dots, M + 1$ ; insert edges from  $w'_{(j-2)(f(n)-1)+1}, w'_{(j-2)(f(n)-1)+2}, \dots, w'_{(j-1)(f(n)-1)}$  to  $z_{M+j}$ , for  $j = 3, \dots, N + 2$ . See Fig. 3(b).

It is easy to see that the triangulation  $G'$  obtained from  $G$  after all vertices  $u_1, \dots, u_k$  have been considered is upward planar. We have the following.



**Fig. 3.** (a) Neighbors of a high-degree vertex  $u_i$ . (b) Replacing  $u_i$  with lower-degree vertices, assuming  $f(n) = 3$ .

**Claim 9.**  $G'$  has  $O(n)$  vertices and  $O(f(n))$  degree. Moreover, for every upward vertex ordering  $\sigma'$  of  $G'$ , there exists an upward vertex ordering  $\sigma$  of  $G$  such that  $\sigma$  and  $\sigma'$  restricted to the vertices that are both in  $G$  and in  $G'$  coincide.

We now describe how to compute a book embedding of  $G$  in  $O(g(n) + \frac{n}{f(n)})$  pages. First, construct the upward planar triangulation  $G'$  as above. Second, construct a book embedding of  $G'$  into  $O(g(n))$  pages. Such a book embedding exists by hypothesis, since  $G'$  has  $O(n)$  vertices and  $O(f(n))$  degree (by Claim 9). Denote by  $\sigma'$  the total ordering of the vertices of  $G'$  in the constructed book embedding. Construct any total ordering  $\sigma$  of the vertices of  $G$  such that  $\sigma$  and  $\sigma'$  restricted to the vertices that are both in  $G$  and in  $G'$  coincide. Such an ordering exists (and can be easily constructed) by Claim 9. The edges of  $G$  can be assigned to pages as follows:  $O(g(n))$  pages suffice to accommodate all the edges that are both in  $G$  and in  $G'$ ; moreover, one page can be used to accommodate all the edges incident to vertex  $u_i$ , for  $i = 1, \dots, k \in O(\frac{n}{f(n)})$ . It follows that  $G$  has a book embedding in  $O(g(n) + \frac{n}{f(n)})$  pages, thus proving Theorem 3.

**Corollary 4.** Every  $n$ -vertex upward planar triangulation has  $o(n)$  page number if and only if every  $n$ -vertex upward planar triangulation with degree  $O(\sqrt{n})$  has  $o(n)$  page number.

## 6 Conclusions

In this paper we studied the relationship between the page number of an upward planar triangulation  $G$  and three important parameters of  $G$ : The connectivity, the diameter, and the degree. It would be interesting, in our opinion, to understand whether the statements of Theorems 1 and 2 can be referred to the page number rather than to the maximum twist size. That is: (1) Is it true that any upward planar triangulation  $G$  has page number  $O(k)$  if and only if every maximal 4-connected subgraph of  $G$  has page number  $O(k)$ ? (2) Is it true that any  $n$ -vertex upward planar triangulation  $G$  with diameter  $D$  has page number  $p(n)$  satisfying  $p(n) = p(\frac{n}{2}) + aD + b$ , for some constants  $a$  and  $b$ ?

Determining whether every  $n$ -vertex upward planar DAG has  $o(n)$  page number and whether there exist upward planar DAGs with  $\omega(1)$  page number remain among the most important problems in the theory of linear graph layouts.

**Acknowledgments.** The first author would like to thank Patrizio Angelini, Giuseppe Di Battista, and Stefano Sarauili for very useful discussions.

## References

1. Alzohairi, M., Rival, I.: Series-Parallel Planar Ordered Sets Have Pagenumber Two. In: North, S.C. (ed.) GD 1996. LNCS, vol. 1190, pp. 11–24. Springer, Heidelberg (1997)
2. Buss, J.F., Shor, P.W.: On the pagenumber of planar graphs. In: Symposium on Theory of Computing (STOC 1984), pp. 98–100. ACM (1984)
3. Cerný, J.: Coloring circle graphs. Elec. Notes Discr. Math. 29, 457–461 (2007)
4. Chung, F.R.K., Leighton, F.T., Rosenberg, A.L.: Embedding graphs in books: A layout problem with applications to VLSI design. SIAM J. Alg. Discr. Meth. 8, 33–58 (1987)
5. Di Battista, G., Tamassia, R.: Algorithms for plane representations of acyclic digraphs. Theor. Comp. Sci. 61, 175–198 (1988)
6. Di Giacomo, E., Didimo, W., Liotta, G., Wismath, S.K.: Book embeddability of series-parallel digraphs. Algorithmica 45(4), 531–547 (2006)
7. Enomoto, H., Nakamigawa, T., Ota, K.: On the pagenumber of complete bipartite graphs. J. Comb. Th. Ser. B 71(1), 111–120 (1997)
8. Ganley, J.L., Heath, L.S.: The pagenumber of  $k$ -trees is  $O(k)$ . Discr. Appl. Math. 109(3), 215–221 (2001)
9. Heath, L.S.: Embedding planar graphs in seven pages. In: Foundations of Computer Science (FOCS 1984), pp. 74–83. IEEE (1984)
10. Heath, L.S., Istrail, S.: The pagenumber of genus  $g$  graphs is  $O(g)$ . J. ACM 39(3), 479–501 (1992)
11. Heath, L.S., Leighton, F.T., Rosenberg, A.L.: Comparing queues and stacks as mechanisms for laying out graphs. SIAM J. Discr. Math. 5(3), 398–412 (1992)
12. Heath, L.S., Pemmaraju, S.V.: Stack and queue layouts of posets. SIAM J. Discr. Math. 10(4), 599–625 (1997)
13. Heath, L.S., Pemmaraju, S.V.: Stack and queue layouts of directed acyclic graphs: Part II. SIAM J. Computing 28(5), 1588–1626 (1999)
14. Heath, L.S., Pemmaraju, S.V., Trenk, A.N.: Stack and queue layouts of directed acyclic graphs: Part I. SIAM J. Computing 28(4), 1510–1539 (1999)
15. Kainen, P.C.: Thickness and coarseness of graphs. Abh. Math. Sem. Univ. Hamburg 39, 88–95 (1973)
16. Kostochka, A.V., Kratochvíl, J.: Covering and coloring polygon-circle graphs. Discr. Math. 163(1-3), 299–305 (1997)
17. Malitz, S.M.: Genus  $g$  graphs have pagenumber  $O(\sqrt{g})$ . J. Algorithms 17(1), 85–109 (1994)
18. Malitz, S.M.: Graphs with  $e$  edges have pagenumber  $O(\sqrt{e})$ . J. Algorithms 17(1), 71–84 (1994)
19. Ollmann, L.T.: On the book thicknesses of various graphs. In: Hoffman, F., Levow, R.B., Thomas, R.S.D. (eds.) Southeastern Conference on Combinatorics, Graph Theory and Computing. Congressus Numerantium, vol. VIII, p. 459 (1973)
20. Rosenberg: The Diogenes approach to testable fault-tolerant arrays of processors. IEEE Trans. Comp. C-32, 902–910 (1983)
21. Tarjan, R.E.: Sorting using networks of queues and stacks. J. ACM 19(2), 341–346 (1972)
22. Yannakakis, M.: Embedding planar graphs in four pages. J. Comp. Syst. Sci. 38(1), 36–67 (1989)