

*Corrigendum*

## Phenotypic recurrent selection for 2n pollen and 2n egg production in diploid alfalfa

S. Tavoletti<sup>1</sup>, A. Mariani<sup>1</sup> & F. Veronesi<sup>2</sup>

<sup>1</sup> *Instituto di Recerche sul Miglioramento Genetico delle Piante Foraggere, C.N.R., Perugia, Italy;*

<sup>2</sup> *Instituto di Produzione Vegetale, Università degli Studi di Udine, Udine, Italy*

*Euphytica* 57: 97–102, 1991.

© 1991 Kluwer Academic Publishers. Printed in the Netherlands.

Owing to an error in the proof-reading process Table 4 of this article has been published incorrectly. The Table with the correct numbers is presented below.

*Table 4.* Selection for 2n egg production: effect of phenotypic recurrent selection on the average seed set per population and on the range of the seed per plant

	2x–4x crosses		
	Unselected population (B <sub>0</sub> )	Population of the 1st cycle (B <sub>1</sub> )	Population of the 2nd cycle (B <sub>2</sub> )
N. of plants utilized	16	139	101
N. of flowers pollinated	563	5347	2469
N. of seeds produced	26	613	652
Average seed set (%) <sup>1</sup>	4.26a	11.46b	26.4c
Seed set per plant (%) (Range)	(0–96)	(0–75)	(0–160)

<sup>1</sup> Average seed set = (N. of seeds produced/N. of flowers pollinated) × 100; means not followed by the same letter differ significantly at P ≤ 0.01.