

References

- Ahmed, Y., Gilmartin, G.M., Hanly, S.M., Nevins, J.R., Greene, W.C. (1991) The HTLV-I rex response element mediates a novel form of mRNA polyadenylation. *Cell* **64**, 727–737
- Barker, D.G., Gallusci, P., Lullien, V., Khan, H., Ghérardi, M., Huget, T. (1988) Identification of two groups of leghemoglobin genes in alfalfa (*Medicago sativa*) and a study of their expression during root nodule development. *Plant Mol. Biol.* **11**, 761–772
- Dean, C., Tamaki, S., Dunsmuir, P., Favreau, M., Katamaya, C., Dooner, H., Bedbrook, J. (1986) mRNA transcripts of several plant genes are polyadenylated at multiple sites in vivo. *Nucleic Acids Res.* **14**, 2229–2240
- Hunt, A.G., MacDonald, M.H. (1989) Deletion analysis of the polyadenylation signal of the pea ribulose-1,5-bisphosphate carboxylase small-subunit gene. *Plant Mol. Biol.* **13**, 125–138
- Hunt, A.G., Chu, N.M., Odell, J.T., Nagy, F., Chua, N.H. (1987) Plant cells do not properly recognize animal gene polyadenylation signals. *Plant Mol. Biol.* **8**, 23–35
- Ingelbrecht, I.L.W., Herman, L.M.F., Dekeyser, R.A., Van Montagu, M.C., Depicker, A.G. (1989) Different 3'end regions strongly influence the level of expression in plant cells. *Plant Cell* **1**, 671–680
- Joshi, C.P. (1987) Putative polyadenylation signals in nuclear genes of higher plants: a compilation and analysis. *Nucleic Acids Res.* **15**, 9627–9640
- Kuhse, J., Pühler, A. (1987) Conserved sequence motifs in the untranslated 3'end of leghemoglobin transcripts isolated from broadbean nodules. *Plant Sci.* **49**, 137–143
- LeMeur, M.A., Galliot, B., Gerlinger, P. (1984) Termination of the ovalbumin gene transcription. *EMBO J.* **3**, 2279–2286
- Malim, M.H., Hauber, J., Le, S.-Y., Maizel, J.V., Cullen, B.R. (1989) The HIV-1 rev trans-activator acts through a structured target sequence to activate nuclear export of unspliced viral mRNA. *Nature* **338**, 254–257
- Manen, J.-F., Simon, P., Van Slooten, J.-C., Osteras, M., Frutiger, S., Hughes, G.J. (1991) A nodulin expressed in senescent nodules of winged bean is a protease inhibitor. *Plant Cell* **3**, 259–270
- Mogen, B.D., MacDonald, M.H., Graybosch, R., Hunt, A.G. (1990) Upstream sequence other than AAUAAA are required for efficient messenger RNA 3'end formation in plants. *Plant Cell* **2**, 1261–1272
- Sánchez, F., Padilla, J.E., Pérez, H., Lara, M. (1991) Control of nodulin genes in root-nodule development and metabolism. *Annu. Rev. Plant Physiol.* **42**, 507–528
- Sanfaçon, H., Brodman, P., Hohn, T. (1991) A dissection of the cauliflower mosaic virus polyadenylation signal. *Genes Develop.* **5**, 141–149
- Sasavage, N.L., Smith, M., Gillam, S., Woychick, R.P., Rottman, F.M. (1982) Variation in the polyadenylation site of bovine prolactin mRNA. *Proc. Natl. Acad. Sci. USA* **79**, 223–227
- Wiborg, O., Hylding-Nielsen, J.J., Jensen, E.O., Paludan, K., Marcker, K.A. (1982) The nucleotide sequence of two leghemoglobin genes from soybean. *Nucleic Acids Res.* **10**, 3487–3495
- Wickens, M. (1990) How the messenger got its tail: addition of poly(A) in the nucleus. *TIBS* **15**, 277–281
- Zuker, M., Stiegler, P. (1981) Optimal computer folding of large RNA sequences using thermodynamic and auxiliary information. *Nucleic Acids Res.* **9**, 133–148

Erratum

Expression of two cloned mRNA sequences during development and germination of Spores of the sensitive fern, *Onoclea sensibilis* L.

V. Raghavan¹* and Joseph C. Kamalay²

¹ Department of Plant Biology and ² Department of Agronomy, The Ohio State University, Columbus, OH 43210, USA

Published in *Planta* (1993) 189:1–9

On page 1, the address listed for J.C. Kamalay should read:

² USDA, Forest Service, Northeastern Forest Experiment Station, 359 Main Road, Delaware, OH 43015, USA