single group whose outcome will be compared to those of the standard treatment group. The same principles of dietary management as outlined in the intervention strategy for the standard treatment group will be followed. Reinforcement of the dietary programme will be carried out by the dietician monthly for the first 6 months and every 3 months thereafter. Self blood glucose monitoring will be performed a minimum of four times a day, to include three pre-prandial and one bedtime sample. A 03.00 h sample will be obtained once a week and repeated the next night if the value is <65 mg/dl (3.6 mmol/l). Initially, patients will be seen every week at the clinic until a stable treatment programme is achieved and at least monthly thereafter. Telephone contact will be made daily for the first week and weekly thereafter. The aim in the experimental treatment group is to achieve and maintain as near normal glycaemic control as possible in the absence of significant hypoglycaemia. The goal for fasting and pre-prandial levels of blood glucose is 70–120 mg/dl (3.9–6.7 mmol/l); for 03.00 h blood glucose, 65 mg/dl (3.6 mmol/l) or above; and for 90–120 min post-prandial levels <180 mg/dl (10 mmol/l). The goal will be to maintain the glycohaemoglobin level within 2 SD of the mean for a non-diabetic sample of persons. Procedures for calculation of these values will be analogous to those used for the standard treatment group (data are not yet available from the questionnaire survey). Hypoglycaemia will be carefully monitored and limited.

During the study, participants will undergo a set of regularly scheduled procedures for patient follow-up and analysis of study end-points. A standardized follow-up history and physical examination will be scheduled yearly for each patient. Glycohaemoglobin, blood glucose profiles, and a standardized ophthalmological examination, including stereo fundus photography, will be determined quarterly; standardized psychological and compliance/adherence assessments will be made annually. Additional visits may be scheduled as needed for clinical care.

Detailed procedures are specified for the management of diabetic hypoglycaemia, ketoacidosis, and pregnancy. The protocol also describes procedures for managing deviations from assigned treatment, for transfers to inactive status, and for protocol changes. Major statistical methods to be used in the analysis of the data are presented.

Mechanisms for assuring quality control and for monitoring the performance of all Diabetes Control and Complications Trial components on a regular basis have been established within the trial structure, including programmes for central laboratory quality control, local laboratory standardization and surveillance of the Clinical Centres and the Coordinating Centre. Two groups are specified which are advisory to the the National Institute of Arthritis, Diabetes, and Digestive and Kidney Diseases and are otherwise external to and independent of the Diabetes Control and Complications Trial: the Policy Advisory Group and the Data, Safety and Quality Review Group.

The outcome of this limited feasibility study will determine whether and how the full-scale trial will be initiated.

A limited number of copies of the complete draft of the Phase II protocol are available upon receipt of a written request enclosing a pre-addressed mailing label. Requests will be honoured on a first-come, first-served basis and should be addressed to: Clinical Trial Coordinator, Division of Diabetes, Endocrinology, and Metabolic Diseases, The National Institute of Arthritis, Diabetes, and Digestive and Kidney Diseases, National Institutes of Health, Westwood Building, Room 607, Bethesda, Maryland, 20205, USA.

Yours sincerely
L. B. Salans

Dr. L. B. Salans
Director, National Institute of Arthritis, Diabetes, and Digestive and Kidney Diseases
National Institutes of Health
Bethesda, Maryland 20205, USA

Therapeutic Trials in Type 1 (Insulin-Dependent) Diabetic Patients on Insulin: Goals and Tools

Dear Sir,

It has been announced that the National Institutes of Health, Bethesda, USA, is beginning a diabetes control trial aimed at assessing the relationship between blood glucose and microvascular complications in Type 1 (insulin-dependent) diabetes. Twenty-one medical centres in the USA and Canada have been selected [1].

Recent articles in both the lay [1] and the medical [2] literature have emphasized the idea that the open-loop delivery system will resolve the controversy over diabetic control and the development of complications.

The utility of any clinical trial can be discussed in two ways: is it going to yield more fundamental knowledge (explanatory approach) or will it help to reach a decision on the choice of the best available treatment (pragmatic approach) [3]? Thus any trial designed to demonstrate the causal relationship between blood glucose levels and microvascular lesions must be an explanatory one [3]. However, such a trial is not easy to design since two goals more or less overlap: (1) what is the effect of elevated blood glucose levels on the microvasculature and (2) is there any definite pathological blood glucose level or some progressive linear relationship between blood glucose and microvascular changes? Moreover, if the design of the protocol also includes an analysis of the tool used to conduct it, pumps for instance, a third goal is added.

Whatever the goal of this kind of explanatory trial (assuming it is unique and well-defined), the setting up of such an investigation implies certain conditions. In particular, it requires the use of a placebo for a double-blind study. Obviously such a requirement cannot be applied to insulin trials. It is also necessary to test a simple parameter and to study a homogenous selected highly motivated population. Even if it yields highly experimental scientific knowledge of wide application, there will be ethical and practical difficulties because of the design; unless the conventionally treated group is maintained frankly hyperglycaemic for many years, failure to find a difference in the rate of development of microangiopathy would be meaningless.

The use of pumps as tools in an explanatory trial on the effect of blood glucose in diabetic patients is not mandatory since a difference in glycaemic control between groups can be achieved using well established method of conventional insulin therapy [4]. If normoglycaemia is required it has been shown that intensified conventional therapy can achieve this goal [5]. If pumps are used it should be stressed (as was done in the Steno Group report which demonstrated a beneficial effect from producing good glycaemic control) that 'the study (is) not designed as a contest between infusion devices and conventional therapy but a comparison between near normal and moderate levels of glycaemic control' [6]. Such confusions between the goals of a study and the tools used are not always avoided [7]. Moreover, insulin treatment with pumps aimed at achieving perfect control on a very large population of patients may be dangerous: 11 deaths have been reported in patients wearing pumps in the USA in 1981 [8]. These are perhaps due, at least in part, to casual and indiscriminate use in patients not sufficiently trained for 'euglycaemic control treatment' [9]. Finally, a wide utilization of pumps may be difficult since a large number of medical staff is needed and patient compliance in the long term is totally unknown and probably poor.

Whatever the tools to be used for conducting the trial, it seems to us that the cost-benefit ratio of such an explanatory trial – assuming it could be done – will be very poor and will raise the question whether it is really needed.

A final point about this kind of explanatory trial is that any study which mimics a laboratory experiment will not necessarily show what the best treatment is under *routine* conditions in daily practice for the majority of diabetic patients. Pragmatic clinical trials [3] may help to discover what is the most effective and safe treatment available and/or degree of blood glucose control which is desirable in the long term. Perhaps it is wise to accept, occasionally, moderately imperfect

control of hyperglycaemia at least at the end of the night instead of running the risk of severe nocturnal hypoglycaemia [10]. Of course 'pragmatic' studies [3] aimed at choosing the most practical treatment for achieving near-normalization of blood glucose levels in diabetic patients who need euglycaemia cannot readily be extrapolated since the treatments studied must be given under routine conditions. Pragmatic trials studying pump treatments are certainly needed. The 'feasibility phase' of the National Institutes of Health trial will thus probably be useful. But, as far as trials on insulin treatment are concerned, we would like to suggest that it would be more rational to spend funds not on explanatory trials but on methods designed to simplify the normalization of glycaemia and to make it safe and available for the majority of diabetic patients.

Yours sincerely

G. Tchobroutsky, D. Job, G. Slama and E. Eschwege

References

1. Siebert C (1982) Diabetes control trial begins at 21 medical centers. *Diabetes Dateline* 3: 1–2
2. Marliss EB (1982) Insulin: sixty years of use. *N Eng J Med* 306: 362–363
3. Schwartz D, Flamant R, Lellouch J (1980) *Clinical trials*. Academic Press, London 1 vol.
4. Job D, Eschwege E, Guyot C, Aubry JP, Tchobroutsky G (1976) Effect of multiple daily insulin injections on the course of diabetic retinopathy. *Diabetes* 25: 463–469
5. Schiffrin A, Belmonte MM (1982) Comparison between continuous subcutaneous insulin infusion and multiple injections of insulin. A one-year prospective study. *Diabetes* 31: 255–264
6. Steno Study Group (1982) Effect of 6 months of strict metabolic control on eye and kidney function in insulin-dependent diabetics with background retinopathy. *Lancet* 1: 121–124
7. Tamborlane WV, Hintz RL, Bergman M, Genel M, Felig P, Sherwin RS (1981) Insulin-infusion pump treatment of diabetes. Influence of improved metabolic control on plasma somatomedin levels. *N Eng J Med* 305: 303–307
8. Anonymous (1982) Death among patients using continuous subcutaneous insulin infusion pumps. *United States. Morbidity Mortality Weekly Report* 31: 80–82
9. Anonymous (1982) Notes and news: deaths in diabetics using insulin infusion pumps. *Lancet* 1, 636
10. Unger RH (1982) Nocturnal hypoglycemia in aggressively controlled diabetes. *N Eng J Med* 306: 1294

Dr. G. Tchobroutsky
Hôtel-Dieu Hospital
1 Place du Parvis Notre Dame
75181 Paris Cedex 04, France

Glucagon Induced Hyperkalaemia in Diabetes

Dear Sir,

Van Gaal and de Leeuw recently reported in a Letter to the Editor [1] their data showing a highly significant positive correlation between potassium and high blood glucose levels (> 20 mmol/l) in 22 admissions for diabetic ketoacidosis. They concluded that 'hyperglycaemia is the single most important cause of hyperkalaemia in these patients'; without evaluating any other factors.

We recently documented a hyperkalaemic role of glucagon in normal subjects [2]. Van Gaal and de Leeuw in their Letter suggest that we concluded from our work in normal volunteers that hyperglucagonaemia is a major cause of hyperkalaemia in decompensated diabetes. In fact we reported that circulating glucagon might be a contributory factor in the change in potassium homeostasis in decompensated diabetes.

In further studies, we documented that in insulin-dependent diabetic patients the increase in potassium following the withdrawal of an insulin infusion, which had maintained the patients euglycaemic overnight, was not correlated with the increase in blood glucose, while it was abolished by somatostatin (which inhibits glucagon) [3]. These data do not exclude the importance of other mechanisms, but suggest again that the increase in glucagon which occurs during non-ketoacidotic decompensation could at least play a part in causing hyperkalaemia. This latter phenomenon could be due to mobilization of potassium from the liver as we have documented in normal subjects [4].

Yours sincerely,

F. Massara, E. Cagliero and G. M. Molinatti

References

1. Van Gaal L, de Leeuw I (1982) Glucagon and hyperkalaemia in decompensated diabetes. *Diabetologia* 22: 492 (Letter)
2. Massara F, Martelli S, Cagliero E, Camanni F, Molinatti GM (1980) Influence of glucagon on plasma levels of potassium in man. *Diabetologia* 19: 414–417
3. Cagliero E, Martina V, Massara F, Molinatti GM (1983) Glucagon-induced increase in plasma potassium levels in Type 1 (Insulin-dependent) diabetic subjects. *Diabetologia* 24: 85–87
4. Massara F, Cagliero E, Martina V, Orzan F, Carini G, Molinatti GM (1982) Glucagon stimulation of potassium efflux from the liver. *Diabetologia* 23: 300 (Abstract)

Professor F. Massara
Cattedra Endocrinologia
Corso Polonia 14
I-10126 Turin
Italy

Virus-Induced Diabetes: Cytomegalovirus and Multiple Environmental Insults

Dear Sir,

Several viruses can cause diabetes in mice [1]. Encephalomyocarditis (EMC) virus infects pancreatic β cells and produces a severe form of diabetes in certain inbred strains of mice [2]. Under some circumstances, reovirus also infects β cells, and in addition, triggers an autoimmune response resulting in a mild form of diabetes characterized by abnormalities in glucose tolerance tests which generally disappear within several weeks [3, 4]. Diabetes may also result from cumulative β cell damage produced by sequential environmental insults. For example, in mice, a sub-diabetogenic dose of streptozotocin reduces the β cell reserve and enhances the capacity of certain viruses (e.g., Coxsackie B viruses) to produce diabetes [5, 6]. The present experiments show that murine cytomegalovirus (CMV) infection leads to mild focal β cell damage, and that multiple environmental insults can increase the severity of diabetes.

Newborn NIH Swiss male and female mice were infected with the Smith strain of CMV [7] or reovirus type 1 that had been passaged more than seven times in mouse pancreatic islet cell cultures to increase their tropism for β cells [4]. Two or four weeks later, the mice were challenged with a sub-diabetogenic dose of streptozotocin (1 mg/mouse) [5, 6], or the D variant of EMC virus [2], or CMV. Beginning one week after the challenge, each mouse was bled four times, and the glucose index and percentage of animals developing diabetes were determined as described previously [2].

Table 1 shows that when NIH Swiss male mice were infected with only CMV or reovirus, or given only streptozotocin, diabetes did not