Erratum

A Practical Approach to the Hepatobiliary Kinetics of ^{99m}Tc -HIDA. Clinical Validation of the Method and a Preliminary Report on its Use for Parametric Imaging

[Eur J Nucl Med (1983) 8:292–298]

G. Galli¹, C.L. Maini¹, M. Salvatori¹, and F. Andreasi²

¹ Istituto di Medicina Nucleare

² Istituto di Fisica Medica, Università Cattolica del Sacro Cuore, Policlinico 'A. Gemelli'
Via Pineta Sacchetti 644, Roma, Italy

1. On page 293, the caption of Figure 1 should read:

Fig. 1. Gamma fit of the hepatogram (dotted line) in 4 different situations – *Upper left*: normal case ($r=0.993$); *upper right*: cirrhosis ($r=0.983$); *bottom left*: cirrhosis ($r=0.981$); *bottom right*: obstructive jaundice ($r=0.996$). The *precordial* curve is also reported (biexponential fitting)

2. On page 295, the last line in the left column should read:

e) *increasing* numerical values of TT .

3. On page 298, Formula (7) should read:

$$B=b^{-1}=\left[1+\frac{c_1 t_1-c_2 t_2}{S_0}-\frac{c_1-c_2}{S_{-1}}\right] / \left(\frac{S_1-S_0}{S_0-S_{-1}}\right). \quad (7)$$

Formula (9) should be

$$\sigma_B^2 = \frac{1}{\left[S_0 \left(\frac{S_1-S_0}{S_0-S_{-1}}\right)\right]^2} \cdot \left[c_1 \left(t_1 - \frac{S_0}{S_{-1}}\right)^2 + c_2 \left(t_2 - \frac{S_0}{S_{-1}}\right)^2\right]. \quad (9)$$

Formula (10) should be

$$\sigma_A^2 = \left(\frac{S_0}{S_{-1}}\right)^2 \cdot \sigma_B^2 + \frac{c_1+c_2}{S_{-1}^2}. \quad (10)$$