

Separate influence of dietary carbohydrate and fibre on the metabolic control in diabetes

G. Riccardi, A. Rivellese, D. Pacioni, S. Genovese, P. Mastranzo and M. Mancini

Institute of Internal Medicine and Metabolic Disease, Second Medical School, University of Naples, Naples, Italy

Summary. To clarify the separate influences of digestible carbohydrate and of dietary fibre on blood glucose control and serum lipoproteins, 14 diabetic patients (six Type 1 and eight Type 2) were submitted to three weight-maintaining diets for 10 days each: (1) low carbohydrate/low fibre diet with 42% carbohydrate and 20 g fibre; (2) high carbohydrate/low fibre diet (carbohydrate 53%, fibre 16 g); (3) high carbohydrate/high fibre diet (carbohydrate 53%, fibre 54 g). In comparison with the low carbohydrate/low fibre diet, the 2-h post-prandial blood glucose and the daily blood glucose profile decreased significantly on the high carbohydrate/high fibre diet, without significant changes during the high carbohydrate/low fibre diet. The diet-induced modifications of blood glucose control were similar in both types of diabetic patients (two-way analysis of variance: $F=5.86$, $p<0.02$ for dietary treatment and $F=2.09$, NS for type of diabetes). Total and low-density lipoprotein cholesterol were also decreased after the high carbohydrate/high fibre diet in comparison with the low

carbohydrate/low fibre diet ($p<0.001$ for both), while they were not significantly modified after the high carbohydrate/low fibre diet. Again the modifications of low density lipoprotein cholesterol induced by diet were similar in both types of diabetic patients ($F=10.02$, $p<0.005$ for dietary treatment and $F=0.14$ for type of diabetes, NS). High-density lipoprotein cholesterol was lower after the two test diets than after the low carbohydrate/low fibre diet. However statistical significance was found only for the high carbohydrate/high fibre diet ($p<0.005$). In conclusion: (1) a simple increase of digestible carbohydrate without a parallel increase of dietary fibre does not help in improving the metabolic control of diabetic patients; (2) the hypoglycaemic and hypolipidaemic effects of high carbohydrate/high-fibre diets are due principally to dietary fibre.

Key words: Diet, Type 1 and 2 diabetes, blood lipids, lipoproteins, dietary fibre, dietary carbohydrate, blood glucose.

The American Diabetes Association Committee on Food and Nutrition recommends that most diabetic patients, especially those with Type 1 (insulin-dependent) diabetes, use a diet containing 50%–60% of the caloric intake in the form of carbohydrate (CHO), limiting the consumption of fats below 30% of the total energy intake [1]. Similar recommendations have been issued by other diabetic associations [2, 3]. This new strategy was adopted with the aims of reducing the rate of atherogenesis and improving blood glucose control in patients with diabetes.

However, criticisms have been raised against these dietary recommendations owing to the lack of sound experimental data on which to establish the new strategy [4]. The available information on the metabolic effects of high CHO diets is controversial [5–8].

Many confounding variables obscure the influence of CHO on lipids and carbohydrate metabolism. In many studies, high CHO diets were used which, at the same time, were high in vegetable fibre. We have demonstrated that a high fibre diet is able, per se, to improve

blood glucose control and to lower low-density lipoprotein cholesterol in diabetic patients [9, 10]. In other studies diets were not physiological, both in the use of non-natural foods (liquid formula diet) and in the choice of a very high CHO content (75%–85%) [5, 6].

This study was therefore undertaken to evaluate the effects of a simple increase of dietary CHO on blood glucose metabolism and on serum lipoproteins, by comparison with a parallel increase of both dietary CHO and fibre. All diets were composed of natural foodstuffs and were tried in patients with both types of diabetes.

Subjects and methods

Subjects

Eight Type 2 (non-insulin-dependent) and six Type 1 (insulin-dependent) diabetic patients participated in the study after giving informed consent. The mean \pm SD age was 47 ± 5 and 33 ± 13 years, respectively. The average body mass index was 28 ± 5 and 22 ± 4 kg/m², and the duration of diabetes was 7 ± 2 and 12 ± 6 years.

Table 1. Composition of the 2200 kcal diet

	Low CHO/low fibre		High CHO/low fibre		High CHO/high fibre	
	(g/day)	(%) ^a	(g/day)	(%) ^a	(g/day)	(%) ^a
<i>Protein</i>	121	21	98	17	98	17
<i>Fat</i>						
Total	93	37	76	30	74	30
Saturated	26		17		18	
Mono-unsaturated	54		50		47	
Poly-unsaturated	13		9		9	
Cholesterol	<0.250		<0.250		<0.250	
<i>Carbohydrate</i>						
Total	252	42	318	53	321	53
Simple	108		111		112	
Complex	144		207		209	
<i>Plant fibre</i>	20		16		54	

^a Contribution to total (100%) available energy content of diet

Table 2. A daily scheme of the fibre-rich diet

<i>Breakfast</i>		<i>Snack</i>	
Skimmed milk	(100 g)	Apple	(150 g)
Wholemeal bread	(40 g)		
Cottage-cheese	(30 g)	<i>Dinner</i>	
		Butter beans	(80 g)
<i>Lunch</i>		Sole fish	(100 g)
Spaghetti	(70 g)	Mushrooms	(300 g)
Beef (lean)	(100 g)	Wholemeal bread	(100 g)
Artichokes	(300 g)	Strawberries	(250 g)
Wholemeal bread	(80 g)		
Orange	(200 g)	<i>Snack</i>	
		Apple	(150 g)

50 g olive oil was consumed daily

Seven of the Type 2 diabetic patients were treated with glibenclamide (5–20 mg), while the remaining patient was on diet alone. The dosage of hypoglycaemic drugs was kept constant throughout the experiment.

All the Type 1 diabetic patients were treated with two daily injections of insulin (isophane + soluble) before breakfast and dinner. The insulin dosage was changed (± 4 units/day) if severe hypoglycaemic attacks or marked glycosuria (> 5 g/day) occurred, but the time of insulin therapy and the type of insulin used were always the same.

Diets

Patients were admitted to a metabolic ward and given three weight-maintaining diets in a random order and for consecutive periods of 10 days. The diets were composed of natural food stuffs easily available at any food-shop. Their composition is outlined in Table 1: (1) a low CHO (42%)/low fibre (20 g) diet which resembled a traditional diet for diabetic patients used in many countries [11]; (2) a high CHO/low fibre diet with an increased proportion of CHO (53%) and a low amount of fibre as in the previous diet (16 g); (3) a high CHO/high fibre diet with increased amounts of both CHO (53%) and fibre (54 g).

The three diets had similar amounts of soluble CHO and an identical polyunsaturated/saturated fat ratio. They were distributed between the two main meals (lunch and dinner), a light breakfast and three snacks. Each meal contained a fixed proportion of nutrients and fibre and was consumed at the same time every day.

The vegetable fibre in the high CHO/high fibre diet was provided mainly by leguminous sources (45%) and by different types of vegetables (37%), the remaining part being wholemeal bread (12%) and

fruit (6%). One of the various daily menus of the fibre-rich diet is given in Table 2. Calories, nutrients and fibres were calculated from tables of food composition [12] prepared using the Southgate method for the analysis of dietary fibre [13].

Experimental procedure

During each experiment, patients were weighed daily. At the end of each dietary period blood glucose levels were measured [14] on samples taken in the fasting state, before lunch, before dinner and 2 h after lunch and dinner. On the same day a fasting serum specimen was taken for the analysis of cholesterol [15] and triglycerides [16] in total serum and in the major lipoprotein classes, which were separated by preparative ultracentrifugation [17].

Statistical analysis

The results of the low CHO/low fibre diet were compared with those of each of the two remaining diets. Statistical significance was tested by Student's paired t-test [18].

A two-way analysis of variance [18] was utilized to test simultaneously the hypothesis that changing the type of diet from a low CHO/low fibre diet to either a high CHO/high fibre or a high CHO/low fibre diet has a similar influence on the various metabolic variables considered in this study (F_2) and that this influence is similar for patients with both types of diabetes (F_1). Moreover this type of analysis evaluates the possibility that there is interaction between the diets and the type of diabetes (F_3).

Results

Blood glucose control

The influence of the three diets on blood glucose levels is shown in Figure 1. Fasting blood glucose did not show any significant modification during the three dietary periods, with average concentrations of 6.9 ± 2.5 mmol/l at the end of the low CHO/low fibre diet, 6.8 ± 2.0 mmol/l after the high CHO/low fibre diet and 6.4 ± 2.2 mmol/l after the high CHO/high fibre diet. The 2-h post-prandial blood glucose was significantly lower after the high CHO/high fibre diet (6.8 ± 2.7 mmol/l) than after either the high CHO/low fibre diet (9.0 ± 2.9 mmol/l, $p < 0.02$) or the low CHO/low fi-




Fig. 1. Blood glucose control at the end of each dietary period. The bar indicates the average concentration + standard deviation for the whole group of patients

Table 3. Variations in blood glucose levels on changing from the low CHO/low fibre diet to either the high CHO/low fibre or to the high CHO/high fibre diet, evaluated for Type 1 and Type 2 diabetic patients

Blood glucose (mmol/l)		Variations seen when changing diets	
		Low CHO to high CHO with low fibre	Low CHO to high CHO + low fibre to high fibre
Fasting	Type 1	-0.38 ± 4.97	-2.09 ± 1.83
	Type 2	1.19 ± 2.57	-0.17 ± 1.18
		F ₁ = 2.55	F ₂ = 1.98 F ₃ = 0.03
2-h Post-prandial	Type 1	-0.15 ± 1.36	-2.83 ± 2.32
	Type 2	1.13 ± 4.10	-1.01 ± 1.21
		F ₁ = 2.09	F ₂ = 5.86 ^b F ₃ = 0.07
Average daily	Type 1	-0.05 ± 1.67	-1.29 ± 1.75
	Type 2	0.68 ± 2.40	-1.05 ± 0.87
		F ₁ = 0.52	F ₂ = 4.86 ^a F ₃ = 0.13

Results expressed as mean ± SD. Two way analysis of variance: F₁ = type of diabetes; F₂ = type of diet; F₃ = interaction. ^ap < 0.05; ^bp < 0.02

bre diet (8.7 ± 3.9 mmol/l, p < 0.01). No significant difference was found between the high CHO/low fibre and the low CHO/low fibre diet.

A similar pattern was seen in the average daily blood glucose which decreased after the high CHO/high fibre diet (6.4 ± 2.2 mmol/l) compared with both the high CHO/low fibre diet (7.8 ± 2.0 mmol/l, p < 0.02) and the low CHO/low fibre diet (7.5 ± 1.5 mmol/l, p < 0.005). Once again no significant change was recorded between the two low fibre diets with low or high CHO content.

Blood glucose level variations, observed when changing from the low CHO/low fibre diet to either the high CHO/low fibre diet or the high CHO/high fibre diet, were then evaluated for each patient and a two-way analysis of variance was performed (Table 3). This type of analysis was chosen in order to test whether the two high CHO diets, with different contents of vegetable fibre, had different influences on blood glucose parameters. At the same time, by this type of analysis it was possible to compare the effects of these diets in Type 1 diabetic patients with those obtained in Type 2 patients.

Fasting blood glucose was not significantly influenced by either the type of diabetes or the type of diet (Table 3).

In the group with Type 1 diabetes, the 2-h post-prandial blood glucose concentration decreased by only 0.2 ± 1.4 mmol/l after the high CHO/low fibre diet and by as much as 2.8 ± 2.3 mmol/l after the high CHO/high fibre diet, by comparison with the low CHO/low fibre diet. In Type 2 diabetic patients the 2-h post-prandial blood glucose concentration increased by 1.1 ± 4.1 mmol/l after the high CHO/low fibre diet. Conversely it fell by 1.0 ± 1.2 mmol/l after the high CHO/high fibre diet. The type of dietary treatment (F₂ = 5.86, p < 0.02) but not the type of diabetes (F₁ = 2.09) had a significant influence on this parameter. No interaction between diets and type of disease (F₃ = 0.07) was statistically significant (Table 3).

The average daily blood glucose concentration showed a similar pattern. In Type 1 patients it showed a slight decrease after the high CHO/low fibre diet




Fig. 2. Composition of the major lipoprotein classes at the end of each dietary period. The bar indicates the average concentration + standard deviation for the whole group of patients

(-0.1 ± 1.7 mmol/l) and a more pronounced reduction after the high CHO/high fibre diet compared with the low CHO/low fibre diet (-1.3 ± 1.7 mmol/l). In Type 2 patients the average daily blood glucose increased by 0.7 ± 2.4 mmol/l after the high CHO/low fibre diet. It decreased, instead, by 1.1 ± 0.9 mmol/l after the high CHO/high fibre diet. Once again only the type of diet ($F_2 = 4.86$, $p < 0.05$) was related to this parameter (Table 3).

Serum lipoproteins

In Figure 2 the concentrations of the major serum lipoproteins are shown after each dietary period (available for six Type 2 and five Type 1 patients).

The LDL cholesterol concentration was significantly lower after the high CHO/high fibre diet (2.8 ± 0.7 mmol/l) than after the low CHO/low fibre diet (3.6 ± 0.8 mmol/l, $p < 0.001$) or the high CHO/low fibre diet (3.8 ± 1.3 mmol/l, $p < 0.01$). No significant difference was found between the two low fibre diets with different amounts of CHO. Total serum cholesterol levels behaved in a similar way. The concentration fell to 4.7 ± 0.7 mmol/l after the high CHO/high fibre diet, from 5.6 ± 0.9 mmol/l after the low CHO/low fibre diet ($p < 0.001$) or 5.9 ± 1.3 mmol/l after the high CHO/low fibre diet ($p < 0.01$). The difference between the two low fibre diets was not significant.

The high-density lipoprotein (HDL) cholesterol was also reduced after the high CHO/high fibre diet compared with the low CHO/low fibre diet (1.3 ± 0.3 versus 1.5 ± 0.4 mmol/l, $p < 0.005$). After the high CHO/low fibre diet the HDL cholesterol concentration of 1.4 ± 0.4 mmol/l was not significantly different from the other two diets.

No significant differences between the three diets were found in the concentrations of total serum triglycerides. The triglyceride composition of the very-low-density lipoprotein class (VLDL) was instead significantly reduced after the high CHO/high fibre diet compared with the high CHO/low fibre diet (0.9 ± 0.5 versus 1.1 ± 0.6 mmol/l, $p < 0.05$). The VLDL triglyceride concentration after the low CHO/low fibre diet (0.8 ± 0.3 mmol/l) was not significantly different from that found after the other two diets.

The ratio VLDL + LDL/HDL cholesterol was lower after the high CHO/high fibre diet (2.6 ± 1.2) than after the high CHO/low fibre diet (3.5 ± 1.5 , $p < 0.02$) or the low CHO/low fibre diet (2.9 ± 1.2). In the latter case the difference did not reach statistical significance. The comparison between the high CHO/low fibre and the low CHO/low fibre diet showed a difference which was significant at the 5% level.

The influence of the two high CHO diets and of both types of diabetes on the composition of the major serum lipoproteins was then checked by two-way analysis of variance performed on the variations between the concentration of serum lipoproteins after each of the two high CHO diets and that found at the end of the low CHO diet (Table 4).

A significant influence of the type of dietary treatment ($F_2 = 10.02$, $p < 0.005$) was found only for LDL cholesterol and not for HDL cholesterol or VLDL triglycerides. Conversely a significant influence of the type of diabetes was found for VLDL triglycerides ($F_1 = 6.55$, $p < 0.02$).

Body weight

Despite the fact that the three diets were designed to keep body weight constant, a slight but significant in-

Table 4. Variations in serum lipoprotein composition changing from the low CHO/low fibre diet to either the high CHO/low fibre or to the high CHO/high fibre diet, evaluated for Type 1 and Type 2 diabetic patients

Lipoproteins (mmol/l)		Variations seen when changing diet		
		Low CHO to high CHO with low fibre	Low CHO to high CHO + low fibre to high fibre	
LDL Cholesterol	Type 1	0.35 ± 0.65	-0.68 ± 0.66	
	Type 2	0.17 ± 0.99	-0.73 ± 0.38	
		F ₁ = 0.14	F ₂ = 10.02 ^b	F ₃ = 0.04
HDL Cholesterol	Type 1	-0.16 ± 0.29	-0.23 ± 0.18	
	Type 2	-0.09 ± 0.29	-0.22 ± 0.15	
		F ₁ = 0.21	F ₂ = 0.98	F ₃ = 0.11
VLDL Triglycerides	Type 1	-0.02 ± 0.29	-0.28 ± 0.45	
	Type 2	0.90 ± 1.10	0.42 ± 0.74	
		F ₁ = 6.55 ^a	F ₂ = 1.37	F ₃ = 0.12

Results expressed as mean ± SD. Two way analysis of variance: F₁ = type of diabetes; F₂ = type of diet; F₃ = interaction. ^a*p* < 0.02; ^b*p* < 0.005

crease in body weight was observed at the end of the high CHO/high fibre diet (64.4 ± 11.6 kg) compared with the low CHO/low fibre diet (64.0 ± 11.5 kg, *p* < 0.02). No significant difference was found after the high CHO/low fibre diet (64.2 ± 11.6 kg).

Treatment

Although Type 2 diabetic patients did not have to modify their drug treatment during the study, four out of six Type 1 patients had to reduce their insulin dosage by an average of 4 units/day during the high CHO/high fibre diet in order to avoid severe hypoglycaemic attacks.

Discussion

The results of this study clearly demonstrate that a simple increase of digestible CHO (up to 53% of the daily caloric intake) gives no benefit at all in terms of blood glucose control, compared with the traditional low fibre (20 g)/low CHO (42%) diet (still used in many countries). Instead, parallel increases of dietary CHO (53%) and fibre (54 g) produced a significant improvement in blood glucose control, whether evaluated as post-prandial blood glucose or as daily blood glucose profile (Fig. 1). Furthermore the high CHO/high fibre diet is equally effective in both Type 1 and Type 2 patients (Table 3).

Other groups have compared a high CHO/high fibre diet with a traditional low CHO/low fibre diet, obtaining a substantial improvement in blood glucose control with the former diet [19, 20]. This positive effect on blood glucose control was ascribed both to the high level of dietary CHO and to vegetable fibre. Our findings clearly demonstrate that increasing the amount of CHO in the diet from 240 to 320 g/day has very little influence on fasting blood glucose concentrations and, if

anything, contributes to the deterioration of blood glucose control in the post-prandial phase.

An increase in the amount of fibre in the diet is more important for diabetic patients. We have shown in this study (Fig. 1), confirming our previous findings, that dietary fibre, per se, can reduce blood glucose concentrations in diabetic patients [9]. Therefore any positive effect of a high CHO/high fibre diet on blood glucose control in diabetes has to be ascribed mainly to the high level of dietary fibre.

A diet for the treatment of diabetic patients has to be judged not only for its effects on blood glucose control but also for its influence on atherogenesis and on the major risk factors for atherosclerosis, such as lipids and lipoproteins. Unfortunately, because of the experimental design of this study, only short-term lipoprotein results are available. Therefore further studies with longer follow-up periods are needed before drawing more definitive conclusions on the influence of dietary fibre on serum lipoprotein composition.

A simple increase of dietary CHO without changes in vegetable fibre did not modify LDL cholesterol concentrations in all the patients (Fig. 2) to any significant extent. This finding was not unexpected even if CHO replaced the lipids in the high CHO/low fibre diet. Any variation in dietary lipids was obtained keeping constant the ratio between polyunsaturated and saturated fats. It is well known that changes in the amount of dietary fats have little effect on serum lipoprotein composition, if this ratio is kept constant [21].

Dietary fibre had a more pronounced effect than CHO on the LDL lipoprotein fraction. A significant reduction of LDL cholesterol was, in fact, recorded after the high CHO/high fibre diet (Fig. 2). This was of the same order of magnitude in both types of diabetes (Table 4). A similar finding was found with a high fibre diet in patients with Type II hyperlipoproteinaemia without diabetes [22].

HDL cholesterol was higher after the low CHO diet than after either of the two high CHO diets, the difference being significant only for the high CHO/high fibre diet (Fig. 2). The two-way analysis of variance, however, did not show any significant difference between the effects of the two high CHO diets on HDL (Table 4), thus suggesting that a low dietary CHO level rather than a low amount of dietary fibre might be able to increase the HDL cholesterol concentration in diabetes. This is confirmed by a recent study by Simpson et al. in which HDL levels were higher after a simple reduction of dietary CHO without any modification of dietary fibre [23]. These changes in HDL cholesterol reflect, in part, parallel modifications in LDL cholesterol concentrations [24] and are probably transient [25, 26]. The VLDL + LDL/HDL cholesterol ratio, a feature of the lipoprotein pattern commonly considered an index of potential atherogenicity, was significantly reduced after the high CHO/high fibre diet compared with the high CHO/low fibre diet.

In Type 2 patients VLDL triglyceride concentrations increased after changing from the low CHO diet to either of the two high CHO diets, as shown by analysis of variance (Table 4). This suggests that the mechanism of the CHO-induced hyper-VLDL triglyceridaemia is working only in Type 2 patients in whom dietary CHO can produce endogenous hyperinsulinism [27, 28] which, in turn, is responsible for accelerating VLDL synthesis [29, 30]. However, dietary fibre can influence VLDL metabolism, reducing VLDL triglyceride concentration (Fig. 2).

In conclusion, increasing the CHO consumption in the diabetic diet gives no benefit in terms of blood glucose control or serum lipoprotein composition. Conversely, increasing at the same time both dietary CHO and fibre improves blood glucose control and reduces LDL cholesterol together with HDL cholesterol. Therefore, according to the present evidence, an increase of available CHO is justified for the dietary treatment of patients with diabetes only when dietary fibres are increased in a parallel fashion. The type of diet we have used (high in fibre and with a moderate amount of CHO) seems to have the therapeutical properties to justify a trial on a larger scale. Moreover, it is composed only of natural food stuffs and is designed to resemble closely the type of diet used in Mediterranean countries a few decades ago.

Acknowledgements. We gratefully acknowledge the cooperation of Mr. A. Gurrieri – the cook of the metabolic unit. This study was supported by Cilag Foundation for Therapeutic Research. This paper was presented, in part, at the 18th Annual Meeting of the EASD.

References

- American Diabetes Association (1979) Principles of nutrition and dietary recommendations for individuals with diabetes. *Diabetes Care* 2: 520–523
- British Diabetic Association (1982) Dietary recommendations for diabetics for the 1980's. *Hum Nutr Appl Nutr* 36A: 378–394
- Special Report Committee (1981) Guidelines for the nutritional management of diabetes mellitus: A special report from the Canadian Diabetes Association. *J Can Dietet Assoc* 42: 110–118
- Reaven GM (1980) How high the carbohydrate? *Diabetologia* 19: 409–413
- Brunzell JD, Lerner RL, Hazzard WR, Porte D jr, Bierman EL (1971) Improved glucose tolerance with high carbohydrate feeding in mild diabetes. *N Engl J Med* 284: 521–524
- Brunzell JD, Lerner RL, Porte D jr, Bierman EL (1979) Effect of a fat free high carbohydrate diet on diabetic subjects with fasting hyperglycemia. *Diabetes* 23: 138–142
- Weiser RL, Seeman A, Herrera G, Assal GP, Soeldner JS, Gleason RE (1974) High and low carbohydrate diets in diabetes mellitus: study of effects on diabetic control, insulin secretion and blood lipids. *Ann Intern Med* 8: 332–341
- Kjehm TG, Anderson JW, Ward K (1976) Beneficial effects of a high carbohydrate, high fibre diet on hyperglycemic diabetic men. *Am J Clin Nutr* 29: 895–899
- Rivellese A, Riccardi G, Giacco A, Pacioni D, Genovese S, Mattioli PL, Mancini M (1980) Effect of dietary fibre on glucose control and serum lipoproteins in diabetic patients. *Lancet* 2: 447–450
- Rivellese A, Riccardi G, Giacco A, Pacioni D, Genovese S, Mattioli PL, Mancini M (1981) A fibre rich diet for the treatment of diabetes. In: Howard AN, McLean Baird I (eds) Recent advances in clinical nutrition I. J Libbey, London, pp 99–100
- Jenswell AS, Thomas BJ, Brown AM (1975) Survey of dietary policy and management in British diabetic clinics. *Br Med J* 2: 7–11
- Fidanza F, Versiglione N (1982) Tabelle di composizione degli alimenti. Idelson, Napoli
- Southgate DAT (1969) Determination of carbohydrates in foods. II. Unavailable carbohydrates. *J Sci Food Agric* 20: 331–335
- Hjelm M (1966) Enzymatic determination of hexoses in blood and urine *Scand. J Clin Lab Invest* 18: (Suppl 192): 85–89
- Klose S, Hagen A, Greif H (1975) Méthode de dosage colorimétrique du cholestérol par voie entièrement enzymatique adaptée à tous les types de auto-analyseur. In: *Biologie prospective*, 111. Colloque de Pont-à-Mousson. L'Expansion Scientifique Française, Paris, pp 505–537
- Wahlefeld AN (1974) Triglycerides determination after enzymatic hydrolysis. In: Bergmeyer HV (ed) *Methods of enzymatic analysis*, Academic Press, New York
- Carlson K (1973) Lipoprotein fractionation. *J Clin Path* 26 (Suppl 5): 32
- Snedecor GW, Cochran WH (1967) *Statistical Methods*. Ames Iowa USA
- Simpson HCR, Simpson RW, Lousley S, Carter RD, Gekh M, Hockaday JDR, Mann JI (1981) A high carbohydrate leguminous fibre diet improves all aspects of diabetic control. *Lancet* 2: 1–5
- Anderson JW, Kyllen W (1979) High carbohydrate-high fiber diets for insulin treated men with diabetes mellitus. *Am J Clin Nutr* 2: 2312–2321
- Report of a WHO Expert Committee (1982) Prevention of coronary heart disease. Technical Report series 678. WHO, Geneva
- Rivellese A, Riccardi G, Giacco A, Postiglione A, Mastranzo P, Mattioli PL (1983) Reduction of risk factors for atherosclerosis in diabetic patients treated with a high fiber diet. *Prev Med* 12: 128–132
- Simpson HCR, Carter RD, Lousley S, Mann JI (1982) Digestible carbohydrate – an independent effect on diabetic control in Type 2 (non-insulin-dependent) diabetic patients? *Diabetologia* 23: 235–239
- Riccardi G, Rivellese A, Romano G, Giacco R, Mancini M (1984) The influence of dietary fibre on serum lipoproteins in diabetic and hyperlipidaemic patients In: Carlson LA, Olsson AC (eds) *Treatment of hyperlipoproteinaemia*. Raven Press, New York (in press)
- Jenkins DJA, Reynolds D, Slavin B, Leeds AR, Jenkins AL, Jepson EM (1980) Dietary fiber and blood lipids: treatment of hypercholesterolemia with guar crispbread. *Am J Clin Nutr* 33: 575–581
- Taskinen M, Nikkilä EA, Ollus A (1983) Serum lipids and lipoproteins in insulin-dependent diabetic subjects during high-carbohydrate, high-fiber diet. *Diabetes Care* 6: 224–230
- Greenfield M, Koltermann O, Olefsky J, Reaven GM (1980) Mechanism of hypertriglyceridaemia in diabetic patients with fasting hyperglycaemia. *Diabetologia* 18: 441–446
- Farquhar JW, Frank A, Gross RC, Reaven GM (1966) Glucose, insulin and triglyceride responses to high and low carbohydrate diets in man. *J Clin Invest* 45: 1641–1656
- Rivellese A, Riccardi G, Farinara E, Capaldo B, Mancini M (1978) Relationship between insulin response to intravenous glucose and VLDL triglycerides in a healthy population. *Diabetes and obesity*. In: Vague J, Vague Ph (eds) *Proceedings of the Vth International Meeting of Endocrinology*. Excerpta Medica, Amsterdam, Oxford, pp 303–306
- Olefsky MJ, Farquhar JW, Reaven GM (1974) Reappraisal of the role of insulin in hypertriglyceridemia. *Am J Med* 57: 552–560

Received: 21 February 1983
and in revised form: 8 September 1983

Dr. G. Riccardi
Clinica Medica 2
2 Facoltà di Medicina e Chirurgia
Nuovo Policlinico
Via S. Pansini, 5
I-80131 Naples, Italy