

## *Erratum*

### **Radioimmunoassay for 2,4-dichlorophenoxyacetic acid**

**Dietmar Knopp<sup>1</sup>, Peter Nuhn<sup>2</sup> and Hans-Joachim Dobberkau<sup>1</sup>**

<sup>1</sup> Research Institute of Hygiene and Microbiology, Department of Environmental Toxicology, Heinrich-Heine-Straße 12, DDR-9933 Bad Elster, German Democratic Republic

<sup>2</sup> Martin-Luther-University Halle – Wittenberg, Section of Pharmacy, Weinbergweg 15, DDR-4020 Halle, German Democratic Republic

Arch Toxicol (1985) 58: 27–32

The legend to Fig. 1 on p 29 should read as follows:

**Fig. 1.** Sensitivity and cross-reactivity of antiserum No. 22, using the iodine tracer and the following derivatives: 2,4-D-tyramine conjugate ( $\Delta-\Delta$ ), 2-(4-Chloro-2-methylphenoxy)propionic acid ( $\bullet-\bullet$ ), 2-(2,4-Dichlorophenoxy)propionic acid ( $\circ-\circ$ ), 2,4-Dichlorophenoxyacetic acid ( $\ominus-\ominus$ ), 2,4-Dichlorophenoxyacetic acid methylester ( $\bullet-\bullet$ ), 4-(4-Chloro-2-methylphenoxy) butyric acid ( $\bullet-\bullet$ ), 4-(2,4-Dichlorophenoxy) butyric acid ( $\ominus-\ominus$ )