

Erratum

Effect of 2,3,7,8-Tetrachlorodibenzo-p-dioxin on the Hepatic Storage of Retinol in Rats with Different Dietary Supplies of Vitamin A (Retinol)

Tuula Thunberg¹, Ulf G. Ahlborg², Helen Håkansson¹, Cilla Krantz²,
and Mats Monier²

¹ Department of Toxicology, Karolinska Institutet,
P.O. Box 60, 400, S-104 01 Stockholm, Sweden

² Toxicology Laboratory National Food Administration,
P.O. Box 622, S-751 26 Uppsala, Sweden

(1980) **45**: 273–285

Page 274 Line 30 should be McConnell et al. (1978a)

Page 275 Line 7 should be McConnell et al. (1978a, b)

Page 276 Line 36 should be 100 µl 0.5 M

Page 276 Line 36 should be pH 7.0

Figures 2 and 3 should be read as follows:




Fig. 2. Retinol concentrations in the liver of control and TCDD-treated (p.o.) rats 4 weeks after administration expressed as µg retinol per g liver. Retinol concentration in food was 1.2, 3.0, and 6.0 mg · kg⁻¹ for the low, normal and high vitamin A diets respectively




Fig. 3. Total amount of retinol in the liver of control and TCDD-treated (p.o.) rats 4 weeks after administration. Retinol concentration in food was 1.2, 3.0, and 6.0 µg · kg⁻¹ for the low, normal and high vitamin A diets respectively