

- Raschke, K., Humble, G. D.: No uptake of anions required by opening stomata of *Vicia faba*: Guard cells release hydrogen ions. *Planta (Berl.)* **115**, 47–58 (1973)
- Sayre, J. D.: Physiology of stomata of *Rumex patientia*. *Science (N.Y.)* **57**, 205–206 (1923)
- Sayre, J. D.: Physiology of stomata of *Rumex patientia*. *Ohio J. Sci.* **26**, 223–266 (1926)
- Scarsh, G. W.: Mechanism of the action of light and other factors on stomatal movement. *Plant Physiol.* **7**, 481–504 (1932)
- Willmer, C. M., Dittrich, P.: Carbon dioxide fixation by epidermal and mesophyll tissue of *Tulipa* and *Commelina*. *Planta (Berl.)* **117**, 123–132 (1974)
- Willmer, C. M., Pallas, J. E., Jr., Black, C. C., Jr.: Carbon dioxide metabolism in leaf epidermal tissue. *Plant Physiol.* **52**, 448–452 (1973)

Erratum

Edwards, K.L., Scott, T.K.: Rapid Growth Responses of Corn Root Segments: Effect of pH on Elongation.

Planta (Berl.) **119**, 27–37 (1974)

Elongation rates in the text and Figs. 2, 3, 4, 5, 7 are multiplied by a factor of five and should read “ $\mu\text{m}/5 \text{ min}/10 \text{ segments}$ ”, the ordinate of Fig. 6 should be expressed as “Increase in Length/10 Segments (0.2 mm)”, and line 22 on p. 36 should read “(1) can be 5–10 times greater than that of shoot tissue”.