

*Erratum*

Eur J Appl Physiol (1981) 46: 121–133

**Lactate After Exercise in Man:  
I. Evolution Kinetics in Arterial Blood**

H. Freund and P. Zouloumian

Centre d'Etudes Bioclimatiques (CNRS),  
21, rue Becquerel, F-67087 Strasbourg Cedex, France

Due to a deplorable technical mishap Table 2 on page 128 was not published correctly. Please find the correct version on the reverse.

**Table 2.** Characteristics of the experimental and fitted arterial blood lactate recovery curves; supine position (Series 3 and 4)

Recovery from exercise at	Series 3						Series 4					
	60% $\dot{V}O_2$ max			90% $\dot{V}O_2$ max			90% $\dot{V}O_2$ max			90% $\dot{V}O_2$ max		
	A <sup>a</sup> (n = 3)	A (A.R.)	A (F.D.)	A (S.M.)	A <sup>a</sup> (n = 3)	A <sup>a</sup> (n = 3)	A <sup>a</sup> (n = 3)	A <sup>a</sup> (n = 6)	A <sup>a</sup> (n = 6)	A <sup>a</sup> (RE) (n = 6)	A <sup>a</sup> (RE) (n = 6)	
$L_a(0)^b$ , $\mu\text{mol} \cdot \text{l}^{-1}$	3,760 $\pm$ 1,175	3,480	4,870	4,770	4,373 $\pm$ 775	6,153 $\pm$ 944	6,642 $\pm$ 1,161	6,153 $\pm$ 944	8,555 $\pm$ 1,156	9,177 $\pm$ 1,789	9,177 $\pm$ 1,789	
$L_a$ max <sup>c</sup> , $\mu\text{mol} \cdot \text{l}^{-1}$	3,834 $\pm$ 1,172	6,980	7,550	8,550	7,693 $\pm$ 795	547 $\pm$ 234	547 $\pm$ 222	547 $\pm$ 222	6,939 $\pm$ 1,320	4,713 $\pm$ 1,504	6,649 $\pm$ 1,621	6,649 $\pm$ 1,621
$L_a(\infty)$ , $\mu\text{mol} \cdot \text{l}^{-1}$	648 $\pm$ 163	407	536	866	610 $\pm$ 234	6,939 $\pm$ 1,320	-10,702 $\pm$ 926	-10,320 $\pm$ 2,095	-11,571	-10,702 $\pm$ 926	-12,606 $\pm$ 2,348	-12,606 $\pm$ 2,348
$A_1$ , $\mu\text{mol} \cdot \text{l}^{-1}$	838 $\pm$ 613	7,734	5,414	7,667	6,939 $\pm$ 1,320	0.486 $\pm$ 0.129	0.626 $\pm$ 0.199	0.507	0.507	0.486 $\pm$ 0.129	0.610 $\pm$ 0.136	0.610 $\pm$ 0.136
$A_2$ , $\mu\text{mol} \cdot \text{l}^{-1}$	-3,950 $\pm$ 1,747	-10,807	-9,728	-10,807	-10,702 $\pm$ 926	0.0741	0.0709 $\pm$ 0.005	0.0741	0.0709 $\pm$ 0.005	0.0511 $\pm$ 0.013	0.0854 $\pm$ 0.041	0.0854 $\pm$ 0.041
$\gamma_1$ , $\text{min}^{-1}$	0.831 $\pm$ 0.374	0.348	0.603	0.603	0.507	67-167	67-167	67-167	67-167	67-167	60-147	60-147
$\gamma_2$ , $\text{min}^{-1}$	0.0832 $\pm$ 0.010	0.0649	0.0736	0.0736	0.0741	27.5	14.5	32	97	32	27.5 <sup>e</sup>	27.5 <sup>e</sup>
SD <sup>d</sup> , $\mu\text{mol} \cdot \text{l}^{-1}$	24-65	56	62	62	62							
$\theta_2$ , min	11.0 <sup>e</sup>	27.5	14.5	14.5	14.5							

A = continuous exercise; (RE) = active recovery

<sup>a</sup> Average of the individual values  $\pm$  SD<sup>b</sup>  $L_a(0)$  = the measured arterial blood lactate concentration at the end of exercise<sup>c</sup>  $L_a$  max = the highest arterial blood lactate concentration during the recovery<sup>d</sup> Standard deviation between the experimental and the fitted data. Accuracy of the estimation (100  $\sigma/\bar{x}$ ) of the parameters  $A_1$  and  $\gamma_1$  is 0.5-4%;  $A_2$  and  $\gamma_2$ , 0.2-2%<sup>e</sup> Measured on the mean arterial - brachial venous blood lactate difference curve