

Erratum

Detection of Lactate Dehydrogenase B₄ in Human Hemolysate of Patients Deficient in Lactate Dehydrogenase B Subunit Activity Using Enzyme Immunoassay¹

Kayoko Sudo^{2,3} Masato Maekawa,² and Takashi Kanno²

On p. 631 of this article, Table III contained some incorrect data. The correct representation of that table appears below.

Table III. LD-B₄ Concentration of Heterozygous Individuals with Deficient in LD-B Subunit Activity in Hemolysate^a

| Case no. | LD-B ₄ concentration (nmol/g Hb) | LD-B ₄ activity (U/g Hb) | Total LD activity (U/g Hb) |
|----------------------------|---|-------------------------------------|----------------------------|
| A-1 | 1.37* | 23.4** | 123.1** |
| A-2 | 1.10* | 14.2** | 100.0** |
| A-3 | 1.54* | 18.7** | 116.1** |
| B-4 | 0.31* | 14.4** | 91.6** |
| C-5 | 0.62* | 11.0** | 94.2** |
| D-6 | 0.35* | 19.7** | 98.3** |
| D-7 | 0.70* | 20.2** | 101.7** |
| Mean | $\bar{X} = 0.86^*$ | $\bar{X} = 17.4^{**}$ | $\bar{X} = 100.6^{**}$ |
| SD | 0.49 | 4.3 | 13.6 |
| normal control ($N = 8$) | | | |
| Mean | $\bar{X} = 1.04^*$ | $\bar{X} = 67.0^{**}$ | $\bar{X} = 181.0^{**}$ |
| SD | 0.60 | 9.3 | 23.0 |

^aFour families (A, B, C and D) were examined. LD-B₄ activity was calculated from total LD activity and LD-B₄ isozyme %.

*Statistically nonsignificant ($P > 0.05$).

**Statistically significant ($P < 0.01$).

¹ *Biochem. Genet.* **24**:625–633 (1986).

² Department of Laboratory Medicine, Hamamatsu University, School of Medicine, Handa-cho 3600, Hamamatsu City, 431-31, Japan.

³ To whom correspondence should be addressed.