- Paul DH, Roberts BL (1976) Responses of cerebellar nuclear neurones in the elasmobranch fish Scyliorhinus canicula. J Physiol 257:43–44 (Abstr)
- Platt CJ, Bullock TH, Czéh G, Kovačević N, Konjević Dj, Gojković M (1974) Comparison of electroreceptor, mechanoreceptor and optic evoked potentials in the brain of some rays and sharks. J Comp Physiol 95:323–355
- Rizzolo A (1929) A study of equilibrium in the smooth dogfish (*Galeus canis* Mitchill) after removal of different parts of the brain. Biol Bull 57:245–249
- Roberts BL (1981) Central processing of acousticolateralis signals in elasmobranchs. In: Tavolga WN, Popper AN,

Fay RR (eds) Hearing and sound communication in fishes. Springer, Berlin Heidelberg New York, pp 357–373

- Ten Cate J (1935) Physiologie des Zentralnervensystems der Fische. Erg Biol 11:335-409
- Tong SL, Bullock TH (1982) Electroreceptive representation and its dynamics in the cerebellum of the catfish, *Ictalurus nebulosus* (Ictaluridae, Siluriformes). J Comp Physiol 145:289–298
- Tong SL, Lee L, Bullock TH (1982) Distribution of responses to visual, mechanical and electrical stimulation in the cerebellum of catfish and ray. Neuroscience 7 (Suppl):S212 (Abstr)

## Erratum

Stein PSG, Robertson GA, Keifer J, Grossman ML, Berenbeim JA, Lennard PR (1982) Motor neuron synaptic potentials during fictive scratch reflex in turtle. J Comp Physiol 146:401–409

The first sentence of the fourth paragraph of the Discussion (p. 407) should read:

The co-activation of HP motor neurons and KE motor neurons during the early A2 phase of the turtle scratch reflex is in contrast to the reciprocal relationship observed between these populations during the flexion reflex demonstrated in the immobilized turtle (Figs. 7 and 8).