Journal of Comparative Physiology B © Springer-Verlag 1994

Erratum

There were several serious errors in the article N. Ohkudo, S. Watabe, T. Oshiro, F. Takashima, H. Nakajima Subunit structures of multiple hemoglobins in carp J Comp Physiol B 163:445-451. Figure 7 is correctly reprinted below: β^1 : $\frac{\text{Val}}{\text{Asp}}$ Glu Trp Thr Asp Ala Glu Arg Ser Ala¹⁰ Ile Ile Ala Leu Trp Gly β^2 : Val Glu Trp Thr Asp Ala Glu Arg Ser Ala¹⁰ Ile Ile Gly Leu Trp Gly β^1 : Lys Leu Asn Pro²⁰ Asp Glu Leu Gly Pro $\frac{Gln}{Glu}$ Ala Leu Ala Arg³⁰

 β^2 : Lys Leu Asn Pro²⁰ Asp Glu Leu Gly Pro $\frac{\text{Gln}}{\text{Glu}}$ Ala Leu Ala Arg³⁰ $\frac{\text{Arg}}{\text{Cys}}$ Leu Ile β^2 : Val Tyr Pro Trp Thr Gln Arg⁴⁰

Fig. 7. N-Terminal a-a sequences of carp Hb β -chains. The β^1 - and β^2 -chains were isolated from carp Hb components CI and CIII, respectively. The sequence of the β^1 -chain from component CII was almost identical except for the heterogeneity at the N-terminal a-a. The one differing residue between the β^1 and β^2 -chains is underlined with *double lines*, while residues in β^1 - and β^2 -chains of this study differing from these reported by Grujic-Injac et al. (1980) are underlined with a single line

In addition, in the legend to Fig. 8, "lanes 9 and 11" should have been "lanes 9, 10 and 11". The legend is reprinted below:

Fig. 8. Exchange of globin subunits between carp Hb components. Hemolyzate (lanes 1, 8 and 12), component CI (lanes 2 and 5), component CII (lanes 9, 10 and 11) component CIII (lanes 3 and 6), and component CI plus CIII (lanes 4 and 7) were applied to PAGE after incubation at 4 °C for 12 h (lanes 1-4) and 24 h (lane 10), or at 37 °C for 12 h (lanes 5-8) and 24 h (lane 11)