

4) Logistic regression analysis is appropriate, as BMI was entered into the equation as a categorical variable, in three categories, where the rate in the two extreme ones is compared to the middle one.

5) Our questionnaire referred to all types of habitual activity, including work time as well as at leisure time. The rate of persons involved in any type of sport was minimal. "High" physical activity was defined as activity score in the highest tertile of the study group. The score consisted of the weekly number of hours spent in moderate activity (heavy activity was not reported by any of the 502 individuals constituting the subsample interviewed for dietary intake and physical activity) + half of the hours spent in light activity defined according to the Committee on Dietary Allowances [2]. Using this crude measure, and accounting for age, BMI and total caloric intake, the risk for Type 2 diabetes for "highly" active individuals of both sexes was 0.67 with 90% confidence limits 0.47–0.96 ($p=0.04$) [3]. This was validated by significantly lower insulin response in the "highly" active individuals in all sex and BMI categories (unpublished data).

Yours sincerely
M. Modan, A. Karasik and H. Halkin

References

1. Modan M, Karasik A, Halkin H, Fuchs Z, Lusky A, Shitrit A, Modan B (1986) Effect of past and concurrent body mass index on prevalence of glucose intolerance and Type 2 (non-insulin-dependent) diabetes and on insulin response: the Israel study of glucose intolerance obesity and hypertension. *Diabetologia* 29: 82–89
2. Committee on Dietary Allowances Food and Nutrition Board (1980) Recommended dietary allowances, 9th revised edn. National Academy of Sciences, p 24
3. Modan M, Lubin F, Shitrit A, Lusky A (1986) Habitual diet, physical activity, body mass index and glucose intolerance. The Israel GOH Study, *Diabetes* 35 [Suppl 1]: 74 A

Dr. M. Modan
Department of Clinical Epidemiology
The Chaim Sheba
Medical Center
Tel Hashomer 52621,
Israel

The editor apologizes for the inadvertent error made during the editing and production process of Dr. Modan's letter [Diabetologia (1986) 29:408]. Instead of "in both Type 1 and Type 2 diabetes" it should have been printed "in both diabetes categories", as originally submitted by Dr. Modan and her associates. The editor deeply regrets this technical error.

Type 1 (insulin-dependent) diabetes and the question of heterogeneity

Dear Sir,

I read with interest the report entitled "HLA-DR3 is associated with a more slowly progressive form of Type 1 (insulin-dependent) diabetes" by Ludvigsson et al. [1]. Since many of those interested in the causes of Type 1 diabetes mellitus believe in the likelihood of genetic heterogeneity, studies attempting to correlate phenotypes with genetic markers are welcome and may shed light on this important question. Unfortunately, most, if not all, claims of such correlations have lacked adequate confirmation. This confusing state of affairs has involved age of onset and other clinical parameters, as well as immunological indicators such as islet cell antibodies, insulin antibodies, etc.

The results of Ludvigsson et al. [1] seem to be in apparent conflict with data we reported recently [2] from cross sectional studies of se-

rum C-peptide in Type 1 diabetic patients and their siblings. We found in this study that DR4+ diabetic patients showed better preservation of their C-peptide blood concentrations than diabetic patients with other HLA types. It is difficult to reconcile these findings with the Ludvigsson et al. [1] results showing that DR3+ diabetic patients have a milder disease. Both our study [2] (not prospective) and the Ludvigsson et al. study [1] (based on questionnaire data and multicenter) have weaknesses. Thus, this subject needs independent confirmation.

The observation by Ludvigsson et al. [1] that 15 patients were DR4–, DR3–, DR2+, presumably mostly from France, is of considerable interest. We have recently shown that there is a third HLA-related susceptibility axis for diabetes (in addition to the DR3– and DR4– related axes) defined by homozygous typing cells (Dw) and restriction fragment length polymorphisms which is part of DR2-LD-MN2 as well as DR1-Dw1 haplotypes [3]. Those 15 patients presumably carried one or both of these haplotypes.

Yours sincerely,
J. Barbosa

References

1. Ludvigsson J, Samuelsson U, Beauforts C, Deschamps I, Dorchy H, Drash A, Francois R, Herz G, New M, Schober E (1986) HLA-DR3 is associated with a more slowly progressive form of Type 1 (insulin-dependent) diabetes. *Diabetologia* 29: 207–210
2. Hoogwerf BJ, Rich SS, Barbosa JJ (1985) Meal-stimulated C-peptide and insulin antibodies in type I diabetic subjects and their nondiabetic siblings characterized by HLA-DR antigens. *Diabetes* 34: 440–445
3. Segall M, Noreen H, Schluender L, Swenson M, Barbosa J, Bach FH (1986) DR2+ haplotypes in insulin-dependent diabetes: analysis of DNA restriction fragment length polymorphisms. *Human Immunology* (in press)

Professor J. Barbosa,
Division of
Endocrinology and Metabolism
Department of Medicine
University of Minnesota
Minneapolis, Minnesota 55455
USA

Hyperkalaemia after interruption of CSII

Dear Sir,

It has recently been suggested that the serum potassium concentrations on admission to hospital with ketoacidosis may be higher in patients treated by continuous subcutaneous insulin infusion (CSII) compared to those receiving conventional injection therapy [1].

In consideration of the possible risks of hyperkalaemia, it may be of interest to reconsider the study which we published in *Diabetologia* in 1982 [2], where CSII was deliberately interrupted for 9 h in 9 resting and fasting Type 1 (insulin-dependent) diabetic patients. We did not record plasma potassium levels at the time of the study, but we have now measured by flame photometry the values on the stored plasma samples (kept at -40°C) from the 7 C-peptide negative patients (all male, mean \pm SD age 33 ± 12 years, duration of diabetes 13 ± 6 years). Figure 1 shows the results, together with the changes in plasma glucose, blood 3-hydroxybutyrate and plasma free insulin concentration, which were assayed at the time of the study. Mean plasma potassium levels increased throughout the 9 h and were significantly different from baseline at all time points from 6 h ($p < 0.01$, t-test) until the end of the experiment ($p < 0.01$).