ERRATA

Siberian Mathematical Journal, $\underline{7}$, 5 (1966)

| Where Found | Was Given As | Should Be |
|---|---|--|
| p. 776, line 22 p. 776, line 26 p. 776, line 27 p. 776, line 28 p. 776, line 37 p. 793, line 8 (from the bottom) p. 815, line 11 p. 843, line 18 p. 894, line 4 p. 894, line 9 p. 906, lines 7 and 8 (from the bottom) | subgroup must $\delta = \pi$ (ms of π (ms of \mathfrak{G}) subgroup contains group for $i > r_1$ and the Sylow class of group. We hyperplane forms the set (G) gen are investigated we have, the associated for, the lines L_1^{k+1} , follows; $L_d^{k,q}$ follows $L_c^{k,q}$ if $0 \le c < d \le +\infty$; $L_b^{k,q}$ | subgroup \mathfrak{M} must $\delta = \pi$ (ms of \mathfrak{M}) \cap π (ms of \mathfrak{G}) subgroup \mathfrak{M} contains group \mathfrak{G} for $i > r_1$ and the Sylow class $\langle \mathfrak{P}_{r_i} \rangle$ of group \mathfrak{G} . We hyperplane $(O, \mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3)$ forms the set \mathfrak{R} (G) \mathfrak{M} gen are investigated we have $dE^1 E^1 $, the associated for $\omega^2 = \omega^3 = 0$, the lines L_1^{k+1}, q follows L_1^{k}, q ; L_0^k, q follows L_1^{k}, q ; L_0^k, q follows L_1^{k}, q if $0 \le c < d \le +\infty$; L_0^{k}, q precedes L_0^{k+1}, q^{-1} if $b > 1$, |
| | precedes if b > 1, c < ∞; | $L_b^{k,q}$ precedes $L_b^{k+1,q}$ if b $c < \infty$; |