

Errata

---

SEX PHEROMONE OF EGYPTIAN COTTON  
LEAFWORM (*Spodoptera littoralis*). ITS  
CHEMICAL TRANSFORMATION UNDER  
FIELD CONDITIONS

A. SHANI and J.T. KLUG

In Ref. 1, the following errors appeared:

Page 876, 6 lines up. Change *N,N*-diethyl-*m*-toluamide to *N*-octyl-*N'*-phenyl-*p*-phenylenediamine.

Page 877, 11 lines down. Change UOP to DEET.

Page 878, 3 lines down. Change (UOP or BHT) to (DEET or BHT).

Page 879, 5 lines up. Change UOP to DEET.

Page 880, 3rd entry and 6th entry in Table 1. Change Prodlure and UOP to Prodlure and DEET.

REFERENCE

SHANI, A., and KLUG, J.T. 1980. *J. Chem. Ecol.* 6:875-881.

TRAPPING THE WESTERN PINE BEETLE AT AND  
NEAR A SOURCE OF SYNTHETIC ATTRACTIVE  
PHEROMONE: EFFECTS OF TRAP SIZE  
AND POSITION

P.E. TILDEN, W.D. BEDARD, D.L. WOOD, K.Q. LINDAHL,  
and P.A. RAUCH

In Ref. 1, the following error appeared:

The first sentence of the second paragraph below Table 4 on p. 528 should read, "When the surrounding traps were 0.19 m<sup>2</sup>, the proportion of males