Errata

SEX PHEROMONE OF EGYPTIAN COTTON LEAFWORM (*Spodoptera littoralis*). ITS CHEMICAL TRANSFORMATION UNDER FIELD CONDITIONS

A. SHANI and J.T. KLUG

In Ref. 1, the following errors appeared:

Page 876, 6 lines up. Change *N,N*-diethyl-*m*-toluamide to *N*-octyl-*N'*-phenyl-*p*-phenylenediamine.

Page 877, 11 lines down. Change UOP to DEET.

Page 878, 3 lines down. Change (UOP or BHT) to (DEET or BHT).

Page 879, 5 lines up. Change UOP to DEET.

Page 880, 3rd entry and 6th entry in Table 1. Change Prodlure and UOP to Prodlure and DEET.

REFERENCE

SHANI, A., and KLUG, J.T. 1980. *J. Chem. Ecol.* 6:875-881.

TRAPPING THE WESTERN PINE BEETLE AT AND NEAR A SOURCE OF SYNTHETIC ATTRACTIVE PHEROMONE: EFFECTS OF TRAP SIZE AND POSITION

P.E. TILDEN, W.D. BEDARD, D.L. WOOD, K.Q. LINDAHL,
and P.A. RAUCH

In Ref. 1, the following error appeared:

The first sentence of the second paragraph below Table 4 on p. 528 should read, "When the surrounding traps were 0.19 m², the proportion of males

caught at the source of attractant was significantly different from the proportions caught 1.5 m away, 1.5 m high and 4.5 m away, 3.0 m high (Table 2). The proportion of males caught 1.5 m away, 1.5 m high was significantly different from the proportion caught 1.5 m away, 3.0 m high." We thank J.A. Byers for bringing this error to our attention.

REFERENCE

- TILDEN, P.E., BEDARD, W.D., WOOD, D.L., LINDAHL, K.Q. and RAUCH, P.A. 1979. *J. Chem. Ecol.* 5:519-531.