

Erratum

In the paper "Serologic Differences Among Nondefective Reticuloendotheliosis Viruses" by P.-Y. Chen *et al.* [Arch Virol 93: 233–245 (1987)] Table 2 on p. 238 was incorrect and should be read as demonstrated on the following page.

Table 2. Cross neutralization tests among 26 reticuloendotheliosis virus isolates

Sub type	Isolate	Virus dose ^a	Homol serum titer	Relative serum titers (\log_2) ^b																			
				DIA	MN 67	MN 81	RU-1	T-A	T-F	T-G	T-Q	T-S	T-T	T-FP	T-E	T-A	RPL	170 A	DIA	SN	MN 2	MN 3	MN 4
1	T-C	8	1:40	0	1	0	2	1	0	0	4	3	2	3	3	3	2	3	2	3	2	3	2
	T-F	32	1:80	1	0	0	1	1	0	2	1	3	3	2	4	3	2	3	2	3	2	3	1
	T-S	16	1:40	1	1	0	1	1	0	2	2	3	3	2	2	2	1	2	1	2	2	2	1
	T-T	16	1:80	1	1	0	0	2	1	0	2	2	3	3	2	3	3	1	1	2	1	3	2
	T-A	8	1:80	0	1	0	1	0	3	3	2	2	2	2	1	1	3	2	3	2	3	2	1
	RPL	16	1:40	0	1	0	2	1	0	0	4	3	2	3	3	3	2	3	2	3	3	2	2
	170 A	16	1:40	0	2	0	1	1	0	2	2	2	2	3	2	0	2	2	1	2	1	2	2
2	DIA	32	1:320	1	3	2	3	2	2	0	0	1	2	1	1	2	2	2	2	2	2	3	2
	SN	32	1:160	2	3	1	0	3	1	1	0	0	0	1	0	2	1	1	2	2	1	1	2
	MN 67	8	1:80	2	2	1	2	2	1	0	0	0	1	1	2	1	1	2	1	1	3	0	2
	MN 81	1	1:160	2	2	1	2	2	1	1	0	0	1	1	2	1	1	3	0	2	1	2	1
	RU-1	12	1:80	2	2	1	2	2	1	0	0	1	2	0	1	2	1	1	3	0	1	2	0
	221 A	12	1:160	2	2	2	2	3	2	1	0	0	1	2	1	0	0	1	2	1	1	2	1
	225 A	4	1:160	2	2	1	2	3	2	1	1	1	2	1	1	0	0	0	1	1	1	2	1
3	339	128	1:80	2	3	1	1	3	2	2	1	1	3	2	2	1	1	1	2	1	1	3	1
	348 G	4	1:80	1	2	1	1	3	2	1	1	0	3	2	1	1	0	0	2	1	0	1	3
	363	15	1:160	2	2	2	1	3	2	1	1	1	2	2	2	2	0	1	0	1	2	1	2
	375 A	32	1:80	1	3	1	1	2	2	1	2	2	2	2	2	1	1	0	1	0	1	1	1
	375 B	15	1:160	1	1	1	3	1	2	0	0	2	1	0	1	0	1	0	1	0	0	1	0
	377 A	16	1:160	2	2	1	1	3	2	1	1	2	3	2	2	1	0	0	1	0	0	0	2
	383 A	32	1:80	2	2	1	1	3	2	1	1	0	3	2	3	2	1	0	0	2	0	0	0
	401	32	1:160	2	3	1	2	3	2	1	1	0	3	1	1	0	2	0	1	2	1	0	1
	488	12	1:80	1	2	1	1	2	1	1	2	2	1	2	2	2	1	1	2	1	1	2	0
	489	32	1:160	1	2	1	1	2	1	1	1	1	2	1	2	2	0	1	2	1	1	2	0
	CS	16	1:80	1	2	1	1	2	1	1	1	1	2	1	1	2	1	0	1	1	2	0	1
	260	32	1:80	1	3	1	1	3	2	2	1	3	1	2	2	1	1	2	2	1	1	0	2

^a Calculated number of tissue culture infectious units per well^b The reciprocal of the titer of a serum against its homologous virus was divided by the reciprocal of the titer against the test virus and expressed as \log_2 . Means of relative titers for each of the 9 blocks are given in Table 3