

## Combined effects of cadmium and salinity on development and survival of herring eggs\*

H. VON WESTERNHAGEN, H. ROSENTHAL & K.-R. SPERLING

*Biologische Anstalt Helgoland (Zentrale);  
Hamburg 50, Federal Republic of Germany*

**KURZFASSUNG:** Kombinierte Wirkungen von Cadmium und Salzgehalt auf Entwicklung und Überleben von Heringseiern. Eier des herbstlaichenden Ostseeherings (*Clupea harengus* L.) wurden in natürlichem und in Cadmium kontaminiertem Seewasser (Konzentrationen: 0,1; 0,5; 1,0; 5,0 ppm Cd) bei verschiedenen Salzgehalten (5 ‰, 16 ‰, 25 ‰, 32 ‰) erbrütet. Die Toxizität von Cd wurde im Hinblick auf folgende Kriterien bewertet: Embryoaktivität und Überlebensraten bis zum Schlupf, Veränderungen der Eimembran, Verschiebung des Schlupfzeitpunktes, Unterschiede in den Schlupfraten, Anteil der lebensfähigen Larven und deren mittlere Totallänge sowie Durchmesser der Augen und Gehörkapseln frischgeschlüpfter Larven. Darüber hinaus wurde die Aufnahme von Cd in Abhängigkeit von Konzentration und Salzgehalt im Verlauf der Embryonalentwicklung bestimmt. Die Beeinträchtigung der Heringsembryonen durch Cd war in brackigem Wasser stärker als in unverdünntem Meerwasser. Auch die konzentrationsabhängige Aufnahme von Cd nahm mit sinkendem Salzgehalt des Erbrütungsmediums zu.

### INTRODUCTION

Cadmium, a biologically non-essential heavy metal, is normally found in minute amounts in sea water (GOLDBERG 1965). In Atlantic offshore waters concentrations are usually not higher than 0.4 ppb with a geometric mean at 0.04 ppb (PRESTON 1973). CHESTER & STONER (1974) found even lower values ranging between 0.02 and 0.18 ppb. In United States continental shelf waters, concentrations range between 0.06 and 0.11 ppb (WINDOM & SMITH 1972). In North Sea waters, concentrations of 0.01 to 0.62 ppb (PRESTON 1973) and 0.024 to 0.25 ppb (MULLIN & RILEY 1956) are considered normal. In the Baltic Sea, equally low mean concentrations of 0.17 to 0.22 ppb have been recorded in open waters (KREMLING 1973). Apart from geological sources, additional cadmium enters coastal areas as a result of industrial activities such as zinc refining and mining, cadmium plating and alloy manufacture, pesticides and fertilizer sources (especially phosphate fertilizers) and paint and pigment manufacture. An increasing number of recent reports indicate that in river estuaries (BUTTERWORTH et al. 1972) and near shore waters of heavily industrialized areas

---

\* This study has been financially supported by the Deutsche Forschungsgemeinschaft (Grant No. Ro 380/2).

(PRESTON et al. 1972, JONES et al. 1973), cadmium content of sea water has risen appreciably (ABDULLAH et al. 1972). In the inner part of the Oslofjord, ANDERSEN et al. (1973) found cadmium concentration to be between 0.67 and 0.92 ppb and cadmium levels in decapitated and eviscerated herring and sprat to vary from 0.2 to 0.7 ppm (dry weight basis). The high cadmium levels in parts of the Severn estuary are also reflected in high accumulation levels in *Littorina* (178 ppm) and *Fucus* (53 ppm) (LEATHERLAND & BURTON 1974).

Of many aquatic poikilothermic organisms, fishes appear to be most resistant to the effects of heavy metals (SCHWEIGER 1957, EISLER 1971). Invertebrates such as molluscs and crustaceans have been shown to be more sensitive to the toxic action of these pollutants (SCHWEIGER 1957, BROWN & AHSANULLAH 1971, EISLER 1971, BIESINGER & CHRISTENSEN 1972, O'HARA 1973b, THURBERG et al. 1973).

Although it is known that larvae of oysters and crustaceans are up to a thousand times more susceptible to heavy metals than adults of the same species (CONNOR 1972), little work has been done on the influence of metals on the early stages of fishes. PICKERING & GAST (1972) found reduced survival in embryos of the fathead minnow *Pimephales promelas* reared in water containing more than 57  $\mu\text{g}$  Cd/l. ROSENTHAL & SPERLING (1974) showed that herring eggs, incubated in sea water of about 16 ‰ salinity and a cadmium concentration of 1.0 ppm, yielded a very low percentage of viable larvae (16.3 ‰) compared to 93 ‰ in the controls. At 5.0 ppm no viable hatch occurred.

The tolerance of aquatic organisms to cadmium varies considerably. In fresh water, tolerance limits for cadmium are to a great extent dependent on the hardness of the water (PICKERING & HENDERSON 1966). In sea water, salinity seems to have a comparable influence on the toxicity of this metal. OLSON & HARREL (1973), using the bivalve *Rangia cuneata*, proved that the toxic action of copper, mercury and chromium was mitigated in water of high salinities.

The present study was established to examine the influence of cadmium on embryonic development of the herring *Clupea harengus* L. from the Baltic Sea under different salinities. The cadmium concentrations employed in this series of experiments were selected in order to obtain visible deviations from normal embryonic development resulting from short time exposure.

## MATERIAL AND METHODS

Eggs of Baltic autumn spawning herring were artificially fertilized in sea water of 5, 16, 25, and 32 ‰ S at a temperature of 10° C. Immediately after fertilization the eggs were transferred into aerated 2 l incubation jars containing natural sea water (controls) and test solutions of 0.1, 0.5, 1.0 and 5.0 ppm Cd<sup>++</sup>. The test media were renewed every other day, while salinity and pH values (7.53–8.10) were determined immediately before and after each renewal. The test solutions were prepared by adding CdCl<sub>2</sub> from a stock solution immediately before each renewal. Temperature readings and water samples for cadmium determination were taken daily. Cadmium working levels maintained in test solutions throughout the experiments were at about 80 ‰

of the initial concentration. Egg samples (5 each) were preserved at different ontogenetic stages. Determination of cadmium uptake in herring embryos was accomplished by means of AAS (Perkin Elmer Type 300) as described by ROSENTHAL & SPERLING (1974).

In order to assess possible changes in the toxic effects of cadmium at different salinities, the following criteria were used: rate of embryonic survival (daily counts); embryonic activity (number of embryo movements per 3 min); effects on egg membrane; incubation time (time from fertilization to 50% hatching); hatching rates (observed in 24 h intervals); percentage of viable hatch; mean total length and yolk sac length of newly hatched larvae, diameter of eyes and otic capsules; cadmium uptake of eggs during the course of embryonic development.




Fig. 1: *Clupea harengus* embryos incubated at different salinity-cadmium concentrations. Mean number of activities per minute measured 24 h prior to hatching. Solid line = complete revolution; broken line = slight movements and "trembling". Incubation temperature 10° C

## RESULTS

## Embryonic activity

At a certain developmental stage herring embryos perform rotations within their egg capsule (HOLLIDAY et al. 1964, ROSENTHAL 1966). This is presumed to circulate the perivitelline fluid thus improving oxygen provision of the embryo. In herring eggs incubated in cadmium contaminated water, changes in this behaviour became apparent. Depending on cadmium concentration, embryos displayed "trembling"-motions of the whole body and fewer rotations occurred. Embryonic activity in contaminated water was also influenced by the salinity of the incubation medium. Figure 1 shows embryonic activity 24 h before hatching. Effects of cadmium on embryonic activity were most pronounced in highly diluted sea water of 5 ‰ S. There was no trembling behaviour in the controls. A 32 ‰ S no activity measurements were conducted.

Thus the detrimental effect of cadmium on the embryonic activity was most apparent in diluted sea water. The action of the metal diminished as salinity of the incubation medium increased.




Fig. 2: *Clupea harengus* eggs incubated at different salinity-cadmium concentrations. Survival rates until hatching in % of fertilized eggs. Incubation temperature 10° C

### Egg membrane

During the course of development, the chorion of eggs incubated in highly contaminated water became dark and less transparent, suggesting that it was directly affected by the cadmium. Also the softening of the chorion under cadmium influence, described by ROSENTHAL & SPERLING (1974), was observed in all salinities investigated. Egg diameter was not altered by the Cd content of the incubating medium. As development proceeded, eggs reared in cadmium contaminated water became less tightly attached to the incubating panes and burst easily when removed.

### Embryonic survival until hatching

Survival of fertilized eggs in cadmium solutions of up to 1.0 ppm was comparable to that in the controls. Rates of survival decreased considerably in 5.0 ppm Cd, being lowest at 5 and 16 ‰ S (21.7 ‰ and 32.8 ‰, respectively) and highest at 25 and 32 ‰ S (54.9 ‰ and 51.0 ‰, respectively). In 5, 16, and 25 ‰ S and in controls, 0.1, 0.5 and 1.0 ppm Cd hatching rates were high, ranging from 85 ‰ to 99 ‰ of fertilized eggs. Survival rates at 32 ‰ S, being generally lower than in less saline water and never exceeding 67 ‰, did not show as spectacular a drop for high cadmium concentrations as found in other salinities (Fig. 2).

### Incubation period

Figure 3 shows hatching distribution at 24 h intervals at different salinities and cadmium concentrations. Cadmium concentrations higher than 0.1 ppm were generally associated with earlier hatch. This was true for all salinities employed, except for the 5 ‰ S/5.0 ppm Cd trial, where 50 ‰ hatching occurred much later than in the other experiments. However, the hatching process in this trial was abnormal. The embryos, having died prior to or immediately after hatching, were only liberated after the membranes disintegrated due to continued action of released hatching enzymes.

### Viable hatch

At 25 ‰ and 32 ‰ S, 94 ‰ to 97.5 ‰ of the total hatch appeared viable in cadmium concentrations of up to 1.0 ppm (Fig. 4). At 16 ‰ S and 1.0 ppm Cd, 61.5 ‰ of the hatched larvae were considered viable, while at 5 ‰ S hatch was already below 90 ‰ in 0.1 ppm Cd and near zero in 0.5 ppm Cd. No viable larvae were obtained in 5.0 ppm Cd in any of the salinities tested. Subdividing the non-viable larvae into bent, crippled and dead specimens (Fig. 5) allows further insight into the effect of cadmium on the developing eggs of the herring. At 5 ‰ S, larvae were most severely damaged (70 ‰ dead at 5.0 ppm Cd, 100 ‰ crippled at 1.00 ppm Cd). At 16 ‰ S and 0.5 ppm Cd, 2 ‰ of the larvae were dead and 98 ‰ were crippled; at 1.0 ppm Cd, 24.5 ‰ were crippled and 14 ‰ displayed a bent longitudinal body axis.




Fig. 3: *Clupea harengus* larvae incubated at different salinity-cadmium concentrations. Hatching distribution (% day) and 50% hatching time (dark bars). Incubation temperature 10° C

### Mean total length of larvae at hatching

Measurements of mean total length of larvae were only possible for straight specimens. The available data show that in all tested salinities larval size decreased




Fig. 4: Percentage viable hatch of herring larvae (*Clupea harengus*) incubated at different salinity-cadmium concentrations at 10° C




Fig. 5: *Clupea harengus*. Rates of malformation and lethal hatch (%) of larvae incubated at different salinity-cadmium concentrations at 10° C

Table 1  
*Clupea harengus* larvae. Total length, diameter of eye and otic capsule at hatching. n = number of larvae measured;  $\bar{x}$  = mean; s = standard deviation;  $s\bar{x}$  = error of the mean. Larvae derived from incubation trials in 32‰ S originated from a second female

S (‰)	Experimental design Cd conc.	Total length (mm)			Otic capsule (mm)			Eye diameter (mm)					
		n	$\bar{x}$	s	$s\bar{x}$	n	$\bar{x}$	s	$s\bar{x}$	n	$\bar{x}$	s	$s\bar{x}$
5	control	80	8.27 ± 0.34		0.04	80	0.327 ± 0.020		0.002	55	0.288 ± 0.018		0.002
5	0.1 ppm	91	7.77 ± 0.54		0.06	89	0.316 ± 0.024		0.002	64	0.280 ± 0.011		0.001
5	0.5 ppm		not measurable			171	0.313 ± 0.026		0.002	125	0.251 ± 0.016		0.001
5	1.0 ppm		not measurable			179	0.314 ± 0.025		0.002	137	0.252 ± 0.018		0.002
5	5.0 ppm		not measurable			33	0.212 ± 0.039		0.007	39	0.220 ± 0.016		0.003
16	control	100	7.78 ± 0.58		0.06	100	0.314 ± 0.023		0.002	76	0.287 ± 0.014		0.002
16	0.1 ppm	62	7.88 ± 0.62		0.08	65	0.311 ± 0.025		0.003	41	0.284 ± 0.017		0.003
16	0.5 ppm	93	7.70 ± 0.51		0.05	93	0.310 ± 0.021		0.002	43	0.285 ± 0.009		0.001
16	1.0 ppm	105	7.13 ± 0.38		0.04	129	0.301 ± 0.027		0.002	129	0.283 ± 0.014		0.001
16	5.0 ppm		not measurable			141	0.244 ± 0.033		0.003	100	0.258 ± 0.020		0.002
25	control	50	7.98 ± 0.17		0.02	49	0.311 ± 0.015		0.002		0.288		
25	0.1 ppm	63	7.90 ± 0.46		0.06	62	0.312 ± 0.021		0.003	63	0.277 ± 0.018		0.002
25	0.5 ppm	72	7.88 ± 0.41		0.05	71	0.315 ± 0.017		0.002	46	0.289 ± 0.010		0.001
25	1.0 ppm	91	7.48 ± 0.60		0.06	91	0.311 ± 0.034		0.004	91	0.293 ± 0.011		0.001
25	5.0 ppm		not measurable			102	0.244 ± 0.062		0.006	82	0.263 ± 0.024		0.003
32	control	50	7.05 ± 0.28		0.04	53	0.279 ± 0.027		0.004	25	0.274 ± 0.008		0.002
32	0.1 ppm	54	7.00 ± 0.34		0.05	56	0.286 ± 0.020		0.003	50	0.265 ± 0.020		0.001
32	0.5 ppm	61	6.94 ± 0.42		0.05	63	0.285 ± 0.027		0.003	38	0.268 ± 0.014		0.002
32	1.0 ppm	74	6.82 ± 0.41		0.05	73	0.288 ± 0.022		0.002	74	0.272 ± 0.018		0.002
32	5.0 ppm		not measurable			35	0.178 ± 0.035		0.006	45	0.234 ± 0.019		0.003



with increasing cadmium concentration. The relative decrease in mean total length was least in 32 ‰ S, showing only 0.24 mm difference between the controls and 5.0 ppm Cd, while at 16 ‰ S and 25 ‰ S the differences were 0.66 and 0.59 mm, respectively (Table 1).

The smaller size of the larvae obtained from the 32 ‰ S experiments can be attributed to the fact that the eggs used in these trials were derived from a different female than in experiments of all other salinities.




Fig. 6: *Clupea harengus*. Mean eye diameter of newly hatched larvae incubated at different salinity-cadmium concentrations at 10°C. Larvae derived from incubation trials in 32 ‰ S originated from a second female. Basic statistics are given in Table 1

### Eye, otic capsule and yolk sac measurements

At all salinities eye diameter decreased with increasing cadmium concentration. The most pronounced reductions in eye diameter were registered at 5 ‰ S (Table 1, Fig. 6). Otic capsule diameter of newly hatched larvae remained essentially the same in 5 ‰, 16 ‰, and 25 ‰ S at cadmium levels of 0.1, 0.5 and 1.0 ppm and the

controls. At 5.0 ppm Cd otic capsule diameter was considerably reduced (Fig. 7), capsules being smaller at 5 ‰ and 32 ‰ S than at 16 ‰ and 25 ‰ S.

Yolk sac length and height of newly hatched larvae were measured. Owing to the drop shape of the yolk sac under cadmium influence, the formula for a cone was used to approximate yolk sac volume (V):

$$V = \frac{1}{3} \pi r^2 \cdot h$$

Yolk sac volumes increased with decreasing salinity (Fig. 8), as found by ROSENTHAL & MANN (1973). In addition, at all four salinities yolk sac volume increased at higher cadmium concentrations. This was particularly evident in 5 ‰ S.




Fig. 7: *Clupea harengus*. Mean diameter of the otic capsule of newly hatched larvae incubated at different salinity-cadmium concentrations at 10°C. Larvae derived from incubation trials in 32 ‰ S originated from a second female. Basic statistics are given in Table 1

### Cadmium uptake during embryonic development

Amount of cadmium accumulated by herring eggs under exposure to cadmium ions is depicted in Figure 9. Initial cadmium uptake during the first hours after

fertilization was fairly rapid at all salinities. Highest cadmium concentrations were attained during this period. Further exposure to cadmium during development led either to declining cadmium levels in the eggs, or to stability in cadmium values as shown for 0.1 ppm (Fig. 9). Eggs incubated in 5 ‰ S showed highest initial contamination (Fig. 9a) when compared with the respective trials at 16 ‰, 25 ‰ and 32 ‰ S. Maximum levels attained depended on the initial cadmium concentrations




Fig. 8: *Clupea harengus*. Mean yolk sac volume of newly hatched larvae incubated at different salinity-cadmium concentrations at 10° C. Larvae derived from incubation trials in 32 ‰ S originated from a second female

employed in the respective tests. At 16 ‰ (Fig. 9b) cadmium concentrations in eggs, in general, were lower than those found at 5 ‰, but considerably higher than those determined for eggs incubated in 25 ‰ or 32 ‰ S (Fig. 9c, d). Lowest contamination values were recorded for eggs incubated at 32 ‰ S.

Newly hatched larvae at all salinity-cadmium combinations contained only minute amounts of the metal, suggesting that cadmium accumulated primarily in the chorion.

## DISCUSSION

The experiments indicate that toxic effects of cadmium on developing herring eggs vary considerably with the salinity of the incubating medium. In diluted sea water deleterious effects of cadmium-contaminated water on herring embryos were more pronounced than in slightly brackish or natural sea water. Prehatching mortality (Fig. 2) and malformation rates (Fig. 5) were highest in the 5 ‰ S trials. The strong detrimental effects of cadmium on herring eggs incubated in highly diluted sea water




Fig. 9: a, b *Clupea harengus*. Mean cadmium content of eggs ( $\mu\text{g}/\text{individual}$ ) during incubation in different salinity-cadmium concentrations. Each value is based upon 3 to 8 determinations




Fig. 9: c, d *Clupea harengus*. Mean cadmium content of eggs ( $\mu\text{g}/\text{individual}$ ) during incubation in different salinity-cadmium concentrations. Each value is based upon 3 to 8 determinations

are supported by the results of the activity measurements; the least active embryos occurred at 5 ‰ S and 1.0 and 5.0 ppm Cd.

The occasional high rate of survival of eggs in the incubating trial with low cadmium concentrations cannot be satisfactorily explained. EATON (1973) reported similar findings with eggs of the fathead minnow (*Pimephales promelas*). It might be suggested that low cadmium levels slightly inhibit the growth of bacteria, which impede gaseous exchange at the egg membrane.

The softening of the egg membrane under the influence of cadmium indicates changes in the properties of the chorion. EATON (1973) observed that spawned eggs from parental fish of the fathead minnow which had been chronically exposed to cadmium were more loosely attached to the substrate than those from control fish. Changes in the physico-chemical properties of the chorion may have detrimental secondary effects. Further investigations are planned in order to evaluate the physiological and ecological importance of these effects.

From Figure 8 it becomes apparent that yolk sacs of newly hatched larvae were generally larger in high cadmium concentrations than those of larvae incubated in less contaminated water. Whether this effect was caused by the shortened incubation period, as reported by ROSENTHAL & MANN (1973), or was due to the repressed activity of cadmium exposed embryos (Fig. 1) and resulting lower energy requirements cannot as yet be determined. In order to properly evaluate activities of embryos it is necessary to obtain data not only on the frequency but also on the duration of active phases. However, activity measurements seem to be useful to detect sublethal effects of pollutants. In relatively low cadmium concentrations the effects of the metal could be observed even when no morphological malformations occurred.

Not only was the deleterious effect of cadmium on herring eggs mitigated in water of higher salinities, but also the actual cadmium content of analyzed eggs dropped considerably with increasing salinity (Fig. 9). Similar findings were reported by O'HARA (1973 a). Working with the fiddler crab *Uca pugilator*, he found that individuals exposed to cadmium would accumulate more of the metal in low (10 ‰ S) than in high (30 ‰ S) salinities. The author attributed this phenomenon to active uptake of salts by crabs kept in brackish water. However, as reported by VERNBERG & O'HARA (1972) for the same species, a similar pattern for uptake of mercury apparently does not exist.

We can assume that in unpolluted water most of the cadmium added to the test solution as  $\text{CdCl}_2$  was available as uncomplexed  $\text{Cd}^{++}$  in the water (ZIRINO & YAMAMOTO 1972, GARDINER 1974). Thus the fact that Cd content of herring eggs exposed to identical contamination levels decreased with increasing salinity might suggest a mechanism of interaction between available metal ions. As reported by ROSENTHAL & SPERLING (1974), the simultaneous presence of Zn and Cd (Zn being less toxic than Cd, DANIELLI & DAVIES 1951) increased the viable hatch of herring eggs from zero ( $\text{Cd}^{++}$ -contaminated sea water without addition of Zn) to 11.4 ‰ (Cd-Zn solution, Cd concentration: 5.0 ppm).

According to their theory and as suggested by DANIELLI (1944) accumulation of metal takes place only on the surface where cadmium is fixed by formation of chemical compounds with mucopolysaccharides of the chorion which acts as an unspecific ion-

exchange medium. Thus the presence of another bivalent ion seems to trigger competition between the two lowering the amount of either fixed to or introduced into the egg. Therefore the availability of more  $\text{Ca}^{++}$  or other metal ions in higher salinities might be responsible for lower Cd content of the eggs, mitigating toxic effects of cadmium. The apparent interaction between  $\text{Ca}^{++}$  and  $\text{Cd}^{++}$  ions has been pointed out by KOBAYASHI (1971). The extremely high cadmium susceptibility of freshwater teleosts such as *Lepomis macrochirus* and *Micropterus salmoides* (CEARLEY & COLEMAN 1974), *Pimephalus promelas* and *Lepomis macrochirus* (PICKERING & HENDERSON 1966), and *Salmo gairdnerii* (BALL 1967) compared to that of saltwater fishes such as *Fundulus heteroclitus* (EISLER 1971) also suggests competitive interaction between calcium and cadmium.

The question arises as to why cadmium was not accumulated in measurable amounts inside the eggs during the incubation period. Cadmium content of the eggs during embryogenesis declined slightly under most salinity regimes (Fig. 9). The same findings have been reported by ROSENTHAL & SPERLING (1974) incubating eggs of Baltic herring. In contrast to this, marine fish larvae do accumulate cadmium (VON WESTERNHAGEN, DETHLEFSEN & ROSENTHAL unpublished) from the surrounding water, as do crustaceans (O'HARA 1973a), molluscs (YAGER & HARRY 1966, NICKLESS et al. 1972, PEDEN et al. 1973), and teleosts (EISLER et al. 1972, CEARLEY & COLEMAN 1974). A possible explanation lies in the assumption that the chorion, which obviously accumulates most of the cadmium in herring eggs (ROSENTHAL & SPERLING 1974), acts as an ion-exchange medium having the capacity of complexing or binding only a certain amount of bivalent metal ions until saturation, which could occur shortly after exposure. Thus high salinities with more competitive ions would cause lower cadmium levels in the egg membrane due to high concentrations of bivalent metals in the test medium.

Higher cadmium concentrations in the chorion would increase the chance of metal eventually seeping through the membrane into the egg as suggested by DANIELLI & DAVIES (1951) for the reaction of heavy metals on cell surfaces. This would directly affect the embryo.

#### SUMMARY

1. Eggs of autumn spawning Baltic herring (*Clupea harengus* L.) were incubated in cadmium-contaminated water (0, 0.1, 0.5, 1.0, 5.0 ppm) at four salinities (5 ‰, 16 ‰, 25 ‰, 32 ‰) in order to evaluate possible changes in toxicity of Cd.
2. Effects of Cd on embryonic survival were found to be dependent on salinity of the incubating water. Deleterious effects of Cd on developing herring embryos were more pronounced in brackish water than in sea water.
3. Embryonic activity, as a measure of viability of developing embryos, decreased in Cd concentrations with decreasing salinity.
4. In none of the trials was egg diameter altered by the Cd content of the incubation water.

5. In all salinities, incubation time appeared to be shortened with increasing Cd content of the test medium.
6. At 5 ‰, 16 ‰, 25 ‰ and 0, 0.1, 0.5 and 1.0 ppm, hatching rate was not significantly altered by Cd. High hatching rates between 85 to 99 % occurred in all salinity-Cd combinations. At high Cd levels (5.0 ppm), there was greater survival of embryos at high salinities (32 ‰ and 25 ‰) than at low salinities (16 ‰ and 5 ‰).
7. Percentage viable hatch was unaffected at 32 ‰, 25 ‰ and 16 ‰ S and 0, 0.1 and 0.5 ppm Cd. In low salinities (5 ‰), only 1 % viable hatch occurred at 0.5 ppm; in 16 ‰, 61.5 % viable hatch occurred at 1.0 ppm Cd. No viable larvae were obtained in any tests at 5.0 ppm Cd.
8. In all salinities examined, mean total length of newly hatched larvae decreased with increasing Cd concentration of the rearing medium. Relative decrease in mean total length was minimum at 32 ‰ S.
9. In all four test concentrations yolk sac volumes of newly hatched larvae increased with rising Cd concentrations, probably associated with declining embryo activity.
10. The Cd content of eggs was found to be generally higher in lower salinities than in more saline water at comparable Cd concentrations.

## LITERATURE CITED

- ABDULLAH, M. J., ROYLE, L. G. & MORRIS, A. W., 1972. Heavy metal concentration in coastal waters. *Nature*, Lond. **235**, 158-160.
- ANDERSEN, A. T., DOMMASNES, A. & HESTHAGEN, I. H., 1973. Some heavy metals in sprat (*Sprattus sprattus*) and herring (*Clupea harengus*) from the inner Oslofjord. *Aquaculture* **2**, 17-22.
- BALL, I. R., 1967. The toxicity of cadmium to rainbow trout (*Salmo gairdnerii* RICHARDSON). *Wat. Res.* **1**, 805-806.
- BIESINGER, K. E. & CHRISTENSEN, G. M., 1972. Effects of various metals on survival, growth, reproduction, and metabolism of *Daphnia magna*. *J. Fish. Res. Bd Can.* **29**, 1691-1700.
- BROWN, B. & AHSANULLAH, M., 1971. Effect of heavy metals on mortality and growth. *Mar. Pollut. Bull.* **2**, 182-187.
- BUTTERWORTH, J., LESTER, P. & NICKLESS, G., 1972. Distribution of heavy metals in the Severn estuary. *Mar. Pollut. Bull.* **3**, 72-74.
- CEARLEY, J. E. & COLEMAN, R. L., 1974. Cadmium toxicity and bioconcentration in large-mouth bass and bluegill. *Bull. environ. Contam. Toxicol.* **11**, 146-151.
- CHESTER, R. & STONER, J. H., 1974. The distribution of zinc, nickel, manganese, cadmium, copper, and iron in some surface waters from the world ocean. *Mar. Chem.* **2**, 17-32.
- CONNOR, P. M., 1972. Acute toxicity of heavy metals to some marine larvae. *Mar. Pollut. Bull.* **3**, 190-192.
- DANIELLI, J. F., 1944. The biological action of ions and the concentration of ions at surfaces. *J. exp. Biol.* **20**, 167-176.
- & DAVIES, J. T., 1951. Reactions at interfaces in relation to biological problems. *Adv. Enzymol.* **11**, 35-89.
- EATON, J. G., 1973. Chronic toxicity of a copper, cadmium and zinc mixture to the fathead minnow (*Pimephales promelas* RAFINESQUE). *Wat. Res.* **7**, 1723-1736.
- EISLER, R., 1971. Cadmium poisoning in *Fundulus heteroclitus* (Pisces: Cyprinodontidae) and other marine organisms. *J. Fish. Res. Bd Can.* **28**, 1225-1234.



- ZAROOGIAN, G. E. & HENNEKEY, R. J., 1972. Cadmium uptake by marine organisms. *J. Fish. Res. Bd Can.* **29**, 1367–1369.
- GARDINER, J., 1974. The chemistry of cadmium in natural water. 1. A study of cadmium complex formation using the cadmium specific-ion electrode. *Wat. Res.* **8**, 23–30.
- GOLDBERG, E. D., 1965: Minor elements in sea water. In: *Chemical Oceanography*. Ed. by J. P. RILEY & G. SKIRROW. Academic Press, New York, **1**, 163–196.
- HOLLIDAY, F. G. T., BLAXTER, J. H. S. & LASKER, R., 1964. Oxygen uptake of developing eggs and larvae of the herring (*Clupea harengus*). *J. mar. biol. Ass. U.K.* **44**, 711–723.
- JONES, P. G. W., HENRY, J. L. & FOLKARD, A. R., 1973. The distribution of selected trace metals in the water of the North Sea 1971–1973. *Coun. Meet. int. Coun. Explor. Sea* (= C.M.—I.C.E.S.) **5**.
- KOBAYASHI, J., 1971. Relation between the "itai-itai" disease and the pollution of river water by cadmium from a mine. *Adv. Wat. Pollut. Res.* **1**, 1–32.
- KREMLING, K., 1973. Voltametrische Messungen über die Verteilung von Zink, Cadmium, Blei und Kupfer in der Ostsee. *Kieler Meeresforsch.* **29**, 77–84.
- LEATHERLAND, T. M. & BURTON, J. D., 1974. The occurrence of some trace metals in coastal organisms with particular reference to the Solent region. *J. mar. biol. Ass. U.K.* **54**, 457–468.
- MULLIN, J. B. & RILEY, J. P., 1956. The occurrence of cadmium in seawater and in marine organisms and sediment. *J. mar. Res.* **15**, 103–122.
- NICKLESS, G., STENNER, R. & TERRILLE, N., 1972. Distribution of cadmium, lead and zinc in the Bristol Channel. *Mar. Pollut. Bull.* **3**, 188–190.
- O'HARA, J., 1973a. Cadmium uptake by fiddler crabs exposed to temperature and salinity stress. *J. Fish. Res. Bd Can.* **30**, 846–848.
- 1973b. The influence of temperature and salinity on the toxicity of cadmium to the fiddler crab, *Uca pugnator*. *Fish. Bull. U.S.* **71**, 149–153.
- OLSON, K. R. & HARREL, R. C., 1973. Effect of salinity on acute toxicity of mercury, copper and chromium for *Rangia cuneata* (Pelecypoda, Mactridae). *Contr. mar. Sci.* **17**, 9–13.
- PEDEN, J. D., CROTHERS, J. H., WATERFALL, C. E. & BEASLEY, J., 1973. Heavy metals in Somerset marine organisms. *Mar. Pollut. Bull.* **4**, 7–9.
- PICKERING, Q. H. & HENDERSON, C., 1966. The acute toxicity of some heavy metals to different species of warm water fishes. *Int. J. Air Wat. Pollut.* **10**, 453–463.
- & GAST, M., 1972. Acute and chronic toxicity of cadmium to the fathead minnow (*Pimephales promelas*). *J. Fish Res. Bd Can.* **29**, 1099–1106.
- PRESTON, A., 1973. Heavy metals in British waters. *Nature, Lond.* **242**, 95–97.
- JEFFRIES, D. F., DUTTON, J. W. R., HARWEY, B. B. & STEELE, E. K., 1972. British Isles coastal waters: The concentrations of selected heavy metals in sea water, suspended matter and biological indicators – a pilot survey. *Environ. Pollut.* **3**, 69–82.
- ROSENTHAL, H., 1966. Leaving the egg shell. A film on the hatching of herring larvae. *Coun. Meet. int. Coun. Explor. Sea* (C.M.—I.C.E.S.) **3**, 1–3.
- & MANN, H., 1973. Wirkung eines proteolytischen Enzyms (Maxatase-P) auf Embryonen des Herings (*Clupea harengus*) bei unterschiedlichen Temperaturen und Salzgehalten. *Arch. FischWiss.* **24**, 217–236.
- & SPERLING, K. R., 1974. Effects of cadmium on development and survival of herring eggs. In: *The early life history of fish*. Ed. by J. H. S. BLAXTER. Springer, Berlin, 383–396.
- SCHWEIGER, G., 1957. The toxic action of heavy metal salts on fish and organisms on which fish feed. *Arch. FischWiss.* **8**, 54–78.
- THURBERG, F. P., DAWSON, M. A. & LOLLIER, R. S., 1973. Effects of copper and cadmium on osmoregulation and oxygen consumption in two species of estuarine crabs. *Mar. Biol.* **23**, 171–175.
- VERNBERG, W. B. & O'HARA, J., 1972. Temperature-salinity stress and mercury uptake in the fiddler crab, *Uca pugnator*. *J. Fish. Res. Bd Can.* **29**, 1491–1494.
- WINDOW, H. L. & SMITH, R. G., 1972. Distribution of cadmium, cobalt, nickel and zinc in southeastern United States continental shelf waters. *Deep Sea Res.* **19**, 727–730.

- YAGER, C. M. & HARRY, H. W., 1966. Uptake of heavy metal ions by *Taphius glabratus*, a snail host of *Schistosoma mansoni*. *Expl. Parasit.* **19**, 174–182.
- ZIRINO, A. & YAMAMOTO, S., 1972. A pH-dependent model for the chemical speciation of copper, zinc, cadmium, and lead in sea water. *Limnol. Oceanogr.* **17**, 661–671.

First author's address: Dr. H. VON WESTERNHAGEN  
2 Hamburg 50  
Biologische Anstalt Helgoland  
Zentrale Hamburg  
Palmaille 9  
Federal Republic of Germany