

Gram positive bacilli, without leucocytes, isolating *C. septicum* and *Staph. epidermidis*. The lack of leucocytes and *C. septicum* isolation were also observed in the following samples.

Primary myonecrosis presents with three types of lesions [2, 3]: a) visceral anaerobic cellulitis (VAC), with sepsis, b) VAC, with spreading to adjacent muscles and c) metastatic myonecrosis, of extremely poor prognosis [4] and present in our case. It is a devastating infection, generally appearing in patients with digestive or hematologic malignancy, and caused by *C. septicum* in 60–80% of the cases [1]. *C. septicum*, an aerotolerant, motile and spore forming organism, is an infrequent pathogen (1.3–20% of clostridial infections) [2, 4–6] and is not a normal inhabitant of the gastrointestinal tract. It is 300-fold more virulent than *C. perfringens* [1] due to its toxins: α -toxin (lecithinase), two hemolysins, deoxyribonuclease, hyaluronidase and fibrinolysin [2–4]. The portal of entry for bacteremia is usually through disruptions of bowel mucosa (72–88% of the cases) [1–5] in colorectal malignancy or “enterocolitis” in hematologic malignancy, due to leukemic infiltration, lack of neutrophils or chemotherapy [1, 4, 5]. It later colonizes and proliferates in tissues with favorable local factors [1], such as the bilateral phlebitis in our patient, releasing toxins that produce necrosis and edema [2]. Primary myonecrosis by *C. septicum* presents with: extreme pain of the affected muscles, erythema, hemorrhagic phlictenas and soft tissue crepitation, although it may not be specific sign [1, 4, 5], with the finding of gas in X-rays [2, 4]. Progression is rapid and mortality is high (67–100%) [1, 2, 4, 5], some cases developing multiorgan failure and death in less than 24 h [1, 3, 5]. Dermoeidermic lesions shows Gram positive bacilli and a typical lack of neutrophils [1, 4], as in our case. This finding aided to differential diagnosis with streptococcal myonecrosis, pyomyositis and synergistic necrotizing cellulitis [4]. Incision of muscle may release large amount of gas and exudate, showing a pale, edematous muscle tissue that does not bleed or contract when cut [4, 6]. Poor prognosis factors are shock, hemorrhage, pneumonia, diminution of neutrophils and hemolysis [3]. Treatment should include antibiotics and surgery. The antibiotics of choice is penicillin [1, 4]; other suggested antibiotics are clindamycin, ceftriaxone, cefoxitin, erythromycin, vancomycin, tetracycline, chloramphenicol, metronidazole and rifampin [1, 4]. Surgery ranges from debridement to amputation, but is always associated to antibiotic therapy [2, 4]. Hyperbaric oxygen therapy can be adju-

vant [1] but its efficacy has not been demonstrated [1–4].

References

1. Stevens DL, Musher DM, Watson DA, Eddy H, Hamill RJ, Gyorkey F, Rosen H, Mader J (1990) Spontaneous, nontraumatic gangrene due to *Clostridium septicum*. *Rev Infect Dis* 12:286–296
 2. Sjolín SU, Hansen AK (1991) *Clostridium septicum* gas gangrene and an intestinal malignant lesion. A case report. *J Bone Joint Surg Am* 9: 11–12
 3. Bodey GP, Rodriguez S, Fainstein V, Elting LS (1991) Clostridial bacteremia in cancer patients. A 12-year experience. *Cancer* 67:1928–1942
 4. Kornbluth AA, Danzig JB, Berstein LH (1989) *Clostridium septicum* infection and associated malignancy. *Medicine* 68:30–37
 5. Ray D, Cohle SD, Lamb P (1992) Spontaneous clostridial myonecrosis. *J Forensic Sci, JFSCA* 37:1428–1432
 6. Swartz MN (1990) Myositis. In: Mandell GL, Douglas RG Jr, Bennett JE (eds) *Principles and practice of infectious diseases*, 3d edn. Churchill Livingstone, pp 812–817
- F. Hernandez Hazañas (✉)
C. Perez Torres · G. Dominguez Tornay
A. Herrera Saval
Intensive Care Unit,
Hospital Infanta Elena,
Huelva, Spain

Principles and Practice of Mechanical Ventilation. M. Tobin. New York: McGraw-Hill 1994. 1300 pp. (ISBN 0-070-64943-X).

Martin Tobin, also editor of this journal, just produced for McGraw-Hill, Inc., publishing company, the most recent and comprehensive book on mechanical ventilation.

This big volume – 1300 pages – with a forty page-index, thousands of references and hundreds of figures, tables and photos, provides the best presently available update on mechanical ventilation, and actually on acute respiratory failure. The book goes from history (a nicely illustrated chapter by G.L. Colice) to the most recent concepts, such as microprocessing, barotrauma, and modes (PAV, pressure-controlled ventilation). It gives the basics of mechanical ventilation (gas exchange, heart-lung interaction, neuromuscular function...) and also answers very practical questions such as the strategy of weaning, the management of the airway, and patient-ventilator fighting. All the modes currently provided by ventilators are explained and compared on the basis of published reports and the experience of renowned experts.

It is a common pitfall of such big textbooks that they are already obsolete the very first day they are printed, due to the large number of authors – all very busy and very much in demand – and the quasi-impossibility for any editor to collect all the manuscripts in time. The miracle in this case is that all the newer concepts in the field are presented here, with no lack of relevant information, from barotrauma (D. Dreyfuss) and permissive hypercapnia (D. Tuxen) to healthcare economics (B. Krieger) and the ethics of withholding life-sustaining therapy (J. Luce).

I am also grateful to Dr. Tobin to have invited such a large number of European authors, who indeed have contributed to improve our knowledge in this particular field. Due to the comprehensiveness, balance and broadness, and also due to the quality of authors, this book will certainly remain a landmark in the field of mechanical ventilation and respiratory intensive care. It has to be recommended to all trainees and practitioners in ICUs, and should be available in any personal, departmental and university library.

F. Lemaire, Créteil