

Metabolism of the Sulphur Amino-Acids in *Stegobium paniceum* L. and *Lasioderma serricorne* F.

That the intracellular yeast symbionts of the drug-store beetle, *Stegobium paniceum* L., supply the host with vitamins of the B-complex was reported by KOCH in 1933¹. Since then his findings in this Anobiid have been enlarged upon by his students and others²⁻⁵. Recent reviews of the literature describe the vital role of intracellular symbionts in many other host organisms^{6,7}. HENRY and BLOCK⁸ showed that the cockroach, *Blattella*

of utilizing inorganic sulphate in the synthesis of methionine and cystine. Larvae, artificially freed of their symbionts, are not able to synthesize these sulphur amino-acids.

Additional experiments are being carried out on *Stegobium* and *Lasioderma* with glucose-C¹⁴-U to determine the role of the yeast symbionts in the carbon synthesis of these and other amino acids, and will be reported later⁹.

Specific activity of S³⁵-amino-acids from larvae of *Stegobium paniceum* and *Lasioderma serricorne* after 3 weeks on diet labelled with Na₂S³⁵O₄

Amino acid	<i>Stegobium paniceum</i> ^a				<i>Lasioderma serricorne</i> ^a			
	Aposymbiotic		Normal		Aposymbiotic		Normal	
	Extract	Hydrolysate	Extract	Hydrolysate	Extract	Hydrolysate	Extract	Hydrolysate
Cystine	8	20	16	26	12	71	32	598
Methionine	0	10	4	18	16	62	23	3663

^a Counts per minute per μ mole of sulphur.

germanica, infected with its bacterial symbionts, is capable of utilizing inorganic sulphate for the synthesis of methionine and cystine, while the aposymbiotic roach cannot do so. The following investigations were carried out to determine the role of the symbionts of *S. paniceum* L. and of the closely related tobacco beetle, *Lasioderma serricorne* F., in the sulphur amino-acid metabolism of the hosts.

Both aposymbiotic and normally infected *Stegobium* and *Lasioderma* larvae, freshly hatched, were placed on a diet consisting of wheat grit and yeast extract (90:10 v/v), which was labelled with Na₂S³⁵O₄. After 3 weeks the larvae were homogenized and extracted, and the proteinaceous residues were hydrolysed. The extracts and hydrolysates were chromatographed in 2 directions on paper. The methionine and cystine spots were eluted from the paper and the radioactivity was measured in a liquid scintillation counter.

The results are listed in the Table. In *Stegobium* the specific activity for methionine and cystine was very low in the extracts and hydrolysates of both aposymbiotic and normally infected larvae. The specific activity of extracts from aposymbiotic and normal *Lasioderma* larvae was also low. However, in hydrolysates of normally infected larvae the specific activity of both methionine and cystine was found to be much greater than in aposymbiotic larvae.

These results show that larvae of *Lasioderma serricorne* F., infected with their normal yeast symbionts, are capable

Résumé. Les larves de *Lasioderma serricorne* F., infectées avec leurs symbiontes intracellulaires normales, ont pu utiliser du sulfate inorganique dans la synthèse de méthionine et cystine. Les larves libérées artificiellement de leurs symbiontes ne furent pas capables de synthétiser ces acides aminés sulfuriques. Les larves de *Stegobium paniceum* L. infectées normalement et les larves de *Stegobium paniceum* L. libérées des symbiontes n'utilisèrent pas de sulfate inorganique dans la synthèse de méthionine et cystine.

F. H. FOECKLER

Paul-Buchner-Institut für Symbioseforschung,
D-8031 Stockdorf bei München (West Germany),
30 June 1969

- 1 A. KOCH, Biol. Zbl. 53, 199 (1933).
- 2 K. E. GRAEBNER, Z. Morph. Ökol. Tiere 41, 471 (1954).
- 3 F. H. FOECKLER, Z. Morph. Ökol. Tiere 50, 119 (1961).
- 4 N. C. PANT and G. FRAENKEL, Biol. Bull 107, 420 (1954).
- 5 H. KÜHLWEIN and G. JURZITZA, Arch. Microbiol. 40, 247 (1961).
- 6 A. KOCH, in *Intracellular Symbiosis in Insects* (Ed. S. M. HENRY; Academic Press, New York 1967), vol. II, p. 1.
- 7 P. BUCHNER, *Endosymbiosis of Animals with Plant Microorganisms* (John Wiley & Sons, New York 1955).
- 8 S. M. HENRY and R. J. BLOCK, Contr. Boyce Thompson Inst. Pl. Res. 20, 317 (1960).
- 9 This work was supported by the Deutsche Forschungsgemeinschaft (German Research Foundation).

Biochemical Characteristics of Yeast Respiration-Deficient Mutants Differing in Buoyant Densities of Mitochondrial DNA

Mitochondrial DNA (M-DNA) has been assumed to represent the extrachromosomal genetic factor whose modification determines cytoplasmically inherited respiratory deficiency in yeast¹. A number of cytoplasmic (ρ^-) mutants have been prepared² which differed from each other and also from the original wildtype strain in buoyant densities of their respective M-DNA reflecting considerable differences in base composition³.

It will be shown in this paper that despite these differences in the M-DNA all the mutants display uniform biochemical deficiencies lacking cytochromes *a*, *a*₃, *b* and *c*₁, oligomycin-sensitivity of mitochondrial ATPase, and the ability to incorporate amino acids by isolated mitochondria. In addition they show a uniform pattern of

mitochondrial 'structural protein' in disc electrophoresis.

Material and methods. Haploid strains of cytoplasmic respiration-deficient mutants of *Saccharomyces cerevisiae*, listed in the Table, were kindly provided by Dr. H. Jakob (Centre de génétique moléculaire, Gif-sur-Yvette, France). Methods of culture, of isolation of mitochondria and of ATPase determination⁴ and assay of amino acids incorporation by isolated mitochondria⁵ have been described previously. Cytochrome spectra were measured in a SF 10 spectrophotometer equipped with an integrating sphere. Mitochondrial 'structural protein' was isolated⁷ and its electrophoresis in polyacrylamide gel performed⁸ according to published procedures.