

5. Jugdutt, B. I., G. M. Hutchins, B. H. Buckley, L. C. Becker: Infarct size reduction by prostacyclin after coronary occlusion in conscious dogs. *Clin. Res.* **27**, 177A (1979).
6. Ledingham, I. McA., T. I. McBride, J. R. Parratt, J. P. Vance: The effect of hypercapnia on myocardial blood flow and metabolism. *J. Physiol.* **210**, 87-105 (1970).
7. Marshall, R. J., J. R. Parratt: The antidysrhythmic aminosteroid ORG 6001 reduces the ST-segment elevation produced by coronary occlusion in the dog. *Brit. J. Pharmacol.* **61**, 315-317 (1977).
8. Marshall, R. J., J. R. Parratt, I. McA. Ledingham: Changes in blood flow and oxygen consumption in normal and ischaemic regions of the myocardium following acute coronary artery ligation. *Cardiovasc. Res.* **8**, 204-215 (1974).
9. Ogletree, M. L., A. M. Lefer, J. B. Smith, K. C. Nicolaou: Studies on the protective effect of prostacyclin in acute myocardial ischaemia. *Eur. J. Pharmacol.* **56**, 95-103 (1979).
10. Ribeiro, L. G. T., L. A. Reduto, T. A. Brandon, A. A. Taylor, B. G. Hopkins, R. R. Miller: Effects of prostacyclin on haemodynamics, regional myocardial blood flow, infarct size and mortality in experimental myocardial infarction. *Clin. Res.* **27**, 199A (1979).

Authors' address:

Dr. S. T. Coker, Dept. of Physiology and Pharmacology, Univ. of Strathclyde, George Street, Glasgow G1 1XW (Scotland, U.K.)

ERRATUM

T. Takala. Protein synthesis in the isolated perfused rat heart. Effects of mechanical work load, diastolic ventricular pressure and coronary pressure on amino acid incorporation and its transmural distribution into left ventricular protein. *Basic Res. Cardiol.* **76**, 44-61 (1981). The last three lines in the legend of Table 1 on page 50 should read:

oxygen consumption at time 60 min are shown. P (versus group II) *) < 0.05, **) < 0.02, ***) < 0.001, P (versus group VI), ²) < 0.05, ³) < 0.01.