

antimycin A and 0.5 mM KCN blocked mitochondrial swelling (curve 2). If, however, 0.1 mM PMS were added to the above system swelling occurred (curve 3). Addition of 1.8 mM amytal or 0.06 μ M piericidin A⁷ did not eliminate the swelling supported by the bypass created with PMS (curves 4 and 5, respectively). Addition of amytal or piericidin A in the absence of PMS to the system represented by curve 2 did not induce swelling (curves 6 and 7, respectively). In the absence of PMS, swelling could be supported by addition of 60 μ M each of ATP and Mg²⁺. (Figure 1). Although not shown here, induction of swelling by APT and M²⁺ was also observed for the systems represented by Figure 2. Curve 8 in Figure 2 represents the swelling mediated by all the 3 energy conserving sites in mitochondria in the presence of 1 mM arsenite without the addition of phosphate and thyroxine.

It is generally accepted that large amplitude swelling of mitochondria is an energy-linked process⁸. In our experimental system energy changes were restricted to the first site specifically by using a combination of antimycin A and cyanide and channelling the electrons from the substrates to an artificial acceptor, namely PMS¹. That sites II and III are not operative under these conditions was established. No swelling took place if PMS was not included (curve 2) in the system indicating that flux of electrons from substrates to PMS is necessary. Any possibility that PMS per se might induce swelling is ruled out by the fact that in the absence of swelling agents, phosphate plus thyroxine, there was no swelling. Addition of amytal and piericidin A did not eliminate the swelling (curves 4 and 5). Under conditions when substrate oxidation does not take place addition of ATP and Mg²⁺

induced swelling indicating the energy-requiring nature of the process (curves 2, 6, 7). These results indicate that site I energy conservation is on the substrate site of amytal and piericidin A sensitive sites associated with NADH-dehydrogenase. Our observations⁹ are consistent with those of GUTMAN et al.³.

Zusammenfassung. Es gelingt, die Schwellung der Rattenlebermitochondrien durch Energiekonservierung an jedem der 3 Phosphorylierungsorte herbeizuführen. Bei Gegenwart von Piericidin A oder Amytal wird die energieabhängige Schwellung nicht unterdrückt, insofern Energieänderungen auf den ersten Phosphorylierungsort beschränkt bleiben, was dafür spricht, dass die Wirkungsorte von Piericidin A und Amytal auf der Sauerstoffseite der Energiekonservierung am Phosphorylierungsort I liegen.

C. BHUVANESWARAN and K. DAKSHINAMURTI

Department of Biochemistry, Faculty of Medicine, University of Manitoba, 770 Bannatyne Avenue, Winnipeg 3 (Manitoba, Canada), 20 December 1971.

⁷ We are grateful to Professor K. FOLKERS for the kind gift of piericidin A.

⁸ A. L. LEHNINGER, *Physiol. Rev.* 42, 467 (1962).

⁹ This work was supported by a grant from the Medical Research Council of Canada.

Identification of Alexandrin

Centaurea alexandrina a biennial or often perennial pubescent herb, occurs only in Egypt¹ and is common in the Mediterranean coastal strip. SALEH and GHARBO² reported the isolation of a crystalline principle named 'alexandrin' from the leaves of *C. alexandrina* and stated that the substance (m.p. 261–262), is neither alkaloidal nor glycosidal in nature.

Extraction of the dried powdered leaves with ether², followed by concentration of the extract yielded a crystalline deposit, which after dissolving in alcohol and decolorising with activated charcoal afforded the substance corresponding to the so-called alexandrin. Thin-layer chromatography (adsorbent: silica gel G, solvents^{3,4}: carbon tetrachloride-ethanol-water 10:8:2, hexane-ethyl acetate-ethanol-water 4:6:5:5, benzene-methanol 8:2) of the substance revealed, upon spraying with *p*-anisaldehyde⁵, the presence of two components; the major one has the same R_f as β -sitosterol- β -D-glucoside, while the other is present in traces. Repeated recrystallization of the substance from methanol afforded the major constituent in pure form, m.p. 298–300° (undepressed), and proved to be β -sitosterol- β -D-glucoside. The identity was proven by chemical and physicochemical means (IR, NMR, MS, tetraacetate, hydrolysis).

The reportedly new constituent thymelol, also isolated by SALEH et al.⁶ from *Thymelea hirsuta*, has been proved by RIZK and RIMPLER⁷ to be a mixture of daphnoretin and β -sitosterol- β -D-glucoside.

Zusammenfassung. Die als Alexandrin beschriebene Substanz, die als Bestandteil von *Centaurea alexandrina* isoliert worden ist, wurde als β -sitosterol- β -D-glucosid identifiziert⁸.

A. M. RIZK, F. M. HAMMOUDA and S. I. ISMAIL

Pharmaceutical Sciences Laboratory, National Research Centre, Dokki, Cairo (Egypt), 8 November 1971.

¹ R. A. MUSCHLER, *A Manual Flora of Egypt* (Ed. Karl Trause, R. Friedländer & Sohn, Berlin 1912).

² M. R. I. SALEH and S. GHARBO, *J. pharm. Sci. U.A.R.*, 4, 17 (1963).

³ G. SCHÖPFLIN, Ph. D. Thesis, Freie Universität Berlin (1967).

⁴ F. M. HAMMOUDA, A. M. RIZK, H. GHALEB and M. M. ABDEL-GAWAD, *Planta med.*, in press (1972).

⁵ E. STAHL, *Dünnschicht-Chromatographie* (Springer-Verlag, Berlin 1967).

⁶ M. R. I. SALEH, D. Y. HADDAD and T. M. SARG, *J. pharm. Sci. U.A.R.*, 4, 49 (1963).

⁷ A. M. RIZK and H. RIMPLER, *Phytochemistry* 11, 473 (1972).

⁸ The authors are greatly indebted to Late Prof. Z. F. AHMED for suggesting the subject and to Prof. H. RIMPLER (Institut für Pharmakognosie der FU Berlin) and Prof. J. SCHMIDT-THOMÉ (Farbwerke Hoechst AG, Frankfurt) for their kind help in executing IR, NMR, MS and elementary analyses.