

phagocytic activity even after a long in vitro culturing period of 2 wk. Our procedure will therefore be useful for investigating research problems involving macrophages.

V. REFERENCES

- Nelson, D. S. Macrophages: progress and problems. *Clin. Exp. Immunol.* 45: 225–233; 1981.

- Harris, C. C.; Ih, C. H.; Stoner, G. D.; Trump, B. F.; Selkirk, J. K. Human pulmonary alveolar macrophages metabolize benzo(a) pyrene to proximate and ultimate mutagens. *Nature* 272: 633–634; 1978.

- Brain, J. D. The effects of increased particles on the number of alveolar macrophages. In Walton, W. H., ed. *Inhaled particles*. London: Unwin Brothers Ltd; 1971: 209–225.

¹ K. C. Biological, Inc., Lenexa, KS.

² Flow Laboratories, Inglewood, CA.

³ Abbott Laboratories, North Chicago, IL.

⁴ McGaw Laboratories, Irving, CA.

⁵ Sigma Chemical Co., St. Louis, CA.

⁶ American Scientific Products, Denver, CO.

⁷ Becton Dickinson and Company, Rutherford, NJ.

⁸ Moore Push Pin Co., Wyndmoor, PA.

⁹ Ethicon, Inc., Sommerville, NJ.

¹⁰ American Sterilizer, Erie, PA.

¹¹ Forma Scientific, Inc., Marietta, OH.

¹² Dremel Electric Co., Racine, WI.

¹³ Dupont Company, Instruments Products, Biomedical Division, Newton, CT.

¹⁴ Inbred colony obtained from Inhalation Toxicology Research Institute, Albuquerque, N.M.

ERRATUM

Preparation of Primary Monolayer Cultures of Adult Rat Hepatocytes

G. L. Engelmann and J. A. Fierer

Volume 7, No. 4, pages 169–173; 1982. (Procedure Number 41147). In III PROCEDURE, item A. 2. b. (page 170, column 2) should read:

b. To 225 ml add 140 mg CaCl₂ (Solution A).