

9. VANBREUSEGHEM, R. & BORGERS, G. 1951: A propos d'une souche de *Sabouraudites (Microsporium) gypseus* isolée au Congo belge. Ann. Soc. belge Med. trop. **31**: No. 3: 377—382.
10. EMMONS, C. W. 1934: Dermatophytes. Natural grouping based on the form of the spores and accessory organs. Arch. Derm. & Syph. **30** No. 3: 337—362.
11. SABOURAUD, R. 1910: Les Teignes. Masson & Cie. Paris.
12. KAPLAN, W., GEORG, L. K. & FOSNAUGH, C. J. 1956: Isolation of the dermatophyte, *Microsporium gypseum*, from a horse with ringworm. J. A. V. M. A. **129**, No. 8: 381—383.
13. FORSTER, R. A. 1956: A useful method for the addition of antibiotic to culture media. J. Med. Lab. Tech. **13**: 417—418.
14. CONANT, N. F., SMITH, D. T., BAKER, R. D., CALLAWAY, J. I. & MARTIN, D. S.: Manual of clinical mycology. 2nd. ed. **336**, 1954. W. B. Saunders Co. Philadelphia.

Addenda

Part I (in vol. XI)

References

37. MEWBORN, A. 1903: Report of a case of favus of scrotum coexisting with ringworm of the thigh giving identical Trichophyton-like cultures. J. Cut. Diseases. **21**, No. 244: 11—17.
38. BLANK, F. 1957: Favus of Mice. Canad. J. Microbiol. **3**, 885—896.

Errata

Part I (in Vol. XI)

Read magnifications as follows:

- Page 265. Plate II. Fig. 1. approx. $\frac{3}{4}$ nat. size. Fig. 2. approx. nat. size.
 Page 272. Plate III. Fig. 1. approx. $1\frac{1}{2}$ times nat. size. Fig. 2. approx. $1\frac{1}{2}$ times nat. size.
 Page 263. Table I. read across: J. H. male. 14 yr. Strain No. 2 scalp 1, face 1.

N.B. Dr. G. C. AINSWORTH (personal communication) has pointed out that MACLEOD & MUENDE (24) mention the binomial *T. quinckeanum* on p. 361.