

Addendum

"Studies on Liver Phosphatidyl Cholines: I. Effects of Fatty Liver Induction on Phosphatidyl Cholines From Liver Mitochondria and Microsomes", James E. Miller and W. E. Cornatzer, The Gudy and Bertha Ireland Research Laboratory, University of North Dakota School of Medicine, Grand Forks, North Dakota 58201, *Lipids* 4, 19-27. In Table V, F-1 control value and F-1 choline deficient value for Fatty Acid 22:6 were misstated. Table should read as follows:

TABLE V
Fatty Acid Composition of Mitochondrial Phosphatidyl Choline Fractions
From Livers of Female Rats Accumulating Fat

Fatty acid	Control ^a				Choline deficient ^a			
	F-1	F-2	F-3	F-4	F-1	F-2	F-3	F-4
14:0	0.1	0.3	0.2	0.1	0.2	0.4	0.2	0.4
15:0	0.1	0.1	t	0.1	0.1	0.2	0.2	0.2
16:0	17.4	12.9	13.3	16.8	20.5	14.9	16.9	29.2
16:1	t	1.8	0.8	1.0	0.5	1.4	1.2	2.6
17:0	0.6	0.5	0.4	0.5	0.2	0.5	0.9	1.9
18:0	28.3	23.6	33.1	31.5	17.6	19.9	22.8	16.3
18:1	1.4	10.8	4.1	19.7	4.2	9.3	5.8	26.2
18:2	3.5	7.5	9.8	24.7	1.8	5.4	5.7	19.5
18:3	0.3	0.3	0.3	1.5	0.7	0.6
20:3	t	t	0.2	1.6	0.9	4.8	1.2	1.0
20:4	3.8	24.0	38.1	3.4	2.9	12.4	43.4	1.9
20:5	1.0	10.3	1.4	10.9
22:5	6.2	10.2
22:6	43.7	1.8	t	47.5	8.6	1.2	0.3
Others	0.1	1.8	t	2.3	8.6	1.2	0.3

^aPer cent of total fatty acid by weight.