

Total Muon Capture Rates and Mean Neutrino Energy.

P. CHRISTILLIN, A. DELLAIORE and M. ROSA-CLOT

Scuola Normale Superiore - Pisa, Italia

Istituto Nazionale di Fisica Nucleare - Sezione di Pisa, Italia

(*Nuovo Cimento*, **34 A**, 272 (1976))

Some trivial misprints appear in sect. 3.

Equation (25) should read

$$\dots + \frac{1}{128} \frac{1}{x^3} \gamma^3 \left. \vphantom{\frac{1}{128}} \right\}.$$

Equation (36)

$$R = \frac{1}{Z} \int v^2 S \, dn_{\mathbf{z}} \, dv \, \delta(\mu - \varepsilon - v - E).$$

Equation (37)

$$R = \int R(E) \, dE = \frac{1}{Z} \int_a^b v^2 S_1 \frac{M^*}{v} \, dE + \frac{1}{Z} \int_b^c v^2 S_2 \frac{M^*}{v} \, dE.$$

Isotope Effect in Total Muon Capture Rates on Nuclei and Isotensor Effective Potential.

P. CHRISTILLIN and M. ROSA-CLOT

Scuola Normale Superiore - Pisa, Italia

Istituto Nazionale di Fisica Nucleare - Sezione di Pisa, Italia

A. DELLAIORE

Department of Theoretical Physics, Oxford University - Oxford, Great Britain

(*Nuovo Cimento*, **34 A**, 296 (1976))

Equation (7) should read

$$\dots + \frac{kx}{1 - kx} \frac{2Z}{A(A + x)}.$$