Bookreviews

JOHN L. HARPER

POPULATION BIOLOGY OF PLANTS

Academic Press, London, New York et San Francisco 1977, XXIV + 892 pp.

The last 20 years have been characterized by an increasing number of papers concerning the population ecology of plants. The end of this period is marked by the publication of the book under consideration. HARPER's book is the first one devoted to population biology of plants and is divided into six parts. The first part, with a single chapter "I. Experiments, analogies, models" gives a concise survey of the theory of population growth.

The second part, "Dispersal, dormancy and recruitment", contains four chapters, namely: 2. The seed rain, 3. Dormancy, 4. The seed bank, 5. The recruitment of seedling populations, describing crucial factors influencing the rise and maintenance of populations.

The third part (The effects of neighbors) is divided into six chapters: 6. The influence on density and mortality, 7. The influence of density on form and reproduction, 8. Mixtures of species: I. Space and proportions, 9. Mixtures of species: II. Changes with time, 10. The limiting resources of the environment, 11. Mechanisms of interaction between species. Questions dealing with the mutual influence of particular plants are discussed in this chapter. These relations are very important in the management of plant populations -- mainly in agriculture and forestry.

The fourth part is called "The effects of predators". The chapters contained in this part are the following: 12. Defoliation, 13. Seasonality, search and choice, 14. The role of grazing animals, 15. The predation of seeds and fruits, 16. Pathogens, 17. The role of predation in vegetation.

"The natural dynamics of plant populations" is the name of next part: plant dynamics is described in three chapters here: 18. Introduction: annuals and biennials, 19. Herbaceous perennials, 20. Woody plants.

The last part, "Plants, vegetation and evolution", has four chapters: 21. Reproduction and growth, 22. Reproduction -- life cycles and fertility schedules, 23. Community structure and diversity, 24. Natural selection and the population biology of plants.

It can be seen from the names of particular chapters that only one part of population biology i.e. population ecology, is the subject of this book.

Population biology of plants is a very young branch of science and is growing rapidly. For this reason, population ecology lacks a number of the laws which are so typical of long-established disciplines. Theory consists in hypotheses to be tested during continuing research. The author solves this situation by a great number of examples. The book is rich in very informative figures. He thus gives a clear picture of the present state of population biology of plants in the world. KROPÁČ's paper on the estimation of seeds in soil is the only paper by a Czechoslovak author which is given in HARPER's references, a fact which accurately reflects the situation in population biology of plants in Czechoslovakia.

The importance of HARPER's book also lies in another field. As the first book dealing with this topic, it compounds rather dispersed information from the literature and, in this way, forms an important starting point for the further development of the young discipline of population ecology.

FRANTIŠEK KRAHULEC

R. H. WHITTAKER and S. E. LEVIN (eds.)

NICHE: THEORY AND APPLICATION

Benchmark Papers in Ecology 3, — Dowden, Hutchinson and Ross, Inc., Stroudsburg, Pennsylvania 1975, XV + 448 pp., 137 Figs.

The term "ecological niche", shortly "niche", embraces various ecological concepts. It has become one of the most frequent terms in contemporary ecological literature. In European literature the term was seldom used except in the architectural sense. Many European ecologists still work without this concept, which suggests that a similar idea or at least parts of the niche concept are contained in other ecological terms. Czech ecologists, for example, have only recently begun to consider the appropriate spelling for Czech (niche, niše, niš or nika?). The question arose: Is the term "niche" actually needed?

The 3rd volume of the series "Benchmark Papers in Ecology" edited by R. H. WHITTAKER and S. A. LEVIN brings together a collection of the most important American papers dealing with or applying the niche concept. The editors themselves took part in the elaboration of the niche theory and its linkage to other ecological concepts. Covering all the major aspects of the problem their selection of papers appears highly representative. Besides an introduction, each part of the book is introduced by the "Editor's Comments on Papers" summarizing the substance and merits of the papers concerned. For the non-American reader, these editorial comments will prove a great help in understanding certain purely American approaches to the use of the concept.

In the introduction, the authors elucidate essential aspects of the niche concept. Its major features are expressed in formulations like "the place of each species in the functional whole", "the role of a species in the system", "the way a species population fits into a given community", "part of the whole set of relationships of the species to the environment", "the way the species relates to other species of the same community". Right at the beginning, the editors separate the term from the closely related, yet differently conceived term "habitat" ("range of environments in which a species lives"). Simple as it may sound, many European authors repeatedly misinterpret the niche concept and stress more its environmental content. For example, the expression "empty niche" often conforms to the idea of a shallow and empty hollow in the wall, derived from architecture.

The origin of the niche concept is described in the first part of the book where excerpts of works by GRINNELL, ELTON and GAUSE are reprinted. Very interesting reading indeed. One can see the gradual development of the concept starting with the rather vague approach of GRINNELL, which included habitat. Part II contains papers and excerpts from papers dealing with the competitive exclusion principle, sometimes called "GAUSE's principle". This idea is strongly linked to the niche concept, and supports the usefulness of this concept even within European terminological systems. In the editors' words "the brilliant young Russian scientist GAUSE set out to give the theory experimental teeth" — a note which suggests that fundamental biological approaches are not geographically determined.

Part III contains comments, papers and excerpts about niche axes, i.e. critical variables that affect the species composition of a community, such as resource availability and pressure of competitors, predators and parasites. Altogether nine authors and two editors show in both theoretical terms and zoological and botanical examples the background of spatiotemporal differences deciding species coexistence.

Part IV called "Niche and Habitat Dimensions" includes a variety of ideas and notions referring to the relationship between the niche and habitat concepts, gradient analysis, ordination of habitat relationships, niche overlap, niche breadth, etc.

Part V contains two excerpts and two papers discussing the evolutionary and distributional variation of niches. In this chapter the notion of niche is extended to the "geographical" and "paleobiological" aspects. The theory of island biogeography is presented as a good example of a niche-related notion. In Part VI four case studies utilizing the niche-concept are presented: two of them refer to birds, one to protozoa, and another to insects. Part VII is the concluding chapter containing three complete papers by (1) HUTCHINSON, (2) MACARTHUR and (3) WHITTAKER, LEVIN et ROOT. The last one published in 1973 in *American Naturalist* is the first summary of the subject.

All European ecologists will appreciate this volume, particularly because of the selection

of many papers which are scattered in less accessible journals, such as "Auk", containing the fundamental papers by GRINNELL. This assemblage of excerpts and full-length articles is also highly useful because of the existing separation between ecologists in zoological and botanical institutions — a frequent obstacle to ready access to literature referring to animals or plants, respectively. It is clear that the niche concept is both inspiring and important for development of future ecological theory. The book will satisfy European ecologists and help them in more adequate application of this concept. European ecologists will have to find out the historical roots of similar terms developed in the past and substitute the niche concept in the present-day literature. Some concepts, such as "life-form", "synusia", "ecological constitution", "behaviour", "cenotype", "cenological relationship", express similar features of life. The best interpretation of the niche concept in the Central-european terminological system will be "the totality of the relationships of the organism (population, species) and the environment", i. e. both its role in the community and the conditions provided by the environment. In this sense this concept will occupy a very useful place in ecological theory.

JAN JENÍK

R. G. WIEGERT (ed.)

ECOLOGICAL ENERGETICS

Benchmark Papers in Ecology 4. — Dowden, Hutchinson and Ross, Inc., Stroudsburg, Pennsylvania, 1976, XV + 456 pp. (Distributed by Halsted Press, a division of John Wiley and Sons, Inc.).

This is another book in the valuable set of "Benchmark Papers in Ecology". This time, it is a collection of altogether 38 classical papers in production ecology and ecological energetics, brought together by RICHARD G. WIEGERT, Professor of Zoology at the University of Georgia, a well-known specialist on the subject. The selected papers were published between 1942 and 1971, but most of them originate from the 'sixties, a period of rapid development of ecological energetics.

The papers are divided into four parts, and within each part, the editor's comments introduce groupings of papers on closely related subjects.

Part I. is entitled "History and Theory". The first paper is A. MACFADYEN'S (1948) essay on definitions of productivity in biological systems. It is followed by 4 papers on thermodynamics in ecological systems by R. G. WIEGERT (1968), D. SCOTT (1965), D. C. SPANNER (1963) and H. T. ODUM et R. C. PINKERTON (1955).

Part II., entitled "Levels of Storage and Efficiencies of Transfer", is introduced by the editor's comment on the distinction between the concepts of trophic levels and species food webs, both describing energy transfers within ecosystems. First in this part is the famous paper by LINDEMAN (1942) followed by G. E. HUTCHINSON'S prophetic postscript written after LINDEMAN'S premature death, at the age of 27 when the paper was in press. The further development of ecology has proved how right HUTCHINSON then was. The next four papers, by PAINE (1971), GOLLEY (1961), MARSHALL et MARPLES (1965) and SLOBODKIN et RICHMAN (1961) could be placed under the title of "ecological calorimetry": they define the energy contents in various biological materials. The next three papers deal with ecological efficiencies of energy transfers between trophic levels and populations. L. B. SLOBODKIN (1960) defines three different concepts of efficiency, and F. E. TURNER (1970) and D. G. KOZLOVSKY (1968) introduce further refinement into the definitions of efficiency. Most of these definitions are applicable to energy transfers between heterotroph populations rather than to the use of solar energy by the plant cover.

Part III. is called "Energy: Rates of Transfer". The five papers included mostly deal with the energetics of metabolism, each paper presenting a specific example: J. H. ZAR (1968) and E. A. LE FEBVRE (1964) — birds, two papers by J. PHILLIPSON (1962, 1963) and one by D. E. REICHLE (1968) deal with forest floor arthropods. The editor's comments draw some general conclusions from all five papers. Further on, nine papers deal with primary (autotrophic) pro-

duction, i. e., that of plants and their communities, and with its role in ecosystems. D. M. GATES (1965) briefly and clearly describes, in energy terms, the complex plant-environment relationships. G. M. WOODWELL and R. H. WHITTAKER (1968) give a survey of approaches to and methods of assessing primary production. E. P. ODUM's (1960) paper contains results of net production estimates in relation to secondary succession in abandoned fields. This paper is of particular interest to geobotanists. L. C. BLISS et E. B. HADLEY (1964) evaluate, in energy terms, the metabolism of plants, alpine lichens, in an extreme environment. Aquatic primary production, its role in lakes at various stages of their development, and the principal methods of its measurement, are evaluated in a clear and concise paper by CH. R. GOLDMAN (1968). A brief communication by R. J. BEYERS et H. T. ODUM (1959) points to the importance of exactly assessing the pH-dependent $\text{CO}_2 : \text{HCO}_3^- : \text{CO}_3^{2-}$ equilibria for any carbon budgets in water. The experimental approach is represented by the paper by N. R. KEVERN et R. C. BALL (1965), who measured algal production in an artificial stream in which the ecological conditions could be manipulated. A biological photoelectric system composed of a blue-green algal mat and a bacteria layer, is described in a brief note by N. E. ARMSTRONG et H. T. ODUM (1964). The next two papers, by S. MC NEIL et J. H. LAWTON (1970) and by F. B. GOLLEY (1968), evaluate the concept of secondary (heterotroph) production in ecosystems.

Part IV, introduced by the editor's relatively detailed remarks, is called "Energetics of Ecosystems". But the first five papers are energy-flow studies of single species populations of heterotrophs: three papers by E. H. BATTLE (two 1960, 1971) — *Saccharomyces cerevisiae*; S. RICHMAN (1958) — *Daphnia pulex*; A. E. SMALLEY (1960) and E. P. ODUM et A. E. SMALLEY (1959) — grasshoppers and molluscs in a salt marsh, depending, respectively, on live and dead *Spartina* as their main food sources. It is the last three papers in both Part IV and the whole book that treat energy flow at the community or ecosystem level: J. M. TEAL (1957) assesses community metabolism in a cold spring, an ecosystem depending to 77 % on allochthonous energy sources, and only to 23 % on its own primary production. The total energy input is, however, used quite efficiently (0.27 %), which is explained by the energy-saving effect of the rather constant and favourable spring environment. J. M. TEAL (1962) summarizes the energy flow in a coastal salt marsh dominated by *Spartina*. The producers (mainly *Spartina*) are also the most important consumers in the ecosystem, by respiring 77 % of their own gross production. Yet, the net efficiency of the primary production is quite high (1.4 % of total incident radiation) so that large amounts of energy are eventually stored in the *Spartina* detritus which is used as the most important food source by heterotrophs both in the marsh and in the adjacent estuary (about 45 % of the marsh production is exported there). M. D. ENGELMANN's (1968) paper on the role of soil arthropods in the energetics of an old-field community points to the inadequacy of a simplified trophic level concept when tracing the energy flow in a particular ecosystem.

In Part V, entitled "Future Directions", the editor outlines his own ideas about the usefulness of studies in ecological energetics and about the way of conducting them. The future of ecological energetics is seen in the predictive modelling of energy flow at the population, community and ecosystem levels. The models should be tested, of course, against experimental evidence. The minimum information required for the construction of a predictive model is stated quite clearly. Any degree of complexity or realistic detail can then be incorporated. Yet, although energy is the most easily defined ecological "currency", one should bear in mind that energy flow, however important it is, represents just one aspect of ecosystem functioning. Progress in ecological theory depends on a meaningful combination of more aspects.

At the end of the book, the reader will find a list of references to papers quoted in the editor's comments, the author citation index and a subject index. The whole book provides an instructive picture of the development of ecological energetics, demonstrated predominantly on the basis of the American literature. Ecologists from other countries have, indeed, worked much less in the field but the inclusion of a few more references to their papers would have broadened the readers' horizons. The energetics of primary production process in plants and plant stands appears relatively little represented. The development from plant physiological laboratory studies to those of the efficiency of interception and utilization of incident solar radiation by stands of both crops and wild-growing plants, could have been traced in a few carefully selected papers. This is, however, perhaps the only critical remark that can be made by a botanist on R. G. WISGERT's selection of Benchmark papers on "Ecological Energetics", a most useful book, in all respects, for any ecologist.

D. L. JAMESON (ed.)

EVOLUTIONARY GENETICS

Benchmark Papers in Genetics 8. — Dowden, Hutchinson and Ross, Inc. 1977, 332 pp., 28 Tab., 13 Figs.

This volume represents a selection of the basic literature from the time when the genetic bases of the evolutionary process was being identified, developed, and established. The volume starts with DARWIN and WALLACE and traces development to the early 1930's when FISHER, HALDANE and WRIGHT established integrated and complementary theories on the basis of which work has since proceeded. This volume is not meant to cover all the history of the early days of genetics. Large areas of genetic investigation are omitted, such as cytological, biochemical, mutational, physiological and microbiological studies, etc. Some of these questions were discussed in previously published volumes of this series. Here, a part of genetics is described that can be clearly identified as "evolutionary genetics" because the main questions concern the evolutionary process. After the 1930's the various parts of genetics became more specialized and many diverse aspects were considered for their own value, for their application to human welfare and/or for their interest to other disciplines and only peripherally did they consider evolutionary topics.

The volume is divided into four main parts. Each of them has the editor's comments on published papers followed by reprints of important original articles by the authors involved. The discussion is centred on the papers selected and the references to these studies in other papers.

In the first part of the volume (The Beginnings) DARWIN, GALTON, BALDWIN and MENDEL are mentioned. The selection from this early period includes DARWIN's and WALLACE's (Paper 1) proposition of the theory of natural selection where inherited variation was emphasized as the component which contributed most to the evolutionary process. GALTON's work (Paper 2) was an attempt to demonstrate his theory of inheritance. BALDWIN's paper (Paper 3) introduced the concept of inheritance of the ability to respond to changing conditions, thus linking developmental processes and inheritance to evolutionary studies. Finally, MENDEL's paper (Paper 4) provided rules for the particulate basis of the hereditary process.

The second part (Conflicts and Resolutions) covers the work of CASTLE, YULE, HARDY, WEINBERG, PEARSON and EAST. CASTLE (Paper 5) compared GALTON's and MENDEL's theories and described the selection of a Mendelian character. YULE (Paper 6) attempted to explain the inheritance of quantitative characters by Mendelism. HARDY (Paper 7) and WEINBERG (Paper 8) identified equilibrium principles and PEARSON (Paper 9 and 10) resolved the two (biometric and Mendelian) approaches except for the universality of the principle of dominance. EAST (Paper 11) provided an experimental demonstration of multiple Mendelian factors in inheritance of a quantitative character.

In the third part (Hammer and Tongs) theoretical papers are included. NORTON et PUNNET (Paper 12) produced a table of the rate change frequency by selection. FISHER's paper (Paper 13) dealt with the dominance ratio and one of WRIGHT's early papers (paper 14) provided a way to measure inbreeding. Papers 15, 16 and 17 of HALDANE flesh out the initial theoretical skeleton. CHETVERIKOV's summary (paper 18 and 19) of the state of evolutionary studies in the middle of the 1920's serves as an introduction to the development of a number of synthetic studies which later integrated various fields and strengthened our understanding of the evolutionary process.

In the early 1930's HALDANE, FISHER and WRIGHT presented their views on the evolutionary process and made them available in a form more accessible than in their numerous separate contributions. In the fourth part (How Nature Works) their articles represent modern studies of how populations change in nature, particularly in the fields of plant and animal breeding, quantitative genetics, behavioural genetics, and speciations.

The study of any discipline assumes a mastery of the literature of the subject: in many branches of science, even one as new as genetics, the expansion of knowledge has been so rapid that there is little hope of learning of the development of all phases of the subject. Benchmark Volume Editors are knowledgeable in specific aspects of the literature of genetics and selected papers to demonstrate both knowledge and the atmosphere in which the knowledge was developed. There is no substitute for reading great papers. Here you can learn how questions are asked, how they are approached, and how difficult and essential it is to obtain definitive answers and clear writing. That is why Mr. JAMESON's compilation is not only worth buying but why it should be read thoroughly.

DAGMAR JIČÍNSKÁ

P. BANARESCU et N. BOŞCAIU

BIOGEOGRAPHIE

Fauna und Flora der Erde und ihre geschichtliche Entwicklung

VEB Gustav Fischer Verlag, Jena 1978, 392 pp., 48 Fig.

Ergebnisse mehrerer Fachbereiche zu synthetisieren, ist nicht leicht und auch keine dankbare Aufgabe. Im rezensierten Buch haben es jedoch beide Autoren, ein Zoologe und ein Botaniker, verstanden, die Ergebnisse der Zoogeographie und der Phytogeographie in Einklang zu bringen. Prof. Dr. PETRU BANARESCU vom Institut für Biologie in Bukarest brachte in das gemeinsame Werk eine Reihe origineller Erkenntnisse aus seinen ichthyologischen Forschungen. Dr. NICOLAE BOŞCAIU, Hauptbiologe bei der Kommission für Naturdenkmäler aus Cluj-Napoca, hat den Text vor allem durch Beispiele aus der Studie der Flora und der Vegetation Rumäniens bereichert.

Dieses Handbuch erschien zum erstenmal i. J. 1973 rumänisch und nach kaum fünf Jahren erscheint es in deutscher Sprache. Das allein ist schon ein deutlicher Beweis dafür, dass die Autoren ein gutes Werk, reich belegt mit konkreten Beispielen, vorgelegt haben. Vorzug dieser Arbeit ist, dass der Inhalt modern, mit der Orientierung auf die Evolutions- und die historische Biogeographie erfasst worden ist. Grosser Nachdruck wird hier auf Fragen der Artbildung, der Populationsgenetik und der natürlichen Selektion gelegt. Im Gegenteil dazu, schweift die traditionell in der Literatur vorgelegte Problematik niemals allzusehr in Details ab. Das rezensierte Handbuch schöpft nicht alle Probleme der Biogeographie aus. So haben die Autoren vollkommen die ökologische Biogeographie ausgelassen, was sie damit begründen, dass auf dieses Thema z. B. in der Zoogeographie eine Arbeit von HESSE und DAHL existiert; das allerdings ist nicht stichhältig, denn seit der Ausgabe dieser Arbeit ist bereits ein halbes Jahrhundert vergangen.

Ein umfassendes Kapitel ist der Artbildung, der Aufgabe der Population im Artbildungsprozess, der Problematik allopatrischer und sympatrischer Artbildung und den polyploiden Reihen besonders bei den Pflanzen gewidmet, wo Polyploidie viel häufiger ist als bei Lebewesen. Mit zahlreichen, gut gewählten Beispielen ist auch das Kapitel über die Zytogeographie belegt. Gründliche Aufmerksamkeit ist weiter dem Begriff Artenareal, den Methoden ihrer Bestimmung, der Grösse und inneren Struktur gewidmet. Beispiele disjunkter Areale sollten nur natürliche Verbreitung erfassen, denn künstliche könnten bei so mancher Art ein diskontinuierliches Areal bilden. Das Beispiel der Gemse auf S. 93 schliesst in der Tschechoslowakei neben den einzigen ursprünglichen Arealen im Gebirgszug Vysoké Tatry (Hohe Tatra) auch die künstliche Verbreitung im Gebirge Lužické hory (Lausitzer Gebirge) und Hrubý Jeseník (Hohes Gesenke) ein.

Einen sehr guten Überblick gewähren die Autoren über die Mechanismen der Flora- und Faunagenese, der faunistischen und der floristischen Elemente sowie der Inselflora und -fauna. Eingehend beachten sie alle bedeutenderen paleogeographischen Hypothesen und ihre Bestätigung, eventuell auch ihre Ablehnung vom Gesichtspunkt der Biogeographie. Vom Gesichtspunkt der Entwicklung unserer gegenwärtigen Flora und Fauna wird die grösste Aufmerksamkeit der ausführlichen Analyse der einzelnen Quartärperioden gewidmet.

Ein umfassendes Kapitel widmen die Autoren auch der Aufteilung der Kontinente in biogeographische Regionen. Im wesentlichen halten sie die konventionelle Gliederung mit kleineren Abweichungen ein; so z. B. lehnen sie die Anerkennung einer selbständigen zirkumatlantischen Region ab, erkennen bloss zirkumatlantische Elemente im Rahmen vor allem einer neotropischen Region an. Wenn es sich auch um ein relativ kleines Gebiet handelt, ist es doch nicht möglich, Flora und Fauna des Südzipfels von Südamerika und der anliegenden Inseln in die gleiche Region einzugliedern wie die Flora und Fauna der Amazon-Tropenurwälder, besonders deswegen, da sie als selbständig z. B. die Region von Madagaskar oder Neuseeland anerkennen.

Was die Abbildungen betrifft, so können wir das Handbuch als eine mit Landkartenmaterial mittelmässig ausgestattete Arbeit ansehen. Schade, dass von den 48 Abbildungen, zumeist Landkarten, dem ersten Autor nur 6, dem zweiten sogar keine zugeschrieben werden kann — die meisten der übrigen Abbildungen wurden von der zitierten Literatur, zu 50 % aus dem Handbuch von G. DE LATTIN „Grundriss der Zoogeographie“ übernommen.

Die Bedeutung der Biogeographie wächst gegenwärtig, und das nicht nur vom Gesichtspunkt der Theorie. Der wachsende Bedarf an neuen Pflanzen- und Tierrohstoffquellen einerseits und die Bedrohung des Genofondes unseres Planeten andererseits, erfordern immer tiefergehendere Kenntnisse über die Gesetzmässigkeit der Organismenverbreitung. Die Weltliteratur hat vorläufig einige gute Handbücher der Biogeographie, wie z. B. der Autoren P. DANSEREAU, H. ELHAI,

H. FREITAG und G. LEMÉE. Die Tatsache, dass die rumänische Biologie in dieser deutschen Übersetzung der biogeographischen Weltliteratur ein weiteres gutes Werk gegeben hat, kann man nur begrüßen und wünschen, dass dieses Beispiel bald auch anderswo Nachfolger findet.

BOHUMIL SLAVÍK

S. G. SHETLER and L. E. SKOG

A PROVISIONAL CHECKLIST OF THE SPECIES FOR FLORA NORTH AMERICA

(Revised)

Missouri Botanical Garden, St. Louis 1978, 19 + 199 p.

The Programm Flora North America (sic!) was broken off for financial reasons in 1973; however, in connection with the international programm "Man and Biosphere", it seems to be reviving now under the guidance of P. H. RAVEN from Missouri Botanical Garden. The edition of the publication reviewed here gives evidence of this revival even though it was prepared already before 1973. At that time S. G. SHETLER and L. E. SKOG gathered materials for this checklist and they submit them now as a basis for preparation of the work being planned and at the same time they present them to a broader botanical public.

The introductory chapter gives information on the history of the work and on the method of its elaboration. The proper checklist is arranged accordingly to the adapted Englerian system. The checklist includes the whole continental U.S.A. (incl. Alaska), Canada and Greenland. Taxonomic units used are family (233), genus (2 350) and species (16 274); the authors of plant names are given only for the names of species. No synonyms are mentioned. Short data refer to the life-duration of the plant (perennial or annual), the character of the occurrence (native or introduction) and some morphobiological characters (tree, shrub, parasite etc.); they are given very briefly by means of one- or two-letter signs. Further data refer to the distribution; the area studied is divided into 17 regions designated by letters or numerals; the distribution area of each species is described by listing the signs of single regions. The amplest genera of North American flora are *Carex* (435 species), *Astragalus* (328), *Eriogonum* (215), *Panicum* — incl. *Dichanthelium* (171), *Crataegus* (167), *Pentstemon* (141), *Quercus* (131), *Phacelia* (116), *Lupinus* (113), *Aster* (111), *Salix* (109), *Erigeron* (108) and *Juncus* (106).

The material gathered in this book certainly represents valuable information on the flora of the North American subcontinent (north of Mexico) and on its geographical distribution. The concept of taxonomic units (especially of genera) is rather broad (e.g. *Delphinium*, *Lycopodium*, *Rumex*, *Succisa*); a broad concept of family is kept also in circumscription of *Polypodiaceae*. In view of the preliminary character of the publication, it is hardly possible to blame the authors for some simplified approaches; however, in many cases the material presented in their publication seems to be only roughly gathered, not processed in any way. This is evident especially from the fact that the same species is given twice in the list, with its correct name and under a synonym, whether in different genera or in the same genus, and again in cases where the species content of a certain genus is incomprehensibly divided into two parts, one of them being retained in the original genus, the other transferred to its segregant.

Examples of the inclusion of the same species into two genera are for instance: *Cheilanthes parryi* — *Notholaena p.* (3); *Dryopteris hexagonoptera* — *Thelypteris h.* (3, 4); *Dryopteris thelypteris* — *Thelypteris palustris* (3, 4); *Fagopyrum esculentum* — *Polygonum fagopyrum* (13, 14); *Cerastium cerastioides* — *Provancheria c.* (18, 19); *Lychnis alpina* — *Viscaria a.* (19, 21); *Saponaria vaccaria* — *Vaccaria pyramidata* (20, 21); *Anemone occidentalis* — *Pulsatilla o.* (26, 28); *Nasturtium officinale* — *Rorippa nasturtium-aquaticum* (37); *Buglossoides arvensis* — *Lithospermum a.* (101, 103); *Hackelia californica* — *Lappulla c.* and *H. deflexa* — *L. d.* (102, 103); *Lycopersicum esculentum* — *Solanum lycopersicum* (109, 110); *Agropyron repens* — *Elytrigia r.* (158, 164); *A. boreale* — *Roegneria b.* and *A. violaceum* — *R. v.* (158, 172); *Andropogon annulatus* — *Dichanthium a.* and *A. sericeus* — *D. s.* (159, 163); *Avena hookeri* — *Helictotrichon h.* and *A. pubescens* — *H. p.* (160, 165) etc. Mentioning the same species under various names in one genus may be shown by the following examples: *Isoetes echinospora* — *I. setacea* (1); *Vaccaria* (three names for one species — 21); *Pyrola chlorantha* — *P. virens* (86); *Veronica catenata* — *V. comosa* (117) etc.

Data on the distribution usually differ in these cases. Unclear generic concepts may be found in the following couples of genera: *Anemone*—*Pulsatilla* (26, *Anemone patens* belongs to *Pulsatilla*); *Chamaesyce* — *Euphorbia* (64, several species of the latter genus belong to the former, e.g. especially *E. chamaesyce*); *Rhodiola* — *Sedum* (40, the type species of *Rhodiola* — *Rh. rosea* — is included in *Sedum*); *Toxicodendron* — *Rhus* (68); *Calystegia* — *Convolvulus* (98); *Dactylorhiza* — *Orchis* (183, 184) etc. An especially confused case of classification is that of *Gentiana* s. l. (91); segregants of *Gentianella* — *Comastoma* and *Gentianopsis* — are accepted here as separate genera, but the members of *Gentianella* proper are still included in *Gentiana*; at the same time several species of *Gentianopsis* are retained in *Gentiana*.

These often repeated defects are due to the fact that single taxa taken over from various sources (flora manuals) under various names were not compared and identified. A quick revision of the material by a few specialists would certainly have eliminated these defects to a great extent, if not entirely.

Further critical comments may be evoked by the taxonomically unsuitable concept of *Montia* (17) including a greater part of the species of *Claytonia*, and by the incorrect classification of *Radiola* to *Myrtaceae* (78). In several cases correct names could be used for some taxa (*Bilderdykia*, *Agrostis alba*, *Anthriscus scandicina*, *Filago germanica*, *Lappula echinata*, *Oenothera strigosa*, *Oxalis europaea* etc.); specific epithets of the type "*hookerianus*" could be corrected to the correct version "*hookeranus*".

Errors may be found among the data on the life-duration of plants (species of *Rhinanthus* are not perennials, just as *Veronica incana* and species of *Succisa* s. l. are not annuals) or in data on the native character of the occurrence (cf. *Cuscuta epithymum*). Some unification is necessary in abbreviations of authors' names (at species names), in which (as in names of species) relatively many printer's errors occur.

Despite the provisional character of this publication (mentioned directly in its title) and the defects mentioned in this review, the checklist by SHETLER and SKOG is an important source giving useful information on the flora of the whole of North America. It is therefore necessary to welcome it. The printing of the checklist itself by computer is not of the first quality. With regard to the provisional character of this work and to its basic importance a really revised edition of this publication should certainly be published soon, using better printing methods

JOSEF HOLUB

Erratum

In der Besprechung des Buches von E. OBERDORFER: „Süddeutsche Pflanzengesellschaften“, zweite Auflage, in Folia Geobotanica et Phytotaxonomica 14/4: 437—439, 1979 sind leider einige Fehler in die Tab. 1 auf S. 438 unterlaufen. Die richtigen Werte in den betreffenden Zeilen sind folgende:

	1957			1977—1978		
	Ord. *	Verb.	Ass.	Ord.	Verb.	Ass.
<i>Asplenietea rupestris</i>	2	2	7	2	5	16
<i>Phragmitetea</i>	1	3	15(+1?)	1	3	24(+10?)
<i>Montio-Cardaminetea</i>	1	2	9	1	2	8(+8?)
<i>Scheuchzerio-Caricetea fuscae</i>	3	4	20(+2?)	3	5	18(+7?)
<i>Nardo-Callunetea</i>	2	4	15	2	5	15
<i>Betulo-Adenostyletea</i>	2	3	4	1	3	7
Insgesamt: Klassen: 23	27	48	175(+4?)	38	70	283(+25?)