

consumption.<sup>3</sup> Others have shown a rapid decline of endothelial ATP especially in the presence of heparin and this depletion occurs within a minute.<sup>4</sup> These effects on intracellular oxygen consumption and ATP availability may account for its direct effect on the cardiovascular system especially on the myocardium. Furthermore, blockade of all potential pathways that may cause hypotension and bradycardia, excluding the direct route, have been unsuccessful in blunting the cardiovascular effect of protamine.<sup>5</sup> These findings strongly indicate a direct influence of this drug on the myocardium.

Finally, protamine is also known to cause both anaphylactoid and anaphylactic reactions through activation of the complement system. Rapid boluses are contributory and should not exceed 50 mg over 10 min.<sup>6</sup> It is possible that the bolus administration of 20 mg was enough to induce such reactions although other indications confirming this as a mechanism of action are not offered.

A.J. Olufolabi MB BS FRCA  
Durham, North Carolina

#### REFERENCES

- 1 Backman SB, Gilron I, Robbins R. Protamine-induced hypotension and bradycardia in a cardiac transplant patient. *Can J Anaesth* 1997; 44: 520–4.
- 2 Kosekova GP, Mitovska MI, Dancheva KI. Protamine inhibition of the oxidative phosphorylation in intact, cytochrome c-depleted and restored mitochondria. *Acta Biologica et Medica Germanica* 1975; 34: 539–47.
- 3 Wakefield TW, Ucros I, Kresawik TF, Hinshaw DB, Stanley JC. Decreased oxygen consumption as a toxic manifestation of protamine sulphate reversal of heparin anticoagulation. *J Vasc Surg* 1989; 9: 772–7.
- 4 Wakefield TW, Hinshaw DB, Burger JM, Burkel WE, Stanley JC. Protamine-induced reductions of endothelial cell ATP. *Surgery* 1989; 106: 378–85.
- 5 Fadali MA, Papacostas CA, Duke JJ, Ledbetter M, Osbakken M. Cardiovascular depressant effect of protamine sulphate: experimental study and clinical implications. *Thorax* 1976; 31: 320–3.
- 6 Horrow JC. Adverse reactions to protamine. *Int Anesthesiol Clin* 1985; 23: 133–44.

#### REPLY

*We assessed possible mechanisms to explain the sudden onset of bradycardia in a cardiac transplant patient following a period of hypotension subsequent to the administration of protamine. We concluded that the change in heart rate could not be explained by cardiac reinnervation, as the patient had undergone transplantation only eight weeks*

*previously. Rather, we suggested that the change was mediated by mechanisms intrinsic to the transplanted heart and extrinsic to the CNS. It was argued that withdrawal of adrenal sympathetic output secondary to stimulation of mechanoreceptors, while possible, was unlikely because of the precipitous change in heart rate. Other possibilities, which we could not discount, included cardiac ischaemia, activation of an intrinsic stretch-related mechanisms in the transplanted heart, and entrainment of the donor to recipient (innervated) atria. Dr. Olufolabi proposes an additional mechanism, which involves a direct effect of protamine on the myocardium of the transplanted heart. This possibility warrants consideration. We thank Dr. Olufolabi for his interest in our report, and for his insight into the possible mechanisms mediating the cardiovascular response produced by protamine in a cardiac transplant patient.*

S.B. Backman MDCM PhD FRCPC  
I. Gilron MD FRCPC  
R. Robbins MD FRCPC  
Montréal; Quebec

#### *Falsely reassuring pulse oximetry in the presence of severe hypoxia*

To the Editor:

Pulse oximetry is a common means of objectively assessing the adequacy of oxygenation in the neonatal intensive care unit. We report the case of a one-day-old neonate with pulse oximeter saturation readings of >95% despite severe hypoxaemia.

This neonate experienced respiratory failure due to perinatal pneumonia (post-mortem diagnosis), and required mechanical ventilation, fluid and inotropic support. Monitoring included ECG, direct arterial blood pressure measurement and pulse oximetry (Nellcor N25). Clinically, the neonate was cyanosed despite ventilation with 100% oxygen, and a discrepancy was noted between the ABG PO<sub>2</sub> and the pulse oximeter saturation readings, the pulse oximeter persistently reading 95–100% despite ABG PO<sub>2</sub> levels in the 2.1–4.63 kPa range (see Table on page 1324). Pre and post ductal saturations were the same. The probe was attached sequentially to all four limbs with the same readings. Pulse oximeter and ECG heart rates always correlated and the pleythsmograph trace was of good quality. A second probe and pulse oximeter was tried and agreed with the first. Both pulse oximeters were used simultaneously and agreed. Subsequently both machines were found to be faultless by the hospital Bio-engineers.

TABLE This table illustrates the values obtained from the six arterial blood gas samples taken during the resuscitation and the corresponding pulse oximeter reading at the same time.

| ABG                            | 1    | 2     | 3     | 4     | 5     | 6     |
|--------------------------------|------|-------|-------|-------|-------|-------|
| pH                             | 7.27 | 6.98  | 6.65  | 6.81  | 6.65  | 6.72  |
| PCO <sub>2</sub>               | 6.69 | 10.92 | 15.49 | 15.69 | 16.2  | 16.73 |
| PaO <sub>2</sub>               | 4.63 | 2.1   | 2.77  | 3.91  | 3.49  | 2.22  |
| BES                            | -4.3 | -16.5 | -27.5 | -21   | -27.6 | -23.1 |
| SATS %                         | 77   | 15.6  | 17.3  | 31    | 24.1  | 13.8  |
| SaO <sub>2</sub> %<br>Oximeter | 100  | 95    | 96    | 98    | 98    | 96    |

The causes of falsely elevated pulse oximetry include, ambient light, elevated carboxyhaemoglobin levels and partial probe detachment.<sup>1</sup> Oximeters overestimate the saturation when it falls below 75%, but the overestimation is minimal.<sup>2</sup> Lighting conditions and carboxyhaemoglobin levels were normal. Probe placement was checked and the patient was pharmacologically paralysed for ventilation.

Neither the manufacturer nor ourselves have satisfactorily explained this observation.

Connail McCrory MB FFARCSI  
Margaret Ryan MB FFARCSI  
Patrick Doherty MB FFARCSI  
Dublin, Ireland

#### REFERENCES

- 1 *Poets C, Southall D.* Non-invasive monitoring of oxygen in infants and children. Practical considerations and areas of concern. *Paediatrics* 1994; 93: 737-44.
- 2 *Grace RF.* Pulse oximetry. Gold standard or false sense of security? *Med J Aust* 1994; 160: 638-43.