Bruce Dick, University of Alberta, Edmonton, AB, Canada;

Saifee Rashid, University of Alberta;

Julie Zhang, University of Alberta;

A Ohinmaa, University of Alberta;

INTRODUCTION: Previous studies have shown that chronic pain has negative effects on a person's quality of life, level of functional disability, and mental health. Best medical practice standards for the management of chronic pain include a variety of treatment options. The biopsychosocial model of chronic pain management is increasingly being used as a part of treatment plan formulation during medical interventions. This study sought to investigate factors that are associated with decreased quality of life in a consecutive sample of patients from a general clinic population of a Multidisciplinary Pain Centre.

METHODS: Local IRB approval was obtained for this study. A consecutive sample of 100 patients with chronic pain were included in this study. In addition to demographic information, general health information, and subjective symptom ratings, patients were asked to provide information via the 15D, a validated measure of quality of life (QOL). Participants' medications were rated within three categories according to strength ('none,' 'mild,' 'moderate,' 'strong'). Chi-square tests and multiple linear regression analyses using a backward selection method were carried out to ascertain associations between background factors and predictors of QOL.

RESULTS: This sample was primarily Female (65%). The chronicity of pain varied such that nearly half of patients reported chronic pain for more than 5 years. The majority of patients (54%) reported more than one co-morbid medical condition. The two primary factors that were found to be significantly predictive of QOL in this sample were whether or not patients were being treated with 'strong' medications and whether they had co-morbid medical problems. Other significant predictors of QOL were gender (Females reported higher QOL), age (poorer QOL in older patients), anxiety, depression, and number of pain sites.

SUMMARY: Medical factors including strength of medication and additional co-morbid medical problems over-and-above chronic pain were the best predictors of QOL in this sample. Females, younger patients, patients with lower depression and anxiety scores, and patients with fewer pain sites all reported higher quality of life in this sample. Data collection is ongoing in order to investigate additional factors that may be associated with QOL in our general clinic population.