

- 5 Bellomo R, Chapman M, Finfer S, Hickling K, Myburgh J. Low-dose dopamine in patients with early renal dysfunction: a placebo-controlled randomised trial. Australian and New Zealand Intensive Care Society (ANZICS) Clinical Trials Group. *Lancet* 2000; 356: 2139–43.

Accidental intravascular injection of levobupivacaine and lidocaine during the transarterial approach to the axillary brachial plexus

To the Editor:

Racemic bupivacaine, when injected intravascularly, is associated with serious cardiac complications¹ such as ventricular fibrillation resistant to successful resuscitation. No such serious outcome was reported hitherto with levobupivacaine. The present case reports the accidental intravascular injection of a combination of levobupivacaine and lidocaine used for axillary brachial plexus blockade.

A 35-yr-old patient was admitted to the hospital for orthopedic surgery. Following premedication with midazolam (4 mg) and placement of all monitors, he received an axillary plexus block by the transarterial approach using a mixture of lidocaine 2% (20 mL) and levobupivacaine 0.75% (20 mL). Twenty-five millilitres of the local anesthetic mixture was deposited posterior and 15 mL anterior to the axillary artery. Briefly after deposition and without showing signs of light central nervous system (CNS) toxicity (lightheadedness, tinnitus, metallic taste), the patient exhibited three interrupted episodes of tonic-clonic seizures, each lasting for about three seconds and eventually resulting in unconsciousness. The patient's heart rate (HR) showed a sinus tachycardia of 160 beats·min⁻¹, the blood pressure (BP) increased to 180/120 mmHg and the SPO₂ decreased to 40% within one minute. For seizure control, the patient was given 5 mg of midazolam and 100 mg of propofol *iv*. Following mask ventilation with 100% oxygen, he was intubated and brought to the postanesthesia care unit. His vital signs stabilized within 30 min (BP 103/74, HR 84, SPO₂ 97%) without further pharmacologic support and he was extubated. Two hours after extubation, he was alert and oriented and discharged to home. The patient did not show signs of sensory and motor blockade.

A recent case report of an accidental intravascular injection following epidural anesthesia with 19 mL of levobupivacaine 0.75% resulted in only minor CNS side effects (drowsiness, slurred speech) and, most

importantly, no cardiac sequelae.² Plasma levels were not taken until 14 min after epidural injection, still they revealed a toxic range of levobupivacaine that, most likely, was substantially higher immediately after its intravascular administration. Though the severity of side effects remains unknown had racemic bupivacaine been administered in this patient, previous reports hint at a more serious outcome after racemic bupivacaine 0.75%.¹ Similarly, no cardiac effects other than a sinus tachycardia occurred in this young and otherwise healthy patient.

It appears that levobupivacaine is a safer drug than racemic bupivacaine, still vigilance and the laws of regional anesthesia (slow and intermittent injections, frequent aspirations) need to be practiced to take advantage of levobupivacaine's wider margin of safety.

Humayon Khan MD
Peter G. Atanassoff MD
New Haven, Connecticut

References

- 1 Albright GA. Cardiac arrest following regional anesthesia with etidocaine or bupivacaine. *Anesthesiology* 1979; 51: 285–7.
- 2 Kopacz DJ, Allen HW. Accidental intravenous levobupivacaine. *Anesth Analg* 1999; 89: 1027–9.

Skin analgesia with lidocaine tape prior to epidural blockade

To the Editor:

Lidocaine tape (Penles®, Japan Lederle, Tokyo, Japan) is a self-adhesive poultice for local anesthesia containing 18 mg of lidocaine at a concentration of 60% in a 30.5 × 50.0 mm polyester film. It has been reported that lidocaine tape provides effective skin analgesia, minimizing the pain caused by percutaneous cannulation, stellate ganglion block, and propofol injection.^{1–3} Eutectic mixture of local analgesics has also been used to alleviate cutaneous pain in children and adults.⁴ However, for optimal analgesic effects, the correct amount of the drug must be applied and the skin should be properly dressed for an effective absorption.⁴ In this regard, lidocaine tape has advantages and is frequently used because of easier application. However, although the tape is clinically useful, elevation of the pain threshold as measured by depth of needle insertion and the optimal duration of application remain unclear.




FIGURE Depth at which pain is elicited by insertion of a needle at the lidocaine site (■) and the placebo site (□). Lidocaine tape increased depth compared to control. ($*P < 0.05$). No differences were observed at control sites.

The present study was undertaken to examine the effects of lidocaine tape on the sensitivities to pain at the sites of epidural block. Thirteen outpatients receiving a series of epidural blocks at the pain clinic of the university hospital were subjected to a double-blind study. Lidocaine or placebo tapes were applied to the block site on the back (paramedian approach) or to a symmetrical position opposite to the block site (control site) for 30, 60, 90, or 120 min. Then, to test the degree of analgesia, a 23-gauge needle was inserted vertically into the skin. The depth at which patients first felt pain was measured. Depth was 3.38 ± 2.60 mm at 30 min, 9.31 ± 6.25 mm at 60 min, 11.54 ± 5.72 mm at 90 min, and 11.69 ± 6.18 mm at 120 min after application of the lidocaine tape (Figure). Depth at which pain was elicited were significantly greater in the lidocaine tape sites than with placebo ($P < 0.05$). No differences were observed at control sites. Our results indicate that the application of lidocaine tape for 90 min is the shortest optimal period to minimize the pain on needle insertion in patients receiving epidural blocks and that it can provide analgesia to a mean depth of 12 mm.

Kazuya Sobue MD
Takako Tsuda MD
Miho Yumoto MD
Takashi Nakagawa MD
Motomi Nakano MD
Hirotada Katsuya MD
Nagoya, Japan

References

- 1 Jin T, Iida T, Isozaki K. Continuous pain-relieving effect of a local anesthetic tape (Japanese). *J Clin Anesth* 1996; 20: 519–22.
- 2 Inada T, Uesugi F, Kawachi S, Inada K. Lidocaine tape relieves pain due to needle insertion during stellate ganglion block. *Can J Anaesth* 1997; 44: 259–62.
- 3 Yokota S, Komatsu T, Komura Y, et al. Pretreatment with topical 60% lidocaine tape reduces pain on injection of propofol. *Anesth Analg* 1997; 85: 672–4.
- 4 Bjerring P, Arendt-Nielsen L. Depth and duration of skin analgesia to needle insertion after topical application of EMLA cream. *Br J Anaesth* 1990; 64: 173–7.