

tored, particularly during the first month of combined therapy. Also, the risk of gastrointestinal haemorrhage with any NSAID must always be remembered.

References

- Hoffbrand BI, Kininmonth DA. Potentiation of anticoagulants. *British Medical Journal* 2: 838-839, 1967
- Koch-Weser J. Hemorrhagic reactions and drug interactions in 500 warfarin-treated patients. *Clinical Pharmacology and Therapeutics* 14: 139, 1973
- Landefeld CS, Goldman L. Major bleeding in outpatients treated with warfarin: incidence and prediction by factors known at the start of outpatient therapy. *American Journal of Medicine* 87: 144-152, 1989
- Landefeld CS, Rosenblatt MW. Bleeding in outpatients treated with warfarin: relation to the prothrombin time and important remediable lesions. *American Journal of Medicine* 87: 153-159, 1989
- Launbjerg J, Egeblad H, Heaf J, Nielsen NH, Fugleholm AM, et al. Bleeding complications to oral anticoagulant therapy: multivariate analysis of 1010 treatment years in 551 outpatients. *Journal of Internal Medicine* 229: 351-355, 1991
- O'Callaghan JW, Thompson RN, Russell AS. Combining NSAIDs with anticoagulants: yes and no. *Canadian Medical Association Journal* 131: 857-859, 1984
- Pullar T, Capell HA. Interaction between oral anti-coagulant drugs and non-steroidal anti-inflammatory agents: a review. *Scottish Medical Journal* 28: 42-47, 1983
- Self TH, Evans WE, Ferguson T. Drug enhancement of warfarin activity. *Lancet* 2: 557, 1975
- Vessel ES, Passananti GT, Johnson AO. Failure of indomethacin and warfarin to interact in normal human volunteers. *Journal of Clinical Pharmacology* 15: 486-495, 1975
- Yip ASB, Chow WH, Tai YT, Cheung KL. Adverse effect of topical methylsalicylate ointment on warfarin anticoagulation: an unrecognized potential hazard. *Postgraduate Medical Journal* 66: 367-369, 1990

Correspondence and reprints: Dr *Thomas Y.K. Chan*, Department of Clinical Pharmacology, The Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, N.T., Hong Kong.

Erratum

Vol. 9, No. 6, 1993, page 447: In figure 1, the middle boxes should be replaced as follows:

<p>No clinical signs of liver disease * and ALT < 3-5xULN (ALP < 2xULN)</p>

<p>Clinical signs of liver disease and/or ALT > 3-5xULN (ALP > 2xULN)</p>
