

116. Jones RM, Cashman JN, Foster JMG, et al. Comparisons of infusions of morphine and lysine acetyl salicylate for the relief of pain following thoracic surgery. *Br J Anaesth* 1984; 57: 259-63
117. Mattila MK, Ahlstrom-Bengs E, Pekkola P. Intravenous indomethacin in prevention of post-operative pain. *BMJ* 1993; 287: 1026
118. Campbell WI, Kendrick R, Patterson C. Intravenous diclofenac sodium. *Anaesthesia* 1990; 45: 763-6
119. Hovorka J, Kallela H, Kortilla K. Effect of intravenous diclofenac on pain and recovery profile after day-case laparoscopy. *Eur J Anaesthiol* 1993; 10: 105-8
120. Laitinen J, Nuutinen L, Kiiskial E-L, et al. Comparison of intravenous diclofenac, indomethacin and oxycodone as post-operative analgesics in patients undergoing knee surgery. *Eur J Anaesthiol* 1992; 9: 29-34
121. Murphy DF, Medley C. Preoperative indomethacin for pain relief after thoracotomy: comparison with postoperative indomethacin. *Br J Anaesth* 1993; 70: 298-300
122. Kavanagh BP, Katz J, Sandler AN, et al. Multimodal analgesia before thoracic surgery does not reduce postoperative pain. *Br J Anaesth* 1994; 73: 184-9

Correspondence and reprints: Dr *Jeremy Cashman*, Consultant Anaesthetist, Department of Anaesthesia, St George's Hospital, Blackshaw Road, London SW17 0QT, England.

Errata

Vol. 47, No. 5, p. 827: In the summary of Dosage and Administration, the second sentence should read, 'In adults with normal renal function, a loading dose of 400mg (approximately 6 mg/kg) every 12 hours for the first 3 doses, followed by 400mg once daily, is recommended.'

[*Brogden RN, Peters DH. Teicoplanin: a reappraisal of its antimicrobial activity, pharmacokinetic properties and therapeutic efficacy. Drugs* 1994 May; 47 (5): 823-54]

Vol. 48, No. 3, p. 389: The first sentence in the Clinical Tolerability summary should read, 'PLS administration has been well tolerated in multicentre clinical trials (including approximately 3200 patients), in which it did . . . '.

p. 397: The first sentence in section 4 Clinical Tolerability should read, 'PLS has been well tolerated in large multicentre clinical trials to date (including a total of approximately 3200 infants), in which it did . . . '.

p. 397: In section 3.2 Prophylactic Use, the sentence beginning on line 7 of the second column should read, 'The findings of a recent meta-analysis of 3 clinical studies, including the trial discussed above^[49] and evaluating a total of 671 neonates, indicate, however, that PLS prophylaxis does significantly improve clinical outcome versus rescue treatment.^[50]'

[*Wiseman LR, Bryson HM. Porcine-derived lung surfactant: a review of the therapeutic efficacy and clinical tolerability of a natural surfactant preparation (Curosurf®) in neonatal respiratory distress syndrome. Drugs* 1994 Sept; 48 (3): 386-403]