Boundary Hamiltonian theory for gapped topological phases on an open surface

Yuting Hu,^{a,1} Zhu-Xi Luo,^b Ren Pankovich,^b Yidun Wan^{a,c,1} and Yong-Shi Wu^{d,a,c,b,1}

^a*Department of Physics and Center for Field Theory and Particle Physics, Fudan University, Shanghai 200433, China*

^b*Department of Physics and Astronomy, University of Utah, Salt Lake City, Utah, 84112, U.S.A.*

^c*Collaborative Innovation Center of Advanced Microstructures, Nanjing 210093, China*

^d*State Key Laboratory of Surface Physics, Fudan University, Shanghai 200433, China*

E-mail: yuting.phys@gmail.com, mercuryzx.luo@gmail.com,
ren.pankovich@utah.edu, ydwan@fudan.edu.cn, yswu@fudan.edu.cn

ABSTRACT: In this paper we propose a Hamiltonian approach to gapped topological phases on open surfaces. Our setting is an extension of the Levin-Wen model to a 2d graph on an open surface, whose boundary is part of the graph. We systematically construct a series of boundary Hamiltonians such that each of them, when combined with the usual Levin-Wen bulk Hamiltonian, gives rise to a gapped energy spectrum which is topologically protected. It is shown that the corresponding wave functions are robust under changes of the underlying graph that maintain the spatial topology of the system. We derive explicit ground-state wavefunctions of the system on a disk as well as on a cylinder. For boundary quasiparticle excitations, we are able to construct their creation, annihilation, measuring and hopping operators etc. Given a bulk string-net theory, our approach provides a classification scheme of possible types of gapped boundary conditions by Frobenius algebras (modulo Morita equivalence) of the bulk fusion category; the boundary quasiparticles are characterized by bimodules of the pertinent Frobenius algebras. Our approach also offers a set of concrete tools for computations. We illustrate our approach by a few examples.

KEYWORDS: Topological States of Matter, Quantum Groups, Topological Field Theories

ARXIV EPRINT: [1706.03329](https://arxiv.org/abs/1706.03329)

¹Corresponding author.

Contents

1	Introduction and summary	1
2	Review of the Levin-Wen model	4
2.1	Topological feature	8
3	The boundary Hamiltonian	9
3.1	Frobenius algebra as input data	9
3.2	Boundary Hamiltonian	11
3.3	Graphical presentation	12
4	Ground state on a disk	14
5	Topological feature of ground states	17
6	Ground states on a cylinder	18
7	Boundary excitations	19
7.1	Measure quasiparticles	20
7.2	Creation operators	21
7.3	Hopping operators	21
7.4	Fluxons	21
8	Examples	22
8.1	Charge boundary	22
8.2	LW $\mathbb{Z}_2$ model	22
8.3	LW Fibonacci model	24
8.4	LW Ising model	25
9	Equivalent boundary conditions	25
10	Relation to the Kitaev-Kong formulation	26
A	Some proofs and details	29
A.1	Topological feature of the graph	29
A.2	Frobenius algebra	30
A.3	Derivation of equation (3.19)	32
A.4	Property of boundary terms	32
A.5	Ground state on a disk	34
A.6	Proof of equation (6.7)	36
A.7	$W_M \Phi\rangle$ is an eigenvector of $\Pi_M = 1$	36
A.8	Bimodules with multiplicity	37

1 Introduction and summary

Two important characteristic properties of a 2d matter phase with an intrinsic topological order [1], crucial for topological quantum computation, are a finite set of topologically protected, degenerate ground states [2–4] and the corresponding anyon excitations obeying braid statistics [5]. While the former furnishes robust quantum memories [6], the latter delineates a logical Hilbert space that supports topological quantum computation via unitary braiding of anyonic quasiparticles [7–10]. It is also hopeful to realize or simulate Abelian [11, 12] and non-Abelian anyons [13, 14]. On a closed spatial 2-surface, the genus number and the fusion rules between anyon excitations determine the ground state degeneracy (GSD) [1, 4, 10, 15, 16]. In particular, on a torus, the GSD equals the number of anyon species. Nevertheless, realizing closed-surface material with topological order is difficult in experiments; it is much more natural to make finite open systems. Yet, any boundary massless modes that often appear must be gapped to have a well defined GSD. The gapping conditions of Abelian topological orders have recently been understood in terms of the concept of Lagrangian subsets [17–24], and subsequently the GSD of these Abelian phases on open surfaces with multiple boundaries were computed [23, 25], based on the idea of anyon transport across boundaries. Experiments detecting and utilizing the topological degeneracy with gapped boundaries were proposed in [26, 27]. The gapping conditions of non-Abelian topological orders have recently been tackled by the mechanism of anyon condensation [28] and equivalently by solving certain algebraic equations [29]. Gapped boundaries of topological orders can also be classified by Frobenius algebras [30], using the mechanism of anyon condensation.

Nevertheless, the rich studies and classifications of gapped boundaries of topological orders are, unfortunately, not practical enough, because they are based on abstract mathematical theories rather than explicit Hamiltonian models consisting of both bulk and boundary terms. This causes consequent issues. For example, ref. [28] offers a closed-form formula of computing the GSD of a topological order on a n -hole surface, in terms of the condensed anyons at the holes and their fusion rules. This formula, though mathematically complete and beautiful, cannot tell us how to realize the boundary conditions on a given Hamiltonian model of the topological order. Although it is known that a gapped boundary of a topological order corresponds to certain anyon condensation, it is not clear how bulk anyons interact with the boundary excitations. Hence, the proposal of using the generalized Laughlin-Tao-Wu charge-pumping argument [2, 31] to braid anyons and realize topological quantum computation in ref. [28] would be impossible unless there is a concrete Hamiltonian model in which pumping a boundary excitation from one boundary to another can be studied in terms of physical operators.

More broadly, a dynamical theory in which one is not able to specify boundary conditions is not a complete dynamical theory. Therefore, Hamiltonian models of topological orders such as the Levin-Wen model [32], the Kitaev Model [7], and the twisted quantum double model [33] as a generalization of the Kitaev model are not complete dynamical models on open surfaces because they do not include boundary terms.

In this paper, we develop a new systematic approach — in the framework of discrete models on a 2d graph with boundary — to gapped topological phases on open surfaces. Our approach leads to the following main results.

1. A large family of exactly solvable boundary Hamiltonian H_{bdry} , using the data of a Frobenius algebra in the unitary fusion category (UFC) (or the fusion algebra associated with it), that defines the bulk Levin-Wen model.
2. A gapped energy spectrum for each boundary Hamiltonian together with its bulk Hamiltonian, that is protected by the spatial topology, as in the original Levin-Wen model.
3. The topological ground-state wavefunctions on a disk and a cylinder in closed form, in terms of solely the input data (fundamental degrees of freedom of the model, whose generalization to surfaces with more boundaries is straightforward).
4. Explicit creation, measuring, and hopping operators for boundary excitations.
5. A classification scheme of possible types of gapped boundary conditions by Morita-equivalent Frobenius algebras of the bulk fusion algebra, as well as that of boundary quasiparticles by bimodules of the pertinent Frobenius algebra.

These results enable one to study explicitly various topological properties, such as the GSDs, the topological entanglement entropy, excitations, bulk boundary correspondence, and so on, of topological orders on open surfaces.

Our approach is based on three physical ansatzes as follows.

1. *Locality*: the boundary Hamiltonian is a local one.
2. *Asymptotics*: the boundary (interacting) theory is the asymptotics of the bulk (interacting) theory; hence, the boundary degrees of freedom should come from the bulk degrees of freedom.
3. *Topological invariance*: the ground-state Hilbert space is invariant under topology-preserving mutations of the graph. This leads to Frobenius algebras characterizing boundary interactions, as emergent structures of the bulk degrees of freedom.

The Locality ansatz is natural. The Asymptotics ansatz and the emergence of Frobenius algebras deserve more justification, as we now elaborate.

It is well-known that for a continuum Chern-Simons gauge theory on a bounded spatial region in a plane, besides the usual Chern-Simons bulk term, the action contains an additional term on the boundary [34], to ensure the gauge invariance of the total action. The physical interpretation for adding a boundary action term is holography, i.e. a holographic correspondence between the bulk and the boundary, respected in a topological phase in two dimensions. Motivated by the desired holography in the discrete framework, the central idea of ours is to construct explicitly and systematically an appropriate boundary Hamiltonian, to be added to the Levin-Wen bulk Hamiltonian that was originally designed for a

Figure 1. Trivalent graph with boundary.

closed surface. Thus, our new approach features the new Hamiltonian below, defined on a trivalent graph as in figure 1:

$$H = H_{\text{LW}} + H_{\text{bdry}}, \quad (1.1)$$

where H_{LW} is the usual Levin-Wen model Hamiltonian in the bulk of the graph, while H_{bdry} is our boundary Hamiltonian defined along the boundary of the graph. As in the original Levin-Wen model, the bulk degrees of freedom, i.e., the string types labeled on the bulk edges, are simple objects of a unitary fusion category (UFC), such as irreducible representations of a finite group or a quantum group. The boundary Hamiltonian in our new approach however are associated with a Frobenius algebra that is a composite object in the bulk UFC, so that in the ground states the boundary degrees of freedom are restricted to be the string types appearing only in the pertinent Frobenius algebra. This consideration of Frobenius algebra is again motivated by the demand of exact solvability and by holography. First, when we restrict the boundary degrees of freedom to live in a subset of those in bulk described by the input UFC, and require the commutativity between the boundary operators and the bulk operators, we find that the boundary degrees of freedom would have to form a Frobenius algebra — an associative algebra with non-degenerate inner product. Second, restricted to bulk ground states only, the boundary theory of a topological order may be thought of as a $(1+1)$ -d topological quantum field theory (TQFT) associated with the bulk $(2+1)$ -d TQFT. It has been shown that $(1+1)$ -d TQFTs are in one-to-one correspondence with commutative Frobenius algebras [35]. Third, in the case with finite groups, the Levin-Wen model is dual to the Kitaev quantum double model. Shor et al. have shown that the boundary degrees of freedom in the Kitaev model defined by a finite group G live in a subgroup of G [37], which in the dual Levin-Wen model corresponds to a Frobenius algebra in the UFC of the representations of G .

Compared to the existing approaches to gapped topological phases on open surfaces, our boundary Hamiltonian approach has the following advantages. First, the usual classification of topological boundaries by Lagrangian subsets and/or by anyon condensation, as mentioned above, is based on the data of the anyon species of excitations in topological orders, which is the output of the dynamics of the system. In contrast, our classification is characterized by the input data of the dynamical model. This is more in line with the spirit

of the usual Hamiltonian dynamics. Secondly, for a given Levin-Wen bulk Hamiltonian, the boundary terms of our model may not be unique, and the gapped energy spectrum of the whole bounded system depends on the choice of the boundary term accordingly. Hence, it is obvious that the boundary Hamiltonians can be used to characterize and classify the boundary conditions that give rise to gapped topological boundaries. Moreover, by solving the total Hamiltonian (bulk plus boundary terms) we can obtain the explicit wave functions of the ground and excited states, all in the form of tensor network states. This will provide us a very detailed, dynamic understanding of the stationary topological states of the whole bounded system, especially of what happens on and near the boundary. For example, our model enables us to study the boundary excitations explicitly. Also anyon condensation may be understood at more microscopic scales. These studies will be reported later separately.

We organize our paper as follows. In section 2 we review the core ingredients of the Levin-Wen model and set up notations. We extract some basic topological features of the Levin-Wen model on a closed surface, that are often overlooked except the original paper, and promote them into basic principles crucial to our development for open surfaces. Section 3 is devoted to the construction of the boundary Hamiltonian in our approach, with a graphical presentation developed along the way. Then the ground state on a disk of the total (bulk plus boundary) Hamiltonian is studied in section 4, while those on a cylinder are studied in section 6. The topological nature of the ground states is examined in section 5. We devote the next section 7 to the construction of relevant operators for studying boundary excitations, such as their creation, annihilation, measuring, and hopping operators etc. Section 8 presents a few explicit examples. In section 9 we discuss in what sense two apparently different boundary conditions are equivalent, resulting in a classification scheme of gapped boundary conditions by Morita equivalence of Frobenius algebras.

It is worth noting that there has been a few studies of the boundary Hamiltonians in the Kitaev model [36–39], as well as a study of the Levin-Wen model [17] with boundary in the language of module categories. While our approach is systematic and easier to access by the condensed matter community, we shall discuss in section 10 the relation between our approach and the one taken by Kitaev and Kong [17]. The appendices collect certain details and proofs.

Two of us also reported in a companion paper [40] a similar approach of constructing the boundary Hamiltonian of the twisted quantum double model. The present paper is a much expanded version, with many details and more results, of a short paper of three of us [41] that briefly reports the main ideas and some results.

2 Review of the Levin-Wen model

Let us start with a brief review of the Levin-Wen model [32] on a closed surface without boundary, setting up our notations and conventions. This model is a lattice Hamiltonian model defined by a set of input data — a unitary fusion category (UFC) $\mathcal{C}$ — that specifies the Hilbert space of the model. In this work, we will use the tensor description of $\mathcal{C}$ in terms of $6j$ -symbols. (For a physicist-readable introduction see, e.g., ref. [42].)

Figure 2. A configuration of string types on a directed trivalent graph. The configuration (b) and hence the associated Hilbert space is regarded the same as (a), with some of the directions of some edges reversed and the corresponding labels j conjugated j^* .

The model is defined on a trivalent graph embedded in a closed, oriented surface. The Hilbert space is spanned by the degrees of freedom on edges (see figure 2), which are the simple objects (called string types) in $\mathcal{C}$ and are labeled by j that runs over a finite set of integers $L = \{j = 0, 1, \dots, N\}$. (For example, string types could be irreducible representations of a finite or quantum group.) The Hilbert space is spanned by all configurations of the labels on edges. Each label j has a “conjugate” j^* , which is also an integer and satisfies $j^{**} = j$. If we reverse the direction of one edge labeled by j and replace the label by j^* , we require the state remains the same. See figure 2. There is a unique “trivial” label $j = 0$ with $0^* = 0$.

As simple objects in a UFC $\mathcal{C}$, the string types are subject to fusion rules. A *fusion rule* on L is a function $N : L \times L \times L \rightarrow \mathbb{N}$ such that for $a, b, c, d \in L$,

$$N_{0a}^b = N_{a0}^b = \delta_{ab}, \quad (2.1)$$

$$N_{ab}^0 = \delta_{ab^*}, \quad (2.2)$$

$$\sum_{x \in L} N_{ab}^x N_{xc}^d = \sum_{x \in L} N_{ax}^d N_{cb}^x. \quad (2.3)$$

A fusion rule is multiplicity-free if $N_{ab}^c \in \{0, 1\}$ for all $a, b, c \in L$. We restrict to the multiplicity-free case throughout this paper unless stated otherwise. We define $\delta_{abc} := N_{ab}^{c^*}$, with the symmetric properties: $\delta_{abc} = \delta_{bca}$ and $\delta_{abc} = \delta_{c^*b^*a^*}$. A triple (a, b, c) is admissible if $\delta_{abc} = 1$.

Given a fusion rule on L , a *quantum dimension* is a map $d : L \rightarrow \mathbb{R}$ such that $d_{a^*} = d_a$ and

$$\sum_{c \in L} d_c \delta_{abc^*} = d_a d_b. \quad (2.4)$$

In particular, $d_0 = 1$. Let $\alpha_j = \text{sgn}(d_j)$, which take values of ± 1 for each label j and satisfy

$$\alpha_i \alpha_j \alpha_k = 1, \quad \text{if } \delta_{ijk} = 1. \quad (2.5)$$

Given fusion rules and quantum dimensions on L , we may define $6j$ -symbols, often denoted as G . A *tetrahedral symmetric unitary* $6j$ -symbol is a map $G : L^6 \rightarrow \mathbb{C}$ satisfying

these conditions:

$$\begin{aligned}
 G_{kln}^{ijm} &= G_{nk^*l^*}^{mij} = G_{ijn^*}^{klm^*} = \alpha_m \alpha_n \overline{G_{l^*k^*n}^{j^*i^*m^*}}, \\
 \sum_n d_n G_{kp^*n}^{mlq} G_{mns^*}^{jip} G_{lkr^*}^{js^*n} &= G_{q^*kr^*}^{jip} G_{mls^*}^{riq^*}, \\
 \sum_n d_n G_{kp^*n}^{mlq} G_{pk^*n}^{l^*m^*i^*} &= \frac{\delta_{iq}}{d_i} \delta_{mlq} \delta_{k^*ip},
 \end{aligned} \tag{2.6}$$

where the second equation above is the *pentagon identity*.

The input data of the Levin-Wen model is such a set: $\{d_j, \delta_{ijk}, G_{klm}^{ijm}\}$ that can be derived from the representation theory of a finite group or a quantum group, and more generally, such a set of data is from a UFC. For instance, we may take the labels j to be the irreducible representations of a finite group H . The trivial label 0 is the trivial representation. The fusion rules indicate whether the tensor product $j_1 \otimes j_2 \otimes j_3$ contains the trivial representation or not. Each number α_j is the Frobenius-Schur indicator telling whether the representation j is real, complex, or pseudoreal. The relation $d_j = \alpha_j \dim(j)$ holds, where $\dim(j)$ is the dimension of the corresponding representation space. The 6j-symbols G_{kln}^{ijm} are identified with the (symmetrized) Racah 6j-symbols of the group H . In this example, the Levin-Wen model can be mapped to the Kitaev quantum double model.

One important property of the 6j-symbols is that

$$G_{kln}^{ijm} = G_{kln}^{ijm} \delta_{ijm} \delta_{klm} \delta_{lin} \delta_{nk^*j^*}. \tag{2.7}$$

To prove this, one can rewrite the orthogonality condition as

$$\sum_n \left(v_n v_q G_{kp^*n}^{mlq} \right) \overline{\left(v_n v_i G_{kp^*n}^{mlq} \right)} = \delta_{iq} \delta_{mli} \delta_{k^*ip}, \tag{2.8}$$

where $\bar{c}$ stands for the complex conjugate of a complex number c . When $q = i$, the above equality implies that $G_{kp^*n}^{mli}$ must vanish unless $\delta_{mlq} \delta_{k^*ip} = 1$. By the tetrahedral symmetry, one arrives at eq. (2.7), where $v_j = \sqrt{d_j}$ is a choice of a square root of the quantum dimension. The number v_j is either real or pure imaginary, depending on the $\alpha_j = \text{sgn}(d_j)$, and is determined up to a sign that can be fixed as follows. From the conditions (2.6), we have $(G_{0kj}^{ijk} v_j v_k)^2 = \delta_{ijk}$, and it is possible to fix the sign of v_j such that $G_{0kj}^{ijk} v_j v_k = \delta_{ijk}$. We define

$$v_j := \frac{1}{G_{00j}^{j^*j0}}. \tag{2.9}$$

In particular, $v_0 = 1$ because $d_0 = 1$ (from eq. (2.4)), and thus $G_{000}^{000} = 1$ from eq. (2.6). Indeed, we can verify $v_j^2 = d_j$ directly from the orthogonality condition in eq. (2.6) together with $d_0 = 1$. The definition in eq. (2.9) also implies that

$$G_{0kj}^{ijk} v_j v_k = \delta_{ijk}, \tag{2.10}$$

which is due to the pentagon identity $d_0 G_{0kj}^{ijk} G_{00j}^{j^*j0} G_{00k}^{k^*k0} = G_{0kj}^{ijk} G_{k^*0j}^{j^*i^*k^*}$ and the orthogonality $d_j G_{0kj}^{ijk} G_{k^*0j}^{j^*i^*k^*} = \frac{1}{d_k} \delta_{ijk}$.

There are two types of local operators, Q_v defined at a vertex v and B_p^s (indexed by the string types $s = 0, 1, \dots, N$) on a plaquette p . On a trivalent graph, a Q_v acts on the labels j_1, j_2 , and j_3 on three edges incident at the vertex v , such that

$$Q_v \left| \begin{array}{c} j_3 \quad j_2 \\ \diagdown \quad \diagup \\ j_1 \end{array} \right\rangle = \delta_{j_1 j_2 j_3} \left| \begin{array}{c} j_3 \quad j_2 \\ \diagdown \quad \diagup \\ j_1 \end{array} \right\rangle, \quad (2.11)$$

where the tensor $\delta_{j_1 j_2 j_3}$ determines whether the triple $\{j_1, j_2, j_3\}$ is admissible or not at v . Since $\delta_{j_1 j_2 j_3}$ is invariant under permutations of its three indices, the ordering in the triple $\{j_1, j_2, j_3\}$ is irrelevant.

An operator B_p^s acts on the boundary edges of the plaquette p , and has the following matrix elements on a triangular plaquette.

$$\begin{aligned} & \left\langle \begin{array}{c} j_5 \quad j_3 \quad j_6 \\ \diagdown \quad \diagup \quad \diagup \\ j_1 \quad j_2 \\ \diagup \quad \diagdown \\ j_4 \end{array} \right| B_p^s \left| \begin{array}{c} j'_5 \quad j'_3 \quad j'_6 \\ \diagdown \quad \diagup \quad \diagup \\ j'_1 \quad j'_2 \\ \diagup \quad \diagdown \\ j'_4 \end{array} \right\rangle \\ &= v_{j_1} v_{j_2} v_{j_3} v_{j'_1} v_{j'_2} v_{j'_3} G_{s j'_3 j'_1}^{j_5 j_1^* j_3} G_{s j'_1 j'_2}^{j_4 j_2^* j_1} G_{s j'_2 j'_3}^{j_6 j_3^* j_2}. \end{aligned} \quad (2.12)$$

The action of B_p^s on a quadrangle, a pentagon, or a hexagon, etc, is similar. Note that the matrix of B_p^s is non-diagonal only on the labels of the boundary edges (such as j_1, j_2 , and j_3 in the above graph). The operators B_p^s have the properties

$$B_p^{s\dagger} = B_p^{s*}, \quad (2.13)$$

$$B_p^r B_p^s = \sum_t \delta_{rst^*} B_p^t, \quad (2.14)$$

which can be verified by using conditions (2.6).

The operators defined above comprise the Hamiltonian of the model:

$$H = - \sum_v Q_v - \sum_p B_p, \quad B_p = \frac{1}{D} \sum_s d_s B_p^s \quad (2.15)$$

where the sum run over vertices v and plaquettes p of the trivalent graph, and $D = \sum_j d_j^2$ is the total quantum dimension.

It turns out that all Q_v and B_p involved are mutually-commuting projectors. Namely, (1) $[Q_v, Q_{v'}] = 0 = [B_p, B_{p'}]$, $[Q_v, B_p] = 0$; (2) $Q_v Q_v = Q_v$ and $B_p B_p = B_p$. Thus the Hamiltonian is exactly soluble. The elementary energy eigenstates are given by common eigenvectors of all these projectors. The ground states satisfies the constraints $Q_v = B_p = 1$ for all v and p , while the excited states violate these constraints for certain plaquettes and/or vertices.

In cases where the input data $\{d, \delta, G\}$ arises from the representations of groups or quantum groups, we have $\delta_{rst^*} = \delta_{srt^*}$. Then the operators B_p^s also meet the following commutation relation,

$$[B_{p_1}^r, B_{p_2}^s] = 0, \quad (2.16)$$

which can be verified by the conditions (2.6), with p_1 and p_2 two neighboring plaquettes, and by eq. (2.14), together with $\delta_{rst^*} = \delta_{srt^*}$ when $p_1 = p_2$.

2.1 Topological feature

We briefly review the topological nature of the bulk ground state(s). Any two given trivalent graphs $\Gamma^{(1)}$ and $\Gamma^{(2)}$ can be mutated into each other by a composition of three elementary moves, call Pachner moves. There are unitary linear maps [42] associated to these moves:

$$\begin{aligned}
 T_{2 \rightarrow 2} \left| \begin{array}{c} \nearrow j_1 \quad \nearrow j_4 \\ \searrow j_2 \quad \searrow j_3 \\ \text{---} j_5 \end{array} \right\rangle &= \sum_{j'_5} G_{j_3 j_4 j'_5}^{j_1 j_2 j_5} v_{j_5} v_{j'_5} \left| \begin{array}{c} \nearrow j_1 \quad \nearrow j_4 \\ \searrow j_2 \quad \searrow j_3 \\ \text{---} j'_5 \end{array} \right\rangle \\
 T_{1 \rightarrow 3} \left| \begin{array}{c} \nearrow j_1 \quad \nearrow j_3 \\ \searrow j_2 \end{array} \right\rangle &= \sum_{j_4, j_5, j_6} \frac{v_{j_5} v_{j_6} v_{j_4}}{\sqrt{D}} G_{j_5 j_6 j_4}^{j_1 j_2 j_3} \left| \begin{array}{c} \nearrow j_1 \quad \nearrow j_6 \quad \nearrow j_3 \\ \searrow j_4 \quad \searrow j_5 \end{array} \right\rangle \\
 T_{3 \rightarrow 1} \left| \begin{array}{c} \nearrow j_1 \quad \nearrow j_6 \quad \nearrow j_3 \\ \searrow j_4 \quad \searrow j_5 \\ \text{---} j_2 \end{array} \right\rangle &= \frac{v_{j_5} v_{j_4} v_{j_6}}{\sqrt{D}} G_{j_5 j_4 j_6}^{j_1 j_3 j_2^*} \left| \begin{array}{c} \nearrow j_1 \quad \nearrow j_3 \\ \searrow j_2 \end{array} \right\rangle
 \end{aligned} \tag{2.17}$$

This provides a linear transformation $\mathcal{H}^{(1)} \rightarrow \mathcal{H}^{(2)}$ between the two Hilbert spaces. Instead of $T_{3 \rightarrow 1}$, one may use another move: squeezing a “bubble”,

$$T'_{2 \rightarrow 0} \left| \begin{array}{c} \uparrow k' \\ \text{---} i \quad \text{---} j \\ \downarrow k \end{array} \right\rangle = \delta_{kk'} \frac{v_i v_j}{v_k \sqrt{D}} \left| \begin{array}{c} \uparrow k \end{array} \right\rangle, \tag{2.18}$$

which is a composition of $T_{3 \rightarrow 1}$ and $T_{2 \rightarrow 2}$.

The topological nature of the bulk ground states is described as follows. The ground-state Hilbert space is invariant under arbitrary composition of $T_{2 \rightarrow 2}$, $T_{1 \rightarrow 3}$, and $T_{3 \rightarrow 1}$. Moreover, given initial and final trivalent graphs there are multiple ways to compose $T_{2 \rightarrow 2}$, $T_{1 \rightarrow 3}$, $T_{3 \rightarrow 1}$, but different ways result in one and the same transformation on the ground-state Hilbert space.

The topological feature is similar in the Hilbert space $\mathcal{H}^{Q=1}$ of simultaneous eigenvectors of $Q = 1$ at all vertices. All states in $\mathcal{H}^{Q=1}$ are invariant under any transformation composed of $T_{2 \rightarrow 2}$. An example that yields the Pentagon identity (A.1) is presented in appendix A.1.

Transformations involving $T_{2 \rightarrow 2}$, $T_{1 \rightarrow 3}$ and $T_{3 \rightarrow 1}$ that moves from one given graph to another are generally not unique; however, there are subsets of all possible sequences of $T_{1 \rightarrow 3}$ and $T_{3 \rightarrow 1}$ moves, each of which leads to a unique transformation. Such a subset is obtained by specifying (using a ‘ $\times$ ’) in the initial graph the plaquettes to be killed by $T_{3 \rightarrow 1}$ moves, and in the final graph the plaquettes (using a ‘ $\cdot$ ’) to be created by $T_{1 \rightarrow 3}$ moves. In the rest of the paper, we will follow the above convention. That is, we suppress the choice of sequence of Pachner moves behind any transformation, and denote the transformation by $\mathcal{T}$ hereafter. In this convention, the B_p operator is compactly expressed by

$$\mathcal{T} \left(\left| \begin{array}{c} \text{---} \text{---} \text{---} \\ \times \end{array} \right\rangle \rightarrow \begin{array}{c} \text{---} \text{---} \text{---} \\ \cdot \end{array} \right\rangle \right). \tag{2.19}$$

3 The boundary Hamiltonian

In this section, we explicitly construct the boundary Hamiltonian for the Levin-Wen model on open surfaces. To understand our systematic construction, we will first introduce certain necessary mathematical structures.

3.1 Frobenius algebra as input data

The topological feature in the bulk is a consequence of the conditions (2.6) on the $6j$ -symbols. We expect the gapped boundaries to have similar topological feature; hence, it is natural to look for ingredients in the input data that may play a role similar to that of the $6j$ -symbols in the bulk.

We shall need a mathematical structure — Frobenius algebra of simple objects in a UFC — to construct the boundary terms to be added to the Levin-Wen bulk Hamiltonian. Let G be a symmetric $6j$ -symbol over the label set L . A *Frobenius algebra* is a subset $L_A \subset L$ equipped with a multiplication f , i.e. a set of multiplication coefficients f_{ijm} ($i, j, k \in L_A$), defined by $i \otimes j = \oplus_k f_{ijk} k$, satisfying

$$\begin{aligned} \text{(associativity)} \quad & \sum_c f_{abc} f_{cde} G_{de}^{abc*} v_c v_g = f_{age} f_{bdg}, \\ \text{(non-degeneracy)} \quad & f_{bb*0} \neq 0, \quad \forall b \in L_A. \end{aligned} \quad (3.1)$$

We can normalize (as described in appendix A.2) the non-degeneracy condition to

$$f_{bb*0} = 1, \quad \forall b \in L_A. \quad (3.2)$$

Due to the symmetry conditions (2.6) of symmetric $6j$ -symbols [42], the multiplication meets the following defining properties.

$$\begin{aligned} \text{(unit)} \quad & f_{bb*0} = f_{b0b*} = f_{0bb*} = 1, \\ \text{(cyclic)} \quad & f_{abc} = f_{cab}, \\ \text{(strong)} \quad & \sum_{ab} f_{abc} f_{c*b*a*} v_a v_b = d_A v_c, \end{aligned} \quad (3.3)$$

where $d_A = \sum_{a \in L_A} d_a$ is the quantum dimension of A .

If we set $g = 0, d = b^*, e = a$ in eq. (3.1), we get

$$\sum_c f_{abc} f_{cb*a*} v_c = v_a v_b. \quad (3.4)$$

Define

$$\Delta_{abc} = f_{abc} f_{c*b*a*}. \quad (3.5)$$

Then the above equation is expressed by

$$\sum_c \Delta_{abc} v_c = v_a v_b, \quad (3.6)$$

which, when compared with eq. (2.4), implies that symbols Δ_{abc} play a role similar to that of the fusion coefficients δ_{ijk} . The definition above can be illustrated graphically by the Pachner moves. The associativity condition (3.1) graphically reads

$$\begin{aligned} \mathcal{T} \left(\sum_c f_{bac^*} f_{cde^*} \left| \begin{array}{c} b \quad e \\ \swarrow \quad \searrow \\ a \quad c \quad d \end{array} \right. \right\rangle \right) \\ = \sum_c f_{abc^*} f_{cde^*} \sum_g G_{de^*g}^{abc^*} v_c v_g \left| \begin{array}{c} a \quad e \\ \swarrow \quad \searrow \\ b \quad d \end{array} \right. \left. \begin{array}{c} g \\ \updownarrow \end{array} \right. \right\rangle = \sum_g f_{bdg^*} f_{age^*} \left| \begin{array}{c} a \quad e \\ \swarrow \quad \searrow \\ b \quad d \end{array} \right. \left. \begin{array}{c} g \\ \updownarrow \end{array} \right. \right\rangle, \end{aligned} \quad (3.7)$$

where in the initial and final states, each vertex is associated with a multiplication f .

The strong condition (3.3) may also be understood graphically as

$$\mathcal{T} \left(\frac{\sqrt{D}}{d_A} \sum_{a,b} f_{abc^*} f_{cb^*a^*} \left| \begin{array}{c} c \\ \updownarrow \\ a \quad b \\ \updownarrow \\ c \end{array} \right. \right\rangle \right) = \left| \begin{array}{c} k \\ \updownarrow \end{array} \right. \right\rangle, \quad (3.8)$$

or, equivalently, as

$$\begin{aligned} \mathcal{T} \left(\frac{\sqrt{D}}{d_A} \sum_{a_2, a_4, a_5} f_{a_4^* a_5 a_6^*} f_{a_1 a_2^* a_4} f_{a_2 a_3 a_5^*} \left| \begin{array}{c} a_6 \\ \swarrow \quad \searrow \\ a_4 \quad a_5 \\ \swarrow \quad \searrow \\ a_1 \quad a_2 \quad a_3 \end{array} \right. \right\rangle \right) \\ = \frac{\sqrt{D}}{d_A} \sum_{a_2, a_4, a_5} f_{a_4^* a_5 a_6^*} f_{a_1 a_2^* a_4} f_{a_2 a_3 a_5^*} \\ \times \frac{v_{a_2} v_{a_4} v_{a_5}}{D^{1/2}} G_{a_2^* a_4 a_5^*}^{a_6 a_3^* a_1^*} \left| \begin{array}{c} a_6 \\ \updownarrow \\ a_1 \quad a_3 \end{array} \right. \right\rangle \\ = f_{a_6^* a_1 a_3} \left| \begin{array}{c} a_6 \\ \updownarrow \\ a_1 \quad a_3 \end{array} \right. \right\rangle. \end{aligned} \quad (3.9)$$

To save writing, we can suppress the coefficients f and related summations. For example, let us express the associativity and strong conditions compactly as:

$$\mathcal{T} \left(\left| \begin{array}{c} i \quad l \\ \swarrow \quad \searrow \\ j \quad k \end{array} \right. \right\rangle \right) = \left| \begin{array}{c} i \quad l \\ \swarrow \quad \searrow \\ j \quad k \end{array} \right. \right\rangle \quad (3.10)$$

$$\mathcal{T} \left(\frac{\sqrt{D}}{d_A} \left| \begin{array}{c} k \\ \updownarrow \\ \text{thick dot} \\ \updownarrow \\ k \end{array} \right. \right\rangle \right) = \left| \begin{array}{c} k \\ \updownarrow \end{array} \right. \right\rangle \quad (3.11)$$

$$\mathcal{T} \left(\frac{\sqrt{D}}{d_A} \left| \begin{array}{c} j_6 \\ \swarrow \quad \searrow \\ \text{thick dot} \\ \swarrow \quad \searrow \\ j_1 \quad j_3 \end{array} \right. \right\rangle \right) = \left| \begin{array}{c} j_6 \\ \swarrow \quad \searrow \\ j_1 \quad j_3 \end{array} \right. \right\rangle \quad (3.12)$$

The rule is to put a thick dot at any vertex associated with an f and draw an unlabeled thick line indicating a summation. We shall call this rule the thick-line convention and

Figure 3. Boundary is a wall carrying tails. j 's are bulk labels and a are tail labels.

follow it hereafter. It is sometimes natural and handy to refer to the multiplication f of a Frobenius algebra A as the Frobenius algebra without causing any confusion.

To summarize, a Frobenius algebra f determines a state (on a trivalent graph) with an f at each vertex, and a factor $\sqrt{D}/d_A$ at an internal plaquette. Such a state is invariant under Pachner moves $T_{2 \rightarrow 2}$, and $T_{3 \rightarrow 1}$ but not invariant under $T_{1 \rightarrow 3}$ that creates plaquettes. In other words, it is invariant under moves from bigger/dense graphs to smaller/sparse graphs.

For computational convenience, we set $f_{ijk} = 0$ for any i, j, k in $L \setminus L_A$, so that f is defined for all labels.

3.2 Boundary Hamiltonian

A section of a generic boundary of our model is depicted in figure 3. The boundary is a domain wall separating the bulk (in gray in the figure) and the vacuum. The bulk edges are labeled by $j_1, j_2, \dots$, which take value in L , the set of objects of the input UFC. The boundary degrees of freedom, also taking value in L , inhabit the tails, i.e. open (or dangling) edges, labeled by $a_1, a_2, \dots$. In ground states the boundary degrees of freedom are required to be restricted to a Frobenius algebra $L_A \subseteq L$; this will be implemented by projection operators comprising the boundary Hamiltonian, to be explained shortly. The Hilbert space of the model thus consists of all possible configurations of the bulk and boundary degrees of freedom.

The boundary Hamiltonian comprises two sets of operators as follows.

$$H_{\text{bdry}} = - \sum_n \bar{Q}_n - \sum_p \bar{B}_p. \quad (3.13)$$

Here, $\bar{Q}_n$ is a boundary edge operator acting on open edge n , which projects the boundary degrees of freedom to $L_A \subseteq L$:

$$\bar{Q}_n \left| \begin{array}{c} j_2 \\ \hline j_1 \end{array} \leftarrow a_n \right\rangle = \delta_{a_n \in L_A} \left| \begin{array}{c} j_2 \\ \hline j_1 \end{array} \leftarrow a_n \right\rangle. \quad (3.14)$$

And $\bar{B}_p$ is an operator comprised of operators $\bar{B}_p^t$:

$$\bar{B}_p = \frac{1}{d_A} \sum_{t \in L_A} v_t \bar{B}_p^t, \quad d_A = \sum_{t \in L_A} d_t, \quad (3.15)$$

where $\overline{B}_p^t$ acts on a boundary open plaquette between two nearest neighboring open edges:

$$\overline{B}_p^t : \left| \begin{array}{c} j_4 \\ \nearrow \\ j_3 \xrightarrow{a_2} \\ \leftarrow j_5 \\ \searrow \\ j_2 \xrightarrow{a_1} \\ \leftarrow j_1 \end{array} \right\rangle \mapsto \sum_{a'_1, a'_2, j'_2, j'_3} f_{t^* a'_2^* a_2} f_{a_1 t a'_1^*} u_{a_1} u_{a_2} u_{a'_1} u_{a'_2} \times$$

$$G_{t^* a'_2^* j'_3}^{j_4 j_3 a_2^*} G_{t^* j'_3^* j'_2}^{j_5 j_2 j_3^*} G_{j_1 a'_1^* a_1}^{t^* j'_2^* j_2} v_{j_2} v_{j_3} v_{j'_2} v_{j'_3} \left| \begin{array}{c} j_4 \\ \nearrow \\ j'_3 \xrightarrow{a'_2} \\ \leftarrow j'_5 \\ \searrow \\ j'_2 \xrightarrow{a'_1} \\ \leftarrow j_1 \end{array} \right\rangle. \quad (3.16)$$

If one of the neighboring open edges a_n or $a_{n+1} \notin L_A$, then we have $f_{t^* a'_2^* a_2} f_{a_1 t a'_1^*} = 0$, and hence

$$\overline{B}_p^t = 0. \quad (3.17)$$

Note that our definition of the boundary Hamiltonians needs a Frobenius algebra A with $L_A \subset L$ as input data.

3.3 Graphical presentation

We have reviewed unitary transformations associated with Pachner moves on 2D graphs on a closed surface, in terms of $6j$ -symbol. These transformations quantitatively describe the topological feature in the bulk. Likewise, we can use Frobenius algebras to associate unitary transformations to Pachner moves on 1D boundary part of the graphs on an open surface.

Similar to transformations (2.17), we can use the Frobenius algebra A to define unitary transformations associated with 1D Pachner moves on the boundaries of a graph: (with $u_a = \sqrt{v_a}$)

$$T_{1 \rightarrow 2} \left| \begin{array}{c} j \\ \uparrow \\ i \xrightarrow{a_1} \end{array} \right\rangle = \sum_{a_2, a_3} \frac{u_{a_1} u_{a_2} u_{a_3}}{\sqrt{d_A}} \sum_k v_k f_{a_2^* a_3^* a_1} G_{a_2^* a_3^* k}^{j^* i a_1^*} \left| \begin{array}{c} j \\ \uparrow \\ k \xrightarrow{a_3} \\ \leftarrow i \xrightarrow{a_2} \end{array} \right\rangle,$$

$$T_{2 \rightarrow 1} \left| \begin{array}{c} j \\ \uparrow \\ k \xrightarrow{a_3} \\ \leftarrow i \xrightarrow{a_2} \end{array} \right\rangle = \sum_{a_1} \frac{u_{a_1} u_{a_2} u_{a_3}}{\sqrt{d_A}} v_k f_{a_2 a_3 a_1^*} G_{a_2^* k^* a_3}^{j a_1 i^*} \left| \begin{array}{c} j \\ \uparrow \\ i \xrightarrow{a_1} \end{array} \right\rangle. \quad (3.18)$$

where $u_a = \sqrt{v_a}$ (sign of square root may be arbitrarily chosen but if fixed once then for all).

Alternative to $T_{1 \rightarrow 2}$ and $T_{2 \rightarrow 1}$, the boundary Pachner moves can be defined as

$$\begin{aligned}
 T'_{1 \rightarrow 2} \left| \begin{array}{c} \text{thick line} \\ \uparrow a_1 \\ \text{thick line} \end{array} \right\rangle &= \sum_{a_2, a_3} \frac{f_{a_2^* a_3^* a_1} u_{a_2} u_{a_3}}{u_{a_1} \sqrt{d_A}} \left| \begin{array}{c} \text{thick line} \\ \uparrow a_1 \\ \text{thick line} \end{array} \right\rangle, \\
 T'_{2 \rightarrow 1} \left| \begin{array}{c} j_3 \\ \uparrow a_1 \\ j_2 \\ \uparrow a_2 \\ j_1 \end{array} \right\rangle &= \sum_{a_3} \frac{f_{a_2 a_1 a_3^*} u_{a_3} \sqrt{D}}{u_{a_2} u_{a_1} \sqrt{d_A}} \left| \begin{array}{c} j_3 \\ \uparrow a_1 \\ j_2 \\ \uparrow a_2 \\ j_1 \end{array} \right\rangle,
 \end{aligned} \tag{3.19}$$

or

$$T''_{2 \rightarrow 1} \left| \begin{array}{c} \text{thick line} \\ \uparrow a_1 \\ \text{thick line} \end{array} \right\rangle = \frac{f_{a_2 a_3 a_1^*} u_{a_2} u_{a_3}}{u_{a_1} \sqrt{d_A}} \left| \begin{array}{c} \text{thick line} \\ \uparrow a_1 \\ \text{thick line} \end{array} \right\rangle. \tag{3.20}$$

The $T_{1 \rightarrow 2}$ and $T_{2 \rightarrow 1}$ can be derived by composing the bulk Pachner moves and these alternative T' moves.

The action of $\bar{B}_p$ can be expressed in terms of T moves: $\bar{B}_p = T_{2 \rightarrow 2} \circ \dots \circ T_{2 \rightarrow 2} \circ T'_{2 \rightarrow 1} \circ T_{1 \rightarrow 2}$. To see this, we expanded in terms of f and G :

$$\begin{aligned}
 \bar{B}_p \left| \begin{array}{c} j_4 \\ \uparrow a_2 \\ j_3 \\ \uparrow j_5 \\ j_2 \\ \uparrow a_1 \\ j_1 \end{array} \right\rangle &= \mathcal{T} \left(\frac{\sqrt{D} u_{a_1'} u_{a_2'}}{d_A u_{a_1} u_{a_2}} \left| \begin{array}{c} j_4 \\ \uparrow a_2 \\ j_3 \\ \uparrow j_5 \\ j_2 \\ \uparrow a_1 \\ j_1 \end{array} \right\rangle \right) \\
 &= \sum_{t, a_2'} \frac{f_{t^* a_2'^* a_2} u_t u_{a_2'}}{u_{a_2} \sqrt{d_A}} \sum_{a_1'} \frac{f_{a_1 t a_1'^*} u_{a_1'} \sqrt{D}}{u_{a_1} u_t \sqrt{d_A}} \sum_{j_3'} G_{t^* a_2'^* j_3'}^{j_4 j_3 a_2^*} v_{a_2} v_{j_3'} \sum_{j_2'} G_{t^* j_3'^* j_2'}^{j_5 j_2 j_3^*} v_{j_3} v_{j_2'} G_{j_1 a_1'^* a_1'}^{t^* j_2'^* j_2^*} v_{j_2} v_{a_1'} \\
 &\quad \times \frac{v_t v_{a_1}}{v_{a_1'} \sqrt{D}} \left| \begin{array}{c} j_4 \\ \uparrow a_2' \\ j_3' \\ \uparrow j_5' \\ j_2' \\ \uparrow a_1' \\ j_1 \end{array} \right\rangle.
 \end{aligned} \tag{3.21}$$

Recall the thick-line convention that a thick dot stands for an f , an unlabeled thick line stands for a summation, and a $\times$ marks the plaquette to be killed by Pachner moves. The detailed derivation of the above equation can be found in appendix A.3. More compactly, we can write $\bar{B}_p$ as

$$\bar{B}_p \left| \begin{array}{c} j_4 \\ \uparrow a_2 \\ j_3 \\ \uparrow j_5 \\ j_2 \\ \uparrow a_1 \\ j_1 \end{array} \right\rangle = \mathcal{T} \left(T'_{1 \rightarrow 2} T'_{2 \rightarrow 1} \left| \begin{array}{c} j_4 \\ \uparrow a_2 \\ j_3 \\ \uparrow j_5 \\ j_2 \\ \uparrow a_1 \\ j_1 \end{array} \right\rangle \right), \tag{3.22}$$

where by eq. (3.19) the sequence of moves, $T'_{1 \rightarrow 2} T'_{2 \rightarrow 1}$, would generate an extra bulk plaquette, which would then be killed by the transformation $\mathcal{T}$ due to eq. (3.12), so that

the final graph would be the one in the last row of eq. (3.21). This sequence of moves and transformations would generate all the coefficients and summations in the last row of eq. (3.21). On a simple open plaquette,

$$\overline{B}_p \left| \begin{array}{c} \uparrow j_3 \\ \leftarrow a_2 \\ \uparrow j_2 \\ \leftarrow a_1 \\ \uparrow j_1 \end{array} \right\rangle = T_{1 \rightarrow 2} T_{2 \rightarrow 1} \left| \begin{array}{c} \uparrow j_3 \\ \leftarrow a_2 \\ \uparrow j_2 \\ \leftarrow a_1 \\ \uparrow j_1 \end{array} \right\rangle. \quad (3.23)$$

On a generic open plaquette, the formula above should be sandwiched between a sequence of moves $T_{2 \rightarrow 2} \dots$ that turns the generic open plaquette to a simple plaquette and another sequence of moves $T_{2 \rightarrow 2} \dots$ that turns the simple plaquette back to the shape of the original generic plaquette. That is, we have $\overline{B}_p = T_{2 \rightarrow 2} \dots T_{1 \rightarrow 2} T_{2 \rightarrow 1} T_{2 \rightarrow 2} \dots$ on a generic open plaquette.

One can use the graphical representation of the boundary terms to prove that: (i) the boundary terms $\overline{Q}_n$ and $\overline{B}_p$ commute with bulk terms Q_v and B_p , and (ii) the boundary plaquette operators $\overline{B}_p$ are mutual commuting projections. Appendix A.4 records the details of the proof. Therefore, the total Hamiltonian — the sum of the bulk and boundary Hamiltonians — is exactly solvable.

4 Ground state on a disk

Now we proceed to consider the sector of states on a disk without quasiparticles in the bulk, which contains the ground state of the whole system. Effectively, on such a disk, we can apply the $\mathcal{T}$ transformation to shrink the bulk graph to a single plaquette, bounded by a circle with outward open edges, as in the equation below. We label the wall edges by l 's taking value in L and the open edges by a 's taking value in $L_A \subseteq L$, with A a Frobenius algebra. Denote by N the length of the boundary.

Restricting to the Hilbert space $H^{Q=1}$, the bulk Hamiltonian reads $H_{\text{bulk}} = -\frac{1}{D} \sum_{s \in L} d_s B_p^s$ with

$$\left\langle \begin{array}{c} \uparrow a_3 \\ \leftarrow l'_2 \\ \uparrow a_2 \\ \leftarrow l'_1 \\ \uparrow a_1 \end{array} \right\rangle \left| B_p^s \right| \left\langle \begin{array}{c} \uparrow a_3 \\ \leftarrow l_2 \\ \uparrow a_2 \\ \leftarrow l_1 \\ \uparrow a_1 \end{array} \right\rangle \quad (4.1)$$

$$= v_{l_1} v_{l'_1} v_{l_2} v_{l'_2} \dots v_{l_N} v_{l'_N} G_{sl'_1 l'_2}^{a_2^* l_2^* l_1} G_{sl'_2 l'_3}^{a_3^* l_3^* l_2} \dots G_{sl'_N l'_1}^{a_1^* l_1^* l_N}.$$

The boundary Hamiltonian takes the form $H_{\text{bdry}} = -\sum_n \overline{B}_{(n,n+1)}$, where $(n,n+1)$ labels the boundary plaquette p sandwiched by the links (a_n, a_{n+1}) , and $\overline{B}_{(n,n+1)} = \frac{1}{d_A} \sum_t \overline{B}_{(n,n+1)}^t$,

$$\left\langle \begin{array}{c} \uparrow l_{n+1} \\ \leftarrow a'_{n+1} \\ \uparrow l'_n \\ \leftarrow a'_n \\ \uparrow l_{n-1} \end{array} \right\rangle \left| \overline{B}_{(n,n+1)}^t \right| \left\langle \begin{array}{c} \uparrow l_{n+1} \\ \leftarrow a_{n+1} \\ \uparrow l_n \\ \leftarrow a_n \\ \uparrow l_{n-1} \end{array} \right\rangle \quad (4.2)$$

$$= u_{a_n} u_{a_{n+1}} u_{a'_n} u_{a'_{n+1}} v_{l_n} v_{l'_n} f_{a_{n+1} l}^{a'_{n+1}} f_{t a_n a'_n} G_{t a'_{n+1} l'_n}^{l_{n+1}^* l_n^* a_{n+1}} G_{a_n^* t^* l'_n}^{l_n^* l_{n-1}^* a_n}.$$

A topologically ordered system on the disk has exactly one ground state, which is the simultaneous $+1$ eigenvector of B_p and $\prod_{n=1}^N \bar{B}_{(n,n+1)}$. To find the unique ground state, we need to first understand the notion of local ground states on the boundary, which boils down to solving the eigen-problem of $\prod_n \bar{B}_{(n,n+1)} = 1$. It turns out that the local eigenvectors are characterized by A -modules over the Frobenius algebra A , which is defined as follows.

A (right) module over Frobenius algebra A (or, a A -module) is a subset $L_M \subseteq L$ of labels equipped with an action tensor $\rho_{j_1 j_2}^a$, with $a \in L_A$ and $j_1, j_2 \in L_M$. Note that L_M is not anything *ad hoc* but to be obtained by solving the tensor equations of $\rho_{j_1 j_2}^a$. The tensor $\rho_{j_1 j_2}^a$ vanishes outside of these subsets and satisfies the following condition.

$$\sum_{j'} \rho_{j_1 j'}^{a_1} \rho_{j' j_2}^{a_2} G_{a_2 j_2^* a'}^{j_1 a_1 j'^*} v_{j'} v_{a'} = \rho_{j_1 j_2}^{a'} f_{a_2 a'^* a_1}, \quad (4.3)$$

which can be understood in terms of Pachner moves:

$$\begin{aligned} T_{2 \rightarrow 2} \sum_{j'} \rho_{j_1 j'}^{a_1} \rho_{j' j_2}^{a_2} \left| \begin{array}{c} j_2 \uparrow \\ \swarrow a_2 \\ j' \uparrow \\ \swarrow a_1 \\ j_1 \uparrow \end{array} \right\rangle \\ = \sum_{j'} \rho_{j_1 j'}^{a_1} \rho_{j' j_2}^{a_2} \sum_{a'} G_{a_2 j_2^* a'}^{j_1 a_1 j'^*} v_{j'} v_{a'} \left| \begin{array}{c} j_2 \uparrow \\ \swarrow a' a_2 \\ j_1 \uparrow \\ \swarrow a_1 \end{array} \right\rangle = \sum_{a'} \rho_{j_1 j_2}^{a'} f_{a_2 a'^* a_1} \left| \begin{array}{c} j_2 \uparrow \\ \swarrow a' a_2 \\ j_1 \uparrow \\ \swarrow a_1 \end{array} \right\rangle, \end{aligned}$$

where the f factor in the last row should be understood as the tensor of the trivial module to be defined below eq. (4.7). Let us again take the thick-line convention. We also suppress the indices of the action tensor ρ and put it in a box. In this box-notation, the condition (4.3) takes the following compact form:

$$T_{2 \rightarrow 2} \left| \begin{array}{c} j_2 \uparrow \\ \boxed{\rho} a_2 \\ \boxed{\rho} a_1 \\ j_1 \uparrow \end{array} \right\rangle = \left| \begin{array}{c} j_2 \uparrow \\ \boxed{\rho} a_2 \\ j_1 \uparrow \\ \swarrow a_1 \end{array} \right\rangle. \quad (4.4)$$

Here a boxed ρ at a vertex means that the tensor ρ is associated with the vertex (e.g., $\rho_{j_1 j_2}^c$ on the r.h.s., with a thick-line summation). We denote the collection of all modules over A as Mod_A .

The unit condition on the Frobenius algebra A implies the unit condition on A -modules:

$$\rho_{jj}^0 = 1. \quad (4.5)$$

If we set $a' = 0$, $a_1 = a_2^* = a$, $j_2 = j_1 = j$, and $j' = k$ in eq. (4.3), we get

$$\sum_k \rho_{jk}^a \rho_{kj}^{a^*} v_k = v_a v_j, \quad (4.6)$$

which is presented graphically using T move (2.18),

$$\tau \left(\left| \begin{array}{c} \rho \\ \downarrow j \\ \rho \\ \uparrow j \end{array} \right. \right\rangle \rangle = \frac{d_a}{\sqrt{D}} \left| \begin{array}{c} \nearrow j \\ \searrow \end{array} \right\rangle. \quad (4.7)$$

Each Frobenius algebra A has a trivial module $M_0 = A$. The action tensor of M_0 is thus $[\rho_{M_0}]_{jk}^a = f_{ak}^* j$. We shall denote the minimal set of inequivalent A -modules by $\{(M, \rho_M)\}$.

Using ρ , the local eigenvectors of $\bar{B}_p = 1$, i.e., the basis of local ground states on the boundary, are expressed as

$$\sum_{a_1, a_2} u_{a_1} u_{a_2} \left| \begin{array}{c} j_2 \\ \rho \\ a_2 \\ \rho \\ a_1 \\ j_1 \end{array} \right\rangle, \quad (4.8)$$

which can be verified directly as in appendix A.5.

The ground state on a disk has neither bulk nor boundary quasiparticle excitations. It is easily shown to be non-degenerate. Using the local basis found above, the unique ground state on the disk is expressed as

$$|\Phi\rangle = \sum_M \frac{d_M}{d_A \sqrt{D}} |\Phi_M\rangle, \quad (4.9)$$

where $d_M = \sum_{j \in L_M} d_j$ and Φ_M is the wavefunction corresponding to the local eigenvector characterized by the A -module (M, ρ_M) :

$$\Phi_M \left(\begin{array}{c} a_3 \uparrow \\ l_3 \\ a_2 \rightarrow \\ l_2 \\ a_1 \downarrow \\ l_1 \\ a_N \leftarrow \end{array} \right) = u_{a_1} u_{a_2} \dots [\rho_M]_{l_1 l_2}^{a_1} [\rho_M]_{l_2 l_3}^{a_2} \dots \quad (4.10)$$

Here $\{(M, \rho_M)\}$ are all (inequivalent) irreducible modules over the algebra A .

The unique ground state on a disk can also be expressed in terms of f_{ijk} and $6j$ -symbols:

$$\begin{aligned} |\Phi\rangle &= B_p |\Phi_{M_0}\rangle \\ &= \frac{1}{D} \sum_s d_s \sum_{a'_1 a'_2 \dots l'_1 l'_2 \dots} \sum_{a_1 a_2 \dots l_1 l_2 \dots} u_{a_1} u_{a_2} \dots v_{l_1} v_{l_2} \dots v_{l'_1} v_{l'_2} \dots \\ &\quad \times G_{sl'_1 l'_2}^{a_1 l_2^* l_1} G_{sl'_2 l'_3}^{a_2 l_3^* l_2} \dots f_{a_1 l_2^* l_1} f_{a_2 l_3^* l_2} \dots \left| \begin{array}{c} a_3 \uparrow \\ l_3 \\ a_2 \rightarrow \\ l_2 \\ a_1 \downarrow \\ l_1 \\ a_N \leftarrow \end{array} \right\rangle. \end{aligned} \quad (4.11)$$

Here p is the only plaquette of bulk in the simplest reduced diagram.

We now prove that $|\Phi\rangle$ in eq. (4.9) is a ground state on the disk. It suffices to show that $B_p|\Phi\rangle = |\Phi\rangle$. We apply unitary Pachner moves on $|\Phi\rangle$ and get

$$\sum_M \sum_j \frac{d_M}{d_A \sqrt{D}} \sum_{a_1 a_2 \dots} u_{a_1} u_{a_2} \dots \times \left| \begin{array}{c} \text{disk } j \\ \rho_M \end{array} \begin{array}{c} \uparrow a_1 \uparrow a_2 \uparrow a_3 \dots \uparrow a_{N-1} \\ \xrightarrow{a_N} \end{array} \right\rangle \quad (4.12)$$

which can be evaluated using eq. (A.17)

$$\frac{1}{\sqrt{D}} \sum_j d_j \sum_{a_1 a_2 \dots} u_{a_1} u_{a_2} \dots \times \left| \begin{array}{c} \text{disk } j \\ 0 \end{array} \begin{array}{c} \uparrow a_1 \uparrow a_2 \uparrow a_3 \dots \uparrow a_{N-1} \\ \xrightarrow{a_N} \end{array} \right\rangle \quad (4.13)$$

which is a $B_p = 1$ eigenvector. Hence $|\Phi\rangle$ is a ground state on disk.

Certain useful proofs can be found in appendix A.5. We often abuse the notation by referring to M as an A -module.

5 Topological feature of ground states

In this section, we study the topological nature of the ground state. Namely, we show that the ground states of our Hamiltonian are invariant under Pachner moves.

The bulk topological feature in the case with boundaries is the same as that in the case without boundaries. We then need only to show the boundary topological feature via the 1+1D Pachner moves on boundaries, which are defined in eqs. (3.18) through (3.20).

With boundary, the topological feature can be described as follows. The ground state is invariant under any transformation composed by $T_{2 \rightarrow 2}, T_{1 \rightarrow 3}, T_{3 \rightarrow 1}$ in the bulk and $T_{1 \rightarrow 2}, T_{2 \rightarrow 1}$ on the boundary. Moreover, such transformation is unique: different ways to composing T 's results in the same transformation.

To show the uniqueness of the transformation, we consider boundary Pachner moves $T_{1 \rightarrow 2}, T_{2 \rightarrow 1}$. Take example of a transformation from N_1 tails to N_2 tails. The composition of $T_{1 \rightarrow 2}, T_{2 \rightarrow 1}$ amounts a graph structure with N_1 input edges and N_2 output edges, where each trivalent vertex is attached with a multiplication f .

(5.1)

From the Frobenius condition, the transformation presented by the graph in the dashed box is unique.

Figure 4. Effective Hilbert space of on the cylinder.

6 Ground states on a cylinder

A topologically ordered system on a cylinder has two boundaries. We can specify the two boundary Hilbert spaces and define the two boundary Hamiltonians by two Frobenius algebras over L_A and L_B , respectively. The corresponding multiplications are denoted by f_{ij}^k for $i, j, k \in L_A$ and g_{ab}^c for $a, b, c \in L_B$.

If we consider the states without any bulk quasiparticles, we can completely shrink the bulk graph by Pachner moves to make it disappear, so that the cylinder graph becomes a ring with open edges on both sides of the ring, as in figure 4. Consider the Hilbert subspace spanned by all the labels in the graph. The total Hamiltonian contains two boundary Hamiltonians defined by two Frobenius algebras.

The ground states are characterized by the A - B -bimodules, as will be defined shortly. Each bimodule P_M gives rise to a ground-state wavefunction:

$$\Phi_M^{\text{cyl}} \left(\text{Diagram of Figure 4} \right) = \prod_n u_{a_n} u_{b_n} [P_M]_{j_{n-1} j_n k_n}^{a_n b_n} . \quad (6.1)$$

An A - B -bimodule is a subset L_M equipped with an action tensor P_{ijk}^{ab} , satisfying

$$\mathcal{T} \left(\left| \begin{array}{c} \text{Diagram with two } P_M \text{ boxes} \end{array} \right\rangle \right) = \left| \begin{array}{c} \text{Diagram with one } P_M \text{ box} \end{array} \right\rangle . \quad (6.2)$$

Here the tensor P_M is expressed by a box, whose meaning is as follows:

$$\left| \begin{array}{c} \text{Diagram with } P_M \text{ box} \end{array} \right\rangle \equiv \sum_j P_{ijk}^{ab} \left| \begin{array}{c} \text{Diagram with } j \text{ edge} \end{array} \right\rangle . \quad (6.3)$$

The r.h.s. is independent of j which is summed; hence, j does not appear on l.h.s.. Note that in this work modules and bimodules are multiplicity free (see appendix A.8 for details).

The A - B -bimodules are subject to the orthonormality and completeness conditions, respectively as follows.

$$\frac{D}{d_A^2} \mathcal{T} \left(\left| \begin{array}{c} \text{Diagram with } P_M \text{ and } P_N \text{ boxes and indices } a, b, j \end{array} \right. \right\rangle \rangle = \delta_{M,N} \frac{d_j}{d_M} \left| \begin{array}{c} \text{Diagram with } P_M \text{ box and index } j \end{array} \right\rangle. \quad (6.4)$$

$$\sum_M \frac{d_M}{d_j} [P_M]_{jij}^{ab} = \delta_{a,0} \delta_{b,0} \delta_{i,j} d_A^2. \quad (6.5)$$

Both conditions can be proved in a fashion similar to that in the case of A -modules. Now the ground state $|\Phi_M^{\text{cyl}}\rangle$ characterized by the bimodule P_M can be expressed graphically as

$$|\Phi_M^{\text{cyl}}\rangle = \sum_{a_1 \dots b_1 \dots} u_{a_1} u_{a_2} \dots u_{b_1} u_{b_2} \dots \times \left| \begin{array}{c} \text{Diagram of a cylinder with four } P_M \text{ boxes and indices } a_i, b_i \end{array} \right\rangle. \quad (6.6)$$

Let us prove this. First, similar to the disk case, we study the local basis of the ground states on a cylinder. By local we mean a piece of the ring comprising two neighboring tensors as follows.

$$\overline{B}_p \overline{B}_{p'} \sum_{a_1 a_2 b_1 b_2} u_{a_1} u_{a_2} u_{b_1} u_{b_2} \left| \begin{array}{c} \text{Diagram of two } P_M \text{ boxes with indices } a_i, b_i \end{array} \right\rangle = \sum_{a'_1 a'_2 b'_1 b'_2} u_{a'_1} u_{a'_2} u_{b'_1} u_{b'_2} \left| \begin{array}{c} \text{Diagram of two } P_M \text{ boxes with indices } a'_i, b'_i \end{array} \right\rangle. \quad (6.7)$$

Detailed derivation can be found in appendix A.6. Hence, $|\Phi_M^{\text{cyl}}\rangle$ is a ground state for the A - B -bimodule M .

Each Frobenius algebra A also has a trivial bimodule $M_0 = A$. The action tensor of M_0 is thus $[P_{M_0}]_{ijk}^{ab} = f_{aj^*i} f_{bjk^*}$.

7 Boundary excitations

Boundary elementary excitations are $\overline{B}_p = 0$ eigenstates for certain boundary plaquettes p (we only consider $\overline{B}_P$ since $\overline{Q}_n = 0$ implies $\overline{B}_p = 0$). Boundary elementary excitations

support topological quasiparticles. In this section we characterize the excitations and topological quasiparticles, by studying the algebra of local operators $\overline{B}_p^t$. We show that topological quasiparticles are classified by the bimodules over A .

The main result is that topological quasiparticles are classified by A - B -bimodules, which are solutions to eq. (6.2). Particularly, if $A = B$, the topological quasiparticles and cylinder ground states are classified by the A - A -bimodules, which are also solutions to eq. (6.2) for $A = B$. The GSD on the cylinder is identical to the number of quasiparticle species on the boundaries.

There are three kinds of important operators to characterize quasiparticles. One is a set of orthonormal projection operators as measuring operators to identify quasiparticles. Another is the set of creation operators to create quasiparticle pairs (quasiparticles can not be singly created). The third is a set of hopping operators that can hop a quasiparticle along a boundary. We will construct these three kinds of operators in the following three subsections. Then we discuss the topological feature of quasiparticles in terms of hopping operators. We also discuss fluxons as a special subset of quasiparticles.

7.1 Measure quasiparticles

In this subsection we construct a set of orthonormal projection operators as measuring operators to identify quasiparticles.

Given a bimodule M , define the corresponding measuring operator Π_M by

$$\Pi_M \left| \begin{array}{c} j_6 \xrightarrow{a_2} \\ j_5 \xrightarrow{\quad} \\ j_4 \xrightarrow{k} \\ j_3 \xrightarrow{\quad} \\ j_2 \xrightarrow{\quad} \\ j_1 \xrightarrow{a_1} \end{array} \right\rangle = \frac{d_M}{d_k} \sum_{a'_1 a'_2} \frac{u_{a'_1} u_{a'_2}}{u_{a_1} u_{a_2}} \mathcal{T} \left(\left| \begin{array}{c} j_6 \xrightarrow{a_2} \\ j_5 \xrightarrow{\quad} \\ j_4 \xrightarrow{k} \\ j_3 \xrightarrow{\quad} \\ j_2 \xrightarrow{\quad} \\ j_1 \xrightarrow{a_1} \end{array} \right\rangle \begin{array}{c} a'_2 \\ \bullet \\ \downarrow \\ \boxed{P_M} \\ \uparrow \\ \bullet \\ a'_1 \end{array} \left| \begin{array}{c} a'_2 \\ \bullet \\ \downarrow \\ \boxed{P_M} \\ \uparrow \\ \bullet \\ a'_1 \end{array} \right\rangle \right).$$

Using the orthonormal condition (6.4) and completeness condition (6.5), we verify that the set $\{\Pi_{M_j}\}$ is orthonormal

$$\Pi_M \Pi_N = \delta_{M,N} \Pi_M, \quad (7.1)$$

$$\sum_M \Pi_M = \mathbb{1}. \quad (7.2)$$

A boundary elementary excitation is a +1 eigenstate of certain Π_M . In particular, however, when M is the trivial module $M_0 = A$,

$$\overline{B}_p \overline{B}_{p'} = \Pi_{M_0}; \quad (7.3)$$

hence, the eigenstate of $\Pi_{M_0} = 1$ is in fact a state without any quasiparticles in the boundary region it acts on. This verifies that Π_M commutes with $\overline{B}_p$, and hence indeed identifies the good quantum numbers of elementary excitations.

7.2 Creation operators

The elementary boundary excitations are characterized by topological quasiparticles. On an A -boundary component, quasiparticle species are identified with the A - A -bimodules M . Consequently, a boundary elementary excitation with quasiparticles carrying an A - A -bimodule M would be a $+1$ eigenstate of the measuring operator $\prod_M$. We construct a creation operator W_M to create a pair of quasiparticles carrying the bimodule M . Below shows how such an operator acts on a boundary section.

$$W_{Mj} \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right\rangle = \sum_{a'_1 j a'_3 k a'_5} \frac{u_j u_{a'_2} u_k}{u_{a_1} u_{a_2} u_{a_3}} \mathcal{T} \left(\frac{D}{d_A^2} \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right\rangle \right), \quad (7.4)$$

for $M \in \text{Mod}_{A|A}$. In this example, the operator W_M creates an M -type and M^* -type quasiparticles on both neighboring open edges j and k of the middle open edge, which becomes a'_3 . We use wavy lines to indicate the quasiparticles. If $j = k = 0$, the quasiparticles become fluxons (to be defined in section 7.4) residing in the corresponding plaquettes. By acting creation operators on ground states, we get an elementary boundary excitation basis $W_M |\Phi\rangle$.

One can verify that $W_M |\Phi\rangle$ is an eigenvector of $\Pi_M = 1$, as shown in appendix A.7.

7.3 Hopping operators

Quasiparticles can move along the boundary under the hopping operator H_M defined by

$$H_M \left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right\rangle = \frac{\sqrt{D}}{d_A} \sum_{j' a'} \frac{u_{a'} u_{j'}}{u_a u_j} \mathcal{T} \left(\left| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \right\rangle \right), \quad (7.5)$$

that hops an M -type quasiparticle initially at the bottom open edge upward across the edge.

The topological feature of elementary excitations can be described using hopping of quasiparticles.

7.4 Fluxons

We consider a subclass of quasiparticles called fluxons. Thus, we can restrict to the Hilbert subspace of $\prod_v \bar{Q}_v = 1$. We find that the local operators $\bar{B}_p^t$ form an algebra

$$\bar{B}_p^r \bar{B}_p^s = \sum_t \frac{v_r v_s}{d_A v_t} f_{r^* s^* t} f_{t^* sr} \bar{B}_p^t. \quad (7.6)$$

The quasiparticles occupied at plaquette p are then identified by the orthonormal projection operators

$$\bar{n}_p^x = \sum_t \bar{Y}_t^x \bar{B}_p^t, \quad (7.7)$$

$$\bar{n}_p^x \bar{n}_p^y = \delta_{x,y} \bar{n}_p^x, \quad (7.8)$$

where Y_t^x satisfies the following conditions, as can be derived from eq. (7.6):

$$\frac{1}{d_A} \sum_{rs} \frac{v_r v_s}{v_t} f_{r^* s^* t} f_{t^* sr} Y_r^x Y_s^x = Y_t^x. \quad (7.9)$$

Particularly, $\bar{n}_p^{x=0} = \bar{B}_p$ with $Y_t^0 = 1$ for all $t \in L_A$.

For an excitation ψ with $\bar{n}_p^x = 1$ we say ψ supports an x -type fluxon at position p . Fluxons are a subclass of the full set of topological quasiparticles identified by the bimodules. Indeed, let $Y_t^x = [P_M]_{0t0}^{tt*}$, then eq. (7.9) is identified with (6.4). Hence, fluxon is a special type of quasiparticle identified by those modules (M, P_M) in which M contains 0.

8 Examples

8.1 Charge boundary

For any input fusion category, there is always a trivial Frobenius algebra $A_0 = 0$, such that $\bar{B}_p$ is trivial and hence the boundary Hamiltonian reduces to

$$H = - \sum_v \bar{Q}_v, \quad (8.1)$$

$$\bar{Q}_n \left| \begin{array}{c} j_2 \\ \uparrow \\ j_1 \end{array} \begin{array}{c} \leftarrow a_n \end{array} \right\rangle = \delta_{a_n, 0} \left| \begin{array}{c} j_2 \\ \uparrow \\ j_1 \end{array} \begin{array}{c} \leftarrow a_n \end{array} \right\rangle. \quad (8.2)$$

The A_0 -modules (and A_0 - A_0 -bimodules) are the entire label set L , with $[\rho_j]_{jj}^0 = 1$ ($[\rho_j]_{jj}^{00} = 1$), $j \in L$. Boundary quasiparticles are then characterized by labels $j \in L$.

8.2 LW $\mathbb{Z}_2$ model

The input fusion category is $\mathbb{Z}_2$; hence the label set is $L = \{0, 1\}$, with $d_0 = d_1 = 1$, and $0^* = 0, 1^* = 1$. Fusion rules are the $\mathbb{Z}_2$ group multiplication rule $\delta_{011} = 1$, and the $6j$ -symbols are

$$G_{kln}^{ijm} = \delta_{ijm} \delta_{klm^*} \delta_{jkn^*} \delta_{inl}. \quad (8.3)$$

There are two Frobenius algebras, one is the trivial one $A_0 = 0$, which defines a charge boundary condition. Quasiparticles on the charge boundary are identified with 1 and e , with e a $\mathbb{Z}_2$ charge.

The nontrivial Frobenius algebra is $A_1 = 0 \oplus 1$, with $L_A = \{0, 1\} = L$. This is a flux boundary. The boundary quasiparticles are identified with $1, m$ with m a $\mathbb{Z}_2$ -flux. The charge-boundary is referred to z -boundary or smooth boundary (respectively, the flux-boundary to x -boundary or rough boundary) in some literature (e.g., see ref. [36]).

Cylindrical model has $\text{GSD} = 2$ with the charge-charge or flux-flux boundary conditions, and $\text{GSD} = 1$ with the charge-flux boundary condition. We shall explain the flux-flux boundary case in detail below.

Consider the model with the flux boundary condition (with the algebra A_1) on a disk illustrated in figure 5. The total Hamiltonian is

$$H = H_{\text{bulk}} + \epsilon H_{\text{bdry}}, \quad (8.4)$$

Figure 5. A trivalent graph on disk.

Figure 6. $\mathbb{Z}_2$ effective boundary theory.

where ϵ is a positive constant, and

$$H_{\text{bulk}} = - \sum_v A_v - \sum_p B_p, \quad H_{\text{bdry}} = - \sum_{p'} \overline{B}_{p'}. \quad (8.5)$$

Seen in figure 5, examples of these operators are

$$A_v = \sigma_1^z \sigma_2^z \sigma_7^z, \quad B_p = \sigma_1^x \sigma_2^x \sigma_3^x \sigma_4^x \sigma_5^x \sigma_6^x, \quad (8.6)$$

and

$$\overline{B}_{p'} = \sigma_8^x \sigma_9^x \sigma_{10}^x \sigma_{11}^x. \quad (8.7)$$

If we consider states without quasiparticles in the bulk, we can simplify the problem with the effective theory on the disk as defined on a chain, see figure 6. The Hilbert space is spanned by $N + 1$ spins: on N external edges and one internal edge denoted by 0 (the spins on all other internal edges are determined by the fusion rules). We require a global constraint

$$\prod_{n=1}^N \sigma_n^z = 1. \quad (8.8)$$

Violating this constraint results in a nontrivial charge in the bulk. The bulk Hamiltonian is reduced to

$$H_{\text{bulk}} = -\sigma_0^x - \prod_{n=1}^N \sigma_n^z. \quad (8.9)$$

The boundary Hamiltonian is

$$H_{\text{bdry}} = - \sum_{n=1}^{N-1} \sigma_n^x \sigma_{n+1}^x - \sigma_0^x \sigma_N^x \sigma_1^x. \quad (8.10)$$

The extra σ_0^x in last term is due to the nontrivial action of $\overline{B}_{p'}$ on the spin at edge 0.

Figure 7. $\mathbb{Z}_2$ effective boundary theory.

Similarly, we have an effective theory for the cylinder as defined on a chain with the flux-flux boundary condition (figure 7). Suppose we have N external edges on both boundaries of the cylinder. Then the Hilbert space is spanned by the $2N + 1$ spins, with the global constraint

$$\prod_n^N \sigma_n^z \prod_{n'}^N \sigma_{n'}^z = 1. \quad (8.11)$$

Here we denote external edges on one boundary by $n = 1, \dots, N$ and on the other boundary by $n' = 1', \dots, N'$. Violating this constraint results in a nontrivial charge in the bulk.

The plaquettes in the bulk region are completely removed by the Pachner moves. The effective cylindrical Hamiltonian consists of a bulk term, which accounts for the global constraint (8.11), and two flux-boundary Hamiltonians on either side:

$$\begin{aligned} H_{\text{cyl}} = & -\frac{1}{\epsilon} \prod_n^N \sigma_n^z \prod_{n'}^N \sigma_{n'}^z - \sum_{n=1}^{N-1} \sigma_n^x \sigma_{n+1}^x - \sigma_0^x \sigma_N^x \sigma_1^x \\ & - \sum_{n'=1}^{N-1} \sigma_{n'}^x \sigma_{(n+1)'}^x - \sigma_0^x \sigma_{N'}^x \sigma_{1'}^x. \end{aligned} \quad (8.12)$$

One verifies that the GSD = 2, as expected. The two ground states are

$$|\sigma_0^x = 1, \sigma_n^x = 1, \sigma_{n'}^x = 1\rangle + |\sigma_0^x = 1, \sigma_n^x = -1, \sigma_{n'}^x = -1\rangle, \quad (8.13)$$

and

$$|\sigma_0^x = 1, \sigma_n^x = 1, \sigma_{n'}^x = -1\rangle + |\sigma_0^x = 1, \sigma_n^x = -1, \sigma_{n'}^x = 1\rangle. \quad (8.14)$$

Here $\sigma_n^x = \pm 1$ denotes the simultaneous eigenvalues of $\sigma_n^x = \pm 1$ for all n .

8.3 LW Fibonacci model

The input fusion category is the Fibonacci category with string types $L = \{0, 2\}$, also denoted by $\{1, \tau\}$. Let $\phi = \frac{1+\sqrt{5}}{2}$ be the golden ratio. The quantum dimensions of 0 and 2 are $d_0 = 1$ and $d_2 = \phi$. The fusion rules are

$$\delta_{000} = \delta_{022} = \delta_{222} = 1, \delta_{002} = 0, \quad (8.15)$$

and the nonzero independent $6j$ -symbols G are given by

$$\begin{aligned} G_{000}^{000} &= 1, & G_{022}^{022} &= G_{222}^{022} = 1/\phi, \\ G_{222}^{000} &= 1/\sqrt{\phi}, & G_{222}^{222} &= -1/\phi^2. \end{aligned} \quad (8.16)$$

The Fibonacci category has two Frobenius algebras: the trivial one $A_0 = 0$ and the nontrivial one $A_1 = 0 \oplus 2$. The A_0 -modules are $N_0 = 0$ and $N_1 = 2$, with action tensor $[\rho_j]_{jj}^0 = 1$. This defines a charge boundary condition.

For $A_1 = 0 \oplus 2$, set $L_{A_1} = \{0, 2\}$. The only nontrivial multiplication reads

$$f_{222} = \phi^{-3/4}. \quad (8.17)$$

A_1 has two modules: (1) $M_0 = 0 \oplus 2$, i.e., A_1 itself, with action morphism being the multiplication $\rho_{jk}^a = f_{ak}^*j$; (2). $M_1 = 2$, with action morphism given by

$$[\rho_1]_{22}^2 = -\phi^{-1/4}. \quad (8.18)$$

The two algebras A_0 and A_1 are Morita equivalent, hence giving rise to the same boundary condition.

8.4 LW Ising model

The input fusion category is the Ising category, with $L = \{0, 1, 2\}$, also denoted by $\{\mathbf{1}, \sigma, \psi\}$. The quantum dimensions are $d_0 = 1$, $d_1 = \sqrt{2}$, and $d_2 = 1$. The fusion rules are

$$\delta_{000} = 1, \delta_{011} = 1, \delta_{022} = 1, \delta_{112} = 1, \quad (8.19)$$

and the nonzero $6j$ -symbols G are

$$\begin{aligned} G_{000}^{000} &= 1, & G_{111}^{000} &= \frac{1}{\sqrt[4]{2}}, & G_{222}^{000} &= 1, & G_{011}^{011} &= \frac{1}{\sqrt{2}}, \\ G_{122}^{011} &= \frac{1}{\sqrt[4]{2}}, & G_{211}^{011} &= \frac{1}{\sqrt{2}}, & G_{022}^{022} &= 1, & G_{112}^{112} &= -\frac{1}{\sqrt{2}}. \end{aligned} \quad (8.20)$$

There are two Frobenius algebras: the trivial one $A_0 = 0$, giving rise to the charge boundary condition, and $A_1 = 0 \oplus 2$.

The Frobenius algebra $A_0 = 0$ has three modules $N_0 = 0$, $N_1 = 1$, and $N_2 = 2$, which are labels in L .

The Frobenius algebra $A_1 = 0 \oplus 2$ has three modules: (1) $M_0 = 0 \oplus 2$, with $[\rho_0]_{20}^2 = [\rho_0]_{02}^2 = 1$. (2). $M_1 = 1$, with $[\rho_1]_{11}^2 = 1$. (3). $M_2 = 1$, with $[\rho_2]_{11}^2 = -1$.

9 Equivalent boundary conditions

The boundary conditions are classified by A - A -bimodules, in the sense that boundary elementary excitations with good quantum numbers are identified with equivalent bimodules.

In this section, however, we discuss a situation where two different Frobenius algebras in a unitary fusion category give rise to equivalent boundary conditions. Two Frobenius algebras A and B are Morita equivalent if category Mod_A of A -modules is equivalent to Mod_B . [43] By the previous analysis, the local ground state basis is characterized by modules. Hence, Morita equivalent Frobenius algebras define equivalent boundary conditions.

For any A -module M , $k \otimes M$ is also a right module. But $k \otimes M$ is reducible; hence, we can decompose $k \otimes M$ into a direct sum of irreducible modules. To do so, we need to study

the equivalence between $k \otimes M$ and some other irreducible module N . Define a morphism $\eta : k \otimes M \rightarrow N$ as a tensor η satisfying

$$\mathcal{T}\left(\left| \begin{array}{c} j_4 \\ \downarrow \\ \eta \\ \uparrow \\ k \\ \downarrow \\ \rho_M \\ \uparrow \\ j_3 \end{array} \right\rangle \right) = \left| \begin{array}{c} j_4 \\ \downarrow \\ \rho_N \\ \uparrow \\ a \\ \downarrow \\ \eta \\ \uparrow \\ k \\ \downarrow \\ j_3 \end{array} \right\rangle. \quad (9.1)$$

We denote the number of independent solutions of η by N_{jM}^N , called the fusion rule in Mod_A .

In the following, we give a practical check of the Morita equivalence between Mod_A and Mod_B . Frobenius algebra A is equivalent to B if the following two conditions hold. (1) All irreducible A -modules $M_0, M_1, \dots, M_n$ are mapped to irreducible B -modules $M'_0, M'_1, \dots, M'_n$. (2) The fusion rule is preserved by the mapping.

For example, in Fibonacci case, the two Frobenius algebras are Morita equivalent. One can easily verify that the fusion rules

$$2 \otimes M_0 = M_0 \oplus M_1, 2 \otimes M_1 = M_0 \quad (9.2)$$

are equivalent to

$$2 \otimes N_0 = N_1, 2 \otimes N_1 = N_0 \oplus N_1, \quad (9.3)$$

by mapping $M_0 \rightarrow N_1$ and $M_1 \rightarrow N_0$. Hence, the two Frobenius algebras are Morita equivalent and give rise to the same boundary conditions.

In the Ising case, the two Frobenius algebras are also Morita equivalent by mapping $M_1 \rightarrow N_2, M_2 \rightarrow N_1$ and $M_3 \rightarrow N_3$. One verifies the fusion rules

$$\begin{aligned} 1 \otimes M_1 &= M_2 \oplus M_3, & 1 \otimes M_2 &= M_1, & 1 \otimes M_3 &= M_1, \\ 2 \otimes M_1 &= M_1, & 2 \otimes M_2 &= M_3, & 2 \otimes M_3 &= M_2, \end{aligned} \quad (9.4)$$

are preserved under the mapping. Hence, the two Frobenius algebras are Morita equivalent and give rise to the same boundary conditions.

10 Relation to the Kitaev-Kong formulation

We used a Frobenius algebra to define the Boundary theory in this paper. This formulation is closely related to Kitaev and Kong's work [17] that formulates boundary theories using module categories over $\mathcal{C}$. In this section we will discuss the relation between our approach and the Kitaev-Kong (KK) formulation.

In our approach, we take boundary degrees of freedom from the labels of the input UFC — the same degrees of freedom as in the bulk, and we start with local boundary Hamiltonians. To write down a “good” boundary Hamiltonian we examine the (unitary representation of) 1+1D boundary Pachner moves. The desired form of the Hamiltonian will be one such that the ground-state Hilbert space is invariant under all bulk and boundary Pachner moves. This invariance leads to a Frobenius algebra structure appearing in the boundary Hamiltonian operators.

With Hilbert space spanned by labels of the input UFC, all operators are explicitly expressed using these labels. Our approach is convenient for computational purposes yet rigorous in characterizing the topological properties. The reader can compute the ground states and excitations by solving the Hamiltonian eigen-problems without knowledge of categories.

Given the bulk Levin-Wen model with input fusion category $\mathcal{C}$, in the KK formulation, the input data to specify the boundary degrees of freedom and boundary operators is a module category $\mathcal{M}$ over $\mathcal{C}$. The topological feature of the boundary ground states comes from the compatibility conditions between the bulk degrees of freedom in $\mathcal{C}$ on the left side and boundary degrees of freedom in $\mathcal{M}$. Here and after we assume the bulk is on the left of a boundary. This is always possible if one tracks along the boundary clockwise.

In the following we will build up the correspondence between the Hilbert space structures in KK formulation and our formulation, by studying the eigen-problem of $\bar{B}_p = 1$, where p label the boundary plaquettes.

Given a boundary Γ , the local basis of boundary ground states (i.e., the $\prod_p \bar{B}_p = 1$ eigenstates) has been discussed in previous sections and has the form

$$\sum_{a_1, a_2} u_{a_1} u_{a_2} \left| \begin{array}{c} j_2 \uparrow \\ \boxed{\rho} \xleftarrow{a_2} \\ \boxed{\rho} \xleftarrow{a_1} \\ j_1 \uparrow \end{array} \right\rangle \quad (10.1)$$

for some $M \in \text{Mod}_A$. This basis is defined for simple boundary plaquettes but can be generalized to cover the cases with generic boundary plaquettes, where bulk edges must also be taken into account. We write the generic form of potential basis vectors as

$$\sum_{a_1, a_2} u_{a_1} u_{a_2} \left| \begin{array}{c} j_2 \uparrow \\ \boxed{\rho_N} \xleftarrow{a_2} \\ \xrightarrow{k} \boxed{\eta} \\ \boxed{\rho_M} \xleftarrow{a_1} \\ j_1 \uparrow \end{array} \right\rangle, \quad (10.2)$$

where we assume two potentially different modules M and N and a tensor η connected to bulk edge k , to be determined by the condition $\bar{B}_p = 1$.

Acting $\bar{B}_p$ on such states yields

$$\frac{\sqrt{D}}{d_A} \mathcal{T} \left(\sum_{a_1, a_2} u_{a_1} u_{a_2} \left| \begin{array}{c} j_2 \uparrow \\ \boxed{\rho_N} \xleftarrow{a_2} \\ \boxed{\rho_N} \\ \xrightarrow{k} \boxed{\eta} \\ \boxed{\rho_M} \xleftarrow{a_1} \\ \boxed{\rho_M} \xleftarrow{a_1} \\ j_1 \uparrow \end{array} \right\rangle \right) \quad (10.3)$$

Figure 8. (a) The Hilbert space structure in the KK formulation. The boundary degrees of freedom live in a module category. (b) The Hilbert space structure in our formulation. With Mod_A treated as the module category, the local basis of boundary ground states recovers the KK Hilbert space structure.

The above is a $\bar{B}_p = 1$ eigenvector if and only if

$$\frac{\sqrt{D}}{d_A} \tau \left(\left| \begin{array}{c} j_4 \\ \rho_N \\ \eta \\ k \\ \rho_M \\ j_3 \end{array} \right\rangle \right) = \left| \begin{array}{c} j_4 \\ \eta \\ k \\ j_3 \end{array} \right\rangle. \quad (10.4)$$

This condition is equivalent to the defining property (9.1) of the morphisms in $Hom(k \otimes M, N)$ of the category Mod_A .

Therefore, on a generic boundary graph, the local basis of boundary ground states is characterized by modules M 's in Mod_A and morphisms η in Mod_A . It is known that Mod_A is equivalent to a module category. We can use such module category data M and η as input degrees of freedom to describe the ground states. See figure 8. Hence, we build up the mapping between the Hilbert space in our formulation and that in the KK formulation on the level of ground states.

This mapping is two-way, which follows from a mathematical theorem: the category of right modules over an algebra A in $\mathcal{C}$ is equivalent to the right module category over (unitary fusion) $\mathcal{C}$ [43]. The mapping is many to one. Namely, two Frobenius algebras A and B are Morita equivalent if Mod_A is equivalent to Mod_B , and they specify the same boundary condition.

In the KK formulation, the boundary excitations are constructed using module functors $Fun(\mathcal{M}, \mathcal{M})$ of the input module category $\mathcal{M}$. In our formulation, however, the elementary excitations are identified with the bimodules $Mod_{A|A}$. In this paper, we are not going to directly prove the equivalence of elementary excitations in the two formulations. Nevertheless, it is known that [43] if $\mathcal{M}$ is taken to be Mod_A then the category $Fun(\mathcal{M}, \mathcal{M})$ is equivalent to $Mod_{A|A}$. Therefore, we expect our formulation also agrees with the KK formulation on boundary elementary excitations.

By above analysis, we show that our approach with an Frobenius algebra A is equivalent to the KK formulation with input module category $\mathcal{M} = Mod_A$.

Figure 9. Two ways of composing $T_{2 \rightarrow 2}$ moves to transform from the left most graph to the right most graph.

Acknowledgments

We thank Zhenghan Wang, Liang Kong, Davide Gaiotto for helpful discussions. YDW is also supported by the Shanghai Pujiang program.

A Some proofs and details

A.1 Topological feature of the graph

In the Hilbert space $\mathcal{H}^{Q=1}$ of simultaneous eigenvectors of $Q = 1$ at all vertices, all states are invariant under any transformation composed of $T_{2 \rightarrow 2}$.

For example, consider composition of $T_{2 \rightarrow 2}$ illustrated in figure 9 involving 5 external edges. There are two ways to compose $T_{2 \rightarrow 2}$ moves to transform from the left most graph to the right most graph. Explicitly, they are

$$\begin{aligned}
 & T \left(\begin{array}{c} 2 \quad 5 \\ \diagdown \quad \diagup \\ 1 \quad 3 \quad 4 \end{array} \rightarrow \begin{array}{c} 2 \quad 5 \\ \diagdown \quad \diagup \\ 1 \quad 3 \quad 4 \end{array} \rightarrow \begin{array}{c} 2 \quad 5 \\ \diagdown \quad \diagup \\ 1 \quad 3 \quad 4 \end{array} \right) : \\
 & \left| \begin{array}{c} j_2 \quad j_6 \quad j_7 \quad j_5 \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ j_1 \quad j_3 \quad j_4 \end{array} \right\rangle \mapsto \sum_{k_3} G_{j_4 j_5^* k_3}^{j_6 j_3 j_7^*} v_{j_7} v_{k_3} \left| \begin{array}{c} j_2 \quad j_6 \quad j_5 \\ \diagdown \quad \diagup \quad \diagdown \\ j_1 \quad k_3 \quad j_4 \end{array} \right\rangle \\
 & \mapsto \sum_{k_1} G_{j_3 j_7^* k_1}^{j_2^* j_1 j_6^*} v_{j_6} v_{k_1} \left| \begin{array}{c} j_2 \quad j_7 \quad j_5 \\ \diagdown \quad \diagup \quad \diagdown \\ j_1 \quad k_1 \quad j_4 \end{array} \right\rangle \\
 & \mapsto \sum_{k_1} G_{j_3 j_7^* k_1}^{j_2^* j_1 j_6^*} v_{j_6} v_{k_1} \sum_{k_2} G_{j_4 j_5^* k_2}^{j_2^* k_1 j_7^*} v_{j_7} v_{k_2} \left| \begin{array}{c} j_2 \quad j_5 \\ \diagdown \quad \diagup \\ j_1 \quad k_1 \quad j_4 \end{array} \right\rangle \quad (A.1)
 \end{aligned}$$

Figure 10. (a) Initial and final graphs with marked plaquettes. (b) A possible sequence of Pachner moves taking the initial graph to the final.

and

$$\begin{aligned}
 & T \left(\begin{array}{c} 2 \quad 5 \\ 1 \quad 3 \end{array} \begin{array}{c} 5 \\ 4 \end{array} \rightarrow \begin{array}{c} 2 \quad 5 \\ 1 \quad 3 \end{array} \begin{array}{c} 5 \\ 4 \end{array} \rightarrow \begin{array}{c} 2 \quad 5 \\ 1 \quad 3 \end{array} \begin{array}{c} 5 \\ 4 \end{array} \rightarrow \begin{array}{c} 4 \quad 5 \\ 1 \quad 2 \end{array} \begin{array}{c} 5 \\ 3 \end{array} \right) : \\
 & \left| \begin{array}{c} j_2 \quad j_6 \quad j_7 \quad j_5 \\ j_1 \quad j_3 \quad j_4 \end{array} \right\rangle \mapsto \sum_{k_3} G_{j_4 j_5^* k_3}^{j_6 j_3 j_7^*} v_{j_7} v_{k_3} \left| \begin{array}{c} j_2 \quad j_6 \quad j_5 \\ j_1 \quad j_3 \quad j_4 \end{array} \right\rangle \\
 & \mapsto \sum_{k_3} G_{j_4 j_5^* k_3}^{j_6 j_3 j_7^*} v_{j_7} v_{k_3} \sum_{k_2} G_{k_3 j_5^* k_2}^{j_2 j_1 j_6^*} v_{j_6} v_{k_2} \left| \begin{array}{c} j_2 \quad j_5 \\ j_1 \quad j_3 \quad j_4 \end{array} \right\rangle \\
 & \mapsto \sum_{k_3} G_{j_4 j_5^* k_3}^{j_6 j_3 j_7^*} v_{j_7} v_{k_3} \sum_{k_2} G_{k_3 j_5^* k_2}^{j_2 j_1 j_6^*} v_{j_6} v_{k_2} \sum_{k_1} G_{j_3 j_4^* k_1}^{k_2 j_1 k_3} v_{k_3} v_{k_1} \left| \begin{array}{c} j_2 \quad j_5 \\ j_1 \quad j_3 \quad j_4 \end{array} \right\rangle. \quad (\text{A.2})
 \end{aligned}$$

By the pentagon identity (2.6), eq. (A.1) is identified with eq. (A.2). We can simply write both transformations as T . In fact, any transformation composed by $T_{2 \rightarrow 2}$ moves only depends on the initial and final graphs (with the same topology) and can be written as T , without specifying the choice of the sequence of $T_{2 \rightarrow 2}$ moves.

Transformations involving $T_{2 \rightarrow 2}$, $T_{1 \rightarrow 3}$ and $T_{3 \rightarrow 1}$ moves from one graph to another are not unique. So we introduce the cross-dot notation as in section 2.1, see figure 10 for an example. In figure 10(a), the initial graph has two plaquettes marked by $\times$, while the final graph has one plaquette marked by $\cdot$. In this case, all possible sequences of $T_{2 \rightarrow 2}$, $T_{1 \rightarrow 3}$, and $T_{3 \rightarrow 1}$ moves result in the same transformation between the Hilbert spaces associated with the initial and final graphs. one such sequence is shown in figure 10(b).

A.2 Frobenius algebra

Property. The unit and cyclic conditions are consequences of the association condition, via appropriate choice of normalization.

Proof. We present a choice of normalization of f . The f_{abc} is determined up to a continuous transformation. If f_{abc} is a solution, then for any nonzero complex function ξ_a

$$f'_{abc} = f_{abc} \frac{\xi_{c^*}}{\xi_a \xi_b} \quad (\text{A.3})$$

is also a solution.

Since $f_{bb^*0} \neq 0$ for all $b \in L_A$. For each dual pair $a = b^*$, set $\xi_a = \xi_b = \sqrt{f_{ab0}}$ (the order of a, b and the sign of the square root are randomly chosen but fixed once for all). We have

$$f_{jj^*0} = 1. \quad (\text{A.4})$$

Particularly, we have

$$f_{000} = 1. \quad (\text{A.5})$$

Now we prove the property with this normalization choice.

Let $a = 0$ in eq. (3.1). Using $G_{de^*g}^{0bc^*} v_c v_g = \delta_{b,c} \delta_{e,g}$, the equation implies

$$f_{0bb^*} = f_{0gg^*} \quad (\text{A.6})$$

since $f_{000} = 1$ we get

$$f_{0bb^*} = 1. \quad (\text{A.7})$$

Similarly, setting $c = 0$ in eq. (3.7) results in

$$f_{a0a^*} = 1. \quad (\text{A.8})$$

By setting $e = 0$ in eq. (3.7), together with that $G_{d0g}^{abc^*} v_c v_g = \delta_{a,g^*} \delta_{d,c^*}$, we obtain the cyclic symmetry condition

$$f_{abc^*} = f_{bc^*a}. \quad (\text{A.9})$$

A.3 Derivation of equation (3.19)

$$\begin{aligned}
 \overline{B}_p \left| \begin{array}{c} j_4 \\ j_3 \xrightarrow{a_2} j_2 \\ j_1 \xrightarrow{a_1} \end{array} \right\rangle &= \mathcal{T} \left(\frac{\sqrt{D} u_{a'_1} u_{a'_2}}{d_A u_{a_1} u_{a_2}} \left| \begin{array}{c} j_4 \xrightarrow{a_2} a'_2 \\ j_3 \xrightarrow{j_2} \times \\ j_1 \xrightarrow{a_1} a'_1 \end{array} \right\rangle \right) \\
 &= \mathcal{T} \left(\sum_{t, a'_2} \frac{f_{t^* a'_2^* a_2} u_t u_{a'_2}}{u_{a_2} \sqrt{d_A}} \sum_{a'_1} \frac{f_{a_1 t a'_1^*} u_{a'_1} \sqrt{D}}{u_{a_1} u_t \sqrt{d_A}} \left| \begin{array}{c} j_4 \xrightarrow{a_2} a'_2 \\ j_3 \xrightarrow{j_2} t \\ j_1 \xrightarrow{a_1} a'_1 \end{array} \right\rangle \right) \\
 &= \mathcal{T} \left(\sum_{t, a'_2} \frac{f_{t^* a'_2^* a_2} u_t u_{a'_2}}{u_{a_2} \sqrt{d_A}} \sum_{a'_1} \frac{f_{a_1 t a'_1^*} u_{a'_1} \sqrt{D}}{u_{a_1} u_t \sqrt{d_A}} \sum_{j'_3} G_{t^* a'_2^* j'_3}^{j_4^* j_3 a_2^*} v_{a_2} v_{j'_3} \left| \begin{array}{c} j_4 \xrightarrow{a'_2} j'_3 \\ j_3 \xrightarrow{j'_2} t \\ j_1 \xrightarrow{a'_1} \end{array} \right\rangle \right) \\
 &= \mathcal{T} \left(\sum_{t, a'_2} \frac{f_{t^* a'_2^* a_2} u_t u_{a'_2}}{u_{a_2} \sqrt{d_A}} \sum_{a'_1} \frac{f_{a_1 t a'_1^*} u_{a'_1} \sqrt{D}}{u_{a_1} u_t \sqrt{d_A}} \sum_{j'_3} G_{t^* a'_2^* j'_3}^{j_4^* j_3 a_2^*} v_{a_2} v_{j'_3} \sum_{j'_2} G_{t^* j'_3^* j'_2}^{j_5 j_2 j_3^*} v_{j_3} v_{j'_2} \left| \begin{array}{c} j_4 \xrightarrow{a'_2} j'_3 \\ j_3 \xrightarrow{j'_2} j'_2 \\ j_5 \xrightarrow{j_2} t \\ j_1 \xrightarrow{a'_1} \end{array} \right\rangle \right) \\
 &= \sum_{t, a'_2} \frac{f_{t^* a'_2^* a_2} u_t u_{a'_2}}{u_{a_2} \sqrt{d_A}} \sum_{a'_1} \frac{f_{a_1 t a'_1^*} u_{a'_1} \sqrt{D}}{u_{a_1} u_t \sqrt{d_A}} \sum_{j'_3} G_{t^* a'_2^* j'_3}^{j_4^* j_3 a_2^*} v_{a_2} v_{j'_3} \sum_{j'_2} G_{t^* j'_3^* j'_2}^{j_5 j_2 j_3^*} v_{j_3} v_{j'_2} G_{j_1 a_1^* a'_1}^{t^* j_2^* j_2} v_{j_2} v_{a'_1} \\
 &\quad \times \frac{v_t v_{a_1}}{v_{a'_1} \sqrt{D}} \left| \begin{array}{c} j_4 \xrightarrow{a'_2} j'_3 \\ j_3 \xrightarrow{j'_2} j'_2 \\ j_5 \xrightarrow{j_2} t \\ j_1 \xrightarrow{a'_1} \end{array} \right\rangle. \tag{A.10}
 \end{aligned}$$

A.4 Property of boundary terms

We state that the *Boundary terms* $\overline{Q}_n$ and $\overline{B}_p$ commute with bulk terms Q_v and B_p . The proof is straightforward though somewhat tedious. Here we offer only a sketch of the proof below.

Sketchy proof. We first consider the nontrivial case of B_p and $\overline{B}_{p'}$ for neighboring plaquettes p and p' . See figure 11. Since both B_p and $\overline{B}_{p'}$ can be expressed as a composition of T moves, the composition of B_p and $\overline{B}_{p'}$ does not depend on the order of the T moves involved:

$$\begin{aligned}
 B_p \overline{B}_{p'} &= \mathcal{T} \left(\begin{array}{c} \text{Diagram 1} \end{array} \rightarrow \begin{array}{c} \text{Diagram 2} \end{array} \rightarrow \begin{array}{c} \text{Diagram 3} \end{array} \right) \\
 &= \overline{B}_{p'} B_p. \tag{A.11}
 \end{aligned}$$

Figure 11. A bulk plaquette p and a boundary plaquette p' neighboring to each other.

Alternatively, We can compare $\bar{B}_{p'}$ with B_p and use the known properties of B_p to argue for the their commutativity. Observe that the matrix element of $\bar{B}_p^t$ can be divided into two parts: f and u factors acting only on open edge labels a 's, while G and v factors on labels j 's and a 's. The part involving G and v factors is exactly the same as those of B_p^t acting on j labels. Loosely speaking, $\bar{B}_p^t$ is a combination of half of B_p^t operator acting on half plaquette involving bulk labels, and the action on the tails determined by f . Since B_p^s commutes with $B_{p'}^s$ for any neighboring plaquette p, p' , $\bar{B}_p^t$ should commute with all $B_{p'}^s$.

Two boundary plaquette operators $\bar{B}_p$ and $\bar{B}_{p'}$ on two different boundary plaquettes p and p' also commute, i.e., $\bar{B}_p \bar{B}_{p'} = \bar{B}_{p'} \bar{B}_p$. This commutativity obviously holds if p and p' are far away, since $\bar{B}_p$ is defined locally. If p and p' are neighboring to each other, we have

$$\begin{aligned} \bar{B}_p \bar{B}_{p'} &= \mathcal{T} \left(\text{Diagram 1} \rightarrow \text{Diagram 2} \rightarrow \text{Diagram 3} \right) \\ &= \mathcal{T} \left(\text{Diagram 4} \rightarrow \text{Diagram 5} \rightarrow \text{Diagram 6} \right) = \bar{B}_{p'} \bar{B}_p, \end{aligned} \quad (\text{A.12})$$

The diagrams represent the sequence of transformations for the operator product. Diagram 1 shows two adjacent boundary plaquettes p and p' . Diagram 2 shows the application of $\bar{B}_p$, which introduces a loop on the shared edge. Diagram 3 shows the application of $\bar{B}_{p'}$. Diagram 4 shows the application of $\bar{B}_{p'}$ first, followed by $\bar{B}_p$ in Diagram 5, leading to Diagram 6 where the operators are swapped.

where the second equality follows from the association condition.

We also state that *boundary plaquette operators $\bar{B}_p$ are mutual commuting projections*:

$$\bar{B}_p \bar{B}_p = \bar{B}_p, \quad (\text{A.13})$$

which is verified by directly computation:

$$\begin{aligned} \bar{B}_p \bar{B}_p \left| \begin{array}{c} j_4 \\ j_3 \\ j_2 \\ j_1 \end{array} \right| \begin{array}{c} a_2 \\ a_1 \end{array} \right\rangle &= \mathcal{T} \left(\frac{D}{d_A^2} \sum_{a'_1, a'_2} \frac{u_{a'_1} u_{a'_2}}{u_{a_1} u_{a_2}} \left| \begin{array}{c} j_4 \\ j_3 \\ j_2 \\ j_1 \end{array} \right| \begin{array}{c} a_2 \\ a_1 \end{array} \right\rangle \right) = \mathcal{T} \left(\frac{D}{d_A^2} \sum_{a'_1, a'_2} \frac{u_{a'_1} u_{a'_2}}{u_{a_1} u_{a_2}} \left| \begin{array}{c} j_4 \\ j_3 \\ j_2 \\ j_1 \end{array} \right| \begin{array}{c} a_2 \\ a_1 \end{array} \right\rangle \right) \\ &= \mathcal{T} \left(\frac{\sqrt{D}}{d_A} \sum_{a'_1, a'_2} \frac{u_{a'_1} u_{a'_2}}{u_{a_1} u_{a_2}} \left| \begin{array}{c} j_4 \\ j_3 \\ j_2 \\ j_1 \end{array} \right| \begin{array}{c} a_2 \\ a_1 \end{array} \right\rangle \right) = \bar{B}_p \left| \begin{array}{c} j_4 \\ j_3 \\ j_2 \\ j_1 \end{array} \right| \begin{array}{c} a_2 \\ a_1 \end{array} \right\rangle, \end{aligned} \quad (\text{A.14})$$

The diagrams in the equation show the evolution of the state under the operator $\bar{B}_p$. The first diagram shows the initial state with labels j_1, j_2, j_3, j_4 and a_1, a_2 . The second diagram shows the application of the operator, which involves a sum over intermediate labels a'_1, a'_2 . The third diagram shows the result of the operator application, which is the same as the initial state, demonstrating the projection property.

where in the second equality uses the association condition, and third the strong condition.

A.5 Ground state on a disk

We first prove (4.8) and then state the proof for orthonormality and completeness for ρ_M 's.

$$\begin{aligned}
 & \overline{B}_p \sum_{a_1, a_2} u_{a_1} u_{a_2} \left| \begin{array}{c} j_2 \\ \rho \\ a_2 \\ \rho \\ a_1 \\ j_1 \end{array} \right\rangle \\
 &= \mathcal{T} \left(\frac{\sqrt{D}}{d_A} \sum_{a'_1, a'_2} u_{a'_1} u_{a'_2} \left| \begin{array}{c} j_2 \\ \rho \\ a'_2 \\ \rho \\ a'_1 \\ j_1 \end{array} \right\rangle \right) \\
 &= \mathcal{T} \left(\frac{\sqrt{D}}{d_A} \sum_{a'_1, a'_2} u_{a'_1} u_{a'_2} \left| \begin{array}{c} j_2 \\ \rho \\ \begin{array}{c} \nearrow a'_2 \\ \searrow a'_1 \end{array} \\ j_1 \end{array} \right\rangle \right) \\
 &= \mathcal{T} \left(\sum_{a'_1, a'_2} u_{a'_1} u_{a'_2} \left| \begin{array}{c} j_2 \\ \rho \\ \begin{array}{c} \nearrow a'_2 \\ \searrow a'_1 \end{array} \\ j_1 \end{array} \right\rangle \right) \\
 &= \sum_{a'_1, a'_2} u_{a'_1} u_{a'_2} \left| \begin{array}{c} j_2 \\ \rho \\ a'_2 \\ \rho \\ a'_1 \\ j_1 \end{array} \right\rangle. \tag{A.15}
 \end{aligned}$$

Orthonormality and completeness.

$$\frac{1}{d_A} \sum_a [\rho_M]_{jk}^a [\rho_M]_{kj}^{a*} \frac{v_a v_k}{v_j} = \frac{d_k}{d_M}, \tag{A.16}$$

$$\sum_M \frac{d_M}{d_j} [\rho_M]_{jj}^a = \delta_{a,0} d_A. \tag{A.17}$$

Orthonormality condition (A.16) is expressed graphically as

$$\frac{\sqrt{D}}{d_A} \mathcal{T} \left(\left| \begin{array}{c} j \\ \rho_N \\ k \\ \rho_M \\ j \end{array} \right\rangle \right) = \delta_{M,N} \frac{d_k}{d_M} \left| \begin{array}{c} j \\ j \end{array} \right\rangle. \tag{A.18}$$

Sketch proof. Given a minimal set of modules $\{\rho_M\}$, we have

$$\mathcal{T}\left(\left|\begin{array}{c} j \\ \downarrow \\ \rho_N \\ \uparrow k \\ \rho_M \\ \uparrow j \end{array}\right\rangle\right) = \delta_{M,N} \beta \left| \begin{array}{c} j \\ \downarrow \end{array} \right\rangle \quad (\text{A.19})$$

for $\beta \in \mathbb{C}$ independent of j . To compute β , we evaluate

$$\begin{aligned} \sum_j \mathcal{T}\left(\left|\begin{array}{c} j \\ \downarrow \\ \rho_M \\ \uparrow k \\ \rho_M \\ \uparrow j \end{array}\right\rangle\right) &= \sum_j \mathcal{T}\left(\left|\begin{array}{c} 0 \\ \downarrow \\ \rho_M \\ \uparrow k \\ \rho_M \\ \uparrow k \end{array}\right\rangle\right) \\ &= \beta \sum_{j \in L_M} \mathcal{T}\left(\left|\begin{array}{c} 0 \\ \downarrow \\ \rho_M \\ \uparrow j \\ \rho_M \\ \uparrow j \end{array}\right\rangle\right). \end{aligned} \quad (\text{A.20})$$

Using eq. (4.7), the last equality is evaluated explicitly as

$$d_k d_A / \sqrt{D} = \beta d_M. \quad (\text{A.21})$$

Hence we proved eq. (A.16).

We now prove completeness condition. using eq. (A.16) and (4.4), we have

$$\sum_M \frac{d_M}{d_j} \frac{\sqrt{D}}{d_A} \mathcal{T}\left(\left|\begin{array}{c} j \\ \downarrow \\ \rho_N \\ \uparrow j \\ \rho_M \\ \uparrow j \end{array}\right\rangle\right) = \left| \begin{array}{c} j \\ \downarrow \\ \rho_N \\ \uparrow j \end{array} \right\rangle. \quad (\text{A.22})$$

Introduce tensor $\mathbb{1}$ with $\mathbb{1}_{jj}^a = \sqrt{D} \delta_{a,0}$, which satisfies

$$\mathcal{T}\left(\left|\begin{array}{c} j \\ \downarrow \\ \rho_N \\ \uparrow j \\ \mathbb{1} \\ \uparrow j \end{array}\right\rangle\right) = \left| \begin{array}{c} j \\ \downarrow \\ \rho_N \\ \uparrow j \end{array} \right\rangle. \quad (\text{A.23})$$

Compare eq. (A.23) and eq. (A.22), we arrive at eq. (A.17).

A.6 Proof of equation (6.7)

In this part we sketch the graphical proof of (6.7).

$$\overline{B}_p \overline{B}_{p'} \sum_{a_1 a_2 b_1 b_2} u_{a_1} u_{a_2} u_{b_1} u_{b_2} \left| \begin{array}{c} \text{diagram} \end{array} \right\rangle = \sum_{a'_1 a'_2 b'_1 b'_2} u_{a'_1} u_{a'_2} u_{b'_1} u_{b'_2} \left| \begin{array}{c} \text{diagram} \end{array} \right\rangle. \quad (\text{A.24})$$

A.7 $W_M |\Phi\rangle$ is an eigenvector of $\Pi_M = 1$

We now verify that $W_M |\Phi\rangle$ is an eigenvector of $\Pi_M = 1$ in the following. It suffices to verify that

$$\begin{aligned} & \Pi_N \mathcal{T} \left(\sum_{a'_2 j k a_4} u_{a'_2} u_j u_k u_{a_4} \left| \begin{array}{c} \text{diagram} \end{array} \right\rangle \right) \\ &= \frac{D}{d_A^2} \mathcal{T} \left(\sum_{a'_2 j k a'_4} \frac{d_N}{d_k} u_{a'_2} u_j u_k u_{a'_4} \left| \begin{array}{c} \text{diagram} \end{array} \right\rangle \right) \\ &= \frac{D}{d_A^2} \sum_{a'_2 j k a'_4} \frac{d_N}{d_k} u_{a'_2} u_j u_k u_{a'_4} \mathcal{T} \left(\left| \begin{array}{c} \text{diagram} \end{array} \right\rangle \right) \\ &= \delta_{M,N} \mathcal{T} \left(\sum_{a'_2 j k a'_4} u_{a'_2} u_j u_k u_{a'_4} \left| \begin{array}{c} \text{diagram} \end{array} \right\rangle \right), \end{aligned} \quad (\text{A.25})$$

where the last equality is due to the completeness condition (6.5). Hence $W_M |\Phi\rangle$ is a $\Pi_M = 1$ eigenvector.

A.8 Bimodules with multiplicity

In general, the action tensor of modules ρ and bimodules P carries extra indices, say, α and β . The action tensor of a bimodule P_M is now expressed by

$$\left| \begin{array}{c} \text{thick line} \\ \text{with } P_M \\ \text{and indices } \alpha, \beta \end{array} \right\rangle \equiv \sum_j P_{ijk}^{ab, \alpha\beta} \left| \begin{array}{c} \text{thin line} \\ \text{with indices } i, j, k \end{array} \right\rangle. \quad (\text{A.26})$$

In defining property (6.2), for a thick line on the l.h.s. we should also sum over appropriate α indices of the two action tensors. Similarly, in general we need to put extra indices to a module tensor action and follow the same convention. Nevertheless, the discussion and derivation throughout the paper remains true when we add the extra indices to tensor actions and add the corresponding summation rule to the thick line convention. Therefore, in the rest of paper, we suppress the α indices for simplicity.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] X.G. Wen, *Topological Order in Rigid States*, *Int. J. Mod. Phys. B* **4** (1990) 239 [[INSPIRE](#)].
- [2] R. Tao and Y.-S. Wu, *Gauge invariance and fractional quantum Hall effect*, *Phys. Rev. B* **30** (1984) 1097 [[INSPIRE](#)].
- [3] Q. Niu, D.J. Thouless and Y.-S. Wu, *Quantized hall conductance as a topological invariant*, *Phys. Rev. B* **31** (1985) 3372 [[INSPIRE](#)].
- [4] X.G. Wen and Q. Niu, *Ground-state degeneracy of the fractional quantum Hall states in the presence of a random potential and on high-genus Riemann surfaces*, *Phys. Rev. B* **41** (1990) 9377 [[INSPIRE](#)].
- [5] Y.-S. Wu, *General Theory for Quantum Statistics in Two-Dimensions*, *Phys. Rev. Lett.* **52** (1984) 2103 [[INSPIRE](#)].
- [6] E. Dennis, A. Kitaev, A. Landahl and J. Preskill, *Topological quantum memory*, *J. Math. Phys.* **43** (2002) 4452 [[quant-ph/0110143](#)] [[INSPIRE](#)].
- [7] A.Yu. Kitaev, *Fault tolerant quantum computation by anyons*, *Annals Phys.* **303** (2003) 2 [[quant-ph/9707021](#)] [[INSPIRE](#)].
- [8] M. Freedman, A.Y. Kitaev, J. Preskill and Z. Wang, *Topological Quantum Computation*, *Bull. Amer. Math. Soc.* **40** (2003) 31 [[quant-ph/0101025](#)].
- [9] A. Stern and B.I. Halperin, *Proposed Experiments to Probe the Non-Abelian $\nu = 5/2$ Quantum Hall State*, *Phys. Rev. Lett.* **96** (2006) 016802.
- [10] C. Nayak, S.H. Simon, A. Stern, M. Freedman and S. Das Sarma, *Non-Abelian anyons and topological quantum computation*, *Rev. Mod. Phys.* **80** (2008) 1083 [[INSPIRE](#)].

- [11] Z. Luo et al., *Experimental Preparation of Topologically Ordered States via Adiabatic Evolution*, [arXiv:1608.06963](#).
- [12] Z. Luo et al., *Experimentally Probing Topological Order and Its Breakdown via Modular Matrices*, *Nature Phys.* (2017) [[arXiv:1608.06978](#)].
- [13] I. Lesanovsky and H. Katsura, *Interacting Fibonacci anyons in a Rydberg gas*, *Phys. Rev. A* **86** (2012) 041601.
- [14] K. Li et al., *Experimental Identification of Non-Abelian Topological Orders on a Quantum Simulator*, *Phys. Rev. Lett.* **118** (2017) 080502.
- [15] X.G. Wen and A. Zee, *Quantum Statistics and Superconductivity in Two Spatial Dimensions*, *Nucl. Phys. Proc. Suppl.* **15** (1990) 135 [[INSPIRE](#)].
- [16] X.-G. Wen, *Topological orders and Chern-Simons theory in strongly correlated quantum liquid*, *Int. J. Mod. Phys. B* **5** (1991) 1641 [[INSPIRE](#)].
- [17] A. Kitaev and L. Kong, *Models for gapped boundaries and domain walls*, *Commun. Math. Phys.* **313** (2012) 351 [[arXiv:1104.5047](#)].
- [18] M. Levin, *Protected edge modes without symmetry*, *Phys. Rev. X* **3** (2013) 021009 [[arXiv:1301.7355](#)] [[INSPIRE](#)].
- [19] L. Kong, *Anyon condensation and tensor categories*, *Nucl. Phys. B* **886** (2014) 436 [[arXiv:1307.8244](#)] [[INSPIRE](#)].
- [20] M. Barkeshli, C.-M. Jian and X.-L. Qi, *Classification of Topological Defects in Abelian Topological States*, *Phys. Rev. B* **88** (2013) 241103 [[arXiv:1304.7579](#)] [[INSPIRE](#)].
- [21] M. Barkeshli, C.-M. Jian and X.-L. Qi, *Theory of defects in Abelian topological states*, *Phys. Rev. B* **88** (2013) 235103 [[arXiv:1305.7203](#)] [[INSPIRE](#)].
- [22] T. Lan and X.-G. Wen, *Topological quasiparticles and the holographic bulk-edge relation in $(2+1)$ -dimensional string-net models*, *Phys. Rev. B* **90** (2014) 115119 [[arXiv:1311.1784](#)] [[INSPIRE](#)].
- [23] J. Wang and X.-G. Wen, *Boundary Degeneracy of Topological Order*, *Phys. Rev. B* **91** (2015) 125124 [[arXiv:1212.4863](#)] [[INSPIRE](#)].
- [24] L.-Y. Hung and Y. Wan, *K matrix Construction of Symmetry-Enriched Phases of Matter*, *Phys. Rev. B* **87** (2013) 195103 [[arXiv:1302.2951](#)] [[INSPIRE](#)].
- [25] T. Iadecola, T. Neupert, C. Chamon and C. Mudry, *Accessing topological order in fractionalized liquids with gapped edges*, *Phys. Rev. B* **90** (2014) 205115 [[arXiv:1407.4129](#)] [[INSPIRE](#)].
- [26] M. Barkeshli, Y. Oreg and X.-L. Qi, *Experimental Proposal to Detect Topological Ground State Degeneracy*, [arXiv:1401.3750](#) [[INSPIRE](#)].
- [27] M. Barkeshli, E. Berg and S. Kivelson, *Coherent Transmutation of Electrons into Fractionalized Anyons*, *Science* **346** (2014) 6210 [[arXiv:1402.6321](#)] [[INSPIRE](#)].
- [28] L.-Y. Hung and Y. Wan, *Ground State Degeneracy of Topological Phases on Open Surfaces*, *Phys. Rev. Lett.* **114** (2015) 076401 [[arXiv:1408.0014](#)] [[INSPIRE](#)].
- [29] T. Lan, J.C. Wang and X.-G. Wen, *Gapped Domain Walls, Gapped Boundaries and Topological Degeneracy*, *Phys. Rev. Lett.* **114** (2015) 076402 [[arXiv:1408.6514](#)] [[INSPIRE](#)].

- [30] L.-Y. Hung and Y. Wan, *Generalized ADE classification of topological boundaries and anyon condensation*, *JHEP* **07** (2015) 120 [[arXiv:1502.02026](#)] [[INSPIRE](#)].
- [31] R.B. Laughlin, *Anomalous quantum Hall effect: An Incompressible quantum fluid with fractionally charged excitations*, *Phys. Rev. Lett.* **50** (1983) 1395 [[INSPIRE](#)].
- [32] M.A. Levin and X.-G. Wen, *String net condensation: A physical mechanism for topological phases*, *Phys. Rev. B* **71** (2005) 045110 [[cond-mat/0404617](#)] [[INSPIRE](#)].
- [33] Y. Hu, Y. Wan and Y.-S. Wu, *Twisted quantum double model of topological phases in two dimensions*, *Phys. Rev. B* **87** (2013) 125114 [[arXiv:1211.3695](#)] [[INSPIRE](#)].
- [34] S. Elitzur, G.W. Moore, A. Schwimmer and N. Seiberg, *Remarks on the Canonical Quantization of the Chern-Simons-Witten Theory*, *Nucl. Phys. B* **326** (1989) 108 [[INSPIRE](#)].
- [35] J. Kock, *Frobenius Algebras and 2-D Topological Quantum Field Theories*, Cambridge University Press (2004), 1st edition, ISBN: 978-0521540315.
- [36] S.B. Bravyi and A.Yu. Kitaev, *Quantum codes on a lattice with boundary*, [quant-ph/9811052](#) [[INSPIRE](#)].
- [37] S. Beigi, P.W. Shor and D. Whalen, *The Quantum Double Model with Boundary: Condensations and Symmetries*, *Commun. Math. Phys.* **306** (2011) 663.
- [38] I. Cong, M. Cheng and Z. Wang, *Topological Quantum Computation with Gapped Boundaries*, [arXiv:1609.02037](#).
- [39] I. Cong, M. Cheng and Z. Wang, *Defects between gapped boundaries in two-dimensional topological phases of matter*, *Phys. Rev. B* **96** (2017) 195129 [[arXiv:1703.03564](#)] [[INSPIRE](#)].
- [40] A. Bullivant, Y. Hu and Y. Wan, *Twisted quantum double model of topological order with boundaries*, *Phys. Rev. B* **96** (2017) 165138 [[arXiv:1706.03611](#)] [[INSPIRE](#)].
- [41] Y. Hu, Y. Wan and Y.-s. Wu, *Boundary Hamiltonian Theory for Gapped Topological Orders*, *Chin. Phys. Lett.* **34** (2017) 077103.
- [42] Y. Hu, S.D. Stirling and Y.-S. Wu, *Ground State Degeneracy in the Levin-Wen Model for Topological Phases*, *Phys. Rev. B* **85** (2012) 075107 [[arXiv:1105.5771](#)] [[INSPIRE](#)].
- [43] V. Ostrik, *Module categories, weak Hopf algebras and modular invariants*, [math/0111139](#).