Displaced B_c^- mesons as an inclusive signature of weakly decaying double beauty hadrons

T. Gershon^a and A. Poluektov^{a,b}

^a*Department of Physics, University of Warwick,
Gibbet Hill Road, Coventry, United Kingdom*

^b*Aix Marseille Université, CNRS/IN2P3, CPPM,
163, avenue de Luminy, Marseille, France*

E-mail: T.J.Gershon@warwick.ac.uk, anton.poluektov@cern.ch

ABSTRACT: The recent discovery, by the LHCb collaboration, of the Ξ_{cc}^{++} doubly charmed baryon, has renewed interest in the spectroscopy of doubly heavy hadrons. Experimentally, however, searches for such states appear highly challenging. The reconstructed final states tend to involve multiple heavy flavoured (beauty or charm) hadrons, so the yield for any exclusive decay mode will be suppressed to unobservably low levels by the product of several branching fractions, each of which is typically 10^{-3} – 10^{-2} . Noting that decays of double beauty hadrons are the only possible source of B_c^- mesons that are displaced from the primary vertices of proton-proton collisions at the LHC, a more promising inclusive search strategy is proposed.

KEYWORDS: Spectroscopy, B physics, Hadron-Hadron scattering (experiments)

ARXIV EPRINT: [1810.06657](https://arxiv.org/abs/1810.06657)

The possible existence of doubly heavy baryons, i.e. bound states that contain two or more beauty or charm quarks, has been predicted since the discoveries of those quarks, not long after the postulation of the quark model. However, such states have relatively low production cross-sections, and require sophisticated detectors to be able to distinguish their decays from backgrounds. The first definitive observation of a doubly heavy baryon was reported by the LHCb collaboration in 2017, with a clear signal of $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ decays obtained from analysis of a data sample of high-energy proton-proton collisions [1]. Subsequently, the $\Xi_c^+ \pi^+$ decay mode has also been observed [2], and the lifetime of the Ξ_{cc}^{++} state has been found to be consistent with expectation for decays mediated by the weak interaction [3].

This discovery has prompted various theoretical predictions for the production and decay properties of other doubly heavy states. Interestingly, these have not been confined to baryonic states. Various recently discovered “exotic” hadrons, which do not fit into the conventional scheme of $q\bar{q}$ mesons and qqq baryons, have led to several different interpretations including models based on tightly bound tetraquarks (see, for example, refs. [4–6] for reviews). Several authors have predicted the existence of a $bb\bar{u}\bar{d}$ tetraquark state with mass below the threshold for decay to a pair of beauty mesons [7–13].¹ It is possible that the $bb\bar{u}\bar{s}$ and $bb\bar{d}\bar{s}$ tetraquarks are also stable against strong and electromagnetic decays. Since these tetraquarks can only decay weakly, they are expected to have comparable lifetimes to the ground state double beauty baryons, i.e. $\sim 0.4\text{--}0.8\text{ ps}$ [14–16].

Double beauty hadrons bbx , where x is either a quark or an anti-diquark, can be produced in high energy collisions such as those at the LHC where, assuming non-negligible lifetimes, they will travel a measurable distance before decaying. Typical estimates for the production cross-sections are $\mathcal{O}(1\text{ nb})$ [17]. Since the accumulated data sample at LHCb (ATLAS and CMS) already exceeds 8 (150) fb^{-1} , with increases of over an order of magnitude anticipated in the HL-LHC era, it might be assumed that prospects for discovery are good. Unfortunately, branching fractions for heavy flavour decays to even the most abundant final states are typically $10^{-2}\text{--}10^{-3}$. Exclusive reconstruction of double beauty hadrons will inevitably involve a chain of such decays. It is also relevant that the final states will involve many particles, leading to lower reconstruction efficiencies. Consequently, the yields that can be obtained are expected to be suppressed to unobservably low levels.

This motivates the development of inclusive search strategies. An important realisation, in this respect, is that weak decays of double beauty hadrons are the only possible source of B_c^- mesons that are displaced from, i.e. do not originate directly from, the primary vertices of the LHC collisions. This is a simple consequence of the facts that: (i) all fundamental Standard Model particles heavier than the b quark decay instantaneously, and do not produce a displaced vertex; (ii) any hadron containing both b and $\bar{b}$ constituent quarks will decay via strong and electromagnetic interactions and hence not have a measurable lifetime. Displaced B_c^- mesons can therefore only be produced² when one of the beauty quarks in a bbx hadron decays via a $b \rightarrow \bar{c}$ transition and the produced $\bar{c}$ antiquark hadronises with the remaining b quark as shown in figure 1. The $b \rightarrow c\bar{c}s$ transition will be dominant, since amplitudes for $b \rightarrow c\bar{c}d$, $b \rightarrow u\bar{c}s$ and $b \rightarrow u\bar{c}d$ processes involve smaller CKM matrix elements. The inclusion of charge conjugate processes is implied throughout this paper.

The signature of displaced B_c^- mesons is quite attractive experimentally, especially for the LHCb experiment which can exploit its precision vertex detector to separate signal

¹Some of these predictions predate the Ξ_{cc}^{++} discovery, in two cases quite considerably so. The experimental data have, however, put the theory on much firmer ground.

²Another conceivable source of displaced B_c^- mesons would be from weak decays of triply heavy $bq\bar{q}\bar{c}$ tetraquark states, where $q \in u, d$. However, even if such a state is stable against strong decays, its production rate will be tiny, its branching fraction for decays to final states involving B_c^- mesons will be small, and its lifetime will be very short. This potential contribution is therefore considered to have negligible impact on the proposed measurement strategy.

Figure 1. Diagram for production of a B_c^- meson from a double beauty hadron decay.

from (prompt) background [18, 19]. Moreover, when the $B_c^- \rightarrow J/\psi \pi^-$, $J/\psi \rightarrow \mu^+ \mu^-$ decay chain is used there are only three final state particles to be reconstructed, and the $J/\psi \rightarrow \mu^+ \mu^-$ signal helps to suppress backgrounds, so that high efficiency can be achieved. The branching fraction for this B_c^- decay is also relatively large, typically expected to be a few percent [20]. Its exact value is not known since the B_c^- production cross-section is unmeasured. The ratio

$$\frac{f_c}{f_u} \frac{\mathcal{B}(B_c^- \rightarrow J/\psi \pi^-)}{\mathcal{B}(B^- \rightarrow J/\psi K^-)} = (0.683 \pm 0.018 \pm 0.009)\%$$

has been measured by LHCb in the fiducial range $p_T(B) < 20 \text{ GeV}$ and $2.0 < y(B) < 4.5$ [21] (units in which $c = 1$ are used throughout this paper). Here, f_c and f_u are the fragmentation fractions for B_c^- and B^- mesons, respectively, while p_T and y denote the component of momentum transverse to the beam direction and the rapidity. This result and the prediction for $\mathcal{B}(B_c^- \rightarrow J/\psi \pi^-)$ are consistent with the expectation that $\frac{f_c}{f_u} \sim \mathcal{O}(10^{-3})$. In spite of this low production rate, a yield of 3325 ± 73 $B_c^- \rightarrow J/\psi \pi^-$ decays has been obtained in 2 fb^{-1} of proton-proton collision data at centre-of-mass energy $\sqrt{s} = 8 \text{ TeV}$, recorded at LHCb [22]. This is significantly larger than the yields reconstructed in other decay modes, and the $B_c^- \rightarrow J/\psi \pi^-$ channel will therefore be the focus of this study. The $B_c^- \rightarrow J/\psi \pi^- \pi^+ \pi^-$ mode [23], with a yield about half as large as that for $B_c^- \rightarrow J/\psi \pi^-$, may also contribute useful sensitivity to an experimental analysis.

The expected yield of displaced B_c^- mesons is given by the product of the following factors, where order-of-magnitude estimated (indicated by “ $\sim$ ”) or known values are given in parentheses: the bbx hadron production cross-section ($\sim 1 \text{ nb}$), the branching fractions for its inclusive decay to final states containing B_c^- mesons ($\sim 10\%$) and for the subsequent $B_c^- \rightarrow J/\psi \pi^-$ ($\sim 2\%$) and $J/\psi \rightarrow \mu^+ \mu^-$ (6%) transitions, the detection efficiency ($\sim 10\%$) and the integrated luminosity. Thus, around 10 displaced B_c^- mesons per fb^{-1} may be reconstructed. To put it another way, $\mathcal{O}(1\%)$ of all B_c^- mesons detected in experiments at the Large Hadron Collider may originate from decays of double beauty hadrons.

The signature of displaced meson production has been used in various ways to determine heavy flavour hadron production cross-sections. For example, displaced J/ψ mesons are commonly used as a signature of b hadron decays — all LHC experiments have ex-

Figure 2. Illustration of the non-zero IP associated with B_c^- production from weakly decaying bbx hadrons. Additional particles produced in the bbx decay are not shown. Reconstruction of the B_c^- decay vertex position and momentum vector from the $J/\psi \pi^-$ final state particles alone is sufficient to determine the IP.

exploited this approach to measure the $b\bar{b}$ production rate in different collision environments. In these analyses, the prompt and displaced J/ψ mesons are typically separated by fitting the distribution of a pseudo-proper decay time variable,

$$t_z = \frac{(z_{J/\psi} - z_{PV}) \times M_{J/\psi}}{p_z J/\psi},$$

where $z_{J/\psi}$ and z_{PV} are the respective positions of the J/ψ decay vertex and the primary vertex, and $p_z J/\psi$ is the component of the J/ψ momentum, along the beam direction. This approach cannot however be used to study displaced B_c^- mesons, as the non-negligible B_c^- lifetime of ~ 0.5 ps [24, 25] causes the production and decay vertices to be separated.

An alternative variable, the impact parameter (IP) can be used to identify displaced B_c^- mesons, as illustrated in figure 2. Due to the non-negligible lifetime of the bbx hadron, the momentum vector of the produced B_c^- meson will not point back directly at the primary vertex, i.e. the IP will be non-zero. The use of the IP to separate displaced and prompt production has previously been used to identify displaced charm hadrons produced in b hadron decays [26–28].³ As shown in figure 3, the displaced component is clearly visible at higher IP values, despite the relative yields of displaced and prompt D^0 mesons being approximately 1 : 20. This signal-to-background ratio is somewhat more favourable than expected for displaced B_c^- mesons, however better alignment has led to improvement of the LHCb IP resolution since the analysis shown in figure 3 was performed, and further improvement is expected with the upgraded vertex locator to be installed before data taking in 2021 [19, 29].

In addition to the lifetime and momentum of the weakly decaying double beauty hadron, the shape of the IP distribution for displaced B_c^- mesons depends on the mo-

³In refs. [26–28], but not in figure 3, the large branching fraction for $b \rightarrow c\mu\nu$ semileptonic decays is exploited and a muon is required to be associated to the charm hadron, which significantly suppresses prompt charm. This approach would not be possible in analysis of displaced B_c^- mesons, which are produced through $b \rightarrow \bar{c}$ transitions.

Figure 3. Separation of prompt and displaced D^0 mesons using the IP variable [26]. The data correspond to 2.9 nb^{-1} of pp collisions, and contain a large prompt component (dashed line) and a smaller displaced component (dotted line). The sum of the two components is shown by a histogram.

momentum released in the decay (i.e. the value Q , which is the difference between the decaying hadron mass and the sum of the final state particle masses). In the limit of small Q value, the momentum vectors of the B_c^- meson and its parent particle will be collinear, and hence small IP values will be obtained. For double beauty baryon decays producing a B_c^- meson through a $b \rightarrow c\bar{c}s$ transition, the final state must contain a charm-strange baryon ($m_{\Xi_c} \sim 2.47 \text{ GeV}$), while for double beauty tetraquark decays a DK combination ($m_D + m_K \sim 2.36 \text{ GeV}$) must be present. Thus, a similar maximum Q value is obtained in both cases: assuming a bbx mass of 10.5 GeV , $Q_{\text{max}} \sim 1.8 \text{ GeV}$. This is sufficient to obtain a significantly non-zero IP value.

In order to understand how the unknown properties of the bbx hadrons, in particular their masses and lifetimes, affect the IP distribution and hence the feasibility of the proposed analysis approach, a study based on a simple simulation of $bb\bar{u}\bar{d}$ tetraquarks has been performed. The kinematic properties, within the LHCb acceptance, are assumed to be the same as those for B mesons, i.e. approximately flat in pseudorapidity and having an exponential distribution in p_T with mean 5 GeV [30]. Decay to the $B_c^- D^+ K^-$ final state, uniform over phase space, is assumed, with subsequent $B_c^- \rightarrow J/\psi \pi^-$, $J/\psi \rightarrow \mu^+ \mu^-$ transitions. Transverse momentum requirements, similar to those in other studies of B_c^- mesons at LHCb [22], are imposed on the final-state particles.

Within an experimental analysis, the B_c^- IP would be determined from a full vertex fit of the three tracks, accounting for the covariance matrices of their position and momentum measurements, which is beyond the scope of the current study. Instead, the IP resolution is assumed to be dominated by the coordinate resolution of the B_c^- vertex, so that the IP resolution can be obtained as the uncertainty of the weighted average of the IP values for the three tracks. The per-track IP resolution is taken to be that obtained by LHCb with data collected in 2012, i.e. $\sigma(\text{IP}_{x,y}) [\mu\text{m}] = 11.6 + 23.4/(p_T [\text{GeV}])$ [19]. The shapes

Figure 4. IP distributions obtained from simplified simulations for different $b\bar{b}x$ hadron masses (and thus Q values) and lifetimes.

obtained for a range of relevant parameters are shown in figure 4. It can be seen that, for $b\bar{b}x$ lifetimes above about 0.5 ps, the distribution for displaced B_c^- hadrons extends to significantly higher IP values than that for prompt production.

The shape of the IP distribution for prompt B_c^- mesons is given purely by the resolution function, since the underlying true distribution is a delta function. Given the expected signal-to-background ratio of around 1 : 100, the sensitivity of the analysis will depend critically on the detector IP resolution — it must be as precise as possible to separate prompt and displaced production, and must be well-known to avoid potentially large systematic uncertainties. It should be stressed that the IP distribution for prompt B_c^- mesons shown in figure 4 is based on a simple study containing several approximations, and therefore is not expected to provide an accurate description of the real experimental situation, particularly in the tails. It will be crucial in an experimental analysis that a suitable control mode can be used to study the resolution function. Fortunately, the IP resolution for $B_c^- \rightarrow J/\psi \pi^-$ decays can be studied in data using the $B^- \rightarrow J/\psi K^-$ decay as a control sample. Contributions from non-prompt B^- mesons can be reduced by requiring production in narrow $\bar{B}_s^{*0} \rightarrow B^- K^+$ decays [31, 32] where the associated K^+ track is required to originate from the primary vertex. Therefore, systematic uncertainties due to the tail of the resolution function should be controllable.

While the IP is a powerful variable that illustrates how prompt and displaced B_c^- mesons can be separated, there is also additional information that could potentially be

exploited to improve the sensitivity in an experimental analysis. The better separation of prompt and displaced components at higher transverse momentum could be exploited by studying the two-dimensional distribution of IP and p_T . The per-candidate estimated uncertainty on the IP measurement could alternatively be used in the analysis to enhance the sensitivity; at LHCb, this is usually done by constructing a χ^2 variable. Similarly, better separation is expected at larger values of the distance between the PV and the B_c^- decay vertex. Determining the optimal analysis approach would be an important aspect of experimental study.

It must be noted that whilst the signature of displaced B_c^- mesons would be sufficient to indicate double beauty hadron production, it would provide only limited information about the properties of those hadrons. The yield is related to the production cross-section times the inclusive branching fraction for decay to final states containing B_c^- mesons, summed over all weakly decaying bbx hadrons. Therefore, with improved predictions for these cross-sections and branching fractions, it could be possible to test whether only double beauty baryons are contributing, or if there is evidence for double beauty tetraquark production. The shape of the distribution depends on the mass, lifetime and kinematic properties of the bbx hadrons, averaged over all the contributing species. Hence with predictions for two of these, the third could potentially be inferred. Moreover, an observation of bbx hadron production from an inclusive analysis would be likely to lead to insights as to how further information could be obtained, for example on which exclusive decays may be most accessible.

Finally, it is worthwhile to consider if any other inclusive signatures of displaced heavy flavour hadrons can be used similarly. Secondary B mesons can only be produced in decays of doubly heavy hadrons, with beauty charm hadron decays likely to be the dominant source. Indeed, the $B_c^- \rightarrow \bar{B}_s^0 \pi^-$ decay has already been observed [33]. However, the Q value for B meson production in B_c^- or Ξ_{bc} decays is relatively small, making it harder to separate signal from prompt production through the IP distribution. This effect also exists, but is less pronounced, for b baryon production, so a search for displaced Λ_b^0 baryons as an inclusive signature of beauty charm hadrons could potentially be interesting. (Secondary Λ_b^0 production through $\Xi_b^- \rightarrow \Lambda_b^0 \pi^-$ is possible [34] but has a very low Q value and hence no significant contribution to the IP.) Another exciting possibility for the long term is that displaced Ξ_{cc} baryons could also be used to search for these states. In these cases a reconstructed muon produced through semileptonic decay could potentially be used to tag the signal and reduce the prompt background significantly.

In conclusion, displaced B_c^- mesons provide a distinctive signature of the production of double beauty hadrons. There appear to be reasonable prospects for the detection of such a signal with the large data samples accumulated by the LHCb detector and its planned upgrades, since excellent IP resolution is a feature of this experiment.

Acknowledgments

The authors wish to thank their colleagues on the LHCb experiment for the fruitful and enjoyable collaboration that inspired this study. In particular, they would like to thank Vanya

Belyaev, Steve Blusk, Alex Bondar, Mat Charles, Vladimir Gligorov, Jibo He and Mika Vesterinen for helpful comments. This work is supported by the Science and Technology Facilities Council.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] LHCb collaboration, *Observation of the doubly charmed baryon Ξ_{cc}^{++}* , *Phys. Rev. Lett.* **119** (2017) 112001 [[arXiv:1707.01621](https://arxiv.org/abs/1707.01621)] [[INSPIRE](#)].
- [2] LHCb collaboration, *First observation of the doubly charmed baryon decay $\Xi_{cc}^{++} \rightarrow \Xi_c^+ \pi^+$* , *Phys. Rev. Lett.* **121** (2018) 162002 [[arXiv:1807.01919](https://arxiv.org/abs/1807.01919)] [[INSPIRE](#)].
- [3] LHCb collaboration, *First measurement of the lifetime of the doubly charmed baryon Ξ_{cc}^{++}* , *Phys. Rev. Lett.* **121** (2018) 052002 [[arXiv:1806.02744](https://arxiv.org/abs/1806.02744)] [[INSPIRE](#)].
- [4] S.L. Olsen, T. Skwarnicki and D. Zieminska, *Nonstandard heavy mesons and baryons: Experimental evidence*, *Rev. Mod. Phys.* **90** (2018) 015003 [[arXiv:1708.04012](https://arxiv.org/abs/1708.04012)] [[INSPIRE](#)].
- [5] A. Ali, J.S. Lange and S. Stone, *Exotics: Heavy Pentaquarks and Tetraquarks*, *Prog. Part. Nucl. Phys.* **97** (2017) 123 [[arXiv:1706.00610](https://arxiv.org/abs/1706.00610)] [[INSPIRE](#)].
- [6] R.F. Lebed, R.E. Mitchell and E.S. Swanson, *Heavy-Quark QCD Exotica*, *Prog. Part. Nucl. Phys.* **93** (2017) 143 [[arXiv:1610.04528](https://arxiv.org/abs/1610.04528)] [[INSPIRE](#)].
- [7] J.P. Ader, J.M. Richard and P. Taxil, *Do narrow heavy multi-quark states exist?*, *Phys. Rev. D* **25** (1982) 2370 [[INSPIRE](#)].
- [8] A.V. Manohar and M.B. Wise, *Exotic $QQ\bar{q}\bar{q}$ states in QCD*, *Nucl. Phys. B* **399** (1993) 17 [[hep-ph/9212236](https://arxiv.org/abs/hep-ph/9212236)] [[INSPIRE](#)].
- [9] P. Bicudo, K. Cichy, A. Peters, B. Wagenbach and M. Wagner, *Evidence for the existence of $u\bar{d}b\bar{b}$ and the non-existence of $s\bar{s}b\bar{b}$ and $c\bar{c}b\bar{b}$ tetraquarks from lattice QCD*, *Phys. Rev. D* **92** (2015) 014507 [[arXiv:1505.00613](https://arxiv.org/abs/1505.00613)] [[INSPIRE](#)].
- [10] A. Francis, R.J. Hudspith, R. Lewis and K. Maltman, *Lattice Prediction for Deeply Bound Doubly Heavy Tetraquarks*, *Phys. Rev. Lett.* **118** (2017) 142001 [[arXiv:1607.05214](https://arxiv.org/abs/1607.05214)] [[INSPIRE](#)].
- [11] M. Karliner and J.L. Rosner, *Discovery of doubly-charmed Ξ_{cc} baryon implies a stable $b\bar{b}u\bar{d}$ tetraquark*, *Phys. Rev. Lett.* **119** (2017) 202001 [[arXiv:1707.07666](https://arxiv.org/abs/1707.07666)] [[INSPIRE](#)].
- [12] E.J. Eichten and C. Quigg, *Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$* , *Phys. Rev. Lett.* **119** (2017) 202002 [[arXiv:1707.09575](https://arxiv.org/abs/1707.09575)] [[INSPIRE](#)].
- [13] A. Czarnecki, B. Leng and M.B. Voloshin, *Stability of tetrons*, *Phys. Lett. B* **778** (2018) 233 [[arXiv:1708.04594](https://arxiv.org/abs/1708.04594)] [[INSPIRE](#)].
- [14] V.V. Kiselev and A.K. Likhoded, *Baryons with two heavy quarks*, *Phys. Usp.* **45** (2002) 455 [[hep-ph/0103169](https://arxiv.org/abs/hep-ph/0103169)] [[INSPIRE](#)].
- [15] M. Karliner and J.L. Rosner, *Baryons with two heavy quarks: Masses, production, decays and detection*, *Phys. Rev. D* **90** (2014) 094007 [[arXiv:1408.5877](https://arxiv.org/abs/1408.5877)] [[INSPIRE](#)].

- [16] A.V. Berezhnuy, A.K. Likhoded and A.V. Luchinsky, *Doubly heavy baryons at LHC*, [arXiv:1809.10058](#) [INSPIRE].
- [17] A. Ali, Q. Qin and W. Wang, *Discovery potential of stable and near-threshold doubly heavy tetraquarks at the LHC*, *Phys. Lett. B* **785** (2018) 605 [[arXiv:1806.09288](#)] [INSPIRE].
- [18] LHCb collaboration, *The LHCb Detector at the LHC*, 2008 *JINST* **3** S08005 [INSPIRE].
- [19] R. Aaij et al., *Performance of the LHCb Vertex Locator*, 2014 *JINST* **9** P09007 [[arXiv:1405.7808](#)] [INSPIRE].
- [20] C.-F. Qiao, P. Sun, D. Yang and R.-L. Zhu, *B_c exclusive decays to charmonium and a light meson at next-to-leading order accuracy*, *Phys. Rev. D* **89** (2014) 034008 [[arXiv:1209.5859](#)] [INSPIRE].
- [21] LHCb collaboration, *Measurement of B_c^+ production in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Phys. Rev. Lett.* **114** (2015) 132001 [[arXiv:1411.2943](#)] [INSPIRE].
- [22] LHCb collaboration, *Search for excited B_c^+ states*, *JHEP* **01** (2018) 138 [[arXiv:1712.04094](#)] [INSPIRE].
- [23] LHCb collaboration, *First observation of decay $B_c^+ \rightarrow J/\psi \pi^+ \pi^- \pi^+$* , *Phys. Rev. Lett.* **108** (2012) 251802 [[arXiv:1204.0079](#)] [INSPIRE].
- [24] LHCb collaboration, *Measurement of the B_c^+ meson lifetime using $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$ decays*, *Eur. Phys. J. C* **74** (2014) 2839 [[arXiv:1401.6932](#)] [INSPIRE].
- [25] LHCb collaboration, *Measurement of the lifetime of the B_c^+ meson using the $B_c^+ \rightarrow J/\psi \pi^+$ decay mode*, *Phys. Lett. B* **742** (2015) 29 [[arXiv:1411.6899](#)] [INSPIRE].
- [26] LHCb collaboration, *Measurement of $\sigma(pp \rightarrow b\bar{b}X)$ at $\sqrt{s} = 7$ TeV in the forward region*, *Phys. Lett. B* **694** (2010) 209 [[arXiv:1009.2731](#)] [INSPIRE].
- [27] LHCb collaboration, *Measurement of b hadron production fractions in 7 TeV pp collisions*, *Phys. Rev. D* **85** (2012) 032008 [[arXiv:1111.2357](#)] [INSPIRE].
- [28] LHCb collaboration, *Measurement of the b -quark production cross-section in 7 and 13 TeV pp collisions*, *Phys. Rev. Lett.* **118** (2017) 052002 [[arXiv:1612.05140](#)] [INSPIRE].
- [29] LHCb collaboration, *LHCb VELO Upgrade Technical Design Report*, CERN-LHCC-2013-021 (2013) [INSPIRE].
- [30] LHCb collaboration, *Measurement of the $B^\pm$ production cross-section in pp collisions at $\sqrt{s} = 7$ and 13 TeV*, *JHEP* **12** (2017) 026 [[arXiv:1710.04921](#)] [INSPIRE].
- [31] CDF collaboration, T. Aaltonen et al., *Observation of orbitally excited B_s mesons*, *Phys. Rev. Lett.* **100** (2008) 082001 [[arXiv:0710.4199](#)] [INSPIRE].
- [32] LHCb collaboration, *First observation of the $B_{s2}^*(5840)^0 \rightarrow B^{*+} K^-$ and studies of excited B_s^0 mesons*, *Phys. Rev. Lett.* **110** (2013) 151803 [[arXiv:1211.5994](#)] [INSPIRE].
- [33] LHCb collaboration, *Observation of the decay $B_c^+ \rightarrow B_s^0 \pi^+$* , *Phys. Rev. Lett.* **111** (2013) 181801 [[arXiv:1308.4544](#)] [INSPIRE].
- [34] LHCb collaboration, *Evidence for the strangeness-changing weak decay $\Xi_b^- \rightarrow \Lambda_b^0 \pi^-$* , *Phys. Rev. Lett.* **115** (2015) 241801 [[arXiv:1510.03829](#)] [INSPIRE].