RECEIVED: October 19, 2018

REVISED: December 13, 2018

ACCEPTED: December 24, 2018

PUBLISHED: January 10, 2019

Non-perturbative effects in $\mu \rightarrow e\gamma$

Wouter Dekens, Elizabeth E. Jenkins, Aneesh V. Manohar and Peter Stoffer¹

*Department of Physics, University of California at San Diego,
9500 Gilman Drive, La Jolla, CA 92093-0319, U.S.A.*

E-mail: wdekens@ucsd.edu, ejenkins@ucsd.edu, amanohar@ucsd.edu,
pstoffer@ucsd.edu

ABSTRACT: We compute the non-perturbative contribution of semileptonic tensor operators $(\bar{q}\sigma^{\mu\nu}q)(\bar{\ell}\sigma_{\mu\nu}\ell)$ to the purely leptonic process $\mu \rightarrow e\gamma$ and to the electric and magnetic dipole moments of charged leptons by matching onto chiral perturbation theory at low energies. This matching procedure has been used extensively to study semileptonic and leptonic weak decays of hadrons. In this paper, we apply it to observables that contain no strongly interacting external particles. The non-perturbative contribution to $\mu \rightarrow e$ processes is used to extract the best current bound on lepton-flavor-violating semileptonic tensor operators, $\Lambda_{\text{BSM}} \gtrsim 450 \text{ TeV}$. We briefly discuss how the same method applies to dark-matter interactions.

KEYWORDS: Chiral Lagrangians, Effective Field Theories, Nonperturbative Effects

ARXIV EPRINT: [1810.05675](https://arxiv.org/abs/1810.05675)

¹Corresponding author.

Contents

1	Introduction	1
2	Matching the LEFT to chiral perturbation theory	2
3	Constraints on electromagnetic dipole interactions	7
3.1	Lepton-flavor-violating processes	7
3.2	Constraints on flavor-diagonal dipole moments	9
3.3	Comment on dark-matter interactions	11
4	Conclusion	12
A	$\mu \rightarrow e$ conversion in nuclei	13
B	Anomalous dimensions	14

1 Introduction

The branching ratio for the purely leptonic decay $\mu \rightarrow e\gamma$ has the current upper bound

$$\text{BR}(\mu \rightarrow e\gamma) < 4.2 \times 10^{-13}, \quad (1.1)$$

determined by the MEG collaboration [1, 2]. In the Standard Model (SM) including flavor-changing Majorana neutrino masses, the decay rate is suppressed by $(m_\nu/M_W)^4 \sim 10^{-48}$, so the result eq. (1.1) provides strong constraints on $\mu \rightarrow e$ transitions due to new physics beyond the Standard Model (BSM). The effect of new physics at energies above the muon mass can be described by an effective field theory, and the leading operators that contribute to $\mu \rightarrow e\gamma$ are the dipole operators¹

$$\mathcal{L} = L_{e\gamma} \bar{e}_L \sigma^{\mu\nu} \mu_R F_{\mu\nu} + L_{e\gamma} \bar{\mu}_L \sigma^{\mu\nu} e_R F_{\mu\nu} + \text{h.c.} \quad (1.2)$$

These dimension-five operators result in a decay width

$$\Gamma(\mu \rightarrow e\gamma) = \frac{m_\mu^3}{4\pi} \left(|L_{e\gamma}|^2 + |L_{e\gamma}|^2 \right) + \mathcal{O}(m_e^2), \quad (1.3)$$

so eq. (1.1) places the limit

$$\sqrt{|L_{e\gamma}|^2 + |L_{e\gamma}|^2} < 3.7 \times 10^{-11} \text{ TeV}^{-1}. \quad (1.4)$$

If one assumes that new physics consists of particles with masses well above the electro-weak scale, the Standard Model Effective Field Theory (SMEFT) provides an adequate

¹We use the conventions of refs. [3, 4]. Other commonly used normalization conventions include factors of e and/or m_μ .

description of BSM effects [5, 6]. At the weak scale, the SMEFT is matched to the low-energy effective field theory (LEFT) [3, 4]. In such a scenario, the dimension-five dipole operators in eq. (1.2) derive from dimension-six dipole operators in the SMEFT, so their coefficients are of the same size as those of the other dimension-six operators. In this paper, we will consider the case where $L_{e\gamma}$ is formally of order $1/\Lambda_{\text{BSM}}^2$, and of the same size as other dimension-six operators. In many BSM models, the dipole operators are typically induced with a coefficient $L_{e\gamma} \sim em_\mu/\Lambda_{\text{BSM}}^2$, and so the limit eq. (1.4) is equivalent to $\Lambda_{\text{BSM}} > 930 \text{ TeV}$.

The operators in the low-energy effective theory below the weak scale have been classified in ref. [3]. We can calculate the decay rate $\mu \rightarrow e\gamma$ in terms of LEFT operators renormalized at a scale $\mu \sim 2 \text{ GeV}$ at which perturbation theory is still valid. The heavy quarks can be integrated out of the theory, so the low-energy effective theory contains three dynamical light quark flavors, $q = u, d, s$.

In addition to dipole operators such as eq. (1.2), LEFT has dimension-six q^4 , ℓ^4 , $q^2\ell^2$, and G^3 operators, where q , ℓ , and G are quark, lepton, and gluon fields. The ℓ^4 and $q^2\ell^2$ operators contribute to lepton-flavor-violating processes such as $\mu \rightarrow e\gamma$. The ℓ^4 operators contribute to $\mu \rightarrow e\gamma$ via a perturbative loop graph giving $L_{e\gamma} \sim em_\ell/(16\pi^2)L_{\ell^4}$, where L_{ℓ^4} is a typical ℓ^4 operator coefficient. In this paper, we compute the non-perturbative contribution of semileptonic $q^2\ell^2$ operators to the purely leptonic process $\mu \rightarrow e\gamma$, and show that these are of order $L_{e\gamma} \sim e(F_\pi^2/\Lambda_\chi)L_{q^2\ell^2}$, where $\Lambda_\chi \sim 4\pi F_\pi$ is the scale of chiral symmetry breaking [7]. These contributions are significant, and the limit eq. (1.1) on the $\mu \rightarrow e\gamma$ branching ratio puts interesting limits on semileptonic tensor operators.

The constraints of lepton-flavor-violating processes such as $\mu \rightarrow e\gamma$ and $\mu \rightarrow 3e$ on the Wilson coefficients of both SMEFT and LEFT have attracted a lot of attention in connection with perturbative effects [8–11], while the non-perturbative effects due to lepton-flavor-violating quark operators have been studied mainly in reactions involving hadrons as external states [12–19]. In our analysis, we consider non-perturbative contributions to $\mu \rightarrow e\gamma$, a decay that has no strongly interacting external particles.

Within the LEFT, the perturbative renormalization-group equations can be used to compute the running and mixing of operators from the weak scale down to the hadronic scale around $\sim 2 \text{ GeV}$. Below this scale, non-perturbative QCD effects are present and must be included. At low energies, the interactions of the lightest hadrons are described by chiral perturbation theory (χ PT) [20–22]. In order to express the low-energy constants of χ PT in terms of the LEFT parameters, a non-perturbative matching has to be performed, either using lattice QCD or phenomenological input. In this paper, we study the matching of semileptonic $q^2\ell^2$ operators onto χ PT operators that contribute to processes with no hadrons.

2 Matching the LEFT to chiral perturbation theory

If we consider the leptonic sector of the LEFT, at leading order in the electromagnetic coupling, the quark dipole operators and the q^4 and G^3 operators only affect the hadronic vacuum polarization function. The most interesting non-perturbative effect on the lepton

$(\bar{L}L)(\bar{L}L)$		$(\bar{L}L)(\bar{R}R)$		$(\bar{L}R)(\bar{L}R) + \text{h.c.}$	
$\mathcal{O}_{eu}^{V,LL}$	$(\bar{e}_{Lp}\gamma^\mu e_{Lr})(\bar{u}_{Lw}\gamma_\mu u_{Lt})$	$\mathcal{O}_{eu}^{V,LR}$	$(\bar{e}_{Lp}\gamma^\mu e_{Lr})(\bar{u}_{Rw}\gamma_\mu u_{Rt})$	$\mathcal{O}_{eu}^{S,RR}$	$(\bar{e}_{Lp}e_{Rr})(\bar{u}_{Lw}u_{Rt})$
$\mathcal{O}_{ed}^{V,LL}$	$(\bar{e}_{Lp}\gamma^\mu e_{Lr})(\bar{d}_{Lw}\gamma_\mu d_{Lt})$	$\mathcal{O}_{ed}^{V,LR}$	$(\bar{e}_{Lp}\gamma^\mu e_{Lr})(\bar{d}_{Rw}\gamma_\mu d_{Rt})$	$\mathcal{O}_{ed}^{S,RR}$	$(\bar{e}_{Lp}e_{Rr})(\bar{d}_{Lw}d_{Rt})$
		$\mathcal{O}_{ue}^{V,LR}$	$(\bar{u}_{Lp}\gamma^\mu u_{Lr})(\bar{e}_{Rw}\gamma_\mu e_{Rt})$	$\mathcal{O}_{eu}^{T,RR}$	$(\bar{e}_{Lp}\sigma^{\mu\nu}e_{Rr})(\bar{u}_{Lw}\sigma_{\mu\nu}u_{Rt})$
		$\mathcal{O}_{de}^{V,LR}$	$(\bar{d}_{Lp}\gamma^\mu d_{Lr})(\bar{e}_{Rw}\gamma_\mu e_{Rt})$	$\mathcal{O}_{ed}^{T,RR}$	$(\bar{e}_{Lp}\sigma^{\mu\nu}e_{Rr})(\bar{d}_{Lw}\sigma_{\mu\nu}d_{Rt})$
$(\bar{R}R)(\bar{R}R)$				$(\bar{L}R)(\bar{R}L) + \text{h.c.}$	
$\mathcal{O}_{eu}^{V,RR}$	$(\bar{e}_{Rp}\gamma^\mu e_{Rr})(\bar{u}_{Rw}\gamma_\mu u_{Rt})$			$\mathcal{O}_{eu}^{S,RL}$	$(\bar{e}_{Lp}e_{Rr})(\bar{u}_{Rw}u_{Lt})$
$\mathcal{O}_{ed}^{V,RR}$	$(\bar{e}_{Rp}\gamma^\mu e_{Rr})(\bar{d}_{Rw}\gamma_\mu d_{Rt})$			$\mathcal{O}_{ed}^{S,RL}$	$(\bar{e}_{Lp}e_{Rr})(\bar{d}_{Rw}d_{Lt})$

Table 1. Semileptonic LEFT operators involving a charged-lepton bilinear and a quark bilinear.

sector is given by semileptonic $q^2\ell^2$ operators. We consider the subset of LEFT operators involving a charged-lepton bilinear and a quark bilinear [3], shown in table 1.

Quark operators in the LEFT Lagrangian can be matched onto operators in the chiral Lagrangian, as long as the momentum transfer is well below Λ_χ . Such a matching has been extensively used in hadronic weak decays (e.g. see ref. [23]). The results for scalar and vector quark bilinears can be obtained from the usual χ PT Lagrangian including external sources [21, 22]. The extension to tensor sources has been derived in ref. [24].

We quickly remind the reader of the construction of the chiral Lagrangian with external sources. The massless QCD Lagrangian is supplemented by quark bilinears according to

$$\begin{aligned}\mathcal{L} = \mathcal{L}_{\text{QCD}}^{M=0} + \mathcal{L}_{\text{ext}} = \mathcal{L}_{\text{QCD}}^{M=0} &+ \bar{q}_L \gamma^\mu l_\mu q_L + \bar{q}_R \gamma^\mu r_\mu q_R + \bar{q}_L S q_R + \bar{q}_R S^\dagger q_L \\ &+ \bar{q}_L \sigma^{\mu\nu} t_{\mu\nu} q_R + \bar{q}_R \sigma^{\mu\nu} t_{\mu\nu}^\dagger q_L,\end{aligned}\quad (2.1)$$

where $q = (u, d, s)^T$ is the three-component column vector of the light quarks and the external sources r_μ , l_μ , S , and $t_{\mu\nu}$ are 3×3 matrices in flavor space.² By promoting the sources to spurion fields transforming under chiral rotations, one obtains a QCD Lagrangian which is formally invariant under chiral symmetry, and one then constructs the most general chiral Lagrangian based on this chiral symmetry involving the spurion source fields and the matrix

$$U(x) = \exp\left(i \frac{\pi(x)}{F_0}\right), \quad \pi(x) = \lambda^a \pi^a(x), \quad \text{Tr}(\lambda^a \lambda^b) = 2 \delta^{ab}, \quad (2.2)$$

transforming as $U \rightarrow RUL^\dagger$ under chiral $\text{SU}(3) \times \text{SU}(3)$ rotations. In the above equations, the matrix $\pi(x)$ is defined in terms of the eight Goldstone boson fields $\pi^a(x)$, $a = 1, \dots, 8$, the Gell-Mann matrices λ^a , and the pion decay constant in the chiral limit, F_0 . Green functions of quark bilinears are computed with the chiral Lagrangian via functional derivatives with respect to the sources. Finally, the spurion sources are fixed to the physical values including the contribution of the semileptonic LEFT operators.

Since the semileptonic LEFT contributions to the sources include a factor $1/v^2$ with $v \sim 246 \text{ GeV}$, we are only interested in terms linear in the LEFT sources: quadratic terms

²Compared with the notation in ref. [24], we use $t_{\mu\nu} \rightarrow t_{\mu\nu}^\dagger$ since the LEFT basis uses $\bar{q}_L \sigma^{\mu\nu} q_R$ operators. The usual χ PT scalar source is $\chi = -2B_0 S^\dagger$.

are of the same size as dimension-eight contributions in the LEFT and can be neglected. This applies to weak SM contributions to the LEFT coefficients and even more so to BSM contributions suppressed by $1/\Lambda_{\text{BSM}}^2$. Therefore, we explicitly split the contribution to the spurions into two parts:

$$S \mapsto S + \tilde{S}, \quad r_\mu \mapsto r_\mu + \tilde{r}_\mu, \quad l_\mu \mapsto l_\mu + \tilde{l}_\mu, \quad t_{\mu\nu} \mapsto t_{\mu\nu} + \tilde{t}_{\mu\nu}. \quad (2.3)$$

The spurions S , r_μ , l_μ , and $t_{\mu\nu}$ describe the mass matrix and the coupling to the electromagnetic field. They are fixed in the end to their physical values (in our sign convention the QED covariant derivative is given by $D_\mu = \partial_\mu + ieQA_\mu$):

$$S \mapsto -M_q^\dagger, \quad r_\mu \mapsto -eQA_\mu, \quad l_\mu \mapsto -eQA_\mu, \quad t_{\mu\nu} \mapsto 0, \quad (2.4)$$

where $M_q = \text{diag}(m_u, m_d, m_s)$ is the light-quark mass matrix.

The spurions $\tilde{S}$, $\tilde{r}_\mu$, $\tilde{l}_\mu$, and $\tilde{t}_{\mu\nu}$ contain the contributions from higher-dimensional operators. We only consider terms linear in the dimension-five and dimension-six LEFT operator coefficients, e.g. the first entry in the 3×3 matrix

$$\tilde{S} = \begin{pmatrix} \tilde{S}_{uu} & \tilde{S}_{ud} & \tilde{S}_{us} \\ \tilde{S}_{du} & \tilde{S}_{dd} & \tilde{S}_{ds} \\ \tilde{S}_{su} & \tilde{S}_{sd} & \tilde{S}_{ss} \end{pmatrix} \quad (2.5)$$

is given by

$$\tilde{S}_{uu} = L_{eu}^{S,RL*}(\bar{e}_{Rr}e_{Lp}) + L_{pruu}^{S,RR}(\bar{e}_{Lp}e_{Rr}). \quad (2.6)$$

Note that terms proportional to G_F induced by the SM weak interaction are included in $\tilde{l}_\mu$.

At order p^4 in the chiral counting,³ the matching is easily computed from the chiral Lagrangian by shifting sources as in eq. (2.3), and amounts to the following replacement rules for the LEFT quark bilinears.

Scalar:

$$\begin{aligned} \bar{q}_L \tilde{S} q_R \rightarrow -2B_0 \Bigg[& \frac{1}{4} F_0^2 \langle \tilde{S} U \rangle + L_4 \langle D_\mu U^\dagger D^\mu U \rangle \langle \tilde{S} U \rangle + L_5 \langle \tilde{S} U D_\mu U^\dagger D^\mu U \rangle \\ & + 2L_6 \langle U^\dagger \chi + \chi^\dagger U \rangle \langle \tilde{S} U \rangle - 2L_7 \langle U^\dagger \chi - \chi^\dagger U \rangle \langle \tilde{S} U \rangle + 2L_8 \langle \tilde{S} U \chi^\dagger U \rangle \\ & + H_2 \langle \tilde{S} \chi \rangle \Bigg] + \mathcal{O}(p^6). \end{aligned} \quad (2.7)$$

³We will treat the LEFT sources formally as the same order in chiral counting as the usual chiral sources, $\tilde{S} \sim p^2$, $\tilde{r}_\mu \sim p$, $\tilde{l}_\mu \sim p$, $\tilde{t}_{\mu\nu} \sim p^2$, so that the equations can be compared easily with the χ PT literature.

Vector:

$$\begin{aligned}
 \bar{q}_R \gamma^\mu \tilde{r}_\mu q_R \rightarrow & \frac{i}{2} F_0^2 \langle \tilde{r}^\mu D_\mu U U^\dagger \rangle + 4i L_1 \langle D_\nu U^\dagger D^\nu U \rangle \langle \tilde{r}^\mu D_\mu U U^\dagger \rangle \\
 & + 4i L_2 \langle D^\mu U^\dagger D^\nu U \rangle \langle \tilde{r}_\mu D_\nu U U^\dagger \rangle + 2i L_3 \langle (U^\dagger \tilde{r}_\mu D^\mu U - D^\mu U^\dagger \tilde{r}_\mu U) D_\nu U^\dagger D^\nu U \rangle \\
 & + 2i L_4 \langle \tilde{r}^\mu D_\mu U U^\dagger \rangle \langle U^\dagger \chi + \chi^\dagger U \rangle + i L_5 \langle (U^\dagger \tilde{r}_\mu D^\mu U - D^\mu U^\dagger \tilde{r}_\mu U) (U^\dagger \chi + \chi^\dagger U) \rangle \\
 & + L_9 \left[-\langle \tilde{r}_\mu F_R^{\mu\nu} D_\nu U U^\dagger \rangle - \langle \tilde{r}_\mu U D_\nu U^\dagger F_R^{\mu\nu} \rangle + \langle \tilde{r}_\mu D_\nu U F_L^{\mu\nu} U^\dagger \rangle + \langle \tilde{r}_\mu U F_L^{\mu\nu} D_\nu U^\dagger \rangle \right] \\
 & - i L_9 \langle \tilde{r}^\mu D^\nu (D_\mu U D_\nu U^\dagger - D_\nu U D_\mu U^\dagger) \rangle + 2L_{10} \langle \tilde{r}_\mu D_\nu (U F_L^{\mu\nu} U^\dagger) \rangle \\
 & + 4H_1 \langle \tilde{r}_\mu D_\nu F_R^{\mu\nu} \rangle + \epsilon \text{ terms} + \mathcal{O}(p^6), \tag{2.8}
 \end{aligned}$$

where we have omitted $\mathcal{O}(p^4)$ terms involving $\epsilon_{\alpha\beta\lambda\sigma}$ from the odd-intrinsic-parity (anomalous) sector [25–28].

Tensor:

$$\bar{q}_L \sigma^{\mu\nu} \tilde{t}_{\mu\nu} q_R \rightarrow \Lambda_1 \langle \tilde{t}_{\mu\nu} (U F_L^{\mu\nu} + F_R^{\mu\nu} U) \rangle + i \Lambda_2 \langle \tilde{t}^{\mu\nu} D_\mu U U^\dagger D_\nu U \rangle + \mathcal{O}(p^6). \tag{2.9}$$

Here $F_R^{\mu\nu} = \partial_\mu r_\nu - \partial_\nu r_\mu - i[r_\mu, r_\nu]$ and similarly for $F_L^{\mu\nu}$ with $r_\mu \rightarrow l_\mu$. B_0 , L_i , and H_i are the usual non-perturbative low-energy constants (LECs) that enter the chiral Lagrangian to order p^2 and p^4 . $\Lambda_{1,2}$ are analogous parameters for tensor sources studied in ref. [24]. The matching of $\bar{q}_R S^\dagger q_L$ and $\bar{q}_R t_{\mu\nu}^\dagger q_L$ is given by the Hermitian conjugates of eqs. (2.7), (2.9), and the one of $\bar{q}_L l^\mu \gamma_\mu q_L$ by $r_\mu \rightarrow l_\mu$, $U \leftrightarrow U^\dagger$, $\chi \leftrightarrow \chi^\dagger$ in eq. (2.8). When computing matrix elements to order p^4 , one has to include the contribution from one-loop diagrams with p^2 vertices and tree graphs with p^4 vertices, as usual in χ PT.

The operators on the l.h.s. of eqs. (2.7), (2.8), and (2.9) are renormalized at a scale μ at which perturbation theory is still valid. While the combination $\chi = -2B_0 S^\dagger$ and the LECs L_i are independent of the perturbative scale μ [21], the LECs B_0 and Λ_i are scale dependent [21, 24]. They connect the perturbative scale μ with low-energy hadron dynamics.

We can now match $q^2 \ell^2$ operators in LEFT onto the chiral Lagrangian. Quarks and leptons do not interact if electromagnetic effects are turned off, so we can directly get the matching to lowest order in α from eqs. (2.7)–(2.9). For example, the LEFT operators

$$\mathcal{O}_{eu}^{T,RR} = (\bar{e}_{Lp} \sigma^{\mu\nu} e_{Rr})(\bar{u}_{Lw} \sigma_{\mu\nu} u_{Rt}), \quad \mathcal{O}_{ed}^{T,RR} = (\bar{e}_{Lp} \sigma^{\mu\nu} e_{Rr})(\bar{d}_{Lw} \sigma_{\mu\nu} d_{Rt}) \tag{2.10}$$

are matched onto

$$\begin{aligned}
 \mathcal{O}_{eq}^{T,RR} \rightarrow & \Lambda_1 (\bar{e}_{Lp} \sigma^{\mu\nu} e_{Rr})(U F_L^{\mu\nu} + F_R^{\mu\nu} U)_{tw} \\
 & + i \Lambda_2 (\bar{e}_{Lp} \sigma^{\mu\nu} e_{Rr})(D_\mu U U^\dagger D_\nu U)_{tw} + \mathcal{O}(p^6) \tag{2.11}
 \end{aligned}$$

in χ PT, where $\Lambda_{1,2}$ are the *same* as in eq. (2.9) up to corrections of order α . Since we are matching onto χ PT, eq. (2.11) can be used for lepton flavors $p, r = e, \mu$, and quark flavors $w, t = u, d, s$. Similarly, using eq. (2.7), the matching of the scalar operators

$$\mathcal{O}_{eu}^{S,RR} = (\bar{e}_{Lp} e_{Rr})(\bar{u}_{Lw} u_{Rt}), \quad \mathcal{O}_{ed}^{S,RR} = (\bar{e}_{Lp} e_{Rr})(\bar{d}_{Lw} d_{Rt}), \tag{2.12}$$

is given by

$$\begin{aligned} \mathcal{O}_{eq}^{S,RR} \rightarrow -2B_0(\bar{e}_{Lp}e_{Rr}) & \left[\frac{1}{4}F_0^2 U_{tw} + L_4 \langle D_\mu U^\dagger D^\mu U \rangle U_{tw} + L_5 (U D_\mu U^\dagger D^\mu U)_{tw} \right. \\ & + 2L_6 \langle U^\dagger \chi + \chi^\dagger U \rangle U_{tw} - 2L_7 \langle U^\dagger \chi - \chi^\dagger U \rangle U_{tw} + 2L_8 (U \chi^\dagger U)_{tw} \\ & \left. + H_2 \chi_{tw} \right] + \mathcal{O}(p^6), \quad q = u, d. \end{aligned} \quad (2.13)$$

The matching of $\mathcal{O}_{eu}^{S,RL}$, $\mathcal{O}_{ed}^{S,RL}$ is given by a similar expression with the hadronic part in eq. (2.13) replaced by its hermitian conjugate.

At order α , the $q^2 \ell^2$ operators can no longer be treated as the product of quark and lepton bilinears: e.g. the matching of the tensor operator eq. (2.10) includes operators such as those in eq. (2.13) where the leptonic part is a Lorentz scalar, since both operators transform as $(\bar{\mathbf{3}}, \mathbf{3})$ under chiral $SU(3) \times SU(3)$.

We want to study the effect of $q^2 \ell^2$ operators on $\mu \rightarrow e\gamma$. The matrix element $\langle \gamma(p, \epsilon) | S | 0 \rangle$ of a scalar operator to a physical photon vanishes due to Lorentz and gauge invariance. The matrix element $\langle \gamma(p, \epsilon) | V^\mu | 0 \rangle$ of a vector operator to a physical photon has the form

$$\langle \gamma(p, \epsilon) | V^\mu | 0 \rangle = A(p^2) [p^\mu (p \cdot \epsilon) - p^2 \epsilon^\mu] \quad (2.14)$$

by gauge invariance, and so vanishes for an on-shell photon. Eq. (2.14) has the structure of a penguin amplitude, and does contribute if the virtual photon is attached to external fermion lines, as in $\mu \rightarrow 3e$. The lowest-order version of this argument is a consequence of eq. (2.7) with $U = 1$ having no $F_{\mu\nu}$ term, and eq. (2.8) with $U = 1$ being proportional to $\partial^\nu F_{\mu\nu}$. The absence of a one-photon matrix element for scalar and vector operators holds to all orders in the chiral expansion, but to lowest order in α , since photon exchange between the quark and lepton bilinears can change the Lorentz structure of the bilinears.

Ignoring QED corrections, which can mix Lorentz structure, the only LEFT operators that contribute to $\mu \rightarrow e\gamma$ are tensor operators, which can have a non-zero hadronic matrix element to an on-shell photon:

$$(\bar{e}_{Lp}\sigma^{\mu\nu}e_{Rr})(\bar{q}_L\sigma_{\mu\nu}q_R) \rightarrow -2Q_q e \Lambda_1 (\bar{e}_{Lp}\sigma^{\mu\nu}e_{Rr})F^{\mu\nu} + \mathcal{O}(p^6), \quad q = u, d, s \quad (2.15)$$

where $F^{\mu\nu}$ is the photon field-strength tensor. To study the impact of eq. (2.15), we need the value of the non-perturbative parameter Λ_1 . Naive dimensional analysis (NDA) [7, 29] gives the estimate

$$\Lambda_1 \sim c_T \frac{F_\pi^2}{\Lambda_\chi} = c_T \frac{\Lambda_\chi}{16\pi^2} \quad (2.16)$$

with c_T of order one. Here $F_\pi = 92.3(1)$ MeV is the physical pion decay constant [30], and $F_0 = F_\pi$ to lowest order in the chiral expansion. Eq. (2.16) is the size one would expect from the graph in figure 1 where the loop integral is estimated using the scale Λ_χ . The model estimate of ref. [31] gives $c_T \approx -3.2$. If we use the lattice input from ref. [32] for the pion tensor charge and relate Λ_1 and Λ_2 by vector-meson saturation [33], we obtain

Figure 1. The matrix element $\langle \gamma | \bar{q} \sigma^{\mu\nu} q | 0 \rangle$.

$c_T \approx -1.0(2)$, where the error is due to uncertainties in the vector couplings F_V and G_V and does not include model uncertainties. Similar estimates can be obtained from [34, 35]. From eq. (2.15) we obtain the low-energy matching condition

$$\delta L_{e\gamma}^T = e c_T \frac{F_\pi^2}{\Lambda_\chi} \left[\frac{2}{3} L_{ed}^{T,RR} + \frac{2}{3} L_{e\mu ss}^{T,RR} - \frac{4}{3} L_{e\mu uu}^{T,RR} \right] \quad (2.17)$$

for the additional non-perturbative contribution to the dipole operator, where the LEFT coefficients $L_{eq}^{T,RR}$ are evaluated at a low scale $\mu \sim 2 \text{ GeV}$, and $c_T(\mu)$ is computed from tensor operators renormalized at the same scale. Note that in (2.17), the μ dependence due to gluon corrections cancels in the product of the LEC c_T and the tensor operator coefficients.

Higher-order corrections in the chiral expansion to (2.17) could be calculated [24] but would involve even more low-energy constants. For off-shell photons, the momentum dependence of tensor-current matrix elements can be determined via unitarity and dispersion relations [36].

3 Constraints on electromagnetic dipole interactions

3.1 Lepton-flavor-violating processes

For notational simplicity, we define

$$L_{e\mu}^T = \frac{2}{3} L_{ed}^{T,RR} + \frac{2}{3} L_{e\mu ss}^{T,RR} - \frac{4}{3} L_{e\mu uu}^{T,RR}, \quad L_{\mu e}^T = \frac{2}{3} L_{\mu edd}^{T,RR} + \frac{2}{3} L_{\mu ess}^{T,RR} - \frac{4}{3} L_{\mu euu}^{T,RR}, \quad (3.1)$$

which contribute to $\mu \rightarrow e\gamma$ using eq. (2.17). Eq. (1.4) gives the bound

$$|c_T| \left(|L_{e\mu}^T|^2 + |L_{\mu e}^T|^2 \right)^{1/2} \lesssim 1.65 \times 10^{-5} \text{ TeV}^{-2}, \quad (3.2)$$

using eq. (2.17) as the only contribution to the dipole operator, which translates to $\Lambda_{\text{BSM}} > |c_T|^{1/2} 250 \text{ TeV} \approx 450 \text{ TeV}$, using $|c_T| = 3.2$.

In table 2, we collect the constraints on the dipole and tensor operator coefficients at the hadronic scale $\mu = 2 \text{ GeV}$. Apart from the above constraints from $\mu \rightarrow e\gamma$, the same tensor operators contribute to $\mu \rightarrow e$ conversion in nuclei. Due to their Lorentz structure these operators induce spin-dependent $\mu \rightarrow e$ amplitudes, and do not directly contribute to the A^2 -enhanced spin-independent rate. However, these interactions do indirectly induce a spin-independent amplitude through their contribution to the dipole operators in

eq. (2.17).⁴ As a result, we can compare the limits on the tensor operators from $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion, the resulting bounds from the latter are shown in the last two rows of table 2. Here we used the expressions of refs. [11, 37], which are collected in appendix A, together with the current and projected experimental constraints [38–41]

$$\text{Br}(\text{Au}) \leq 7 \times 10^{-13}, \quad \text{Br}(\text{Al}) \lesssim 10^{-16} \quad (\text{projected}), \quad (3.3)$$

and employed the same hadronic and nuclear input as ref. [11]. The constraints in table 2 are derived by turning on a single operator at a time at the scale $\mu = 2 \text{ GeV}$. In this scenario, the current constraints from $\mu \rightarrow e\gamma$ are stronger than the constraints from $\mu \rightarrow e$ conversion, while the projected bounds are comparable. Note, however, that in a realistic BSM scenario, there is no reason to believe that only a single operator is present at the hadronic scale and in general the constraints have to be applied to the linear combination $L_{e\gamma} + \delta L_{e\gamma}$.

In a second step, we consider the constraints on the operator coefficients at the scale $\mu = M_W$ by taking into account the renormalization-group equations (RGEs) discussed in appendix B. Detailed analyses of these RGEs have been presented in refs. [10, 11, 13, 19, 42–44]. We evaluate the bounds from $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion by turning on single operators at a time at the scale $\mu = M_W$. These constraints almost directly translate to limits on operators in the SMEFT, using the tree-level matching [4],

$$L_{eu}^{S,RR} = -C_{lequ}^{(1)}, \quad L_{eu}^{T,RR} = -C_{lequ}^{(3)}. \quad (3.4)$$

In the case of the tensor operators, the running and mixing from $\mu = M_W$ down to $\mu = 2 \text{ GeV}$ and the non-perturbative matching lead to two effects at the hadronic scale: on the one hand, the tensor operators contribute to the effective dipole operators through the non-perturbative matching effects discussed in section 2, leading to a contribution proportional to c_T . On the other hand, the RGE evolution from $\mu = M_W$ down to $\mu = 2 \text{ GeV}$ induces perturbative mixing contributions to the coefficients of the dipole operators which are independent of c_T . Therefore, the bounds on $\mu \rightarrow e\gamma$ constrain the linear combinations of terms proportional of c_T and independent of c_T as shown in the upper half of table 3. Assuming $c_T = \mathcal{O}(1)$, the non-perturbative contribution dominates for the couplings to up and down quarks, while it is expected to be of the same order of magnitude as the perturbative term for the strange-quark couplings.

In the case of $\mu \rightarrow e$ conversion, the RGEs have more important consequences: the tensor operator coefficients also mix into the coefficients of semileptonic scalar operators $L_{ed,eu}^{S,RR}$ [3, 11]. As discussed in ref. [44], these mixing effects from $\mu = M_W$ down to $\mu = 2 \text{ GeV}$ turn out to be large and give a significant contribution to the spin-independent

⁴It should be noted that the spin-dependent contributions in principle contribute at a similar level as the spin-independent ones induced through the dipole operators. Furthermore, there are additional non-perturbative contributions as discussed in footnote 5. Here we simply take the contributions arising from the dipole operators as an estimate.

$\mu = 2 \text{ GeV}$	$L_{e\mu}^{e\gamma}$	$c_T L_{e\mu uu}^{T,RR}$	$c_T L_{e\mu dd}^{T,RR}$	$c_T L_{e\mu ss}^{T,RR}$
(c) $\text{Br}(\mu \rightarrow e\gamma) \leq 4.2 \times 10^{-13}$	3.7×10^{-11}	1.2×10^{-5}	2.5×10^{-5}	2.5×10^{-5}
(e) $\text{Br}(\mu \rightarrow e\gamma) \lesssim 4 \times 10^{-14}$	1.1×10^{-11}	3.8×10^{-6}	7.6×10^{-6}	7.6×10^{-6}
(c) $\text{Br}(\text{Au}) \leq 7 \times 10^{-13}$	7.6×10^{-10}	2.6×10^{-4}	5.1×10^{-4}	5.1×10^{-4}
(e) $\text{Br}(\text{Al}) \lesssim 1 \times 10^{-16}$	1.1×10^{-11}	3.7×10^{-6}	7.4×10^{-6}	7.4×10^{-6}

Table 2. Current (c) and expected (e) limits from $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion experiments on the coefficients of the dipole and tensor operators at the scale $\mu = 2 \text{ GeV}$. The limits are given in units of TeV^{-2} for the tensor operators, while those for $L_{e\mu}^{e\gamma}$ are in units of TeV^{-1} . They constrain the absolute values of both the listed coefficients as well as the coefficients $L_{\mu e}^{e\gamma}$ and $c_T L_{\mu eq}^{T,RR}$.

amplitude.⁵ As in the case of $\mu \rightarrow e\gamma$, the bounds on $\mu \rightarrow e$ conversion constrain a combination of the couplings at $\mu = M_W$ that are independent of c_T and a non-perturbative contribution proportional to c_T . Due to the large mixing into scalar operators, the non-perturbative pieces are negligible for the couplings to the up and down quarks, while they contribute at the $\mathcal{O}(10\%)$ level for the strange-quark couplings, as shown in the lower half of table 3.

Note that the current constraints from $\mu \rightarrow e$ conversion are of the same order of magnitude as the limits from $\mu \rightarrow e\gamma$, if we use the model estimate $c_T \approx -3$. In this case, the $\mu \rightarrow e$ conversion limits are somewhat stronger for the couplings involving the up and down quarks, while the strange-quark couplings are more stringently constrained by the $\mu \rightarrow e\gamma$ limit. It should be mentioned that these comparisons are currently rather uncertain due to the fact that only NDA and model estimates exist for c_T . In addition, the two observables depend on different linear combinations of the tensor interactions. In particular, the combination of couplings that enters $\mu \rightarrow e\gamma$ only depends on the quark charges, while the spin-independent $\mu \rightarrow e$ conversion rate also involves the matrix elements $\langle N | \bar{q}q | N \rangle$. This implies that the two observables are complementary probes, as one would generally expect BSM physics to generate multiple operators in the LEFT.

3.2 Constraints on flavor-diagonal dipole moments

Analogously to the $\mu \rightarrow e\gamma$ operators in eq. (2.17), flavor-diagonal dipole operators can be induced by $L_{lluu}^{T,RR}$, $L_{lldd}^{T,RR}$, and $L_{llss}^{T,RR}$. These dipole couplings subsequently contribute to the anomalous magnetic moments of the leptons, $\Delta a_l = a_l - a_l^{\text{SM}}$, through their real parts,

⁵ In principle there are additional contributions to the spin-independent amplitude that also appear at $\mathcal{O}(e^2/(4\pi)^2)$. Firstly, a spin-independent $(\bar{N}N)(\bar{\mu}e)$ interaction can be induced in the chiral Lagrangian by the combination of the tensor operator with two electromagnetic currents. In addition, there are chiral loops involving photon exchange around the leading-order spin-dependent $(\bar{N}N)(\bar{\mu}e)$ interaction that is generated by $L_{ed,eu}^{T,RR}$. As these contributions involve unknown LECs we neglect them here and use the RGE-induced contribution as an estimate of the full amplitude.

$\mu = M_W$	$L_{e\mu}^{e\gamma}$	$(c_T - 0.07)L_{e\mu\mu u}^{T,RR}$	$(c_T - 0.15)L_{e\mu dd}^{T,RR}$	$(c_T - 3.1)L_{e\mu ss}^{T,RR}$
(c) $\text{Br}(\mu \rightarrow e\gamma) \leq 4.2 \times 10^{-13}$	3.8×10^{-11}	1.4×10^{-5}	2.8×10^{-5}	2.8×10^{-5}
(e) $\text{Br}(\mu \rightarrow e\gamma) \lesssim 4 \times 10^{-14}$	1.2×10^{-11}	4.4×10^{-6}	8.8×10^{-6}	8.8×10^{-6}
	$L_{e\mu}^{e\gamma}$	$(1 + 5 \times 10^{-3}c_T)L_{e\mu\mu u}^{T,RR}$	$(1 + 5 \times 10^{-3}c_T)L_{e\mu dd}^{T,RR}$	$(1 + 0.1c_T)L_{e\mu ss}^{T,RR}$
(c) $\text{Br}(\text{Au}) \leq 7 \times 10^{-13}$	7.9×10^{-10}	1.6×10^{-6}	3.0×10^{-6}	7.0×10^{-5}
(e) $\text{Br}(\text{Al}) \lesssim 1 \times 10^{-16}$	1.1×10^{-11}	2.0×10^{-8}	4.0×10^{-8}	8.9×10^{-7}

Table 3. Current (c) and expected (e) limits from $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$ conversion experiments on the coefficients of the dipole and tensor operators at the scale $\mu = M_W$. The first and fourth rows indicate the combinations of Wilson coefficients and the LEC c_T that are constrained by each observable. The limits are given in units of TeV^{-2} for the tensor operators, while those for $L_{e\mu}^{e\gamma}$ are in units of TeV^{-1} . The linear combinations $(1 + 5 \times 10^{-3}c_T)$ etc. differ by about 10% between Au and Al.

while the imaginary parts induce electric dipole moments (EDMs), d_l ,

$$\Delta a_l = -4 \frac{m_l}{e} \text{Re } L_{ll}^{e\gamma}, \quad d_l = -2 \text{Im } L_{ll}^{e\gamma}. \quad (3.5)$$

With the most recent measurement of the fine-structure constant [45] a tension of 2.4σ between SM prediction [46] and measurement [47] of the anomalous magnetic moment of the electron is observed:

$$\Delta a_e = -0.88(36) \times 10^{-12}. \quad (3.6)$$

In the case of the muon, there is a long-standing discrepancy of 3–4 σ between the experimental value [48] and the SM prediction, which consists of QED [46, 49, 50], electroweak [51–58], and hadronic [59–65] contributions:⁶

$$\Delta a_\mu = 2.64(0.63)(0.46) \times 10^{-9}. \quad (3.7)$$

Assuming Gaussian distributions, we translate these values to bounds

$$|\Delta a_e| \leq 1.34 \times 10^{-12}, \quad \text{CL} = 90\%, \\ |\Delta a_\mu| \leq 3.64 \times 10^{-9}, \quad \text{CL} = 90\%, \quad (3.8)$$

which imply limits on the tensor operators due to the induced dipole interactions. Assuming $L_i^{T,RR} = 1/\Lambda_{\text{BSM}}^2$ and considering only the contribution of a single operator at $\mu = M_W$, we obtain $\Lambda_{\text{BSM}} \gtrsim 5\text{--}7 \text{ TeV}$ for the tensor couplings to the electron, while for those to the muon we find $\Lambda_{\text{BSM}} \gtrsim 1\text{--}2 \text{ TeV}$. However, these constraints are not competitive with current LHC limits, which, in the case of $L_{eeuu}^{T,RR}$, find $\Lambda_{\text{BSM}} \gtrsim 12 \text{ TeV}$ [80, 81].

⁶A broad effort is being made to corroborate and reduce the hadronic uncertainties in the vacuum-polarization [66–71] and light-by-light-scattering contributions [72–79] using lattice QCD and dispersion relations.

The CP-odd parts of the dipole moments can be constrained by EDM measurements, which, in the case of the muon [82] and tau [83] do not lead to significant limits. The electron EDM is more relevant and is currently most stringently constrained by the recently improved measurements on the paramagnetic ThO molecule [84, 85]. This system receives contributions from the electron EDM as well as from scalar interactions between electrons and nucleons [86–88]

$$\omega_{\text{ThO}} = \left[(120.6 \pm 4.9) \left(\frac{d_e}{10^{-27} e \text{ cm}} \right) + (181.6 \pm 7.3) \times 10^7 \left(\frac{Z}{A} C_S^{(p)} + \frac{A-Z}{A} C_S^{(n)} \right) \right] \text{ mrad/s},$$

$$C_S^{(N)} = -v^2 \text{Im} \left[f_{S,N}^{(u)} \frac{m_N}{m_u} \left(L_{eeuu}^{S,RR} + L_{eeuu}^{S,RL} \right) + \sum_{q=d,s} f_{S,N}^{(q)} \frac{m_N}{m_q} \left(L_{eeqq}^{S,RR} + L_{eeqq}^{S,RL} \right) \right], \quad (3.9)$$

where $f_{S,N}^{(q)}$ are the scalar couplings of the nucleons for which we use the results of refs. [89, 90]. The tensor operators thus contribute to ω_{ThO} non-perturbatively via d_e , while RGE contributions arise through both d_e and C_S (see appendix B for details).⁷ In terms of the couplings at $\mu = M_W$ we have

$$\frac{\omega_{\text{ThO}}}{\text{mrad/s}} = \text{Im} \left[(16 + 4.0 c_T) L_{eeuu}^{T,RR} - (8.0 + 2.0 c_T) L_{eedd}^{T,RR} + (5.7 - 2.0 c_T) L_{eeess}^{T,RR} \right] \times 10^8 \text{ TeV}^2, \quad (3.10)$$

showing that the non-perturbative contributions are of similar size as those induced through the RGEs. Using the experimental upper bound, $\omega_{\text{ThO}} \leq 1.3 \text{ mrad/s}$ [85], we obtain $\Lambda_{\text{BSM}} \gtrsim 5 \times 10^3 \text{ TeV}$ and $\Lambda_{\text{BSM}} \gtrsim 3 \times 10^3 \text{ TeV}$ for the couplings to up and down quarks, respectively, while for the interaction with strange quarks we have $\Lambda_{\text{BSM}} \gtrsim 10^4 \text{ TeV}$.

The EDM of Mercury receives direct contributions from tensor interactions as it is a diamagnetic system. Following the analysis of [91] for the atomic [92] and hadronic [93] input and using the current experimental constraint [94, 95] leads to somewhat stronger constraints on the up and down quark couplings. In particular, for the coupling to the up (down) quark, the mercury EDM leads to $\Lambda_{\text{BSM}} \gtrsim 6(12) \times 10^3 \text{ TeV}$, while it gives rise to a weaker limit on the coupling to the strange quark, $\Lambda_{\text{BSM}} \gtrsim 2 \times 10^3 \text{ TeV}$. This implies that although the tensor interactions contribute to the mercury EDM directly, the contributions to paramagnetic molecular systems can be significant, in particular in the case of strange-quark couplings.

3.3 Comment on dark-matter interactions

A very similar situation to the semileptonic tensor operators arises in the context of dark-matter interactions. For dark matter consisting of weakly interacting massive particles that couple to SM particles only through a heavy mediator, an effective field theory can be used

⁷Analogous to the case of $\mu \rightarrow e$ conversion, there are additional contributions that appear at the same order, $\mathcal{O}(e^2/(4\pi)^2)$. These consist of a spin-independent $(\bar{N}N)(\bar{e}e)$ interaction that is induced by the combination of the tensor operator with two electromagnetic currents, as well as chiral loops involving photon exchange around the leading-order spin-dependent $(\bar{N}N)(\bar{e}e)$ interaction. We again neglect these contributions and use the RGE-induced contribution as an estimate.

to describe the dark matter interaction with SM particles [96–99]. Under the assumption that the dark matter particles carry a conserved quantum number, the dark matter fields always appear in bilinears and give rise to operators that are similar to those in the LEFT. For example, at dimension five the dipole operators

$$\mathcal{L}_{\text{DM}}^{(5)} = C^{(5)} \bar{\chi}_L \sigma^{\mu\nu} \chi_R F_{\mu\nu} + \text{h.c.} \quad (3.11)$$

arise, where χ denotes the dark matter fermion. At dimension six, the EFT contains four-fermion operators $(\bar{\chi}\Gamma\chi)(\bar{q}\Gamma q)$, e.g. the tensor operators

$$\mathcal{L}_{\text{DM}}^{(6)} = C_q^{(6)} (\bar{\chi}_L \sigma^{\mu\nu} \chi_R) (\bar{q}_L \sigma_{\mu\nu} q_R) + \text{h.c.} \quad (3.12)$$

For direct-detection experiments, the interaction of dark matter with nucleons has to be studied. The constraints of chiral symmetry have been analyzed in [97, 100] for the case of vector, axial-vector, scalar, and pseudo-scalar interactions. The tensor operators (3.12) lead to spin-dependent interactions that are not enhanced by A^2 . The leading contribution to the nucleon tensor form factors [101, 102] is given by the tensor charges and requires lattice QCD input [93]. Through perturbative RGE mixing, the tensor operators induce the electromagnetic dipole operators (3.11) [103], which leads to a spin-independent contribution to the cross section [104, 105]. In addition to these perturbative RGE effects, again a non-perturbative contribution is generated in analogy to the semileptonic dipole operators discussed in section 2. Given the similarities to the case of $\mu \rightarrow e\gamma$, the linear combination of the contributions due to the RGEs and those due to c_T is the same as those in the first row of table 3 (where $L_{e\mu}^{\gamma}$ and $L_{ed,eu}^{T,RR}$ are to be identified with $C^{(5)}$ and $C_q^{(6)}$, respectively). As a result, for $c_T = \mathcal{O}(1)$, the contributions of the couplings to up and down quarks will be dominated by the non-perturbative piece, while it is expected to be of similar size as the RGE term for the strange-quark couplings.

Apart from the contribution proportional to c_T , another non-perturbative contribution is given by the coupling of the tensor current to pion exchange between two nucleons or to pion loops at the one-nucleon level.⁸ The coupling of the tensor current to the pion is parametrized by the LEC Λ_2 in (2.9), which can be estimated using the lattice result [32] for the pion tensor charge, $\Lambda_2 = 6.0(3) \text{ MeV}$.

4 Conclusion

In this paper, we have discussed the non-perturbative contributions of semileptonic LEFT operators to the purely leptonic process $\mu \rightarrow e\gamma$. Although scalar and vector operators do not induce this process (without additional QED corrections), $q^2\ell^2$ tensor interactions do generate this decay through a non-perturbative contribution to the leptonic dipole operators, $L_{e\gamma} \sim e(F_\pi^2/\Lambda_\chi)L_{q^2\ell^2}$. To quantify this contribution we have matched the semileptonic tensor operators to χ PT, and estimated the required low-energy constant using NDA and the model estimate of ref. [31]. These estimates show that the non-perturbative

⁸We thank M. Hoferichter for pointing this out.

contribution is expected to dominate over the perturbative RGE effects for the couplings to up and down quarks, while it is of the same order of magnitude for the coupling to the strange quark. The constraints on the scale of BSM physics that can be set using the experimental limit on the $\mu \rightarrow e\gamma$ branching ratio are of the order of $\Lambda_{\text{BSM}} \gtrsim 450 \text{ TeV}$, as discussed in section 3. These limits are of similar size as those obtained from $\mu \rightarrow e$ conversion in nuclei for the couplings to up and down quarks, while they currently provide the most stringent constraints on the couplings to strange quarks.

Acknowledgments

We thank Zvi Bern and the Mani L. Bhaumik Institute at UCLA for a stimulating email announcement that started this investigation, and V. Cirigliano, A. Crivellin, M. Hoferichter, and S. Davidson for useful discussions and comments on the manuscript. This work was supported in part by DOE Grant No. DE-SC0009919. P. S. is supported by a grant of the Swiss National Science Foundation (Project No. P300P2167751).

A $\mu \rightarrow e$ conversion in nuclei

Here we discuss in more detail the contributions of the operators in table 1 to $\mu \rightarrow e$ conversion in nuclei. The rate for this process can be written as [11, eq. (3.5)]

$$\begin{aligned} \Gamma(\mu A \rightarrow e A) = & \frac{m_\mu^3}{4} \left| D_A L_{\mu e}^* \gamma + 4m_\mu \sum_{\substack{q=u,d,s \\ N=n,p}} \left(C_q^{SL} \frac{m_N}{m_q} f_{S,N}^{(q)} S_A^{(N)} + C_q^{VR} f_{V,N}^{(q)} V_A^{(N)} \right) \right|^2 \\ & + \frac{m_\mu^3}{4} \left| D_A L_{e\mu} \gamma + 4m_\mu \sum_{\substack{q=u,d,s \\ N=n,p}} \left(C_q^{SR} \frac{m_N}{m_q} f_{S,N}^{(q)} S_A^{(N)} + C_q^{VL} f_{V,N}^{(q)} V_A^{(N)} \right) \right|^2. \end{aligned} \quad (\text{A.1})$$

Here $f_{S,N}^{(q)}$ and $f_{V,N}^{(q)}$ are hadronic matrix elements, related to the scalar- and vector-charges of the nucleons, respectively. D_A , $S_A^{(N)}$, and $V_A^{(N)}$ involve overlap integrals between the nuclear and lepton wavefunctions and are taken from [11, eq. (3.10)], while $C_q^{SL,SR}$ and $C_q^{VL,VR}$ are the following combinations of couplings:

$$\begin{aligned} C_u^{SR} &= L_{eu}^{S,RR} + L_{e\mu u}^{S,RL}, & C_d^{SR} &= L_{ed}^{S,RR} + L_{e\mu d}^{S,RL}, & C_s^{SR} &= L_{es}^{S,RR} + L_{e\mu s}^{S,RL}, \\ C_u^{VR} &= L_{eu}^{V,RR} + L_{e\mu u}^{V,LR}, & C_d^{VR} &= L_{ed}^{V,RR} + L_{e\mu d}^{V,LR}, & C_s^{VR} &= L_{es}^{V,RR} + L_{e\mu s}^{V,LR}, \\ C_u^{VL} &= L_{eu}^{V,LL} + L_{e\mu u}^{V,LR}, & C_d^{VL} &= L_{ed}^{V,LL} + L_{e\mu d}^{V,LR}, & C_s^{VL} &= L_{es}^{V,LL} + L_{e\mu s}^{V,LR}. \end{aligned} \quad (\text{A.2})$$

The couplings C_q^{SL} can be obtained from C_q^{SR} by replacing the labels $e\mu qq \rightarrow \mu e qq$ and taking the complex conjugate. It is conventional to define a branching ratio through $\text{Br}(A) = \Gamma(\mu A \rightarrow e A) / \Gamma^{\text{capt}}(A)$, where Γ^{capt} is the muon capture rate. We follow ref. [11] for the hadronic [89, 90] and nuclear input [106].

In order to obtain the non-perturbative tensor operator contribution to $\mu \rightarrow e$ conversion, one has to substitute $L_{e\gamma} \mapsto L_{e\gamma} + \delta L_{e\gamma}$ with $\delta L_{e\gamma}$ given in (2.17).

B Anomalous dimensions

The RGEs for the dipole, scalar and tensor operators relevant for $\mu \rightarrow e\gamma$ from refs. [3, 4] are summarized below. We treat $L_{e\gamma} \sim em_\mu/\Lambda_{\text{BSM}}^2$ so that terms quadratic in $L_{e\gamma}$ have been dropped. These equations agree with earlier results in ref. [11]. There are some differences with the results in refs. [10, 42, 43].

$$\begin{aligned}
 \dot{L}_{rs}^{e\gamma} &= \left[(10 - b_{0,e}) e^2 \right] L_{rs}^{e\gamma} + 8e[M_d]_{wv} L_{ed}^{T,RR} - 16e[M_u]_{wv} L_{eu}^{T,RR} - 2e[M_e]_{wv} L_{ee}^{S,RR}, \\
 \dot{L}_{prst}^{S,RR} &= - \left[\frac{26}{3} e^2 + 6g^2 C_F \right] L_{prst}^{S,RR} + 64e^2 L_{eu}^{T,RR}, \\
 \dot{L}_{prst}^{T,RR} &= \frac{4}{3} e^2 L_{prst}^{S,RR} + \left[\frac{26}{9} e^2 + 2g^2 C_F \right] L_{prst}^{T,RR}, \\
 \dot{L}_{prst}^{S,RL} &= - \left[\frac{26}{3} e^2 + 6g^2 C_F \right] L_{prst}^{S,RL}, \\
 \dot{L}_{prst}^{S,RR} &= - \left[\frac{20}{3} e^2 + 6g^2 C_F \right] L_{prst}^{S,RR} - 32e^2 L_{prst}^{T,RR}, \\
 \dot{L}_{prst}^{T,RR} &= - \frac{2}{3} e^2 L_{prst}^{S,RR} + \left[\frac{20}{9} e^2 + 2g^2 C_F \right] L_{prst}^{T,RR}, \\
 \dot{L}_{prst}^{S,RL} &= - \left[\frac{20}{3} e^2 + 6g^2 C_F \right] L_{prst}^{S,RL}, \\
 \dot{L}_{prst}^{S,RR} &= 16e^2 L_{prst}^{S,RR} - 4e^2 L_{prst}^{S,RR}, \tag{B.1}
 \end{aligned}$$

where $\dot{L} = 16\pi^2 \mu \frac{d}{d\mu} L$, $C_F = \frac{4}{3}$, and $b_{0,e} = -\frac{4}{3}(n_e + \frac{1}{3}n_d + \frac{4}{3}n_u)$ is the leading coefficient of the QED β -function. Note that $L_{prst}^{S,RR} = L_{stpr}^{S,RR}$ by symmetry of the operator $\mathcal{O}_{ee}^{S,RR}$. The labels p, r, s, t, w, v in eq. (B.1) run over the flavors active at a given scale μ .

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] MEG collaboration, *New constraint on the existence of the $\mu^+ \rightarrow e^+ \gamma$ decay*, *Phys. Rev. Lett.* **110** (2013) 201801 [[arXiv:1303.0754](https://arxiv.org/abs/1303.0754)] [[INSPIRE](#)].
- [2] MEG collaboration, *Search for the lepton flavour violating decay $\mu^+ \rightarrow e^+ \gamma$ with the full dataset of the MEG experiment*, *Eur. Phys. J. C* **76** (2016) 434 [[arXiv:1605.05081](https://arxiv.org/abs/1605.05081)] [[INSPIRE](#)].
- [3] E.E. Jenkins, A.V. Manohar and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Anomalous Dimensions*, *JHEP* **01** (2018) 084 [[arXiv:1711.05270](https://arxiv.org/abs/1711.05270)] [[INSPIRE](#)].

- [4] E.E. Jenkins, A.V. Manohar and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Operators and Matching*, *JHEP* **03** (2018) 016 [[arXiv:1709.04486](#)] [[INSPIRE](#)].
- [5] W. Buchmüller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl. Phys. B* **268** (1986) 621 [[INSPIRE](#)].
- [6] B. Grzadkowski, M. Iskrzyński, M. Misiak and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085 [[arXiv:1008.4884](#)] [[INSPIRE](#)].
- [7] A. Manohar and H. Georgi, *Chiral Quarks and the Nonrelativistic Quark Model*, *Nucl. Phys. B* **234** (1984) 189 [[INSPIRE](#)].
- [8] A. Crivellin, S. Najjari and J. Rosiek, *Lepton Flavor Violation in the Standard Model with general Dimension-Six Operators*, *JHEP* **04** (2014) 167 [[arXiv:1312.0634](#)] [[INSPIRE](#)].
- [9] G.M. Pruna and A. Signer, *The $\mu \rightarrow e\gamma$ decay in a systematic effective field theory approach with dimension 6 operators*, *JHEP* **10** (2014) 014 [[arXiv:1408.3565](#)] [[INSPIRE](#)].
- [10] S. Davidson, *$\mu \rightarrow e\gamma$ and matching at m_W* , *Eur. Phys. J. C* **76** (2016) 370 [[arXiv:1601.07166](#)] [[INSPIRE](#)].
- [11] A. Crivellin, S. Davidson, G.M. Pruna and A. Signer, *Renormalisation-group improved analysis of $\mu \rightarrow e$ processes in a systematic effective-field-theory approach*, *JHEP* **05** (2017) 117 [[arXiv:1702.03020](#)] [[INSPIRE](#)].
- [12] D. Black, T. Han, H.-J. He and M. Sher, *τ - μ flavor violation as a probe of the scale of new physics*, *Phys. Rev. D* **66** (2002) 053002 [[hep-ph/0206056](#)] [[INSPIRE](#)].
- [13] M. Carpentier and S. Davidson, *Constraints on two-lepton, two quark operators*, *Eur. Phys. J. C* **70** (2010) 1071 [[arXiv:1008.0280](#)] [[INSPIRE](#)].
- [14] A. Celis, V. Cirigliano and E. Passemar, *Lepton flavor violation in the Higgs sector and the role of hadronic τ -lepton decays*, *Phys. Rev. D* **89** (2014) 013008 [[arXiv:1309.3564](#)] [[INSPIRE](#)].
- [15] A. Celis, V. Cirigliano and E. Passemar, *Model-discriminating power of lepton flavor violating τ decays*, *Phys. Rev. D* **89** (2014) 095014 [[arXiv:1403.5781](#)] [[INSPIRE](#)].
- [16] A. Crivellin, M. Hoferichter and M. Procura, *Improved predictions for $\mu \rightarrow e$ conversion in nuclei and Higgs-induced lepton flavor violation*, *Phys. Rev. D* **89** (2014) 093024 [[arXiv:1404.7134](#)] [[INSPIRE](#)].
- [17] A. Crivellin, G. D'Ambrosio, M. Hoferichter and L.C. Tunstall, *Violation of lepton flavor and lepton flavor universality in rare kaon decays*, *Phys. Rev. D* **93** (2016) 074038 [[arXiv:1601.00970](#)] [[INSPIRE](#)].
- [18] D.E. Hazard and A.A. Petrov, *Radiative lepton flavor violating B , D and K decays*, *Phys. Rev. D* **98** (2018) 015027 [[arXiv:1711.05314](#)] [[INSPIRE](#)].
- [19] S. Davidson and A. Saporta, *Constraints on $2\ell 2q$ operators from $\mu \leftrightarrow e$ flavour-changing meson decays*, [arXiv:1807.10288](#) [[INSPIRE](#)].
- [20] S. Weinberg, *Nonlinear realizations of chiral symmetry*, *Phys. Rev.* **166** (1968) 1568 [[INSPIRE](#)].
- [21] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory to One Loop*, *Annals Phys.* **158** (1984) 142 [[INSPIRE](#)].

- [22] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark*, *Nucl. Phys. B* **250** (1985) 465 [INSPIRE].
- [23] H. Georgi, *Weak Interactions and Modern Particle Theory*, Benjamin/Cummings Pub. Co., Menlo Park, U.S.A., (1984).
- [24] O. Catà and V. Mateu, *Chiral perturbation theory with tensor sources*, *JHEP* **09** (2007) 078 [[arXiv:0705.2948](#)] [INSPIRE].
- [25] J. Wess and B. Zumino, *Consequences of anomalous Ward identities*, *Phys. Lett.* **37B** (1971) 95 [INSPIRE].
- [26] E. Witten, *Global Aspects of Current Algebra*, *Nucl. Phys. B* **223** (1983) 422 [INSPIRE].
- [27] A. Manohar and G.W. Moore, *Anomalous Inequivalence of Phenomenological Theories*, *Nucl. Phys. B* **243** (1984) 55 [INSPIRE].
- [28] R. Kaiser and H. Leutwyler, *Large N_c in chiral perturbation theory*, *Eur. Phys. J. C* **17** (2000) 623 [[hep-ph/0007101](#)] [INSPIRE].
- [29] B.M. Gavela, E.E. Jenkins, A.V. Manohar and L. Merlo, *Analysis of General Power Counting Rules in Effective Field Theory*, *Eur. Phys. J. C* **76** (2016) 485 [[arXiv:1601.07551](#)] [INSPIRE].
- [30] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *Phys. Rev. D* **98** (2018) 030001 [INSPIRE].
- [31] V. Mateu and J. Portolés, *Form-factors in radiative pion decay*, *Eur. Phys. J. C* **52** (2007) 325 [[arXiv:0706.1039](#)] [INSPIRE].
- [32] I. Baum, V. Lubicz, G. Martinelli, L. Orifici and S. Simula, *Matrix elements of the electromagnetic operator between kaon and pion states*, *Phys. Rev. D* **84** (2011) 074503 [[arXiv:1108.1021](#)] [INSPIRE].
- [33] G. Ecker, J. Gasser, A. Pich and E. de Rafael, *The Role of Resonances in Chiral Perturbation Theory*, *Nucl. Phys. B* **321** (1989) 311 [INSPIRE].
- [34] O. Catà and V. Mateu, *Novel patterns for vector mesons from the large- N_c limit*, *Phys. Rev. D* **77** (2008) 116009 [[arXiv:0801.4374](#)] [INSPIRE].
- [35] J.A. Miranda and P. Roig, *Effective-field theory analysis of the $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ decays*, *JHEP* **11** (2018) 038 [[arXiv:1806.09547](#)] [INSPIRE].
- [36] M. Hoferichter, B. Kubis, J. Ruiz de Elvira and P. Stoffer, *Nucleon matrix elements of the tensor current*, [[arXiv:1811.11181](#)] [INSPIRE].
- [37] V. Cirigliano, R. Kitano, Y. Okada and P. Tuzon, *On the model discriminating power of $\mu \rightarrow e$ conversion in nuclei*, *Phys. Rev. D* **80** (2009) 013002 [[arXiv:0904.0957](#)] [INSPIRE].
- [38] SINDRUM II collaboration, *A search for muon to electron conversion in muonic gold*, *Eur. Phys. J. C* **47** (2006) 337 [INSPIRE].
- [39] COMET collaboration, *Conceptual design report for experimental search for lepton flavor violating $\mu^- \rightarrow e^-$ conversion at sensitivity of 10^{-16} with a slow-extracted bunched proton beam (COMET)*, KEK-2009-10 (2009).
- [40] R.K. Kutschke, *The Mu2e Experiment at Fermilab*, in *Proceedings, 31st International Conference on Physics in collisions (PIC 2011): Vancouver, Canada, August 28 – September 1, 2011*, [[arXiv:1112.0242](#)] [INSPIRE].

- [41] COMET collaboration, *A search for muon-to-electron conversion at J-PARC: The COMET experiment*, *PTEP* **2013** (2013) 022C01 [[INSPIRE](#)].
- [42] A. Crivellin, S. Davidson, G.M. Pruna and A. Signer, *Complementarity in lepton-flavour violating muon decay experiments*, in *18th International Workshop on Neutrino Factories and Future Neutrino Facilities Search (NuFact16) Quy Nhon, Vietnam, August 21–27, 2016*, 2016, [arXiv:1611.03409](#) [[INSPIRE](#)].
- [43] S. Davidson, Y. Kuno and A. Saporta, “*Spin-dependent*” $\mu \rightarrow e$ conversion on light nuclei, *Eur. Phys. J. C* **78** (2018) 109 [[arXiv:1710.06787](#)] [[INSPIRE](#)].
- [44] V. Cirigliano, S. Davidson and Y. Kuno, *Spin-dependent* $\mu \rightarrow e$ conversion, *Phys. Lett. B* **771** (2017) 242 [[arXiv:1703.02057](#)] [[INSPIRE](#)].
- [45] R.H. Parker, C. Yu, W. Zhong, B. Estey and H. Müller, *Measurement of the fine-structure constant as a test of the standard model*, *Science* **360** (2018) 191.
- [46] T. Aoyama, T. Kinoshita and M. Nio, *Revised and Improved Value of the QED Tenth-Order Electron Anomalous Magnetic Moment*, *Phys. Rev. D* **97** (2018) 036001 [[arXiv:1712.06060](#)] [[INSPIRE](#)].
- [47] D. Hanneke, S. Fogwell and G. Gabrielse, *New Measurement of the Electron Magnetic Moment and the Fine Structure Constant*, *Phys. Rev. Lett.* **100** (2008) 120801 [[arXiv:0801.1134](#)] [[INSPIRE](#)].
- [48] MUON G-2 collaboration, *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)] [[INSPIRE](#)].
- [49] S. Laporta and E. Remiddi, *The analytical value of the electron $g - 2$ at order α^3 in QED*, *Phys. Lett. B* **379** (1996) 283 [[hep-ph/9602417](#)] [[INSPIRE](#)].
- [50] S. Laporta, *High-precision calculation of the 4-loop contribution to the electron $g - 2$ in QED*, *Phys. Lett. B* **772** (2017) 232 [[arXiv:1704.06996](#)] [[INSPIRE](#)].
- [51] R. Jackiw and S. Weinberg, *Weak interaction corrections to the muon magnetic moment and to muonic atom energy levels*, *Phys. Rev. D* **5** (1972) 2396 [[INSPIRE](#)].
- [52] G. Altarelli, N. Cabibbo and L. Maiani, *The Drell-Hearn sum rule and the lepton magnetic moment in the Weinberg model of weak and electromagnetic interactions*, *Phys. Lett.* **40B** (1972) 415 [[INSPIRE](#)].
- [53] I. Bars and M. Yoshimura, *Muon magnetic moment in a finite theory of weak and electromagnetic interaction*, *Phys. Rev. D* **6** (1972) 374 [[INSPIRE](#)].
- [54] K. Fujikawa, B.W. Lee and A.I. Sanda, *Generalized Renormalizable Gauge Formulation of Spontaneously Broken Gauge Theories*, *Phys. Rev. D* **6** (1972) 2923 [[INSPIRE](#)].
- [55] S. Peris, M. Perrottet and E. de Rafael, *Two loop electroweak corrections to the muon $g-2$: A new class of hadronic contributions*, *Phys. Lett. B* **355** (1995) 523 [[hep-ph/9505405](#)] [[INSPIRE](#)].
- [56] A. Czarnecki, B. Krause and W.J. Marciano, *Electroweak Fermion loop contributions to the muon anomalous magnetic moment*, *Phys. Rev. D* **52** (1995) R2619 [[hep-ph/9506256](#)] [[INSPIRE](#)].
- [57] T. Gribouk and A. Czarnecki, *Electroweak interactions and the muon $g-2$: Bosonic two-loop effects*, *Phys. Rev. D* **72** (2005) 053016 [[hep-ph/0509205](#)] [[INSPIRE](#)].

- [58] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, *The electroweak contributions to $(g-2)_\mu$ after the Higgs boson mass measurement*, *Phys. Rev. D* **88** (2013) 053005 [[arXiv:1306.5546](#)] [[INSPIRE](#)].
- [59] J. Prades, E. de Rafael and A. Vainshtein, *The Hadronic Light-by-Light Scattering Contribution to the Muon and Electron Anomalous Magnetic Moments*, *Adv. Ser. Direct. High Energy Phys.* **20** (2009) 303 [[arXiv:0901.0306](#)] [[INSPIRE](#)].
- [60] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, *Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order*, *Phys. Lett. B* **734** (2014) 144 [[arXiv:1403.6400](#)] [[INSPIRE](#)].
- [61] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, *Remarks on higher-order hadronic corrections to the muon $g-2$* , *Phys. Lett. B* **735** (2014) 90 [[arXiv:1403.7512](#)] [[INSPIRE](#)].
- [62] A. Nyffeler, *Hadronic light-by-light scattering in the muon $g-2$* , in *International Workshop on e^+e^- Collisions from Phi to Psi (PHIPSI17) Mainz, Germany, June 26–29, 2017*, [arXiv:1710.09742](#) [[INSPIRE](#)].
- [63] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon $g-2$ and $\alpha(m_Z^2)$ using newest hadronic cross-section data*, *Eur. Phys. J. C* **77** (2017) 827 [[arXiv:1706.09436](#)] [[INSPIRE](#)].
- [64] F. Jegerlehner, *Muon $g-2$ theory: The hadronic part*, *EPJ Web Conf.* **166** (2018) 00022 [[arXiv:1705.00263](#)] [[INSPIRE](#)].
- [65] A. Keshavarzi, D. Nomura and T. Teubner, *Muon $g-2$ and $\alpha(M_Z^2)$: a new data-based analysis*, *Phys. Rev. D* **97** (2018) 114025 [[arXiv:1802.02995](#)] [[INSPIRE](#)].
- [66] B. Ananthanarayan, I. Caprini, D. Das and I. Sentitemsu Imsong, *Precise determination of the low-energy hadronic contribution to the muon $g-2$ from analyticity and unitarity: An improved analysis*, *Phys. Rev. D* **93** (2016) 116007 [[arXiv:1605.00202](#)] [[INSPIRE](#)].
- [67] B. Chakraborty, C.T.H. Davies, P.G. de Oliveira, J. Koponen, G.P. Lepage and R.S. Van de Water, *The hadronic vacuum polarization contribution to a_μ from full lattice QCD*, *Phys. Rev. D* **96** (2017) 034516 [[arXiv:1601.03071](#)] [[INSPIRE](#)].
- [68] M. Della Morte et al., *The hadronic vacuum polarization contribution to the muon $g-2$ from lattice QCD*, *JHEP* **10** (2017) 020 [[arXiv:1705.01775](#)] [[INSPIRE](#)].
- [69] BUDAPEST-MARSEILLE-WUPPERTAL collaboration, *Hadronic vacuum polarization contribution to the anomalous magnetic moments of leptons from first principles*, *Phys. Rev. Lett.* **121** (2018) 022002 [[arXiv:1711.04980](#)] [[INSPIRE](#)].
- [70] RBC and UKQCD collaborations, *Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment*, *Phys. Rev. Lett.* **121** (2018) 022003 [[arXiv:1801.07224](#)] [[INSPIRE](#)].
- [71] G. Colangelo, M. Hoferichter and P. Stoffer, *Two-pion contribution to hadronic vacuum polarization*, [arXiv:1810.00007](#) [[INSPIRE](#)].
- [72] J. Green, O. Gryniuk, G. von Hippel, H.B. Meyer and V. Pascalutsa, *Lattice QCD calculation of hadronic light-by-light scattering*, *Phys. Rev. Lett.* **115** (2015) 222003 [[arXiv:1507.01577](#)] [[INSPIRE](#)].

- [73] A. Gérardin, H.B. Meyer and A. Nyffeler, *Lattice calculation of the pion transition form factor $\pi^0 \rightarrow \gamma^* \gamma^*$* , *Phys. Rev. D* **94** (2016) 074507 [[arXiv:1607.08174](#)] [[INSPIRE](#)].
- [74] T. Blum et al., *Connected and Leading Disconnected Hadronic Light-by-Light Contribution to the Muon Anomalous Magnetic Moment with a Physical Pion Mass*, *Phys. Rev. Lett.* **118** (2017) 022005 [[arXiv:1610.04603](#)] [[INSPIRE](#)].
- [75] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Rescattering effects in the hadronic-light-by-light contribution to the anomalous magnetic moment of the muon*, *Phys. Rev. Lett.* **118** (2017) 232001 [[arXiv:1701.06554](#)] [[INSPIRE](#)].
- [76] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersion relation for hadronic light-by-light scattering: two-pion contributions*, *JHEP* **04** (2017) 161 [[arXiv:1702.07347](#)] [[INSPIRE](#)].
- [77] T. Blum et al., *Using infinite volume, continuum QED and lattice QCD for the hadronic light-by-light contribution to the muon anomalous magnetic moment*, *Phys. Rev. D* **96** (2017) 034515 [[arXiv:1705.01067](#)] [[INSPIRE](#)].
- [78] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S.P. Schneider, *Pion-pole contribution to hadronic light-by-light scattering in the anomalous magnetic moment of the muon*, *Phys. Rev. Lett.* **121** (2018) 112002 [[arXiv:1805.01471](#)] [[INSPIRE](#)].
- [79] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S.P. Schneider, *Dispersion relation for hadronic light-by-light scattering: pion pole*, *JHEP* **10** (2018) 141 [[arXiv:1808.04823](#)] [[INSPIRE](#)].
- [80] S. Alioli, W. Dekens, M. Girard and E. Mereghetti, *NLO QCD corrections to SM-EFT dilepton and electroweak Higgs boson production, matched to parton shower in POWHEG*, *JHEP* **08** (2018) 205 [[arXiv:1804.07407](#)] [[INSPIRE](#)].
- [81] R. Gupta, Y.-C. Jang, B. Yoon, H.-W. Lin, V. Cirigliano and T. Bhattacharya, *Isovector Charges of the Nucleon from 2+1+1-flavor Lattice QCD*, *Phys. Rev. D* **98** (2018) 034503 [[arXiv:1806.09006](#)] [[INSPIRE](#)].
- [82] MUON (G-2) collaboration, *An Improved Limit on the Muon Electric Dipole Moment*, *Phys. Rev. D* **80** (2009) 052008 [[arXiv:0811.1207](#)] [[INSPIRE](#)].
- [83] BELLE collaboration, *Search for the electric dipole moment of the τ lepton*, *Phys. Lett. B* **551** (2003) 16 [[hep-ex/0210066](#)] [[INSPIRE](#)].
- [84] ACME collaboration, *Order of Magnitude Smaller Limit on the Electric Dipole Moment of the Electron*, *Science* **343** (2014) 269 [[arXiv:1310.7534](#)] [[INSPIRE](#)].
- [85] ACME collaboration, *Improved limit on the electric dipole moment of the electron*, *Nature* **562** (2018), 355.
- [86] M. Denis and T. Fleig, *In search of discrete symmetry violations beyond the standard model: Thorium monoxide reloaded*, *J. Chem. Phys.* **145** (2016) 214307.
- [87] L.V. Skripnikov, A.N. Petrov and A.V. Titov, *Communication: Theoretical study of ThO for the electron electric dipole moment search*, *J. Chem. Phys.* **139** (2013) 221103.
- [88] L.V. Skripnikov, A.N. Petrov and A.V. Titov, *Theoretical study of ThO for electron electric dipole moment search*, *J. Chem. Phys.* **139** (2013) 221103 [[arXiv:1308.0414](#)].
- [89] M. Hoferichter, J. Ruiz de Elvira, B. Kubis and U.-G. Meißner, *High-Precision Determination of the Pion-Nucleon σ Term from Roy-Steiner Equations*, *Phys. Rev. Lett.* **115** (2015) 092301 [[arXiv:1506.04142](#)] [[INSPIRE](#)].

- [90] P. Junnarkar and A. Walker-Loud, *Scalar strange content of the nucleon from lattice QCD*, *Phys. Rev. D* **87** (2013) 114510 [[arXiv:1301.1114](#)] [[INSPIRE](#)].
- [91] W. Dekens, J. de Vries, M. Jung and K.K. Vos, *The phenomenology of electric dipole moments in models of scalar leptoquarks*, [arXiv:1809.09114](#) [[INSPIRE](#)].
- [92] T. Fleig and M. Jung, *Model-independent determinations of the electron EDM and the role of diamagnetic atoms*, *JHEP* **07** (2018) 012 [[arXiv:1802.02171](#)] [[INSPIRE](#)].
- [93] R. Gupta, B. Yoon, T. Bhattacharya, V. Cirigliano, Y.-C. Jang and H.-W. Lin, *Flavor diagonal tensor charges of the nucleon from $(2+1+1)$ -flavor lattice QCD*, *Phys. Rev. D* **98** (2018) 091501 [[arXiv:1808.07597](#)] [[INSPIRE](#)].
- [94] B. Graner, Y. Chen, E.G. Lindahl and B.R. Heckel, *Reduced Limit on the Permanent Electric Dipole Moment of ^{199}Hg* , *Phys. Rev. Lett.* **116** (2016) 161601 [Erratum *ibid.* **119** (2017) 119901] [[arXiv:1601.04339](#)] [[INSPIRE](#)].
- [95] W.C. Griffith, M.D. Swallows, T.H. Loftus, M.V. Romalis, B.R. Heckel and E.N. Fortson, *Improved Limit on the Permanent Electric Dipole Moment of ^{199}Hg* , *Phys. Rev. Lett.* **102** (2009) 101601 [[INSPIRE](#)].
- [96] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T.M.P. Tait and H.-B. Yu, *Constraints on Dark Matter from Colliders*, *Phys. Rev. D* **82** (2010) 116010 [[arXiv:1008.1783](#)] [[INSPIRE](#)].
- [97] M. Hoferichter, P. Klos and A. Schwenk, *Chiral power counting of one- and two-body currents in direct detection of dark matter*, *Phys. Lett. B* **746** (2015) 410 [[arXiv:1503.04811](#)] [[INSPIRE](#)].
- [98] F. Bishara, J. Brod, B. Grinstein and J. Zupan, *From quarks to nucleons in dark matter direct detection*, *JHEP* **11** (2017) 059 [[arXiv:1707.06998](#)] [[INSPIRE](#)].
- [99] J. Brod, A. Gootjes-Dreesbach, M. Tamaro and J. Zupan, *Effective Field Theory for Dark Matter Direct Detection up to Dimension Seven*, *JHEP* **10** (2018) 065 [[arXiv:1710.10218](#)] [[INSPIRE](#)].
- [100] V. Cirigliano, M.L. Graesser and G. Ovanessian, *WIMP-nucleus scattering in chiral effective theory*, *JHEP* **10** (2012) 025 [[arXiv:1205.2695](#)] [[INSPIRE](#)].
- [101] S. Weinberg, *Charge symmetry of weak interactions*, *Phys. Rev.* **112** (1958) 1375 [[INSPIRE](#)].
- [102] S.L. Adler et al., *Renormalization Constants for Scalar, Pseudoscalar and Tensor Currents*, *Phys. Rev. D* **11** (1975) 3309 [[INSPIRE](#)].
- [103] U. Haisch and F. Kahlhoefer, *On the importance of loop-induced spin-independent interactions for dark matter direct detection*, *JCAP* **04** (2013) 050 [[arXiv:1302.4454](#)] [[INSPIRE](#)].
- [104] T. Banks, J.-F. Fortin and S. Thomas, *Direct Detection of Dark Matter Electromagnetic Dipole Moments*, [arXiv:1007.5515](#) [[INSPIRE](#)].
- [105] V. Barger, W.-Y. Keung and D. Marfatia, *Electromagnetic properties of dark matter: Dipole moments and charge form factor*, *Phys. Lett. B* **696** (2011) 74 [[arXiv:1007.4345](#)] [[INSPIRE](#)].
- [106] R. Kitano, M. Koike and Y. Okada, *Detailed calculation of lepton flavor violating muon electron conversion rate for various nuclei*, *Phys. Rev. D* **66** (2002) 096002 [Erratum *ibid.* **D 76** (2007) 059902] [[hep-ph/0203110](#)] [[INSPIRE](#)].