RECEIVED: November 4, 2020

REVISED: December 28, 2020

ACCEPTED: January 4, 2021

PUBLISHED: February 17, 2021

Algebra of quantum $\mathcal{C}$ -polynomials

Andrei Mironov^{a,b,c} and Alexei Morozov^{d,b,c}

^a*Lebedev Physics Institute,
Moscow 119991, Russia*

^b*Institute for Theoretical and Experimental Physics (ITEP),
Moscow 117218, Russia*

^c*Institute for Information Transmission Problems,
Moscow 127994, Russia*

^d*Moscow Institute of Physics and Technology (MIPT),
Dolgoprudny, 141701, Russia*

E-mail: mironov@lpi.ru, morozov@itep.ru

ABSTRACT: Knot polynomials colored with symmetric representations of $SL_q(N)$ satisfy difference equations as functions of representation parameter, which look like quantization of classical $\mathcal{A}$ -polynomials. However, they are quite difficult to derive and investigate. Much simpler should be the equations for coefficients of differential expansion nicknamed quantum $\mathcal{C}$ -polynomials. It turns out that, for each knot, one can actually derive two difference equations of a finite order for these coefficients, those with shifts in spin n of the representation and in $A = q^N$. Thus, the $\mathcal{C}$ -polynomials are much richer and form an entire ring. We demonstrate this with the examples of various defect zero knots, mostly discussing the entire twist family.

KEYWORDS: Chern-Simons Theories, Wilson, 't Hooft and Polyakov loops, Topological Field Theories

ARXIV EPRINT: [2009.11641](https://arxiv.org/abs/2009.11641)

Contents

1	Introduction	1
2	Equations on knot polynomials: generalities	4
2.1	Motivation: from equations on characters to $\mathcal{C}$ -polynomials	4
2.2	Non-local difference equations	5
2.3	Local difference equations: $\mathcal{C}$ -polynomials	7
2.4	Genus expansion and the special point	7
3	$\mathcal{C}$-polynomials for twist knots: examples	8
3.1	DE coefficients for twist and double-braid knots	8
3.2	$\text{twist}_{-1} = 4_1$	9
3.3	$\text{twist}_1 = 3_1$	9
3.4	$\text{twist}_{-2} = 6_1$	9
3.5	$\text{twist}_2 = 5_1$ and $k \longleftrightarrow -k$ duality	11
3.6	$\text{twist}_{-3} = 8_1$	11
4	Algebra of $\mathcal{C}$-polynomials	12
5	General formulas for the entire twist family	14
5.1	Twist knots: n -evolution	15
5.2	Twist knots: A -evolution	15
5.3	Twist knots: mixed n - A -evolutions	16
6	Another defect zero knot	16
7	Conclusion	18
A	A-evolution for twist knots	19
A.1	Examples at low k	19
A.2	General answer	20
A.3	Examples of $G_{n,j}^{(k)}$	21
B	Details on knot 9_{46}	23

1 Introduction

Wilson loop averages are the main observables in Yang-Mills theories, and it is very important to understand the equations they satisfy. Best studied are the Ward identities, i.e. direct implications of quantized equations of motion, which in the case of Wilson averages

are known under the name of loop equations [1–13] (in the case of the eigenvalue matrix models, they are nothing but the celebrated Virasoro and W-constraints). However, there is also a complementary set of relations reflecting the representation theory background behind the Yang-Mills theory, and these are studied much worse because they are associated with Wilson loops in non-fundamental representations, which are only starting to attract attention within generic Yang-Mills context [14, 15]. The biggest progress on this way is achieved in the 3d Chern-Simons theory [16, 17], which is topological, and hence does not possess any dynamical loop equations so that the pure representation theory equations are the only ones which remain. Wilson loop averages in this case are known as *knot polynomials* [18–25], and equations in question are called *non-commutative (quantum) $\mathcal{A}$ -polynomials* [26].

The $\mathcal{A}$ -polynomials have a number of applications, basically to and around Kashaev large- r limit [27–36]. However, this is only a beginning of the story. Being *exactly* calculable non-perturbative quantities, both the HOMFLY-PT polynomials and $\mathcal{A}$ -polynomials provide a nice possibility of studying the basic questions in quantum field theory, from integrability [37] to topological expansions [38–42] or the resurgent structure [43, 44].

In fact, the $\mathcal{A}$ -polynomials appear to be very complicated and difficult to study in any systematic way. The way out was proposed almost from the very beginning in [45]: it suggests to look not at the knot polynomials themselves, but at what is now known as the coefficients of their differential expansion (DE) [46–50]. The quantum $\mathcal{C}$ -polynomials made out of these coefficients in a more direct way than the usual knot polynomials (Wilson averages) were first studied in [45] but then attracted surprisingly low attention (most probably because of a limited knowledge about the DE coefficients and the lacking relation to other popular subjects though their classical limit is related by a rational map with the classical $\mathcal{A}$ -polynomials). *In this paper, we return to the abandoned story of $\mathcal{C}$ -polynomials and consider them in the general case of HOMFLY-PT polynomials.* The original considerations [26, 45, 51–54] were mostly restricted to the Jones polynomials, which are functions only of the parameter $q = e^{2\pi i/(\kappa+2)}$, κ being the Chern-Simons coupling constant, and of the representation, and describe Wilson averages in the Chern-Simons theory with the gauge group $SU(2)$. On the contrary, the HOMFLY-PT polynomials are functions of parameters $q = e^{2\pi i/(\kappa+N)}$, $A = q^N$, and of the representation, and describe Wilson averages in the Chern-Simons theory with the gauge group $SU(N)$. Because of an additional parameter A , there emerge additional difference equations with respect to shifts of A (additional $\mathcal{C}$ -polynomials), and we discuss the full ring of these $\mathcal{C}$ -polynomials, which opens new approaches and perspectives. Being polynomials of A and q , HOMFLY-PT are easily analytically continued to arbitrary complex values of these parameters, and we do not address here the interesting subject of specific representations arising when they are roots of unity, in application to $\mathcal{C}$ -polynomials this will be considered elsewhere.

The original quantum $\mathcal{A}$ -polynomial [26] was introduced to describe a difference equation w.r.t. representation label for the unreduced Jones polynomials of knot $\mathcal{K}$

$$J_{[r]}^{\mathcal{K}} = \sum_{n=0}^r \frac{[r+n+1]!}{[r-n]!} \cdot \{q\}^{2n} \cdot F_{[n]}^{\mathcal{K}}(A, q) \Big|_{A=q^2} \quad (1.1)$$

where $F_{[n]}$ are the coefficients of differential expansion of the unreduced HOMFLY-PT polynomials colored with symmetric representation $[r]$,

$$H_{[r]}^{\mathcal{K}}(A, q) = \sum_{n=0}^r Z_{[r]}^{[n]}(A, q) \cdot F_{[n]}^{\mathcal{K}}(A, q)$$

$$Z_{[r]}^{[n]}(A, q) := \text{Bin}_q(r, n) \cdot \prod_{i=1}^n \{Aq^{r+i-1}\} \cdot \{Aq^{i-2}\} \quad (1.2)$$

at the specialization to the Jones polynomials $A = q^2$. The notation used in these formulas can be found at the end of the Introduction.

This linear transformation of the Jones polynomial, (1.1) was first considered in [55] at q being roots of unity, hence the name for the DE coefficients cyclotomic function (though the number of elements in the product is not related to degree of the root of unity). In general, however, the products like $\prod_i \{Aq^{i-2}\}$ do not look like cyclic polynomials at all, instead the factors are nothing but the DGR “differentials” (related also to those in Khovanov-Rozansky calculus) [46]. Therefore the expansion in Z -factors (1.2) was named “differential” in [47]. The knot-independent Z -factors for symmetric representations were found in [56] and [47] (they are actually known for all rectangular representations for defect-zero knots [57–59], for the situation with other defects and non-rectangular case, see [50] and references therein).

Less known than the $\mathcal{A}$ -polynomials, the quantum $\mathcal{C}$ -polynomials [45]¹ describe equations for the DE coefficients $F_{[n]}$ themselves and are not restricted to the Jones polynomials. Instead they should be formulated differently for knots with different defects [60] (which are defined by the powers of the corresponding Alexander polynomials), because the factorization properties of the differential expansion depends on it. With the increase of defect, the Z -factors get smaller (and are halved in the limit of large defects), while the DE coefficients become more involved and acquire additional sub-structures [49], which will probably show up at the level of $\mathcal{C}$ -polynomials as well. In this paper we concentrate on the examples of defect-zero knots only and, more important, on symmetric representations, i.e. on representations associated with the single-line Young diagrams $[r]$.

In [49], we suggested that the DE coefficients can be a better characteristic of knot than the original knot polynomials. They have no direct interpretation in Chern-Simons theory and can instead be a base for alternative effective theory of knots, of which the examples were attempted in [61–63] and [64]. This direction seems also related to the Khovanov calculus [65–69, 72–74] and its far-going generalizations [75–81]. Unfortunately, study of the DE coefficients remains very difficult, it is still based more on particular examples and computer experiments. This is true for quantum $\mathcal{C}$ -polynomials as well: the search for them remains more a piece of art than a regular method.

The symmetric DE coefficients $F_{[n]}^{\mathcal{K}}(A, q)$ for a given knot $\mathcal{K}$ are functions of three variables A , q , and n , and therefore would satisfy three different equations. However, we

¹S. Garoufalidis et al. introduced $\mathcal{A}$ -polynomials [26] and $\mathcal{C}$ -polynomials [45]. We do not know what the name “ $\mathcal{B}$ -polynomials” was reserved for, but clearly the properties of knot polynomials is rich enough to find an appropriate candidate.

restrict ourselves to two equations and treat q as a *parameter* rather than a *variable*. As usual, each equation can be considered as a *quantum polynomial*, and, for every knot, the two non-commuting quantum polynomials generate a non-trivial algebra with operator-valued structure “constants”, which is the main object of our study in this paper. It is quite intriguing that such simple quantities as the DE coefficients for the twist knots are associated with highly non-trivial quantum polynomials and algebras.

The paper is organized as follows. In section 2, we discuss general existing equations for knot invariants. In section 3, we consider first examples of the equations for a few first terms of the family of twist knots. In section 4, we continue considering examples with a discussion of rings that are formed by $\mathcal{C}$ -polynomials. In section 5, we describe the $\mathcal{C}$ -polynomials for the entire twist knot family, while, in section 6, obtain the $\mathcal{C}$ -polynomials in the simplest example of non-twist (and non-double-braid) knot of still zero defect, knot 9_{46} . Section 7 contains some concluding remarks. The most tedious formulas are collected in two appendices.

Notation. First of all, all knot polynomials are unreduced and given in the topological framing.

As soon as throughout the paper we deal with the symmetric representations only, we denote the DE coefficients instead of $F_{[n]}$ just as F_n .

The quantum numbers are defined as $[m] := \frac{\{q^m\}}{\{q\}} = q^{1-m}[[m]]$ with $\{z\} := z - z^{-1}$ and $[[m]] = 1 + q^2 + \dots + q^{2(m-1)}$.

The q -binomial coefficients are denoted as $Bin_q(n, k) = \frac{[n]!}{[k]![n-k]!}$.

With each representation μ , we associate the Young diagram $\mu : \mu_1 \geq \mu_2 \geq \dots \geq \mu_l > 0$, and $|\mu| := \sum_{i=1}^l \mu_i$, where l is the number of lines in the Young diagram μ , which we often denote l_μ .

2 Equations on knot polynomials: generalities

2.1 Motivation: from equations on characters to $\mathcal{C}$ -polynomials

We now present one more a kind of motivation for studying A and $\mathcal{C}$ -polynomials, which is often ignored in the literature.

The irreducible representation of SL_N group associated with the Young diagram R is described by the character $\chi_R\{p\}$, which depends on traces of powers of the SL_N group element in the fundamental representation, $p_n = \text{Tr } g^n$ and is the Schur polynomial. These characters form a closed algebra

$$\chi_{R_1}\{p\} \cdot \chi_{R_2}\{p\} = \sum_{R \in R_1 \otimes R_2} N_{R_1 R_2}^R \cdot \chi_R\{p\} \quad (2.1)$$

where $N_{R_1 R_2}^R$ are known as Littlewood-Richardson coefficients or structure constants. Alternatively, one can say that each χ_R induces a linear operator acting on characters:

$$\hat{\mathcal{O}}_R : \chi_Q \longrightarrow \sum_{Q' \in R \otimes Q} N_{RQ}^{Q'} \cdot \chi_{Q'} \quad (2.2)$$

If the character variables are restricted to the topological locus

$$p_k^* := \frac{\{A^k\}}{\{q^k\}} = \frac{[Nk]}{[k]} \quad (2.3)$$

with $A := q^N$, one gets an action on the space of quantum dimensions $D_R(N) := \chi_R \{p_k^*\}$. If we further restrict ourselves to $N = 2$ (i.e. SL_2), we get an action on the space of symmetric representations only,

$$\hat{\mathcal{O}}_{[r]} : D_{[n]}(2) \longrightarrow \sum_{m=0}^r D_{[n-r+2m]}(2) \quad (2.4)$$

Moving all the terms in (2.1) to the l.h.s., one obtains a system of linear equations for characters: an equation per each R . The simplest of them, associated with $R = [1]$ is just

$$(q + q^{-1})D_{[n]}(2) - D_{[n+1]}(2) - D_{[n-1]}(2) = 0, \quad \text{i.e.} \quad (q + q^{-1})[n+1] - [n+2] - [n] = 0 \quad (2.5)$$

Knot polynomials are non-trivial deformations of characters (or, at least, of quantum dimensions) [37], moreover, they provide an entire *family* of deformations labeled by knots (or, perhaps, better to say, by braids). As time goes we find more and more demonstration of this analogy, which forces us to take it seriously. In particular, quite some time ago, the colored Jones polynomials (the colored HOMFLY-PT polynomials at $N = 2$) were demonstrated to satisfy some linear equations [51–54] nicknamed non-commutative (quantum) $\mathcal{A}$ -polynomials [26], which can be thought of some analogy of the above linear equations for characters. It was also realized that the equations actually simplify if written not on the knot polynomials themselves, but on the coefficients of their differential (also called cyclotomic) expansion, these equations were nicknamed $\mathcal{C}$ -polynomials [45].

An extension to the generic HOMFLY-PT case is pretty difficult, because, as in the case of characters, the equations involve not only symmetric representations but also other Young diagrams with no more than $N - 1$ lines. However, it still can be studied in the case of HOMFLY-PT polynomials colored with symmetric representations for particular knots [56, 82, 83]. Similarly to the $N = 2$ case, the $\mathcal{A}$ -polynomials turns out to be much more involved as compared to $\mathcal{C}$ -polynomials. In particular, while the $\mathcal{A}$ -polynomials for the whole series of twist knots are unknown, it is possible to construct $\mathcal{C}$ -polynomials for them, as we demonstrate it in the present paper.

Since the DE coefficients depend on *two* variables: $[n]$ (representation) and $A = q^N$, one can actually expect *pairs* of equations emerging, and, since they do not necessarily commute, actually a whole algebra of equations: an algebra of $\mathcal{C}$ -polynomials. The simplest examples of the $\mathcal{C}$ -polynomials and of their algebras is the subject of the present paper.

2.2 Non-local difference equations

Knot-independent sum rules. We begin with a kind of trivial or, if one prefers, of universal $\mathcal{C}$ -polynomial, which is almost independent of the knot, and therefore deserve the name of *sum rules*. The point is that the DE coefficients $F_n^{\mathcal{K}}(A, q)$ are Laurent polynomials

in A^2 and q^2 , and their degree in A^2 is often determined by the *special* polynomial,² which is the specialization of the HOMFLY-PT polynomial at $q = 1$ trivially depending on the representation [85]:

$$H_R^K(A, q = 1) = \left(H_{[1]}^K(A, q = 1) \right)^{|R|} = \left(1 + \{A\}^2 F_1^K(A, q = 1) \right)^{|R|}$$

$$\xrightarrow{(1.2)} F_n^K(A, q = 1) = \left(F_1^K(A, q = 1) \right)^n \quad (2.6)$$

If $F_1^K(A, q) = \sum_{m=m_1}^{m_2} c_m(q) A^{2m}$, then $F_n^K(A, q = 1)$ has powers of A^2 in between nm_1 and nm_2 .

Now, since

$$\sum_{j=0}^{M+1} (-x)^j \cdot \frac{[M+1]!}{[j]![M+1-j]!} = \prod_{i=0}^M (q^{M-2i} - x) \quad (2.7)$$

the sum

$$\sum_{j=0}^{M+1} \frac{(-q^{2m})^j}{q^{(2p+1)Mj}} \cdot \frac{[M+1]!}{[j]![M+1-j]!} = \prod_{i=0}^M \left(q^{M-2i} - q^{2m-(2p+1)M} \right) \quad (2.8)$$

vanishes for all m in between $nm_1 = pM$ and $nm_2 = (p+1)M$. These boundaries coincide with nm_1 and nm_2 when $M = n(m_2 - m_1)$ and $2p+1 = \frac{m_2+m_1}{m_2-m_1} := \nu$. This means that whenever ν is odd integer,³

$$\frac{m_2 + m_1}{m_2 - m_1} \in \mathbb{Z}_{\text{odd}} \implies \sum_{j=0}^{M+1} \frac{(-)^j}{q^{(m_2+m_1)nj}} \cdot \frac{[M+1]!}{[j]![M+1-j]!} \cdot F_n(q^j A, q) = 0 \quad (2.9)$$

with $M = n(m_1 - m_2)$

The number of items in these sum rules depends on representation size n , we call such equations *non-local*. An important feature of this kind of equation is that it does not depend on the knot, but only on the minimal and maximal degrees of its special polynomial.

Knot-dependent semi-local equations. We also distinguish *semi-local* equations, which are certainly non-local since the numbers of items in them grow with n , but slower than n , say, like $n/2$: these are already restrictive but not too much as compared to explicit expressions for F_n . However, the semi-local equations essentially depends on knot details. We will consider examples of the both types of non-local equations in the next sections.

²For instance, this is the case for the fundamental HOMFLY-PT polynomial for all 250 knots with up to 10 crossings from the Rolfsen table [84] for exception of 19 knots with non-trivial DE coefficient, 6 knots with $F_1^K(A, q = 1) = 0$ (knots 8_{14} , 10_{82} , 10_{108} , 10_{116} , 10_{118} , 10_{146}) and 38 knots with $F_1^K(A, q = 1) \sim A^k$ (where k is an integer).

³This property is also often fulfilled for knots. For instance, this is the case for all knots with 5 and 6 crossings from the Rolfsen table [84], for 2 knots with 7 crossings, etc. In particular, this is the case for twist knots with negative twist, when $\nu = -1$, while for the positive twist k $\nu = \frac{k+1}{k-1}$. For a generic knot, ν is not odd, and is not even integer.

2.3 Local difference equations: $\mathcal{C}$ -polynomials

More interesting are *local* relations, where the number of terms does not depend on n . There are basically two types of local equations: those involving items at different n at the same A (we call them n -evolution equations), and those involving shifted A at the same n (we call them A -evolution equations). Having these two types of equations, one can certainly generate many other local equations of a mixed type with both n - and A -evolution presented. However, all these new equations are corollaries of the basic two.

n -evolution equations. The n -evolution equations were first considered in [26, 51–54] in the case of Jones polynomials. In [26], it was proposed that a proper classical limit of these equations (for the Jones polynomials) coincide with the classical $\mathcal{A}$ -polynomial (the hypothesis known as AJ-conjecture). This is why these equations have been called non-commutative $\mathcal{A}$ -polynomials. Their generalization to equations for the HOMFLY-PT polynomials colored with symmetric representations (and further to the superpolynomials) were considered in [82, 83, 86].

A -evolution equations. In contrast with the n -evolution, the A -evolution equations were first considered later [56], since they require changing A , and, hence, cannot be formulated for the Jones polynomials, only for HOMFLY-PT ones, while the first example of the generic symmetric HOMFLY-PT polynomial was obtained in [56]. In [56], there is only mixed evolution presented, and, in this paper, we present the pure A -evolution.

In the next section, we demonstrate all these types of equations in the simplest case of first twist knots, and, in section 6, in example of a simple non-twist knot 9_{46} .

2.4 Genus expansion and the special point

Expansions of the HOMFLY polynomials can be performed in a few different ways: one can parameterize $q := e^{\hbar}$, $A := e^{N\hbar}$ and consider the limit of $\hbar \rightarrow 0$ (such an expansion gives rise to the Vassiliev invariants [87]), one can alternatively keep A fixed in the same limit. This expansion is called genus expansion [88–90], we keep the representation in this limit fixed. When the size of the representation infinitely grows consistently with $\hbar^{-1}$, one obtains the volume conjecture limit [27–30] of the Jones polynomial, and, for the equations, this limit gives rise to either the classical $\mathcal{A}$ -polynomial [26], or to the augmentation variety [91], depending on details.

The genus expansion can be performed both in equations and in the polynomials. In particular, the leading term of expansion of the polynomial is, as follows from (2.6),

$$\left(1 + F_1^{\mathcal{K}}(q=1)\{A\}^2\right)^r = \sum_n F_n^{\mathcal{K}}(q=1) \cdot \frac{r!}{n!(r-n)!} \{A\}^{2n} \quad (2.10)$$

which explains a natural appearance of binomial coefficients in their explicit expressions. However, the quantization, i.e. switching on of $q \neq 1$, or taking into account higher genera can be quite nontrivial.

The genus expansion of equations will be discussed later.

3 $\mathcal{C}$ -polynomials for twist knots: examples

In this section, we provide some details about the DE coefficients for first few knots of the twisted family. This is the case when the DE coefficients are best studied, even far beyond the symmetric representations but still not in full generality [47, 50, 92]. It is especially important because it provides exclusive Racah matrices $\bar{S}$ and S , needed for arborescent calculus and its further generalizations [61–63].

3.1 DE coefficients for twist and double-braid knots

For the twist knots, there is a general formula for symmetric DE coefficients [47]:

$$F_n^{\text{twist}_k} = q^{n(n-1)/2} A^n \sum_{j=0}^n (-)^j \frac{[n]!}{[j]![n-j]!} \frac{\{Aq^{2j-1}\} \cdot (Aq^{j-1})^{2jk}}{\prod_{i=j-1}^{n+j-1} \{Aq^i\}} \quad (3.1)$$

It follows from evolution in k and can be rewritten as

$$F_n^{\text{twist}_k} = \sum_m (\mathcal{B}^{k+1})_{[n],[m]} \quad (3.2)$$

i.e. made from elements of the KNTZ matrix [58, 59]

$$\mathcal{B}_{\lambda,\mu}^{-1} = \frac{\chi_{\lambda/\mu}^{(0)} \chi_{\mu}^{(0)}}{\chi_{\lambda}^{(0)}} \cdot \Lambda_{\mu}^{-1} \quad (3.3)$$

evaluated at the locus $p_k^{(0)} = \frac{1}{\{q^k\}}$. A enters entirely through $\Lambda_{\mu} = \left(A^{|\mu|} q^{\kappa(\mu)}\right)^2$, which is proportional to the square of eigenvalue of the corresponding R -matrix. Here $\kappa(\mu) := \sum_{i=1}^{l_{\mu}} \mu_i(\mu_i - 2i + 1)$.

One of the properties of the twist knot family is that their DE coefficients for positive and negative twists are related by a simple formula:

$$\frac{F_n^{\text{twist}_k}(A, q)}{F_n^{\text{twist}_1}(A, q)} = \frac{F_n^{\text{twist}_{-k}}(A^{-1}, q^{-1})}{F_n^{\text{twist}_{-1}}(A^{-1}, q^{-1})} \quad (3.4)$$

Since

$$F_n^{4_1} = F_n^{\text{twist}_{-1}} = 1, \quad F_n^{3_1} = F_n^{\text{trefoil}} = F_n^{\text{twist}_1} = (-)^n A^{2n} q^{n(n-1)} \quad (3.5)$$

formula (3.4) becomes

$$F_n^{\text{twist}_k}(A, q) = (-)^n A^{2n} q^{n(n-1)} \cdot F_n^{\text{twist}_{-k}}(A^{-1}, q^{-1}) \quad (3.6)$$

From any of these expressions, the DE coefficients are easily calculated and described in the third way as multiple q -binomial sums. Indeed, introduce a function

$$\mathcal{P}_1(n) = \sum_{j=0}^n A^{-2j} q^{-j(j+n-2)} \frac{[n]!}{[n-j]![j]!} \quad (3.7)$$

and its iterations

$$\mathcal{P}_m(n) = \sum_{j=0}^n \mathcal{P}_{m-1}(j) \cdot A^{-2j} q^{-j(j+n-2)} \frac{[n]!}{[n-j]![j]!} \quad (3.8)$$

Then

$$F_n^{\text{twist}_{-k}} = \mathcal{P}_{k-1}(n) \quad (3.9)$$

For the double braids, there is a mysterious product formula [57]

$$F_n^{\text{db}_{k,l}} = \frac{F_n^{\text{twist}_k} F_n^{\text{twist}_l}}{F_n^{\text{trefoil}}} \quad (3.10)$$

It would be interesting to understand what this means for the $\mathcal{C}$ -polynomials.

3.2 twist₋₁ = 4₁

For the figure-eight knot 4₁, all the DE coefficients $F_n^{\text{twist}_{-1}}(A, q) = 1$, and all the equations are trivial.

3.3 twist₁ = 3₁

For the trefoil 3₁, the DE coefficients are not quite unities, still they are simple monomials:

$$F_n^{\text{twist}_1}(A, q) = (-)^n A^{2n} q^{n(n-1)} \quad (3.11)$$

hence, the n - and A -evolution equations are rather trivial:

$$\begin{aligned} F_{n+1}^{\text{twist}_1}(A, q) &= -A^2 q^{2n} \cdot F_n^{\text{twist}_1}(A, q) \\ F_n^{\text{twist}_1}(Aq, q) &= q^{2n} F_n^{\text{twist}_1}(A, q) \end{aligned} \quad (3.12)$$

3.4 twist₋₂ = 6₁

Truly non-trivial equations emerge for the first time in the case of 6₁ knot, where

$$F_n^{\text{twist}_{-2}}(A, q) = \sum_{j=0}^n \frac{[n]!}{[j]![n-j]!} \cdot \frac{1}{A^{2j} q^{j(n+j-2)}} = \sum_{j=0}^n \frac{[[n]]!}{[[j]]![[n-j]]!} \cdot \frac{1}{(Aq^{n-1})^{2j}} \quad (3.13)$$

To avoid possible confusions from now on, we *omit the argument* q of $F_n(A, q)$, when it does not matter (is the same in all the terms of equation).

As we discussed in section 2, the DE coefficients $F_n^{\text{twist}_{-2}}(A, q)$ satisfy a set of local and non-local difference equations:

n -evolution:

$$\begin{aligned} F_{n+1}^{\text{twist}_{-2}}(A) &- \left(1 + \frac{1}{A^2 q^{2n}}\right) F_n^{\text{twist}_{-2}}(A) \\ &= q^{-2}(1 - q^{-2n}) \left(\left(1 - \frac{1}{A^2 q^{2n-2}}\right) \cdot F_n^{\text{twist}_{-2}}(A) - F_{n-1}^{\text{twist}_{-2}}(A) \right) \end{aligned} \quad (3.14)$$

A-evolution:

$$F_n^{\text{twist}-2}(A) = \left(1 - \frac{1}{A^2 q^{2n}}\right) F_n^{\text{twist}-2}(Aq) + \frac{1}{A^2} F_n^{\text{twist}-2}(Aq^2) \quad (3.15)$$

In the genus expansion $q = e^h$ with $h \rightarrow 0$, the first equation is reduced to $F_{n+1}^{\text{twist}-2} = (1 + \frac{1}{A^2}) \cdot F_n^{\text{twist}-2}$, while the second one implies $A \frac{\partial F_n^{\text{twist}-2}}{\partial A} = -2n \frac{F_n^{\text{twist}-2}}{A^{2+1}}$. This is obviously true for arbitrary n for the special polynomial $F_n^{\text{twist}-2} \Big|_{q=1} = \left(F_1^{\text{twist}-2} \Big|_{q=1}\right)^n = (1 + \frac{1}{A^2})^n$, which is, indeed, the case for 6_1 knot.

Non-local eq. As follows from (3.13), one has to choose $m_1 = -1$ and $m_2 = 0$ in (2.9) so that the non-local sum rule is

$$\sum_{j=0}^{n+1} (-)^j q^{nj} \cdot \frac{[n+1]!}{[j]![n+1-j]!} \cdot F_n^{\text{twist}-2}(Aq^j) = 0 \quad (3.16)$$

Semi-local eq. In order to obtain a semi-local equation, one has to combine the A -shifts with n -shifts (otherwise, the equations are the above local equations). The leading dependence of F_n on n is typically $\sim A^{\alpha n} q^{\beta n^2}$, i.e. the power of A is linear in n , while that of q is quadratic in n . This means that an infinitesimal/continuous equation would be of the diffusion type $\partial/\partial \log A \sim \partial^2/\partial(\log q)^2$. A finite-difference lift of the diffusion equation is more involved. Still, such equations exist. Looking at the last representation in (3.13), it is clear that one could make a q -shift $q \rightarrow q^2$ and compensate it by an A -shift $A \rightarrow q^{1-n} A$ modulo the change of binomial coefficient $\frac{[n]!}{[j]![n-j]!}$. This change, however, is absent for the “boundary” terms with $j = 0$ and $j = n$, which leads to

$$\begin{aligned} & F_n^{\text{twist}-2}(Aq^{1-n}, q^2) - F_n^{\text{twist}-2}(A, q) \\ &= \sum_{j=1}^{n-1} \left(\frac{[n]!}{[j]![n-j]!} \Big|_{q=q^2} - \frac{[n]!}{[j]![n-j]!} \Big|_{q=q} \right) \cdot \frac{1}{(Aq^{n-1})^{2j}} \end{aligned} \quad (3.17)$$

with the sum at the r.h.s. can be expressed through $F_{n'}^{\text{twist}-2}$ with $n' \leq n-2$. After some algebra, one can finally obtain a semi-local equation

$$F_n^{\text{twist}-2}(A, q) = \sum_{j=0}^{n/2} \frac{(-)^j (q^2 - 1)^j}{A^{2j} q^{3j(j-1)}} \frac{[n]!}{[n-2j]![2j]!!} \cdot F_{n-2j}^{\text{twist}-2}(Aq^{4j+1-n}, q^2) \quad (3.18)$$

It contains an n -dependent sum, but it is two times shorter than n , and hence is a restrictive relation for the DE coefficients F_n . Note that the double factorial appears in denominator so that the weight in the sum deviates from the ordinary q -binomial coefficient.

The local pair (3.14), (3.15) provides a set of independent equations for the particular case of the knot 6_1 . In addition, there are plenty of other *local* equations where both n and A are

shifted. It is natural to assume they are all corollaries of the first two, which is confirmed in section 4. Note that we deal only with two equations (3.14) and (3.15), though the DE coefficients depends on the three parameters n , A and q : it seems more natural to treat q , which specifies the A -shift (or, perhaps better, the N -shift) as a parameter rather than an independent third variable. This is also justified by the fact that the q -shifts satisfy only *semi-local* equations.

3.5 twist₂ = 5₁ and $k \longleftrightarrow -k$ duality

The DE coefficients themselves are analytical functions of k described by the same formulas at positive and negative k . This is a manifestation of $k \longleftrightarrow -k$ duality for the twist family. However, the equations are slightly different for positive and negative k , (3.6). This can be a sign of their relation to superpolynomials and Khovanov calculus [65–74] which are also known to slightly break $k \longleftrightarrow -k$ symmetry [47, 93–95].

In particular, in the case of knot twist₂ = 5₂:

$$\begin{aligned} F_{n+1}^{\text{twist}_2}(A, q) + (q^{2n} + A^2 q^{6n}) \cdot F_n^{\text{twist}_2}(A, q) \\ = (q^{2n} - 1) \left(A^2 q^{2n+2} F_n^{\text{twist}_2}(A, q) + q^{4n} F_{n-1}^{\text{twist}_2}(A, q) \right) \\ F_n^{\text{twist}_2}(A, q) = \left(\frac{1}{q^{2n}} - A^2 \right) F_n^{\text{twist}_2}(Aq, q) + \frac{A^2}{q^{4n}} F_n^{\text{twist}_2}(Aq^2, q) \end{aligned} \quad (3.19)$$

Since $F_n^{\text{twist}_2}(A, q) = A^2(1 + A^2)$, the sum rule (2.9) is

$$\sum_{j=0}^{n+1} \frac{(-)^j}{q^{3nj}} \cdot \frac{[n+1]!}{[j]![n+1-j]!} \cdot F_n^{\text{twist}_2}(Aq^j) = 0 \quad (3.20)$$

3.6 twist₋₃ = 8₁

As an intermediate step from section 3 to generic formulas, we list explicit expressions for one more twist knot, twist₋₃ = 8₁.

n -evolution:

$$\begin{aligned} F_{n+1}^{\text{twist}_{-3}}(A) - \left(1 + \frac{1}{A^2 q^{2n}} + \frac{1}{A^4 q^{4n}} \right) \cdot F_n^{\text{twist}_{-3}}(A) = \\ = q^{-3}(1 - q^{-2n}) \left\{ \left([2] - \frac{q^{2-3n}(q^{n+1} - q^{-n-1})}{A^2} - \frac{q^{3-5n}(q^n + q^{-n})}{A^4} \right) \cdot F_n^{\text{twist}_{-3}}(A) + \right. \\ \left. + \left(-q^{-1}[3] + q^{-2n-1} - \frac{[2]}{q^{4n-2}A^2} \right) \cdot F_{n-1}^{\text{twist}_{-3}}(A) + q^{-3}(1 - q^{2-2n}) \cdot F_{n-2}^{\text{twist}_{-3}}(A) \right\} \end{aligned} \quad (3.21)$$

A -evolution:

$$\begin{aligned} (q^{2n+2}A^2 - 1) \cdot A^6 q^{4n+2} \cdot F_n^{\text{twist}_{-3}}(A) - \\ (q^{2n}A^2 - 1) \left(A^2 q^{2n+2} + q^{2n+2} - 1 \right) \cdot A^4 q^{2n+2} \cdot F_n^{\text{twist}_{-3}}(Aq) - \\ - (q^{2n+2}A^2 - 1) \left(1 + q^{2n}A^2(q^{2n+2} - 1) \right) \cdot F_n^{\text{twist}_{-3}}(Aq^2) + \\ + (q^{2n}A^2 - 1) \cdot q^{4n} \cdot F_n^{\text{twist}_{-3}}(Aq^3) = 0 \end{aligned} \quad (3.22)$$

Non-local eq.: As usual, there is a knot-independent non-local sum rule (2.9), which, in this case, reads

$$\sum_{j=0}^{2n+1} (-)^j q^{2nj} \cdot \frac{[2n+1]!}{[j]![2n+1-j]!} \cdot F_n^{\text{twist}-3}(Aq^j) = 0 \quad (3.23)$$

4 Algebra of $\mathcal{C}$ -polynomials

Twist -1 knot. Introduce two operators $\hat{\Delta}_n, \hat{D}_A$, acting on any function $f_n(A)$ of two variables by the rules

$$\begin{aligned} \hat{\Delta}_n \cdot f_n(A) &= f_{n+1}(A) \\ \hat{D}_A \cdot f_n(A) &= f_n(Aq) \end{aligned} \quad (4.1)$$

Then the equations for $F_n^{\text{twist}-1}(A) = 1$ are just

$$\begin{cases} (\hat{\Delta}_n - 1) \cdot F_n^{\text{twist}-1}(A) = 0 \\ (\hat{D}_A - 1) \cdot F_n^{\text{twist}-1}(A) = 0 \end{cases} \quad (4.2)$$

and the two operators $\hat{\mathcal{O}}_1^{\text{twist}-1} := (\hat{\Delta}_n - 1)$, $\hat{\mathcal{O}}_2^{\text{twist}-1} := (\hat{D}_A - 1)$ obviously commute,

$$[\hat{\mathcal{O}}_1^{\text{twist}-1}, \hat{\mathcal{O}}_2^{\text{twist}-1}] = 0 \quad (4.3)$$

Twist -2 knot. Similarly, the equations (3.14) and (3.15) for twist -2 can be rewritten as

$$\begin{cases} \hat{\mathcal{O}}_1^{\text{twist}-2} \cdot F_n^{\text{twist}-2}(A) = 0 \\ \hat{\mathcal{O}}_2^{\text{twist}-2} \cdot F_n^{\text{twist}-2}(A) = 0 \end{cases} \quad (4.4)$$

$$\hat{\mathcal{O}}_1^{\text{twist}-2} := \hat{\Delta}_n - \left[\left(1 + \frac{1}{A^2 q^{2n+2}} \right) + \frac{1}{q^2} \left(1 - \frac{1}{q^{2n+2}} \right) \left(1 - \frac{1}{A^2 q^{2n}} \right) \right] \cdot \hat{\Delta}_n + \frac{1}{q^2} \left(1 - \frac{1}{q^{2n+2}} \right) \quad (4.5)$$

$$\hat{\mathcal{O}}_2^{\text{twist}-2} := 1 - \left(1 - \frac{1}{A^2 q^{2n}} \right) \cdot \hat{D}_A - \frac{1}{A^2} \cdot \hat{D}_A^2 \quad (4.6)$$

The commutator of these operators is

$$\begin{aligned} [\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_2^{\text{twist}-2}] &= [2] \frac{\{q\}}{A^2 q^{2n+2}} \cdot \hat{\Delta}_n \cdot \hat{D}_A \cdot \left(\hat{\Delta}_n - 1 - \frac{1}{A^2 q^{2n}} \hat{D}_A \right) \\ &= -[2] \frac{\{q\}}{A^2 q^{2n+2}} \cdot \hat{\Delta}_n \cdot \hat{D}_A \cdot \hat{\mathcal{O}}_3^{\text{twist}-2} \end{aligned} \quad (4.7)$$

Thus, we generated a new operator

$$\hat{\mathcal{O}}_3^{\text{twist}-2} := -\frac{A^2 q^{2n-2}}{\{q^2\}} \cdot \hat{\Delta}_n^{-1} \cdot \hat{D}_A^{-1} \cdot [\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_2^{\text{twist}-2}] = \hat{\Delta}_n - 1 - \frac{1}{A^2 q^{2n}} \hat{D}_A \quad (4.8)$$

Its commutation relations

$$\begin{aligned} \left[\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_3^{\text{twist}-2} \right] &= \frac{\{q\}}{A^2 q^{2n+6}} \hat{\Delta}_n \cdot \left(\left[qA^2 + q^4 [2] \hat{D}_A - \frac{[2]}{q^{2n-2}} \right] \hat{\mathcal{O}}_3^{\text{twist}-2} - \frac{[2]}{q^{2n-2}} \hat{\mathcal{O}}_2^{\text{twist}-2} \right) \\ \left[\hat{\mathcal{O}}_2^{\text{twist}-2}, \hat{\mathcal{O}}_3^{\text{twist}-2} \right] &= \frac{\{q\}}{A^2 q^{2n+1}} \hat{D}_A \cdot \left(\hat{\mathcal{O}}_2^{\text{twist}-2} + \hat{\mathcal{O}}_3^{\text{twist}-2} \right) \end{aligned} \quad (4.9)$$

do not involve any new operators, thus the constraint algebra gets closed. Everything what one can generate is a corollary of (3.14) and (3.15).

For instance, the newly generated operator $\hat{\mathcal{O}}_3^{\text{twist}-2}$ gives rise to the equation

$$F_{n+1}^{\text{twist}-2}(A) = F_n^{\text{twist}-2}(A) + \frac{1}{A^2 q^{2n}} F_n^{\text{twist}-2}(Aq) \quad (4.10)$$

while the combination

$$\left(\frac{1}{q^{2n}} - 1 - \frac{1}{q^{2n+2}} \right) \hat{\mathcal{O}}_2^{\text{twist}-2} + (\hat{D}_A - 1) \cdot \hat{\mathcal{O}}_3^{\text{twist}-2} \quad (4.11)$$

produces

$$F_n^{\text{twist}-2}(A) = F_n^{\text{twist}-2}(Aq) + \frac{1 - q^{-2n}}{A^2} F_{n-1}^{\text{twist}-2}(Aq^2) \quad (4.12)$$

These (4.10) and (4.12) imply a whole variety of other *local* equations, where both n and A shifted:

$$F_{n+1}^{\text{twist}-2}(A) = F_n^{\text{twist}-2}(Aq) + \frac{1}{A^2} F_n^{\text{twist}-2}(Aq^2) \quad (4.13)$$

$$F_n^{\text{twist}-2}(Aq^2) - F_{n-1}^{\text{twist}-2}(Aq^2) = \frac{1}{q^{2n}} \left(F_n^{\text{twist}-2}(Aq) - F_{n-1}^{\text{twist}-2}(Aq^2) \right) \quad (4.14)$$

$$F_{n+1}^{\text{twist}-2}(A) = F_n^{\text{twist}-2}(A) + \frac{1}{A^2 q^{2n}} F_n^{\text{twist}-2}(Aq^2) + \frac{1 - q^{-2n}}{A^4 q^{2n+2}} F_{n-1}^{\text{twist}-2}(Aq^3) \quad (4.15)$$

and so on.

The twist -2 analogue of the commutativity (4.3) is now a more sophisticated algebra of D -module constraints

$$\begin{aligned} \left[\hat{\mathcal{O}}_1^{\text{twist}-2}, \left[\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_2^{\text{twist}-2} \right] \right] &= \hat{\alpha}_1 \hat{\mathcal{O}}_1^{\text{twist}-2} + \hat{\beta}_1 \hat{\mathcal{O}}_2^{\text{twist}-2} + \hat{\gamma}_1 \left[\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_2^{\text{twist}-2} \right] \\ \left[\left[\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_2^{\text{twist}-2} \right], \hat{\mathcal{O}}_2^{\text{twist}-2} \right] &= \hat{\alpha}_2 \hat{\mathcal{O}}_1^{\text{twist}-2} + \hat{\beta}_2 \hat{\mathcal{O}}_2^{\text{twist}-2} + \hat{\gamma}_2 \left[\hat{\mathcal{O}}_1^{\text{twist}-2}, \hat{\mathcal{O}}_2^{\text{twist}-2} \right] \end{aligned} \quad (4.16)$$

with operator-valued coefficients (standing at the left of $\hat{\mathcal{O}}$).

Twist $-k$ knots for higher k . It is natural to conjecture that, for the twist $-k$, we will get a closure at the k -th commutator level, but we did not get enough evidence to support it, despite the form of the “generators” $\hat{\mathcal{O}}_1^{\text{twist}-k}$ and $\hat{\mathcal{O}}_2^{\text{twist}-k}$ is explicitly known.

The genus zero limit. Consider the genus expansion of algebra at the vicinity of the special polynomial point. As we emphasized after eqs. (3.14)–(3.15), in this limit,

$$\begin{aligned}\hat{\mathcal{O}}_1^{\text{twist}_{-2}} &:= \hat{\Delta}_n - 1 - \frac{1}{A^2} \\ \hat{\mathcal{O}}_2^{\text{twist}_{-2}} &:= \left(1 + \frac{1}{A^2}\right) \cdot \frac{\partial}{\partial A} + \frac{2n}{A^3}\end{aligned}\quad (4.17)$$

Commutator of these two operators gives

$$\left[\hat{\mathcal{O}}_1^{\text{twist}_{-2}}, \hat{\mathcal{O}}_2^{\text{twist}_{-2}}\right] = \frac{2}{A^3} \left(\hat{\Delta}_n - 1 - \frac{1}{A^2}\right) = \frac{2}{A^3} \hat{\mathcal{O}}_1^{\text{twist}_{-2}} \quad (4.18)$$

which is, indeed, consistent with the genus zero part of operator (4.8)

$$\hat{\mathcal{O}}_3^{\text{twist}_{-2}} = \hat{\Delta}_n - 1 - \frac{1}{A^2} = \hat{\mathcal{O}}_1^{\text{twist}_{-2}} \quad (4.19)$$

Similarly, in the case of $k = -3$, the two operators in the leading order of the genus expansion are

$$\begin{aligned}\hat{\mathcal{O}}_1^{\text{twist}_{-3}} &:= \hat{\Delta}_n - \left(1 + \frac{1}{A^2} + \frac{1}{A^4}\right) \\ \hat{\mathcal{O}}_2^{\text{twist}_{-3}} &:= \left(1 + \frac{1}{A^2} + \frac{1}{A^4}\right) \cdot \frac{\partial}{\partial A} + \frac{2n}{A^3} \left(1 + \frac{2}{A^2}\right)\end{aligned}\quad (4.20)$$

so that

$$\left[\hat{\mathcal{O}}_1^{\text{twist}_{-3}}, \hat{\mathcal{O}}_2^{\text{twist}_{-3}}\right] = \frac{2}{A^3} \left(1 + \frac{2}{A^2}\right) \hat{\mathcal{O}}_1^{\text{twist}_{-3}} \quad (4.21)$$

At generic $k \in \mathbb{Z}_{>0}$,

$$\begin{aligned}\hat{\mathcal{O}}_1^{\text{twist}_{-k}} &:= \hat{\Delta}_n - \frac{1}{A^{2(k-1)}} \frac{A^{2k} - 1}{A^2 - 1} \\ \hat{\mathcal{O}}_2^{\text{twist}_{-k}} &:= \frac{1}{A^{2(k-1)}} \frac{A^{2k} - 1}{A^2 - 1} \cdot \frac{\partial}{\partial A} + \frac{2n}{A^3} \frac{A^{2(k+1)} + k - A^2(1+k)}{A^{2(k-1)}(1 - A^2)^2}\end{aligned}\quad (4.22)$$

so that

$$\left[\hat{\mathcal{O}}_1^{\text{twist}_{-k}}, \hat{\mathcal{O}}_2^{\text{twist}_{-k}}\right] = \frac{2}{A^3} \frac{A^{2(k+1)} + k - A^2(1+k)}{A^{2(k-1)}(1 - A^2)^2} \hat{\mathcal{O}}_1^{\text{twist}_{-k}} \quad (4.23)$$

Thus, one can see that, in the limit of $q \rightarrow 1$, the algebra of $\mathcal{C}$ -polynomials becomes the very simple triangular Lie algebra, however, as we demonstrated above, its q -deformation is quite sophisticated.

5 General formulas for the entire twist family

From examples of twist knots in section 3, we come here to the general structure of twisted $\mathcal{C}$ -polynomials for the entire family.

5.1 Twist knots: n -evolution

Though equations for positive and negative twists are related by the simple formula (3.6), here we write down the both cases to compare with the $A = q^2$ answers of [45]. Let k always be positive, then

$$\begin{aligned}
 F_{n+k}^{\text{twist}-k}(A, q) &= \sum_{j=0}^{k-1} \frac{(-1)^{j+k+1}}{\{q\}^{j-k+1}} \cdot \left(V_{n,j}^{(-,k)} - q^{-2(k+n)} \cdot V_{n+1,j-1}^{(-,k-1)} \right) \cdot F_{n+j}^{\text{twist}-k}(A, q) \\
 &\text{with} \\
 V_{n,j}^{(-,k)} &:= q^{-\frac{1}{2}(k-1)(3k+2n) - \frac{1}{2}j(3j+2k+6n-5)} \cdot \frac{[n+k-1]!}{[n+j]!} \cdot \sum_{i=0}^j \frac{q^{(4n+2k+2j-3)i}}{A^{2(j-i)}} \cdot \frac{[k-i-1]![k-j+i]!}{[k-j-1]![k-j]![j-i]![i]!} \\
 &\text{and} \\
 F_{n+k}^{\text{twist}k}(A, q) &= \sum_{j=0}^{k-1} \frac{(-1)^{j+k}}{\{q\}^{j-k+1}} \cdot \left(V_{n,j}^{(+,k)} - q^{2(k+n)} V_{n+1,j-1}^{(+,k-1)} \right) \cdot F_{n+j}^{\text{twist}k}(A, q) \\
 &\text{with} \\
 V_{n,j}^{(+,k)} &:= q^{\frac{1}{2}(k-1)(5k+2n)+2kn - \frac{1}{2}j(3j+2k+6n-3)} \cdot \frac{[n+k-1]!}{[n+j]!} \\
 &\quad \cdot \sum_{i=0}^j \frac{q^{(4n+2k+2j-3)i}}{A^{2(j-i-k)}} \cdot \frac{[k-i]![k-j+i-1]!}{[k-j-1]![k-j]![j-i]![i]!}
 \end{aligned} \tag{5.1}$$

The number of items in these relations depend on k (i.e. on the knot), but not on representation size n , therefore these are, indeed, *local* difference equations.

5.2 Twist knots: A -evolution

We gave earlier in section 3 examples of A -evolution for $k = -2$ and $k = -3$. The general structure is

$$\begin{aligned}
 0 &= \sum_{j=0}^{k-1} \epsilon_j G_{n,j}^{(k)} \cdot F_n^{\text{twist}-k}(Aq^j) + \epsilon_k \cdot q^{2(k-1)n} \cdot \prod_{i=0}^{\frac{k-3}{2}} \left(A^2 q^{2n+2i} - 1 \right) \cdot F_n^{\text{twist}-k}(Aq^k) \\
 &\text{where } \epsilon_j = \sqrt{2} \cdot \cos\left(\frac{(2j+1)\pi}{4}\right) \text{ and} \\
 G_{n,j}^{(k)} &= \left(A^2 q^{2n+2j-2} - 1 \right) \cdot \prod_{i=0}^{\text{floor}(\frac{j-3}{2})} \left(A^2 q^{2n+2i} - 1 \right) \cdot \prod_{i=\text{floor}(\frac{k+j}{2})}^{k-2} \left(A^2 q^{2n+2i} - 1 \right) \times \\
 &\quad \times A^{k^2 - (j+1)k + 4\text{frac}(\frac{k}{2})\text{frac}(\frac{j}{2})} q^{2(k-j-1)n + \sigma_{k,j}} \cdot g_{n,j}^{(k)}
 \end{aligned} \tag{5.2}$$

Here $g_{n,j}^{(k)}$ is a polynomial⁴ in A^2 , q^{2n} and q^2 . Note that, in variance with the n -evolution equations (5.1), equations (5.2) form two different series for odd and even k , since the polynomials $g_{n,j}^{(k)}$ differ in these series, see examples of these polynomials and of the numerical coefficients σ_k in appendix A.

Equation (5.2) (or (A.1)–(A.4) in appendix A) can be considered as a recursion in m for the coefficients $F_{n|m}^{\text{twist}-k}$ in $F_n^{\text{twist}-k} = \sum_{m \geq 0} F_{n|m}^{\text{twist}-k} A^{-2m}$. For instance, (A.1) implies that

$$F_{n|m+1}^{\text{twist}-2} = \frac{q^{2n-2m} - 1}{q^{2n-2}(q^{2m+2} - 1)} \cdot F_{n|m}^{\text{twist}-2} \tag{5.3}$$

⁴Its degree in A^2 is $\text{floor}(\frac{k-1}{2}) + 2\text{frac}(\frac{j+1}{2})\text{frac}(\frac{k+1}{2})$ and in q^{2n} is j .

5.3 Twist knots: mixed n - A -evolutions

In fact, equations with mixed n - A -evolutions like that first considered in [56], for instance, eqs. (4.12), (4.10), (4.13)–(4.15), etc. can be also generalized to the whole twist series. For instance, eq. (4.15) would look like

$$F_{n+1}^{\text{twist}-k}(A) = F_n^{\text{twist}-k}(A) + \sum_{j=1}^{k-2} \frac{1}{A^{2j}q^{2n}} F_n^{\text{twist}-k}(Aq) + \frac{1-\delta_{k,1}}{A^{2(k-1)}q^{2n}} F_n^{\text{twist}-k}(Aq^2) + \\ + \sum_{j=1}^{\min(k-1,n)} \frac{(-)^{j-1} c_j^{(k)} \{q\}^j}{A^{2kj} q^{(2j+1)(j-1)k - \frac{(j-1)(j-2)}{2} + (2+j)n+2}} \frac{[n]!}{[n-j]!} F_{n-j}^{\text{twist}-k}(Aq^{2j+1}) \quad (5.4)$$

with

$$\sum_{k=2}^{\infty} c_j^{(k)} z^k = - \sum_{i=0}^{\infty} \sum_{k=2i+2}^{\infty} \left(\frac{C_{j,i}^{(k)}(q)}{A^{2i} q^{k+2ji-1}} + (1 - \delta_{k,2i+j+1}) \frac{C_{j,i}^{(k)}(q^{-1})}{(Aq^{j+1})^{2(k-j-1)} A^{-2i} q^{-k-2ji+1}} \right) \cdot z^k \quad (5.5)$$

$$C_{j,i}^{(k)}(q) = \frac{[i+j-1]!}{[i]!([j]!)^2} \cdot \frac{[k-i-2]!}{[k-i-j-1]!} \left([i][k-2j-i-1] + q^{k-i-1}[j][i-j] \right) \quad (5.6)$$

6 Another defect zero knot

Of course, quantum $\mathcal{C}$ -polynomials exist not only for the twist knots. For illustration, we provide an example of defect-zero [60] knot 9_{46} , the smallest one beyond the double-braid family. For the double braid, the coefficients of differential expansion are given by formula (3.10). For higher defects, the coefficients of differential expansion are somewhat reduced, and it is interesting to see what would this mean for the $\mathcal{C}$ -polynomials: in variance with $\mathcal{A}$ -polynomials, their *definition* begins to depend on the defect. We leave these questions for the future, and here we just demonstrate that our discussion above is in no way restricted to the twist family, in fact, one can lift it to other families of knots like those introduced in [63] or [64].

Knot 9_{46} is a very simple knot, with $F_1^{9_{46}} = A^2(A^2 + 1)$. This first DE coefficient is independent of q , which is the sign of the vanishing defect [60]. Note it is almost the same as $F_1^{5_1} = F_1^{\text{twist}_2} = -A^2(A^2 + 1)$ or $F_1^{6_2} = F_1^{\text{twist}_{-2}} = \frac{A^2+1}{A^2}$. However, in general $F_n^{9_{46}}(A) = F_n^{\text{twist}_{-2}} \left(\frac{1}{Aq^{3(n-1)}} \right) + O\left([n][n-1][n-2]\{q\}^2\right)$, and the actual DE coefficients in this case are rather involved as compared to (3.13):

$$F_n^{9_{46}}(A) = A^{2n} \sum_{j=0}^{n/2} (-)^j (q^2 - 1)^j q^{n^2 + (2j-1)n - 3j(j+1)} \frac{[n]!}{[n-2j]![2j]!!} \prod_{i=n-1}^{2n-2j-2} (A^2 q^{2i} + 1) \quad (6.1)$$

Quantum binomial formula (2.7) can be applied to convert the product at the r.h.s. into another sum, but (6.1) itself contains double factorial and is similar in this respect to (3.18).

The equations ($\mathcal{C}$ -polynomials) are also complicated, somewhat unexpectedly for such a simple knot:

n -evolution:

$$\sum_{i=0}^5 \text{Pol}_i(q, A^2) F_{n+i}^{946}(A) = 0 \quad (6.2)$$

where the polynomials $\text{Pol}_i(q, A^2)$ are of the degree at most 6 in A^2 . However, in this case, they are remarkably complicated, explicit formulas being given in appendix B.

A -evolution:

$$\begin{aligned} F_n^{946}(A) = & \left(A^4 q^{4n+2} + \frac{[5]}{q^{2n+4}} \right) F_n^{946}(Aq) + \left(A^4 \left(\frac{[2]}{q} - [3]q^{2n} \right) - \frac{[4][5]}{[2]q^{4n+8}} \right) F_n^{946}(Aq^2) + \\ & + \left(-A^{10}q^{8n+10} + [3]A^4 \left(\frac{1}{q^2} - \frac{[2]}{q^{2n+3}} \right) + \frac{[4][5]}{[2]} \frac{1}{q^{6n+12}} \right) F_n^{946}(Aq^3) + \\ & + \left(A^{10}q^{4n+10} - A^8q^6 + A^4 \left(\frac{[3][2]}{q^{4n+5}} - \frac{1}{q^{2n+4}} \right) - \frac{[5]}{q^{8n+16}} \right) F_n^{946}(Aq^4) + \\ & + \left(\frac{A^8}{q^{2n-6}} - \frac{[2]A^4}{q^{6n+7}} + \frac{1}{q^{10n+20}} \right) F_n^{946}(Aq^5) \end{aligned} \quad (6.3)$$

or

$$\begin{aligned} \sum_{j=0} \frac{(-1)^{j+1}}{q^{2jn}} \left(\frac{\text{Bin}_q(5, j)}{q^{4j}} + \frac{A^4 q^{6n+4}}{q^{2j}} \text{Bin}_q(3, j-1) - \frac{A^4 q^{4n+3} [2]}{q^{2j}} \text{Bin}_q(3, j-2) - \right. \\ \left. - \frac{A^{10} q^{10+20n}}{q^{-2jn}} \text{Bin}_q(1, j-3) + A^8 q^{8n+6} \text{Bin}_q(1, j-4) \right) F_n^{946}(Aq^j) = 0 \end{aligned} \quad (6.4)$$

Note that, for this knot similarly to the twist knots, the order of the both n -evolution and A -evolution equations is the same, though the equation itself is much simpler for the A -evolution.

Non-local eq.: The sum rule (2.9) in this case is just the same as (3.20) because $F_1^{946} = F_1^{51} = A^2(1 + A^2)$ and $m_1 = 1$, $m_2 = 2$:

$$\sum_{j=0}^{n+1} \frac{(-1)^j}{q^{3jn}} \cdot \frac{[n+1]!}{[j]![n+1-j]!} \cdot F_n^{946}(q^j A) = 0 \quad (6.5)$$

Semi-local eq.:

$$\begin{aligned} F_{n+1}^{946}(A) + \frac{A^2(q^{2n}A^2 + 1)}{q^{2n}} F_n^{946}(q^2 A) \\ - \sum_{j=0}^{\frac{n-1}{2}} (q^2 - 1)^{2j+1} \frac{A^{4(j+1)} q^{2j^2 + (2j+3)n - 3(j+1)} [n]!}{[n-2j-1]!} F_{n-2j-1}^{946}(Aq^{2j+2}) \end{aligned} \quad (6.6)$$

As usual, there are a lot of mixed evolution equations, both local equations and non-local ones.⁵ For instance, there are two rather simple local equations

$$\begin{aligned}
F_{n+1}^{946}(A) &= A^4 q^{4n} \cdot F_n^{946}(Aq) + \frac{A^2}{q^{4n}} \cdot F_n^{946}(Aq^2) \\
&\quad - \{q^n\}\{q^{n-1}\} A^6 \left(q^{2n+3} \cdot F_{n-2}^{946}(Aq^3) + \frac{[2]}{q^{2n-8}} \cdot F_{n-2}^{946}(Aq^4) \right) + \\
&\quad + A^{10} q^{20} \{q^n\}\{q^{n-1}\}\{q^{n-2}\}\{q^{n-3}\} \cdot F_{n-4}^{946}(Aq^6)
\end{aligned} \tag{6.9}$$

and

$$F_n^{946}(A) = q^{-2n} F_n^{946}(Aq) - (q^{2n} - 1) A^4 \cdot F_{n-1}^{946}(Aq^2) \tag{6.10}$$

Note a spectacular simplicity of (6.10), especially impressive given the relative complexity of all other quantum C -polynomials. In variance with those, this equation has just three terms, still it is rather informative: (6.10) does *not* determine recursively only the lowest-degree item A^{2n} in each $F_n^{946} = A^{4n} q^{4n(n+1)} + \dots + c_n A^{2n}$.

7 Conclusion

In this paper, we revisited the forgotten subject of quantum C -polynomials [45] taking advantage of the new knowledge about the differential expansion of the HOMFLY-PT polynomials. Most important, it allows us to lift our consideration from the Jones polynomials to the level of HOMFLY-PT polynomials. Though we keep ourselves restricted to the symmetric representations $[n]$ only, this extension allows us to complement the difference equations in n by new equations in $A = q^N$ (or in N , if one prefers). This promotes a set of C -polynomials to the entire non-commutative ring, where the n -evolution and the A -evolution are treated on equal footing and serve as the ring generators. Similar structures should exist for $\mathcal{A}$ -polynomials, but it is much more difficult to study.

The differential expansion and thus the C -polynomials depend on the *defect* of the knot [60], so far we concentrated on the simplest case of defect zero. All the smallest knots with this property (i.e. with the Alexander polynomial of degree one) belong to the families of twisted knots and of double braids (much similar in properties to the twist knots), the first one beyond these families is knot 9_{46} . We make use of an explicit knowledge of the symmetric DE coefficients for the twist family [47] (see [50] for review and references on

⁵An example of an unusual non-local equation is

$$F_{n+1}^{946}(A) = \sum_{j=0}^n (-)^j A^{2j+2} q^{5jn} \frac{[n]!}{[n-j]!} \{q\}^j \left(a_j \cdot q^{8n} A^2 + b_j \cdot q^{4n} \right) \cdot F_{n-j}^{946}(A) \tag{6.7}$$

with

$$\begin{aligned}
a_0 &= 1, & a_1 &= q^{-6}, & a_2 &= q^{-19}[3], & a_3 &= q^{-37}[3]^2, & a_4 &= q^{-60}([3]^2 + q^{-6}[2]^2 + q^{-10}), \dots \\
b_0 &= 1, & b_1 &= q^{-7}[2], & b_2 &= q^{-19}[2]^2, & b_3 &= q^{-37}([3][2]^2 + q^{-6}), \dots
\end{aligned} \tag{6.8}$$

These are quantizations of sequences A063020 and A007858 from [96]. In particular, $y = \sum_{n=0}^{\infty} a_n x^{n+1}$ at $q = 1$ is a solution of the equation $y(y-1)(y^2-1) = x$.

the progress since then) to give a rather general description of the C -polynomials in this case. We consider also the example of 9_{46} to illustrate the relevance of our studies beyond the twist family. For the knots discussed in this paper, it turns out that the difference equations in n and in A are of the same order (of order k for the k -twist knots and of order 5 for knot 9_{46}). It would be interesting to establish if this is a universal feature of knot polynomials.

Surprisingly or not, the quantum C -polynomials appear to be rather sophisticated. Despite our strong belief that they are more fundamental than the A -polynomials, the theory does not yet look elementary. Still it is much more straightforward and explicit. However, new insights are still needed. In particular, we did not truly exploit the powerful KNTZ-matrix description of the DE [58, 59], which can be directly related to the ring structure of C -polynomials. This is one of the many questions for further investigation.

Acknowledgments

Our work is supported in part by the grant of the Foundation for the Advancement of Theoretical Physics “BASIS”, by RFBR grants 19-01-00680 (A.Mir.) and 19-02-00815 (A.Mor.), by joint grants 19-51-50008-YaF-a (A.Mir.), 19-51-53014-GFEN-a, 18-51-05015-Arm-a. The work was also partly funded by RFBR and NSFB according to the research project 19-51-18006.

A A -evolution for twist knots

A.1 Examples at low k

$$\boxed{k = -2}$$

$$A^2 q^{2n} \cdot F_n^{\text{twist}-2}(A) - \left(A^2 q^{2n} - 1\right) \cdot F_n^{\text{twist}-2}(Aq) - q^{2n} \cdot F_n^{\text{twist}-2}(Aq^2) = 0 \quad (\text{A.1})$$

$$\boxed{k = -3}$$

$$\begin{aligned} & \left(q^{2n+2} A^2 - 1\right) \cdot A^6 q^{4n+2} \cdot F_n^{\text{twist}-3}(A) - \\ & - \left(q^{2n} A^2 - 1\right) \left(A^2 q^{2n+2} + q^{2n+2} - 1\right) \cdot A^4 q^{2n+2} \cdot F_n^{\text{twist}-3}(Aq) - \\ & - \left(q^{2n+2} A^2 - 1\right) \left(1 + q^{2n} A^2 (q^{2n+2} - 1)\right) \cdot F_n^{\text{twist}-3}(Aq^2) + \\ & + \left(q^{2n} A^2 - 1\right) \cdot q^{4n} \cdot F_n^{\text{twist}-3}(Aq^3) = 0 \quad (\text{A.2}) \end{aligned}$$

$$k = -4$$

$$\begin{aligned}
 & \left(q^{2n+4} A^2 - 1 \right) \cdot A^{12} q^{6n+10} \cdot F_n^{\text{twist}-4}(A) - \\
 & - \left(q^{2n+4} A^2 - 1 \right) \left(q^{2n} A^2 - 1 \right) (A^2 + 1) \cdot A^8 q^{4n+10} \cdot F_n^{\text{twist}-4}(Aq) - \\
 & - \left(q^{2n+2} A^2 - 1 \right) \left\{ 1 + q^{2n} A^2 \left((q^2 + 1) q^{2n+2} - (q^4 + 1) \right) + A^4 q^{4n+4} \right\} \\
 & \quad \times A^4 q^{2n+8} \cdot F_n^{\text{twist}-4}(Aq^2) + \\
 & + \left(q^{2n+4} A^2 - 1 \right) \left(q^{2n} A^2 - 1 \right) (q^{4n+4} A^2 + 1) \cdot F_n^{\text{twist}-4}(Aq^3) + \\
 & + \left(q^{2n} A^2 - 1 \right) \cdot q^{6n} \cdot F_n^{\text{twist}-4}(Aq^4) = 0 \quad (\text{A.3})
 \end{aligned}$$

$$k = -5$$

$$\begin{aligned}
 & \left(A^2 q^{2n+6} - 1 \right) \left(A^2 q^{2n+4} - 1 \right) \cdot A^{20} q^{8n+26} \cdot F_n^{\text{twist}-5}(A) - \\
 & - \left(A^2 q^{2n+6} - 1 \right) \left(A^2 q^{2n} - 1 \right) \left(A^4 q^{2n+4} + (A^2 + 1)(q^{2n+4} - 1) \right) \cdot A^{16} q^{6n+26} \cdot F_n^{\text{twist}-5}(Aq) - \\
 & - \left(A^2 q^{2n+6} - 1 \right) \left(A^2 q^{2n+2} - 1 \right) \left((q^{2n+4} - 1) A^4 q^{2n+2} + (q^{2n+4} - 1) \right. \\
 & \quad \times (q^{2n} + q^{2n-2} - 1) A^2 q^2 + (q^2 + 1) \left. \right) \cdot A^{10} q^{4n+22} \cdot F_n^{\text{twist}-5}(Aq^2) + \\
 & + \left(A^2 q^{2n+4} - 1 \right) \left(A^2 q^{2n} - 1 \right) \left((q^2 + 1) A^4 q^{6n+10} + (q^{2n+4} - 1)(q^{2n} + q^{2n-2} - 1) \right) \\
 & \quad \times A^2 q^{2n+6} + (q^{2n+4} - 1) \cdot A^6 q^{2n+14} \cdot F_n^{\text{twist}-5}(Aq^3) + \\
 & + \left(A^2 q^{2n+6} - 1 \right) \left(A^2 q^{2n} - 1 \right) \left((q^{2n+4} - 1) A^4 q^{6n+8} + (q^{2n+4} - 1) A^2 q^{2n+2} + 1 \right) \cdot F_n^{\text{twist}-5}(Aq^4) - \\
 & - \left(A^2 q^{2n+2} - 1 \right) \left(A^2 q^{2n} - 1 \right) \cdot q^{8n} \cdot F_n^{\text{twist}-5}(Aq^5) = 0 \quad (\text{A.4})
 \end{aligned}$$

A.2 General answer

The general structure of the answer is

$$\sum_{j=0}^{k-1} \epsilon_j G_{n,j}^{(k)} \cdot F_n^{\text{twist}-k}(Aq^j) + \epsilon_k \cdot q^{2(k-1)n} \cdot \prod_{i=0}^{\frac{k-3}{2}} \left(A^2 q^{2n+2i} - 1 \right) \cdot F_n^{\text{twist}-k}(Aq^k) = 0 \quad (\text{A.5})$$

$$\begin{aligned}
 G_{n,j}^{(k)} = & \left(A^2 q^{2n+2j-2} - 1 \right) \cdot \prod_{i=0}^{\text{floor}\left(\frac{j-3}{2}\right)} \left(A^2 q^{2n+2i} - 1 \right) \cdot \prod_{i=\text{floor}\left(\frac{k+j}{2}\right)}^{k-2} \left(A^2 q^{2n+2i} - 1 \right) \times \\
 & \times A^{k^2 - (j+1)k + 4\text{frac}\left(\frac{k}{2}\right)\text{frac}\left(\frac{j}{2}\right)} q^{2(k-j-1)n + \sigma_{k,j}} \cdot g_{n,j}^{(k)} \quad (\text{A.6})
 \end{aligned}$$

where $\epsilon_j = \sqrt{2} \cdot \cos\left(\frac{(2j+1)\pi}{4}\right)$, and we conjecture that

$$\begin{aligned}
 \sigma_{k,j} = & \text{floor}\left(\frac{k(k-2)(2k-3)}{4}\right) - \frac{j(j-1)}{2}k + \text{floor}\left(\frac{j}{2}\right)^2 + \text{floor}\left(\frac{j}{2}\right) \\
 & - 2(-1)^j \text{frac}\left(\frac{k}{2}\right) \text{floor}\left(\frac{j}{2}\right) \quad (\text{A.7})
 \end{aligned}$$

and

$$g_{n,0}^{(k)} = 1 \quad (\text{A.8})$$

$$g_{n,1}^{(k)} = \begin{cases} k \text{ even} & \xi_k(A) \\ k \text{ odd} & q^{2n+k-1} \cdot \xi_{k+1}(A) - \xi_{k-1}(A) \end{cases} \quad (\text{A.9})$$

$$g_{n,2}^{(k)} = \begin{cases} k \text{ even} & A^2 q^{4n} \sum_{i=1}^{\frac{k}{2}} q^{k-4+2i} \xi_{2i}(A) + \left(1 - A^2 q^{2n}(1 + q^k)\right) \sum_{i=1}^{\frac{k}{2}-1} q^{2(i-1)} \xi_{2i}(A) \\ k \text{ odd} & A^2 q^{4n} \sum_{i=1}^{\frac{k-1}{2}} q^{k-3+2i} \xi_{2i}(A) + \left(1 - A^2 q^{2n}\right) \sum_{i=1}^{\frac{k-1}{2}} q^{2(i-1)} \xi_{2i}(A) + \sum_{i=0}^{\frac{k-3}{2}} q^{2i+4} \xi_{2i}(A) \end{cases} \quad (\text{A.10})$$

$$\dots \quad (\text{A.11})$$

where $\xi_l(A) := \frac{A^l - 1}{A^2 - 1}$ is a polynomial of A only (since j is even in the formulas above).

Similarly, expressions for $g_{n,j}^{(k)}$ at $j > 2$ involve $\frac{(q^m A)^l - 1}{(q^m A)^2 - 1}$ with $m \leq j - 2$.

A.3 Examples of $G_{n,j}^{(k)}$

$$G_{n,0}^{(k)} = \prod_{i=\text{floor}(\frac{k}{2})}^{k-2} \left(A^2 q^{2n+2i} - 1 \right) \cdot A^{k(k-1)} q^{2(k-1)n + \sigma_k} \quad (\text{A.12})$$

$$\begin{aligned} G_{n,1}^{(2)} &= (A^2 q^{2n} - 1) \\ G_{n,1}^{(3)} &= (A^2 q^{2n} - 1) \cdot (q^{2n+2}(A^2 + 1) - 1) \cdot A^4 q^{2n+2} \\ G_{n,1}^{(4)} &= (A^2 q^{2n} - 1) \cdot (A^2 q^{2n+4} - 1)(A^2 + 1) \cdot A^8 q^{4n+10} \\ G_{n,1}^{(5)} &= (A^2 q^{2n} - 1) \cdot (A^2 q^{2n+6} - 1) \cdot (q^{2n+4}(A^4 + A^2 + 1) - (A^2 + 1)) \cdot A^{16} q^{6n+26} \\ G_{n,1}^{(6)} &= (A^2 q^{2n} - 1) \cdot (A^2 q^{2n+6} - 1)(A^2 q^{2n+8} - 1) \cdot (A^4 + A^2 + 1) \cdot A^{24} q^{8n+54} \\ G_{n,1}^{(7)} &= (A^2 q^{2n} - 1) \cdot (A^2 q^{2n+8} - 1)(A^2 q^{2n+10} - 1) \\ &\quad \cdot (q^{2n+6}(A^6 + A^4 + A^2 + 1) - (A^4 + A^2 + 1)) \cdot A^{36} q^{10n+96} \\ &\dots \end{aligned} \quad (\text{A.13})$$

$$\begin{aligned} G_{n,2}^{(3)} &= (A^2 q^{2n+2} - 1)(A^2 q^{2n}(q^{2n+2} - 1) + 1) \\ G_{n,2}^{(4)} &= (A^2 q^{2n+2} - 1) \left\{ A^4 q^{4n+4} + (q^{2n+2}(q^2 + 1) - (q^4 + 1)) A^2 q^{2n} + 1 \right\} \cdot A^4 q^{2n+8} \\ G_{n,2}^{(5)} &= (A^2 q^{2n+2} - 1)(A^2 q^{2n+6} - 1) \\ &\quad \times (A^4 q^{2n+2}(q^{2n+4} - 1) + (q^{2n+4} - 1)(q^{2n+2} + q^{2n} - q^2) \cdot A^2 + (q^2 + 1)) \cdot A^{10} q^{4n+22} \\ G_{n,2}^{(6)} &= (A^2 q^{2n+2} - 1)(A^2 q^{2n+8} - 1) \cdot \left\{ A^6 q^{4n+8} + (q^{4n+8} + q^{4n+6} - q^{2n+8} - q^{2n+2}) A^4 + \right. \\ &\quad \left. + ((q^4 + q^2 + 1) q^{4n+4} - (q^4 + q^2 + q^{-2} + q^{-4}) q^{2n+4} + q^2) A^2 + (q^2 + 1) \right\} \cdot A^{18} q^{6n+50} \end{aligned}$$

$$\begin{aligned}
 G_{n,2}^{(7)} = & \left(A^2 q^{2n+2} - 1\right) \left(A^2 q^{2n+8} - 1\right) \left(A^2 q^{2n+10} - 1\right) \\
 & \cdot \left\{ (q^{2n+6} - 1) \left(A^6 q^{2n+4} + \left((q^2 + 1)q^{2n+2} - q^4\right)A^4 + \left((q^4 + q^2 + 1)q^{2n} - (q^4 + q^2)\right)A^2\right) + \right. \\
 & \left. + (q^4 + q^2 + 1) \right\} \cdot A^{28} q^{8n+90} \\
 & \dots
 \end{aligned} \tag{A.14}$$

$$\begin{aligned}
 G_{n,3}^{(4)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+4} - 1\right) \left(A^2 q^{4n+4} + 1\right) \\
 G_{n,3}^{(5)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+4} - 1\right) \\
 & \times \left(A^4 q^{6n+10} (q^2 + 1) + (q^{2n+4} - 1)(q^{2n} + q^{2n-2} - 1) \cdot A^2 q^{2n+6} + (q^{2n+4} - 1)\right) \cdot A^6 q^{2n+14} \\
 G_{n,3}^{(6)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+4} - 1\right) \left(A^2 q^{2n+8} - 1\right) \cdot (q^2 + 1) \\
 & \times \left(A^4 q^{4n+8} + (q^{4n+4} - q^{2n+4} + q^{2n+2} - q^{2n} + 1)A^2 q^2 + 1\right) \cdot A^{12} q^{4n+38} \\
 G_{n,3}^{(7)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+4} - 1\right) \left(A^2 q^{2n+10} - 1\right) \\
 & \times \left\{ (q^4 + q^2 + 1)A^6 q^{6n+16} + (q^{2n+6} - 1) \left((q^2 + 1) \left((q^2 + 1)q^{4n} - q^{2n+2}\right)A^4 q^8 + \right. \right. \\
 & \left. \left. + \left((q^4 + q^2 + 1)q^{4n+4} - (q^2 + 1)q^{2n+6} - q^{2n} + q^2 + 1\right)A^2 q^2 + q^2 + 1\right) \right\} \cdot A^{22} q^{6n+78} \\
 & \dots
 \end{aligned} \tag{A.15}$$

$$\begin{aligned}
 G_{n,4}^{(5)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+6} - 1\right) \left((A^4 q^{4n+12} + A^2 q^{4n+6} + 1) - (A^2 q^{4n+6} + 1) \cdot A^2 q^{2n+2}\right) \\
 G_{n,4}^{(6)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+6} - 1\right) \left\{ (q^2 + 1)A^6 q^{8n+16} + \right. \\
 & + \left((q^4 + q^2 + 1)q^{4n+2} - (q^4 + q^2 + q^{-2} + q^{-4})q^{2n+2} + 1\right)A^4 q^{4n+10} + \\
 & \left. + \left((q^2 + 1)q^{4n+6} - (q^6 + 1)q^{2n+2}\right)A^2 + 1 \right\} \cdot A^6 q^{2n+24} \\
 G_{n,4}^{(7)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+6} - 1\right) \left(A^2 q^{2n+10} - 1\right) \cdot \left\{ (q^{2n+6} - 1) \left((q^2 + 1)A^6 q^{6n+14} + \right. \right. \\
 & + \left((q^4 + q^2 + 1)q^{6n+10} - (q^2 + 1)q^{4n+12} - q^{4n+6} + (q^2 + 1)q^{2n+6}\right)A^4 + \\
 & \left. \left. + (q^2 + 1) \left((q^2 + 1)q^{2n+2} - q^4\right)A^2\right) + (q^4 + q^2 + 1) \right\} \cdot A^{14} q^{4n+58} \\
 & \dots
 \end{aligned} \tag{A.16}$$

$$\begin{aligned}
 G_{n,5}^{(6)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+2} - 1\right) \left(A^2 q^{2n+8} - 1\right) \left(A^4 q^{8n+16} + A^2 q^{4n+8} + 1\right) \\
 G_{n,5}^{(7)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+2} - 1\right) \left(A^2 q^{2n+8} - 1\right) \\
 & \times \left\{ (q^4 + q^2 + 1)A^6 q^{10n+26} + \left((q^4 + q^2 + 1) \cdot q^{8n+16} - (q^2 + 1) \cdot q^{6n+18}\right) \cdot A^4 + \right. \\
 & \left. + \left((q^2 + 1) \cdot q^{4n+8} - q^{2n+10}\right) \cdot A^2 + (q^{2n+6} - 1) \right\} \cdot A^8 q^{2n+34} \\
 & \dots
 \end{aligned} \tag{A.17}$$

$$\begin{aligned}
 G_{n,6}^{(7)} = & \left(A^2 q^{2n} - 1\right) \left(A^2 q^{2n+2} - 1\right) \left(A^2 q^{2n+10} - 1\right) \\
 & \times \left\{ \left(A^6 q^{8n+20} + A^4 q^{4n+10} + A^2\right) (q^{2n+6} - 1) q^{2n+4} + 1 \right\} \\
 & \dots
 \end{aligned} \tag{A.18}$$

B Details on knot 9_{46}

Explicit expressions for the entries of (6.2):

$$\begin{aligned}
 & \{q^{n+2}\}q^{7n+12}A^6\left([2]^4[3]^2\{q\}^2\alpha_{0,1}^{(1)}-q^{2n-1}[2]^3\alpha_{0,2}^{(1)}-q^{4n+4}[2]^2\alpha_{0,3}^{(1)}-q^{6n+14}[4][2]\alpha_{0,4}^{(1)}+\right. \\
 & +A^2\left[q^{2n+13}[6][4][3]^2[2]^4\{q\}^3-q^{4n}[2]^4\alpha_{1,1}^{(1)}+q^{6n+3}[2]\alpha_{1,2}^{(1)}\right. \\
 & \left.-q^{8n+14}[3]\alpha_{1,3}^{(1)}+q^{10n+33}[2]^3[4]^3(q^8[2]-[4])\right]\Big)F_{n+1}^{9_{46}}(A)- \\
 & -A^4q^{4n+6}\left(q[3][2]^5\{q\}^2\alpha_{0,1}^{(2)}+q^{2n}[2]^2\alpha_{0,2}^{(2)}+q^{4n+4}\alpha_{0,3}^{(2)}+q^{6n+15}[4]\alpha_{0,4}^{(2)}+q^{8n+38}[4]^4[2]^4-\right. \\
 & -A^2\left[q^{2n}[2]^4[3]\{q\}^2\alpha_{1,1}^{(2)}-q^{4n}[2]^2\alpha_{1,2}^{(2)}+q^{6n+4}\alpha_{1,3}^{(2)}+q^{8n+16}[2]^2\alpha_{1,4}^{(2)}+q^{10n+28}[4]^2[2]^2\alpha_{1,5}^{(2)}\right]- \\
 & -A^4\left[q^{4n+16}[6][4][3]^3[2]^3\{q\}^3-q^{6n+5}[3][2]^3\alpha_{2,1}^{(2)}-q^{8n+6}\alpha_{2,2}^{(2)}-q^{10n+13}[2]\alpha_{2,3}^{(2)}-q^{12n+26}[4]^2\alpha_{2,4}^{(2)}\right]- \\
 & -A^6\left[q^{8n+24}[6]^2[3][2]^5\{q\}^3-q^{10n+12}[2]^4\alpha_{3,1}^{(2)}+q^{12n+16}[2]^2\alpha_{3,2}^{(2)}\right. \\
 & \left.+q^{14n+22}\alpha_{3,3}^{(2)}+q^{16n+45}[2]^3[4]^3(q^8[4]-[2])\right]\Big)F_{n+2}^{9_{46}}(A)+ \\
 & +A^2q^{2n+5}\left(q^{-1}[3][2]^4\{q\}^2\alpha_{0,1}^{(3)}-q^{2n}[2]^3\alpha_{0,2}^{(3)}-q^{4n+6}[2]\alpha_{0,3}^{(3)}-q^{6n+29}[2]^3[4]^3([3]+q^{10})+\right. \\
 & +A^2\left[q^{2n+6}[3]^2[2]^4\{q\}^3\alpha_{1,1}^{(3)}+q^{4n+3}[2]^2\alpha_{1,2}^{(3)}+q^{6n+9}\alpha_{1,3}^{(3)}+q^{8n+26}[4]^2[2]\alpha_{1,4}^{(3)}\right]- \\
 & -A^4\left[q^{4n+1}[3][2]^4\{q\}^2\alpha_{2,1}^{(3)}-q^{6n+3}[2]^4\alpha_{2,2}^{(3)}+q^{8n+9}[2]^2\alpha_{2,3}^{(3)}+q^{10n+20}[4]\alpha_{2,4}^{(3)}+q^{12n+41}[4]^4[2]^4\right]- \\
 & -A^6\left[q^{6n+16}[6][4][3]^2[2]^4\{q\}^3+q^{8n+9}[2]^4\alpha_{3,1}^{(3)}-q^{10n+10}[2]\alpha_{3,2}^{(3)}-q^{12n+17}\alpha_{3,3}^{(3)}-q^{14n+39}[4]^3[3][2]\alpha_{3,4}^{(3)}\right]- \\
 & -A^8\left[q^{10n+28}[6][3]^2[2]^5\{q\}^3+q^{12n+21}[2]^4\alpha_{4,1}^{(3)}-q^{14n+28}[2]\alpha_{4,2}^{(3)}-q^{16n+37}[4][2]\alpha_{4,3}^{(3)}\right]- \\
 & -A^{10}\left[q^{14n+39}[6][3]^2[2]^4\{q\}^3-q^{16n+32}[2]^3\alpha_{5,1}^{(3)}-q^{18n+39}\alpha_{5,2}^{(3)}-q^{20n+60}[4]^3[2]^4\right]\Big)F_{n+3}^{9_{46}}(A)- \\
 & -\left(q^8[6][4][3]^2[2]^3\{q\}^3+q^{2n+1}[3][2]^2\{q\}\alpha_{0,1}^{(4)}-q^{4n+9}[4][2]\{q\}\alpha_{0,2}^{(4)}+\right. \\
 & +A^2\left[q^{2n-2}[3][2]^4\{q\}^2\alpha_{1,1}^{(4)}-q^{4n-2}[2]^2\alpha_{1,2}^{(4)}-q^{6n+6}\alpha_{1,3}^{(4)}-q^{8n+27}[4]^3[2]^4\right]- \\
 & -A^4\left[q^{4n+3}[3][2]^5\{q\}^2\alpha_{0,1}^{(2)}-q^{6n+2}[2]^2\alpha_{0,2}^{(2)}-q^{8n+8}\alpha_{0,3}^{(2)}-q^{10n+31}[4]^3[2]^2\alpha_{2,1}^{(4)}\right]- \\
 & -A^6\left[q^{6n+10}[3]^2[2]^4\{q\}^2\alpha_{3,1}^{(4)}-q^{8n+8}[2]^2\alpha_{3,2}^{(4)}-q^{10n+14}\alpha_{3,3}^{(4)}-q^{12n+32}[4]^2[2]^2\alpha_{3,4}^{(4)}\right]- \\
 & -A^8\left[q^{8n+20}[6][3]^2[2]^4\{q\}^3+q^{10n+13}[2]^3\alpha_{4,1}^{(4)}+q^{12n+20}\alpha_{4,2}^{(4)}+q^{14n+37}[4]^2[2]^3\alpha_{4,3}^{(4)}\right]\Big)F_{n+4}^{9_{46}}(A)+ \\
 & -q^8\left([3][2]^4\{q\}^2\beta_1+q^{2n}[2]^2\beta_2+q^{4n+8}\beta_3+q^{6n+27}[4]^3[2]^4\right)\left(F_{n+3}^{9_{46}}(A)\right. \\
 & -q^{2n+4}A^2F_{n+2}^{9_{46}}(A)+\{q^{n+2}\}q^{5n+6}A^4F_{n+1}^{9_{46}}(A)+\{q^{n+1}\}\{q^{n+2}\}q^{8n+7}A^6F_n^{9_{46}}(A))+ \\
 & +q^{10n+22}\left(q^5[6][2]^4[3]^2\{q\}^3-q^{2n}[2]^3\beta_4-q^{4n+1}\beta_5-q^{6n+25}[8][4]^2[2]^3\right) \\
 & \times\left(q^{6n+7}A^{12}F_{n+2}^{9_{46}}(A)+q^{3n+2}[2]\{q^{n+2}\}A^{10}F_{n+1}^{9_{46}}(A)+\{q^{n+1}\}\{q^{n+2}\}A^8F_n^{9_{46}}(A)\right)- \\
 & -\{q\}[2]^2\left([3][2]^2\{q\}\beta_6+q^{2n}[2]\beta_7+q^{4n+14}[3][4]^2(q^7[3]-[2])\right) \\
 & \times\left(\left[q^{6n+14}A^6+q^{4n+8}A^4-q^{2n}A^2-q^{-6}\right]F_{n+5}^{9_{46}}(A)-A^{10}F_{n+4}^{9_{46}}(A)\right)=0
 \end{aligned}$$

where the coefficients involve the following structures:

$$\begin{aligned}
\alpha_{0,1}^{(1)} &= q^{20} - 2q^{14} - q^{12} - 2q^{10} - 3q^8 - 4q^6 - 3q^4 - q^2 - 1 \\
\alpha_{0,2}^{(1)} &= q^{36} + 4q^{34} + 8q^{32} + 11q^{30} + 12q^{28} + 16q^{26} + 28q^{24} + 46q^{22} + 64q^{20} + 74q^{18} + \\
&\quad + 82q^{16} + 76q^{14} + 60q^{12} + 31q^{10} + 11q^8 - q^6 - 5q^4 - 4q^2 - 2 \\
\alpha_{0,3}^{(1)} &= -2q^{38} - 8q^{36} - 19q^{34} - 36q^{32} - 60q^{30} - 94q^{28} - 139q^{26} - 192q^{24} - 242q^{22} - 278q^{20} - \\
&\quad - 287q^{18} - 263q^{16} - 207q^{14} - 137q^{12} - 73q^{10} - 29q^8 - 3q^6 + 5q^4 + 4q^2 + 1 \\
\alpha_{0,4}^{(1)} &= q^{32} + 4q^{30} + 8q^{28} + 14q^{26} + 24q^{24} + 40q^{22} + 57q^{20} + 70q^{18} + 74q^{16} + 71q^{14} + \\
&\quad + 61q^{12} + 47q^{10} + 30q^8 + 13q^6 + q^4 - 2q^2 - 1 \\
\alpha_{1,1}^{(1)} &= q^{36} + 5q^{34} + 8q^{32} + 16q^{30} + 15q^{28} + 23q^{26} + 18q^{24} + 27q^{22} + 25q^{20} + 33q^{18} + \\
&\quad + 29q^{16} + 26q^{14} + 16q^{12} + 8q^{10} + q^8 + 2q^4 + 2q^2 + 1 \\
\alpha_{1,2}^{(1)} &= 2q^{42} + 14q^{40} + 45q^{38} + 108q^{36} + 202q^{34} + 313q^{32} + 409q^{30} + 469q^{28} + 486q^{26} + \\
&\quad + 467q^{24} + 414q^{22} + 333q^{20} + 228q^{18} + 116q^{16} + 24q^{14} - 23q^{12} - 25q^{10} - 4q^8 + 11q^6 + 11q^4 + 5q^2 + 1 \\
\alpha_{1,3}^{(1)} &= q^{36} + 7q^{34} + 23q^{32} + 59q^{30} + 117q^{28} + 183q^{26} + 231q^{24} + 240q^{22} + 201q^{20} + 126q^{18} + \\
&\quad + 35q^{16} - 52q^{14} - 113q^{12} - 134q^{10} - 117q^8 - 77q^6 - 37q^4 - 12q^2 - 2 \\
\alpha_{0,1}^{(2)} &= q^{24} + 2q^{20} + 3q^{16} - q^{14} + 3q^{12} + q^{10} + 4q^8 + 2q^4 + 1 \\
\alpha_{0,2}^{(2)} &= q^{40} + 5q^{38} + 10q^{36} + 15q^{34} + 16q^{32} + 13q^{30} - 2q^{28} - 23q^{26} - 39q^{24} - 42q^{22} - 46q^{20} - \\
&\quad - 57q^{18} - 61q^{16} - 47q^{14} - 24q^{12} - 5q^{10} + 4q^8 + 9q^6 + 9q^4 + 6q^2 + 2 \\
\alpha_{0,3}^{(2)} &= -2q^{44} - 12q^{42} - 34q^{40} - 66q^{38} - 93q^{36} - 85q^{34} + 187q^{30} + 454q^{28} + 730q^{26} + 936q^{24} + \\
&\quad + 1025q^{22} + 992q^{20} + 856q^{18} + 650q^{16} + 417q^{14} + 208q^{12} + 62q^{10} - 12q^8 - 30q^6 - 20q^4 - 7q^2 - 1 \\
\alpha_{0,4}^{(2)} &= q^{36} + 5q^{34} + 11q^{32} + 15q^{30} + 5q^{28} - 35q^{26} - 117q^{24} - 229q^{22} - 336q^{20} - 398q^{18} - \\
&\quad - 397q^{16} - 339q^{14} - 248q^{12} - 154q^{10} - 77q^8 - 25q^6 - q^4 + 3q^2 + 1 \\
\alpha_{1,1}^{(2)} &= q^{28} + 2q^{26} + 4q^{24} + 4q^{22} + 3q^{20} + q^{18} - q^{16} - 2q^{14} - 8q^{12} - 10q^{10} - 11q^8 - 7q^6 - 5q^4 - 2q^2 - 1 \\
\alpha_{1,2}^{(2)} &= q^{42} + 7q^{40} + 19q^{38} + 37q^{36} + 53q^{34} + 64q^{32} + 62q^{30} + 63q^{28} + 78q^{26} + 114q^{24} + \\
&\quad + 149q^{22} + 180q^{20} + 204q^{18} + 210q^{16} + 177q^{14} + 113q^{12} + 44q^{10} - 16q^6 - 14q^4 - 7q^2 - 2 \\
\alpha_{1,3}^{(2)} &= 3q^{46} + 22q^{44} + 78q^{42} + 189q^{40} + 359q^{38} + 579q^{36} + 821q^{34} + 1053q^{32} + 1272q^{30} + 1523q^{28} + 1839q^{26} + \\
&\quad + 2169q^{24} + 2391q^{22} + 2399q^{20} + 2153q^{18} + 1696q^{16} + 1134q^{14} + 610q^{12} + 240q^{10} \\
&\quad + 50q^8 - 12q^6 - 15q^4 - 6q^2 - 1 \\
\alpha_{1,4}^{(2)} &= -3q^{38} - 15q^{36} - 41q^{34} - 85q^{32} - 145q^{30} - 219q^{28} - 298q^{26} - 382q^{24} - 468q^{22} - 562q^{20} - \\
&\quad - 643q^{18} - 686q^{16} - 660q^{14} - 562q^{12} - 414q^{10} - 259q^8 - 135q^6 - 57q^4 - 17q^2 - 3 \\
\alpha_{1,5}^{(2)} &= q^{26} + 3q^{24} + 5q^{22} + 7q^{20} + 8q^{18} + 10q^{16} + 13q^{14} + 18q^{12} + 20q^{10} + 18q^8 + 13q^6 + 8q^4 + 3q^2 + 1 \\
\alpha_{2,1}^{(2)} &= q^{34} + 5q^{32} + 11q^{30} + 17q^{28} + 19q^{26} + 21q^{24} + 20q^{22} + 25q^{20} + 27q^{18} + 30q^{16} + \\
&\quad + 26q^{14} + 24q^{12} + 17q^{10} + 9q^8 + 3q^6 + 1 \\
\alpha_{2,2}^{(2)} &= -q^{44} - 14q^{42} - 69q^{40} - 210q^{38} - 464q^{36} - 813q^{34} - 1193q^{32} - 1527q^{30} - 1754q^{28} - 1853q^{26} - \\
&\quad - 1833q^{24} - 1715q^{22} - 1509q^{20} - 1229q^{18} - 908q^{16} - 601q^{14} - 356q^{12} \\
&\quad - 194q^{10} - 105q^8 - 59q^6 - 30q^4 - 11q^2 - 2 \\
\alpha_{2,3}^{(2)} &= -q^{42} + 5q^{40} + 34q^{38} + 112q^{36} + 252q^{34} + 447q^{32} + 659q^{30} + 836q^{28} + 930q^{26} + 925q^{24} + \\
&\quad + 834q^{22} + 690q^{20} + 525q^{18} + 366q^{16} + 236q^{14} + 144q^{12} + 86q^{10} + 53q^8 + 33q^6 + 17q^4 + 6q^2 + 1 \\
\alpha_{2,4}^{(2)} &= q^{30} - 6q^{26} - 19q^{24} - 33q^{22} - 44q^{20} - 47q^{18} - 41q^{16} - 29q^{14} - 18q^{12} - 8q^{10} - 2q^8 - 2q^6 - 4q^4 - 3q^2 - 1
\end{aligned}$$

$$\begin{aligned}
 \alpha_{3,1}^{(2)} &= q^{36} + 5q^{34} + 9q^{32} + 13q^{30} + 15q^{28} + 21q^{26} + 25q^{24} + 25q^{22} + 21q^{20} + 23q^{18} + \\
 &\quad + 28q^{16} + 26q^{14} + 20q^{12} + 9q^{10} + 9q^8 + 2q^6 + 3q^4 + 1 \\
 \alpha_{3,2}^{(2)} &= 2q^{40} + 13q^{38} + 38q^{36} + 82q^{34} + 138q^{32} + 200q^{30} + 252q^{28} + 285q^{26} + 282q^{24} + \\
 &\quad + 254q^{22} + 214q^{20} + 180q^{18} + 141q^{16} + 99q^{14} + 57q^{12} + 31q^{10} + 19q^8 + 14q^6 + 9q^4 + 4q^2 + 1 \\
 \alpha_{3,3}^{(2)} &= -q^{42} - 10q^{40} - 43q^{38} - 123q^{36} - 270q^{34} - 483q^{32} - 723q^{30} - 922q^{28} - 1013q^{26} - 959q^{24} - \\
 &\quad - 774q^{22} - 522q^{20} - 277q^{18} - 94q^{16} + 6q^{14} + 35q^{12} + 26q^{10} + 8q^8 - 4q^6 - 7q^4 - 4q^2 - 1 \\
 \alpha_{0,1}^{(3)} &= q^{24} + q^{22} + 2q^{20} + q^{18} + q^{16} - q^{14} - 2q^{12} - 3q^{10} - 4q^8 - 5q^6 - 4q^4 - 2q^2 - 1 \\
 \alpha_{0,2}^{(3)} &= q^{36} + 4q^{34} + 7q^{32} + 10q^{30} + 11q^{28} + 13q^{26} + 13q^{24} + 15q^{22} + 20q^{20} + 27q^{18} + \\
 &\quad + 34q^{16} + 36q^{14} + 35q^{12} + 24q^{10} + 12q^8 + q^6 - 3q^4 - 3q^2 - 1 \\
 \alpha_{0,3}^{(3)} &= -2q^{38} - 10q^{36} - 26q^{34} - 49q^{32} - 75q^{30} - 101q^{28} - 127q^{26} - 155q^{24} - 184q^{22} - 214q^{20} - \\
 &\quad - 238q^{18} - 248q^{16} - 234q^{14} - 190q^{12} - 126q^{10} - 64q^8 - 20q^6 + 3q^2 + 1 \\
 \alpha_{1,1}^{(3)} &= q^{12} + q^{10} + q^8 + q^6 + 3q^4 + 2q^2 + 1 \\
 \alpha_{1,2}^{(3)} &= q^{34} + 2q^{32} + q^{30} - 4q^{28} - 6q^{26} - 7q^{24} - 9q^{22} - 21q^{20} - 31q^{18} - 34q^{16} - 34q^{14} - \\
 &\quad - 37q^{12} - 38q^{10} - 29q^8 - 12q^6 - q^4 + 2q^2 + 1 \\
 \alpha_{1,3}^{(3)} &= -2q^{36} - 9q^{34} - 21q^{32} - 30q^{30} - 25q^{28} + 5q^{26} + 65q^{24} + 152q^{22} + 248q^{20} + 324q^{18} + \\
 &\quad + 355q^{16} + 334q^{14} + 271q^{12} + 189q^{10} + 113q^8 + 58q^6 + 24q^4 + 7q^2 + 1 \\
 \alpha_{1,4}^{(3)} &= q^{18} + 2q^{16} + 3q^{14} + q^{12} - 3q^{10} - 9q^8 - 11q^6 - 9q^4 - 5q^2 - 2 \\
 \alpha_{2,1}^{(3)} &= q^{28} + 2q^{26} + 5q^{24} + 6q^{22} + 8q^{20} + 7q^{18} + 9q^{16} + 7q^{14} + 4q^{12} - 2q^{10} - 4q^8 - 4q^6 - 4q^4 - 2q^2 - 1 \\
 \alpha_{2,2}^{(3)} &= q^{38} + 4q^{36} + 8q^{34} + 14q^{32} + 14q^{30} + 16q^{28} + 9q^{26} + 11q^{24} + 4q^{22} + 10q^{20} + \\
 &\quad + 14q^{18} + 33q^{16} + 39q^{14} + 39q^{12} + 24q^{10} + 15q^8 + 4q^6 - 2q^2 - 1 \\
 \alpha_{2,3}^{(3)} &= 2q^{40} + 10q^{38} + 30q^{36} + 63q^{34} + 103q^{32} + 139q^{30} + 167q^{28} + 187q^{26} + 208q^{24} + \\
 &\quad + 238q^{22} + 283q^{20} + 332q^{18} + 361q^{16} + 344q^{14} + 279q^{12} + 187q^{10} + 100q^8 + 41q^6 + 10q^4 - 1 \\
 \alpha_{2,4}^{(3)} &= -q^{34} - 6q^{32} - 20q^{30} - 47q^{28} - 86q^{26} - 133q^{24} - 187q^{22} - 251q^{20} - 319q^{18} - 378q^{16} - \\
 &\quad - 410q^{14} - 399q^{12} - 340q^{10} - 250q^8 - 154q^6 - 74q^4 - 24q^2 - 4 \\
 \alpha_{3,1}^{(3)} &= q^{36} + q^{34} + q^{32} - q^{30} - 6q^{28} - 11q^{26} - 20q^{24} - 18q^{22} - 25q^{20} - 23q^{18} - 32q^{16} - \\
 &\quad - 25q^{14} - 28q^{12} - 23q^{10} - 25q^8 - 15q^6 - 7q^4 - q^2 + 1 \\
 \alpha_{3,2}^{(3)} &= q^{46} + 6q^{44} + 17q^{42} + 31q^{40} + 38q^{38} + 18q^{36} - 51q^{34} - 181q^{32} - 348q^{30} - 508q^{28} - \\
 &\quad - 625q^{26} - 691q^{24} - 700q^{22} - 660q^{20} - 586q^{18} - 495q^{16} - 387q^{14} \\
 &\quad - 268q^{12} - 154q^{10} - 70q^8 - 26q^6 - 10q^4 - 4q^2 - 1 \\
 \alpha_{3,3}^{(3)} &= -2q^{46} - 12q^{44} - 37q^{42} - 77q^{40} - 116q^{38} - 115q^{36} - 22q^{34} + 198q^{32} + 529q^{30} + 900q^{28} + 1212q^{26} + \\
 &\quad + 1391q^{24} + 1419q^{22} + 1326q^{20} + 1153q^{18} + 931q^{16} + 685q^{14} + 449q^{12} \\
 &\quad + 256q^{10} + 125q^8 + 53q^6 + 20q^4 + 6q^2 + 1 \\
 \alpha_{3,4}^{(3)} &= q^{18} + q^{16} - 2q^{12} - 2q^{10} - 2q^8 - q^6 - q^4 - q^2 - 1 \\
 \alpha_{4,1}^{(3)} &= q^{28} + 6q^{26} + 7q^{24} + 13q^{22} + 8q^{20} + 17q^{18} + 15q^{16} + 21q^{14} + 13q^{12} + 11q^{10} + 9q^8 + 7q^6 + 2q^4 - q^2 - 1 \\
 \alpha_{4,2}^{(3)} &= -q^{32} + 11q^{28} + 38q^{26} + 82q^{24} + 138q^{22} + 203q^{20} + 267q^{18} + 308q^{16} + 308q^{14} + \\
 &\quad + 266q^{12} + 199q^{10} + 130q^8 + 71q^6 + 30q^4 + 8q^2 + 1 \\
 \alpha_{4,3}^{(3)} &= q^{26} - 3q^{22} - 9q^{20} - 16q^{18} - 28q^{16} - 42q^{14} - 52q^{12} - 51q^{10} - 38q^8 - 18q^6 - 3q^4 + 2q^2 + 1 \\
 \alpha_{5,1}^{(3)} &= q^{28} + 4q^{26} + 8q^{24} + 13q^{22} + 13q^{20} + 15q^{18} + 15q^{16} + 17q^{14} + 14q^{12} + 11q^{10} + 10q^8 + 7q^6 + 2q^4 - q^2 - 1 \\
 \alpha_{5,2}^{(3)} &= -q^{30} - 8q^{28} - 29q^{26} - 74q^{24} - 143q^{22} - 221q^{20} - 281q^{18} - 304q^{16} - 289q^{14} - \\
 &\quad - 248q^{12} - 193q^{10} - 135q^8 - 81q^6 - 38q^4 - 12q^2 - 2;
 \end{aligned}$$

$$\begin{aligned}
 \alpha_{0,1}^{(4)} &= 2q^{26} + 5q^{24} + 9q^{22} + 8q^{20} + 2q^{18} - 7q^{16} - 14q^{14} - 14q^{12} - 9q^{10} - 5q^8 + 3q^4 + 3q^2 + 1 \\
 \alpha_{0,2}^{(4)} &= q^{24} + 4q^{22} + 10q^{20} + 14q^{18} + 10q^{16} - 3q^{14} - 16q^{12} - 21q^{10} - 15q^8 - 4q^6 + 3q^4 + 3q^2 + 1 \\
 \alpha_{1,1}^{(4)} &= q^{24} + 2q^{22} + 3q^{20} + 3q^{18} + 4q^{16} + 5q^{14} + 3q^{12} + q^{10} - q^8 - q^6 - 2q^4 - q^2 - 1 \\
 \alpha_{1,2}^{(4)} &= 2q^{36} + 7q^{34} + 14q^{32} + 15q^{30} + 10q^{28} + q^{26} - 3q^{24} - 2q^{22} + 8q^{20} + 24q^{18} + \\
 &\quad + 40q^{16} + 46q^{14} + 42q^{12} + 30q^{10} + 18q^8 + 7q^6 - 2q^2 - 1 \\
 \alpha_{1,3}^{(4)} &= -q^{36} - 6q^{34} - 19q^{32} - 38q^{30} - 54q^{28} - 59q^{26} - 61q^{24} - 80q^{22} - 132q^{20} - 209q^{18} - \\
 &\quad - 279q^{16} - 309q^{14} - 289q^{12} - 229q^{10} - 154q^8 - 87q^6 - 39q^4 - 12q^2 - 2 \\
 \alpha_{2,1}^{(4)} &= q^{14} + q^{12} + q^{10} - q^8 - 2 * q^6 - 2 * q^4 - q^2 - 1 \\
 \alpha_{3,1}^{(4)} &= q^{20} + q^{18} + 2q^{16} + 2q^{14} + 3q^{12} + 4q^{10} + 2q^8 + 2q^6 - 1 \\
 \alpha_{3,2}^{(4)} &= q^{38} + 5q^{36} + 12q^{34} + 21q^{32} + 26q^{30} + 26q^{28} + 16q^{26} + q^{24} - 16q^{22} - 22q^{20} - \\
 &\quad - 12q^{18} + 12q^{16} + 35q^{14} + 47q^{12} + 46q^{10} + 36q^8 + 19q^6 + 5q^4 - q^2 - 1 \\
 \alpha_{3,3}^{(4)} &= -2q^{40} - 12q^{38} - 37q^{36} - 78q^{34} - 122q^{32} - 146q^{30} - 129q^{28} - 73q^{26} - 10q^{24} + \\
 &\quad + 11q^{22} - 37q^{20} - 135q^{18} - 234q^{16} - 290q^{14} - 282q^{12} - 223q^{10} - 145q^8 - 76q^6 - 30q^4 - 8q^2 - 1 \\
 \alpha_{3,4}^{(4)} &= q^{20} + 2q^{18} + 3q^{16} + q^{14} - 2q^{12} - 4q^{10} - q^8 + 3q^6 + 6q^4 + 5q^2 + 2 \\
 \alpha_{4,1}^{(4)} &= q^{38} + 2q^{36} + 2q^{34} + q^{32} + q^{30} - 7q^{26} - 15q^{24} - 20q^{22} - 15q^{20} - 12q^{18} - 11q^{16} - \\
 &\quad - 14q^{14} - 13q^{12} - 11q^{10} - 10q^8 - 7q^6 - 2q^4 + q^2 + 1 \\
 \alpha_{4,2}^{(4)} &= -2q^{40} - 11q^{38} - 29q^{36} - 49q^{34} - 58q^{32} - 44q^{30} + 3q^{28} + 84q^{26} + 181q^{24} + \\
 &\quad + 260q^{22} + 299q^{20} + 296q^{18} + 266q^{16} + 231q^{14} + 202q^{12} + 170q^{10} + 128q^8 + 80q^6 + 38q^4 + 12q^2 + 2 \\
 \alpha_{5,3}^{(4)} &= q^{20} + q^{18} + q^{16} - q^{14} - 2q^{12} - 2q^{10} - q^8 - q^6 - q^4 - 2q^2 - 1 \\
 \beta_1 &= q^{14} + 2q^{12} + 2q^{10} + 2q^8 + 2q^6 + 4q^4 + 2q^2 + 1 \\
 \beta_2 &= q^{28} + 2q^{22} + 16q^{20} + 33q^{18} + 46q^{16} + 49q^{14} + 48q^{12} + 40q^{10} + 23q^8 + 5q^6 - 3q^4 - 3q^2 - 1 \\
 \beta_3 &= -q^{28} - 4q^{26} - 14q^{24} - 39q^{22} - 91q^{20} - 173q^{18} - 269q^{16} - 344q^{14} - 365q^{12} - 320q^{10} \\
 &\quad - 229q^8 - 131q^6 - 58q^4 - 18q^2 - 3 \\
 \beta_4 &= q^{26} + 4q^{24} + 7q^{22} + 8q^{20} + 7q^{18} + 8q^{16} + 12q^{14} + 17q^{12} + 18q^{10} + 18q^8 + 13q^6 + 10q^4 + 4q^2 + 1 \\
 \beta_5 &= -2q^{34} - 12q^{32} - 36q^{30} - 74q^{28} - 118q^{26} - 158q^{24} - 190q^{22} - 220q^{20} - 246q^{18} - 258q^{16} - \\
 &\quad - 244q^{14} - 202q^{12} - 143q^{10} - 86q^8 - 44q^6 - 19q^4 - 6q^2 - 1 \\
 \beta_6 &= q^{14} + q^{12} + 2q^{10} + 2q^8 + 4q^6 + 3q^4 + 2q^2 + 1 \\
 \beta_7 &= -2q^{24} - 5q^{22} - 10q^{20} - 13q^{18} - 15q^{16} - 11q^{14} - 3q^{12} + 7q^{10} + 12q^8 + 13q^6 + 9q^4 + 4q^2 + 1
 \end{aligned} \tag{B.1}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A.A. Migdal, *Loop Equations and 1/N Expansion*, *Phys. Rept.* **102** (1983) 199 [[INSPIRE](#)].
- [2] J. Ambjørn, J. Jurkiewicz and Y.M. Makeenko, *Multiloop correlators for two-dimensional quantum gravity*, *Phys. Lett. B* **251** (1990) 517 [[INSPIRE](#)].
- [3] F. David, *Loop Equations and Nonperturbative Effects in Two-dimensional Quantum Gravity*, *Mod. Phys. Lett. A* **5** (1990) 1019 [[INSPIRE](#)].

- [4] J. Ambjørn and Y.M. Makeenko, *Properties of Loop Equations for the Hermitean Matrix Model and for Two-dimensional Quantum Gravity*, *Mod. Phys. Lett. A* **5** (1990) 1753 [[INSPIRE](#)].
- [5] A. Mironov and A. Morozov, *On the origin of Virasoro constraints in matrix models: Lagrangian approach*, *Phys. Lett. B* **252** (1990) 47 [[INSPIRE](#)].
- [6] H. Itoyama and Y. Matsuo, *Noncritical Virasoro algebra of $d < 1$ matrix model and quantized string field*, *Phys. Lett. B* **255** (1991) 202 [[INSPIRE](#)].
- [7] A.S. Alexandrov, A. Mironov and A. Morozov, *Partition functions of matrix models as the first special functions of string theory. 1. Finite size Hermitean one matrix model*, *Int. J. Mod. Phys. A* **19** (2004) 4127 [[hep-th/0310113](#)] [[INSPIRE](#)].
- [8] A. Morozov, *Integrability and matrix models*, *Phys. Usp.* **37** (1994) 1 [[hep-th/9303139](#)] [[INSPIRE](#)].
- [9] A. Morozov, *Matrix models as integrable systems*, in *CRM-CAP Summer School on Particles and Fields '94*, Banff Canada (1994), pg. 127 [[hep-th/9502091](#)] [[INSPIRE](#)].
- [10] A. Morozov, *Challenges of matrix models*, in *NATO Advanced Study Institute and EC Summer School on String Theory: From Gauge Interactions to Cosmology*, Cargèse France (2005), pg. 129 [[hep-th/0502010](#)] [[INSPIRE](#)].
- [11] A. Mironov, *2 – D gravity and matrix models. 1. 2 – D gravity*, *Int. J. Mod. Phys. A* **9** (1994) 4355 [[hep-th/9312212](#)] [[INSPIRE](#)].
- [12] A. Mironov, *Matrix models of two-dimensional gravity*, *Phys. Part. Nucl.* **33** (2002) 537 [[INSPIRE](#)].
- [13] A. Mironov, *Quantum Deformations of τ -functions, Bilinear Identities and Representation Theory*, *Electron. Res. Announc. AMS* **9** (1996) 219 [[hep-th/9409190](#)] [[INSPIRE](#)].
- [14] T.D. Cohen and S. Sen, *Large N_c gauge theory with quarks in high representations*, *Phys. Rev. D* **90** (2014) 085008 [[arXiv:1406.4892](#)] [[INSPIRE](#)].
- [15] A. Shibata, K.-I. Kondo, R. Matsudo and S. Nishino, *Quark confinement in the Yang-Mills theory with a gauge-invariant gluon mass in view of the gauge-invariant BEH mechanism*, *PoS(LATTICE2019)060* [[arXiv:1912.01247](#)] [[INSPIRE](#)].
- [16] S.-S. Chern and J. Simons, *Characteristic Forms and Geometric Invariants*, *Ann. Math.* **99** (1974) 48.
- [17] E. Witten, *Quantum Field Theory and the Jones Polynomial*, *Commun. Math. Phys.* **121** (1989) 351 [[INSPIRE](#)].
- [18] J.W. Alexander, *Topological invariants of knots and links*, *Trans. Am. Math. Soc.* **30** (1928) 275.
- [19] J.H. Conway, *An Enumeration of Knots and Links, and Some of Their Algebraic Properties*, in *Computational Problems in Abstract Algebra, Proceedings of a Conference*, Oxford U.K. (1967), J. Leech eds., Pergamon Press, London U.K. (1970), pg. 329.
- [20] V.F.R. Jones, *Index for subfactors*, *Invent. Math.* **72** (1983) 1.
- [21] V.F.R. Jones, *A polynomial invariant for links via von Neumann algebras*, *Bull. Am. Math. Soc.* **12** (1985) 103.
- [22] V.F.R. Jones, *Hecke algebra representations of braid groups and link polynomials*, *Ann. Math.* **126** (1987) 335.

- [23] L. Kauffman, *State models and the Jones polynomial*, *Topology* **26** (1987) 395.
- [24] P. Freyd, D. Yetter, J. Hoste, W.B.R. Lickorish, K. Millett and A. Ocneanu, *A new polynomial invariant of knots and links*, *Bull. Am. Math. Soc.* **12** (1985) 239.
- [25] J.H. Przytycki and K.P. Traczyk, *Invariants of Conway type*, *Kobe J. Math.* **4** (1987) 115 [[arXiv:1610.06679](#)].
- [26] S. Garoufalidis, , *Geom. Topol. Monogr.* **7** (2004) 291 [[math/0306230](#)].
- [27] R.Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, *Mod. Phys. Lett. A* **39** (1997) 269 [[q-alg/9601025](#)].
- [28] H.Murakami and J.Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, *Acta Math.* **186** (2001) 85.
- [29] S. Gukov and H. Murakami, *SL(2, $\mathbb{C}$) Chern-Simons theory and the asymptotic behavior of the colored Jones polynomial*, *Lett. Math. Phys.* **86** (2008) 79 [[math/0608324](#)] [[INSPIRE](#)].
- [30] H.Murakami, *An Introduction to the Volume Conjecture*, [arXiv:1002.0126](#).
- [31] T. Dimofte, *Quantum Riemann Surfaces in Chern-Simons Theory*, *Adv. Theor. Math. Phys.* **17** (2013) 479 [[arXiv:1102.4847](#)] [[INSPIRE](#)].
- [32] A. Mironov and A. Morozov, *Equations on knot polynomials and 3d/5d duality*, *AIP Conf. Proc.* **1483** (2012) 189 [[arXiv:1208.2282](#)] [[INSPIRE](#)].
- [33] S. Garoufalidis and D. Zagier, *Asymptotics of Nahm sums at roots of unity*, [arXiv:1812.07690](#) [[INSPIRE](#)].
- [34] S. Gukov and C. Manolescu, *A two-variable series for knot complements*, [arXiv:1904.06057](#) [[INSPIRE](#)].
- [35] J. Brown, T. Dimofte, S. Garoufalidis and N. Geer, *The ADO Invariants are a q-Holonomic Family*, [arXiv:2005.08176](#) [[INSPIRE](#)].
- [36] T. Ekholm, A. Gruen, S. Gukov, P. Kucharski, S. Park and P. Sułkowski, *$\widehat{Z}$ at large N : from curve counts to quantum modularity*, [arXiv:2005.13349](#) [[INSPIRE](#)].
- [37] A. Mironov, A. Morozov and A. Morozov, *Character expansion for HOMFLY polynomials. I. Integrability and difference equations*, in *Strings, gauge fields, and the geometry behind: The legacy of Maximilian Kreuzer*, A. Rebhan et al. eds., World Scientific, New York U.S.A. (2013), pg. 101 [[arXiv:1112.5754](#)] [[INSPIRE](#)].
- [38] H. Ooguri and C. Vafa, *Knot invariants and topological strings*, *Nucl. Phys. B* **577** (2000) 419 [[hep-th/9912123](#)] [[INSPIRE](#)].
- [39] J.M.F. Labastida and M. Mariño, *Polynomial invariants for torus knots and topological strings*, *Commun. Math. Phys.* **217** (2001) 423 [[hep-th/0004196](#)] [[INSPIRE](#)].
- [40] J.M.F. Labastida and M. Mariño, *A New point of view in the theory of knot and link invariants*, [math/0104180](#) [[INSPIRE](#)].
- [41] J.M.F. Labastida, M. Mariño and C. Vafa, *Knots, links and branes at large N* , *JHEP* **11** (2000) 007 [[hep-th/0010102](#)] [[INSPIRE](#)].
- [42] M. Mariño and C. Vafa, *Framed knots at large N* , *Contemp. Math.* **310** (2002) 185 [[hep-th/0108064](#)] [[INSPIRE](#)].
- [43] S. Gukov, M. Mariño and P. Putrov, *Resurgence in complex Chern-Simons theory*, [arXiv:1605.07615](#) [[INSPIRE](#)].

- [44] S. Garoufalidis, J. Gu and M. Mariño, *The resurgent structure of quantum knot invariants*, [arXiv:2007.10190](#) [[INSPIRE](#)].
- [45] S. Garoufalidis and X. Sun, *The C-polynomial of a knot*, *Algebr. Geom. Topol.* **6** (2006) 1623 [[math/0504305](#)].
- [46] N.M. Dunfield, S. Gukov and J. Rasmussen, *The Superpolynomial for knot homologies*, *Exp. Math.* **15** (2006) 129 [[math/0505662](#)] [[INSPIRE](#)].
- [47] A. Mironov, A. Morozov and A. Morozov, *Evolution method and “differential hierarchy” of colored knot polynomials*, *AIP Conf. Proc.* **1562** (2013) 123 [[arXiv:1306.3197](#)] [[INSPIRE](#)].
- [48] E. Gorsky, S. Gukov and M. Stosic, *Quadruply-graded colored homology of knots*, [arXiv:1304.3481](#) [[INSPIRE](#)].
- [49] S.B. Arthamonov, A. Mironov and A. Morozov, *Differential hierarchy and additional grading of knot polynomials*, *Theor. Math. Phys.* **179** (2014) 509 [[arXiv:1306.5682](#)] [[INSPIRE](#)].
- [50] L. Bishler and A. Morozov, *Perspectives of differential expansion*, *Phys. Lett. B* **808** (2020) 135639 [[arXiv:2006.01190](#)] [[INSPIRE](#)].
- [51] R. Gelca, *On the relation between the A-polynomial and the Jones polynomial*, *Math. Proc. Cambridge Phil. Soc.* **133** (2002) 311 [[math/0004158](#)].
- [52] R. Gelca and J. Sain, *The noncommutative A-ideal of a $(2,2p+1)$ -torus knot determines its Jones polynomial*, *J. Knot Theory Ramif.* **12** (2003) 187 [[math/0201100](#)].
- [53] S. Gukov, *Three-dimensional quantum gravity, Chern-Simons theory, and the A polynomial*, *Commun. Math. Phys.* **255** (2005) 577 [[hep-th/0306165](#)] [[INSPIRE](#)].
- [54] S. Garoufalidis and T. Le, *The colored Jones function is q-holonomic*, *Geom. Topol.* **9** (2005) 1253 [[math/0309214](#)].
- [55] K. Habiro, *On the quantum sl_2 invariants of knots and integral homology spheres*, *Geom. Topol. Monogr.* **4** (2002) 55 [[math/0211044](#)].
- [56] H. Itoyama, A. Mironov, A. Morozov and A. Morozov, *HOMFLY and superpolynomials for figure eight knot in all symmetric and antisymmetric representations*, *JHEP* **07** (2012) 131 [[arXiv:1203.5978](#)] [[INSPIRE](#)].
- [57] A. Morozov, *Differential expansion and rectangular HOMFLY for the figure eight knot*, *Nucl. Phys. B* **911** (2016) 582 [[arXiv:1605.09728](#)] [[INSPIRE](#)].
- [58] M. Kameyama, S. Nawata, R. Tao and H.D. Zhang, *Cyclotomic expansions of HOMFLY-PT colored by rectangular Young diagrams*, [arXiv:1902.02275](#) [[INSPIRE](#)].
- [59] A. Morozov, *On exclusive Racah matrices $\bar{S}$ for rectangular representations*, *Phys. Lett. B* **793** (2019) 116 [[arXiv:1902.04140](#)] [[INSPIRE](#)].
- [60] Y. Kononov and A. Morozov, *On the defect and stability of differential expansion*, *JETP Lett.* **101** (2015) 831 [[arXiv:1504.07146](#)] [[INSPIRE](#)].
- [61] A. Mironov, A. Morozov, A. Morozov, P. Ramadevi and V.K. Singh, *Colored HOMFLY polynomials of knots presented as double fat diagrams*, *JHEP* **07** (2015) 109 [[arXiv:1504.00371](#)] [[INSPIRE](#)].
- [62] A. Mironov and A. Morozov, *Towards effective topological field theory for knots*, *Nucl. Phys. B* **899** (2015) 395 [[arXiv:1506.00339](#)] [[INSPIRE](#)].

- [63] A. Mironov, A. Morozov, A. Morozov, P. Ramadevi, V.K. Singh and A. Sleptsov, *Tabulating knot polynomials for arborescent knots*, *J. Phys. A* **50** (2017) 085201 [[arXiv:1601.04199](#)] [[INSPIRE](#)].
- [64] A. Mironov, A. Morozov and A. Morozov, *Tangle blocks in the theory of link invariants*, *JHEP* **09** (2018) 128 [[arXiv:1804.07278](#)] [[INSPIRE](#)].
- [65] M. Khovanov, *A categorification of the Jones polynomial*, *Duke Math. J.* **101** (2000) 359 [[math/9908171](#)] [[INSPIRE](#)].
- [66] M. Khovanov, *Patterns in knot cohomology I*, *Exp. Math.* **12** (2003) 365374 [[math/0201306](#)].
- [67] M. Khovanov, *Categorifications of the colored Jones polynomial*, *J. Knot Theory Ramif.* **14** (2005) 111 [[math/0302060](#)].
- [68] M. Khovanov, *$sl(3)$ link homology*, *Algebr. Geom. Topol.* **4** (2004) 1045 [[math/0304375](#)].
- [69] M. Khovanov, *Triply-graded link homology and Hochschild homology of Soergel bimodules*, *Int. J. Math.* **18** (2007) 869885 [[math/0510265](#)].
- [70] M. Khovanov, *Link homology and categorification*, [math/0605339](#).
- [71] M. Khovanov, *Categorifications from planar diagrammatics*, [arXiv:1008.5084](#).
- [72] D. Bar-Natan, *On Khovanov's categorification of the Jones polynomial*, *Algebr. Geom. Topol.* **2** (2002) 337 [[math/0201043](#)].
- [73] D. Bar-Natan, *Khovanov's homology for tangles and cobordisms*, *Geom. Topol.* **9** (2005) 1443 [[math/0410495](#)].
- [74] D. Bar-Natan, *Fast Khovanov Homology Computations*, *J. Knot Theory Ramif.* **16** (2007) 243255 [[math/0606318](#)].
- [75] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology*, *Fund. Math.* **199** (2008) 191 [[math/0401268](#)].
- [76] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology II*, *Geom. Topol.* **12** (2008) 1387 [[math/0505056](#)].
- [77] M. Khovanov and L. Rozansky, *Virtual crossings, convolutions and a categorification of the $SO(2N)$ Kauffman polynomial*, [math/0701333](#).
- [78] N. Carqueville and D. Murfet, *Computing Khovanov-Rozansky homology and defect fusion*, *Algebr. Geom. Topol.* **14** (2014) 489 [[arXiv:1108.1081](#)] [[INSPIRE](#)].
- [79] V. Dolotin and A. Morozov, *Introduction to Khovanov Homologies. I. Unreduced Jones superpolynomial*, *JHEP* **01** (2013) 065 [[arXiv:1208.4994](#)] [[INSPIRE](#)].
- [80] V. Dolotin and A. Morozov, *Introduction to Khovanov Homologies. III. A new and simple tensor-algebra construction of Khovanov-Rozansky invariants*, *Nucl. Phys. B* **878** (2014) 12 [[arXiv:1308.5759](#)] [[INSPIRE](#)].
- [81] S. Nawata and A. Oblomkov, *Lectures on knot homology*, *Contemp. Math.* **680** (2016) 137 [[arXiv:1510.01795](#)] [[INSPIRE](#)].
- [82] H. Awata, S. Gukov, P. Sulkowski and H. Fuji, *Volume Conjecture: Refined and Categorified*, *Adv. Theor. Math. Phys.* **16** (2012) 1669 [[arXiv:1203.2182](#)] [[INSPIRE](#)].
- [83] H. Fuji, S. Gukov and P. Sulkowski, *Super-A-polynomial for knots and BPS states*, *Nucl. Phys. B* **867** (2013) 506 [[arXiv:1205.1515](#)] [[INSPIRE](#)].

- [84] D. Bar-Natan and S. Morrison, <http://katlas.org>.
- [85] P. Dunin-Barkowski, A. Mironov, A. Morozov, A. Sleptsov and A. Smirnov, *Superpolynomials for toric knots from evolution induced by cut-and-join operators*, *JHEP* **03** (2013) 021 [[arXiv:1106.4305](#)] [[INSPIRE](#)].
- [86] S. Garoufalidis, A.D. Lauda and T.T.Q. Lê, *The colored HOMFLYPT function is q-holonomic*, *Duke Math. J.* **167** (2018) 397 [[arXiv:1604.08502](#)] [[INSPIRE](#)].
- [87] S.Chmutov, S.Duzhin and J.Mostovoy, *Introduction to Vassiliev Knot Invariants*, Cambridge University Press, Cambridge U.K. (2012) [ISBN 978-1-107-02083-2] [[arXiv:1103.5628](#)].
- [88] A. Mironov, A. Morozov and A. Sleptsov, *Genus expansion of HOMFLY polynomials*, *Teor. Mat. Fiz.* **177** (2013) 179 [[arXiv:1303.1015](#)] [[INSPIRE](#)].
- [89] A. Mironov, A. Morozov and A. Sleptsov, *On genus expansion of knot polynomials and hidden structure of Hurwitz tau-functions*, *Eur. Phys. J. C* **73** (2013) 2492 [[arXiv:1304.7499](#)] [[INSPIRE](#)].
- [90] A. Mironov, A. Morozov, A. Sleptsov and A. Smirnov, *On genus expansion of superpolynomials*, *Nucl. Phys. B* **889** (2014) 757 [[arXiv:1310.7622](#)] [[INSPIRE](#)].
- [91] M. Aganagic, T. Ekhholm, L. Ng and C. Vafa, *Topological Strings, D-Model, and Knot Contact Homology*, *Adv. Theor. Math. Phys.* **18** (2014) 827 [[arXiv:1304.5778](#)] [[INSPIRE](#)].
- [92] A. Morozov, *KNTZ trick from arborescent calculus and the structure of differential expansion*, *Theor. Math. Phys.* **204** (2020) 863 [[arXiv:2001.10254](#)] [[INSPIRE](#)].
- [93] A. Morozov, *Knot polynomials for twist satellites*, *Phys. Lett. B* **782** (2018) 104 [[arXiv:1801.02407](#)] [[INSPIRE](#)].
- [94] P. Dunin-Barkowski, A. Popolitov and S. Popolitova, *Evolution for Khovanov polynomials for figure-eight-like family of knots*, [arXiv:1812.00858](#) [[INSPIRE](#)].
- [95] A. Anokhina, A. Morozov and A. Popolitov, *Nimble evolution for pretzel Khovanov polynomials*, *Eur. Phys. J. C* **79** (2019) 867 [[arXiv:1904.10277](#)] [[INSPIRE](#)].
- [96] The On-Line Encyclopedia of Integer Sequences, <http://oeis.org>.