

One-loop amplitudes for $\mathcal{N} = 2$ homogeneous supergravities

Maor Ben-Shahar and Marco Chiodaroli

*Department of Physics and Astronomy, Uppsala University,
Uppsala SE-75108, Sweden*

E-mail: benshahar.maor@physics.uu.se, marco.chiodaroli@physics.uu.se

ABSTRACT: We compute one-loop matter amplitudes in homogeneous Maxwell-Einstein supergravities with $\mathcal{N} = 2$ supersymmetry using the double-copy construction. We start from amplitudes of $\mathcal{N} = 2$ super-Yang-Mills theory with matter that obey manifestly the duality between color and kinematics. Taking advantage of the fact that amplitudes with external hypermultiplets have kinematical numerators which do not present any explicit dependence on the loop momentum, we find a relation between the one-loop divergence of the supergravity amplitudes and the beta function of the non-supersymmetric gauge theory entering the construction. Two distinct linearized counterterms are generated at one loop. The divergence corresponding to the first is nonzero for all homogeneous supergravities, while the divergence associated to the second vanishes only in the case of the four Magical supergravities.

KEYWORDS: Extended Supersymmetry, Scattering Amplitudes, Supergravity Models, Supersymmetric Gauge Theory

ARXIV EPRINT: [1812.00402](https://arxiv.org/abs/1812.00402)

Contents

1	Introduction and summary of results	1
2	Double-copy construction for homogeneous supergravities	3
3	One-loop gauge amplitudes with hypermultiplets	6
4	One-loop supergravity amplitudes from the double copy	17
4.1	Examples	20
5	Discussion	21
A	Conventions	24
B	Feynman rules	25
C	Details on the orbifold numerators	26
D	Integral reduction	27

1 Introduction and summary of results

In the past decade, the duality between color and kinematics and the related double-copy construction [1, 2] have produced a breath-taking increase in our ability to conduct multi-loop calculations in gravity and supergravity theories. The double-copy method expresses loop-level gravity integrands in terms of gauge-theory building blocks, which are typically far easier to obtain. Taking advantage of this construction, calculations have been performed up to five loops in maximal supergravity [3–5] and up to four loops in half-maximal supergravity [6–8]. The studies of ultraviolet (UV) behaviors which have been made possible by these calculations have uncovered a set of so-called enhanced cancellations [9, 10] and revealed a connection between UV-divergences and U(1) quantum anomalies [11, 12].

Going beyond very special theories with extended supersymmetry, it has become clear that there exists a large web of double-copy-constructible theories, which includes pure ungauged supergravities [13, 14], ungauged supergravities with various matter contents [15–20], Yang-Mills-Einstein theories [21, 22], spontaneously-broken supergravities [23], gauged supergravities [24] and conformal supergravities [25, 26]. Conversely, color/kinematics duality has been established for large classes of gauge theories, including for example QCD [27, 28]. Various non-gravitational theories, most prominently the Dirac-Born-Infeld theory and the non-linear sigma model, can also be understood in terms of the double copy and can provide examples in which the duality between color and kinematics is particularly

well-understood [29–33]. In addition, the double copy has been instrumental in explaining amplitude relations between Einstein-Yang-Mills and Yang-Mills (YM) theories [34] that were first observed in ref. [35] (see also refs. [36–42]).

In the case of gravitational theories that can be directly obtained from string theory, insight about color/kinematics duality can be obtained from string monodromy relations [43–48]. Duality-satisfying structures can be obtained from the low-energy limit of string-theory amplitudes [49–52]. New sets of double-copy relations that combine string and field-theory building blocks have also become a powerful tool for understanding the structure of various string-theory amplitudes [53–56]. In addition, the double copy is an intrinsic feature of other modern approaches to scattering amplitudes, including the Cachazo-He-Yuan formalism [48, 57–62] and ambitwistor strings [63, 64].

Given the considerable progress in establishing double-copy structures for wider and wider classes of theories, it is natural to seek additional examples in which the double copy can be employed for shedding some light on the loop-level properties of supergravities with matter. A related issue is that the double-copy is by nature a construction at the integrand level, a fact that makes it more difficult to establish a direct relation between the physical properties of a gravity theory and the ones of the gauge theories entering the construction, which become manifest after integration.

In this paper we focus on the double-copy construction for Maxwell-Einstein supergravities with $\mathcal{N} = 2$ supersymmetry in four and five dimensions and homogeneous target spaces, which was first presented in ref. [17]. These theories have been explicitly classified in the supergravity literature [65] and provide a natural testing ground for amplitude methods. Their double-copy construction involves a $\mathcal{N} = 2$ super-Yang-Mills (sYM) theory with one half-hypermultiplet in a pseudo-real representation as one gauge-theory factor, and a non-supersymmetric YM theory with adjoint scalars and matter fermions in the same representation as the other gauge-theory factor.

Loop amplitudes with four external hypermultiplets that obey the duality between color and kinematics were first obtained in ref. [66]. These amplitudes have the striking property that the loop momentum does not appear in the kinematical numerators. Focusing on supergravity amplitudes between four vector multiplets in which the vectors are constructed as the product of two spin-1/2 asymptotic states, the above property allows us to find an explicit relation between supergravity amplitudes and gauge-theory physical quantities which holds at the integrated level.

Our main result is that the one-loop divergence for the (super)amplitudes we are inspecting admits a very compact expression,

$$\mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} = \frac{2is\delta^4(Q)}{\langle 12 \rangle \langle 34 \rangle} \left(\frac{\kappa}{2}\right)^4 \left\{ sA_{s,\phi}^{\text{tree}} \left(1 + \frac{n_F}{4} - \frac{n_S}{2}\right) + sA_{s,A}^{\text{tree}} \left(\frac{11}{6} + \frac{n_F}{6} - \frac{n_S}{12}\right) \right\} \frac{c_\Gamma}{\epsilon} + \text{Perms}, \tag{1.1}$$

where n_F and n_S are the numbers of four-dimensional fermions and scalars in the non-supersymmetric gauge theory entering the double-copy construction, and $c_\Gamma = i/(4\pi)^2$. The presence of the Grassmann delta function in (1.1) makes $\mathcal{N} = 2$ supersymmetry manifest. In turn, the expression (1.1) can be directly related to a linear combination of

various parts of the one-loop beta function of the non-supersymmetric gauge theory,

$$\mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} = -\frac{s\delta^4(Q)}{\langle 12 \rangle \langle 34 \rangle} \left(\frac{\kappa}{2}\right)^4 \left\{ sA_{s,\phi}^{\text{tree}} \left(\beta_\phi \Big|_{T(G)} - \frac{\beta_\phi}{2} \Big|_{T(R)} \right) + sA_{s,A}^{\text{tree}} \left(\beta_A \Big|_{T(G)} - \frac{\beta_A}{2} \Big|_{T(R)} \right) \right\} \frac{c_T}{\epsilon} + \text{Perms}, \tag{1.2}$$

where β_ϕ, β_A are the beta functions for the gauge and Yukawa couplings and $\beta \Big|_{T(G), T(R)}$ are the parts of the relevant beta function which are proportional to the index of the adjoint and matter representations. $A_{s,A}^{\text{tree}}$ and $A_{s,\phi}^{\text{tree}}$ are the contributions to the non-supersymmetric gauge theory’s tree-level s -channel from gluon and scalar exchange, respectively.

This paper is structured as follows. In section 2, we review the construction for homogeneous supergravities and introduce the gauge-theory Lagrangians employed in the calculation. In section 3, we consider one-loop gauge-theory amplitudes with external hypermultiplets and extend the results of ref. [66] to the pseudo-real case by restoring the permutation symmetry at the superamplitude’s level. In section 4, we compute one-loop amplitudes for the non-supersymmetric gauge theory and use the double-copy method for obtaining gravity amplitudes. We then conduct multiple checks and comparisons with previous results in the literature and end with a discussion of future directions.

2 Double-copy construction for homogeneous supergravities

The double-copy construction relies on organizing gauge theory amplitudes at L loops and n points as sums over a set of cubic graphs,

$$\mathcal{A}_n^{(L)} = i^{L-1} g^{n-2+2L} \sum_{i \in \text{cubic}} \int \frac{d^{LD} \ell}{(2\pi)^{LD}} \frac{1}{S_i} \frac{C_i n_i}{D_i}, \tag{2.1}$$

where D_i are products of inverse scalar propagators and S_i are symmetry factors. Each cubic graph is associated to a color factor C_i , which is constructed out of gauge-group invariant tensors (i.e., structure constants, representation matrices and Clebsch-Gordan coefficients with three matter-representation indices). Since the relevant group-theoretical objects obey algebraic relations (e.g. commutation relations and Jacobi identities), there exist triplets of graphs $\{i, j, k\}$ such that $C_i + C_j + C_k = 0$. Graphs in the above amplitude presentation are further associated to numerator factors, which contain theory-specific kinematical information and involve external and loop momenta, as well as Grassmann variables in case of superamplitudes. A theory obeys color/kinematics duality if the amplitude numerators possess the same algebraic properties as the corresponding color factors,

$$n_i - n_j = n_k \quad \Leftrightarrow \quad C_i - C_j = C_k. \tag{2.2}$$

The double-copy construction utilizes duality-satisfying numerators to write (super)gravity amplitudes by replacing the color factors with a second set of numerators,

$$\mathcal{M}_n^{(L)} = i^{L-1} \left(\frac{\kappa}{2}\right)^{n-2+2L} \sum_{i \in \text{cubic}} \int \frac{d^{LD} \ell}{(2\pi)^{LD}} \frac{1}{S_i} \frac{n_i \tilde{n}_i}{D_i}, \tag{2.3}$$

where κ is the gravity coupling. The numerators n_i, \tilde{n}_i may come from different gauge theories which share the same set of color factors. The formula requires that at least one of the two sets of numerators satisfies manifestly color/kinematics duality [1, 2]. If this requirement is satisfied, it is possible to show that the double-copy amplitudes obey the Ward identities corresponding to linearized diffeomorphisms and hence have the interpretation of amplitudes from some gravitational theory [34]. In general, identifying the gravity theory produced by a given double-copy construction (i.e. reconstructing the corresponding Lagrangian) is a highly-nontrivial problem. However, symmetry considerations, combined with minimal information on the supergravity interactions, have been instrumental in applying the double-copy method to very large classes of theories.

Among the double copies formulated to date, the construction for $\mathcal{N} = 2$ homogeneous Maxwell-Einstein supergravities plays a key role. Supergravities in this family have been explicitly classified by the supergravity literature [65] and hence give a natural testing ground for the double-copy method. Their double-copy construction has the following schematic form [17]:

$$\left(\mathcal{N} = 2 \text{ hom. sugras} \right) = \left(\mathcal{N} = 2 \text{ sYM} + \frac{1}{2} \text{ hyper} \right) \otimes (\text{YM}_D + P \text{ fermions}) . \quad (2.4)$$

The right gauge theory entering the construction is a YM theory in D spacetime dimensions with P additional fermions which transform in a pseudo-real gauge group representation. The corresponding Lagrangian is

$$\mathcal{L}_R = -\frac{1}{4} F_{\mu\nu}^{\hat{a}} F^{\hat{a}\mu\nu} + \frac{i}{2} \bar{\lambda} \Gamma^\mu D_\mu \lambda, \quad (2.5)$$

where field strengths and covariant derivatives are defined as

$$F_{\mu\nu}^{\hat{a}} = \partial_\mu A_\nu^{\hat{a}} - \partial_\nu A_\mu^{\hat{a}} + g f^{\hat{a}\hat{b}\hat{c}} A_\mu^{\hat{b}} A_\nu^{\hat{c}}, \quad (2.6)$$

$$D_\mu \lambda = \partial_\mu \lambda - ig T_R^{\hat{a}} A_\mu^{\hat{a}} \lambda. \quad (2.7)$$

Here $\mu, \nu = 0, \dots, D - 1$ are spacetime indices and we use a mostly-minus metric. The D -dimensional gamma matrices obey the Clifford algebra relation

$$\{\Gamma^\mu, \Gamma^\nu\} = 2\eta^{\mu\nu}, \quad (2.8)$$

where all gamma matrices except Γ^0 are taken to be antihermitian (our conventions are summarized in appendix A). $\hat{a}, \hat{b}, \hat{c}$ are adjoint indices of the gauge group.¹ The fermions' indices are not displayed explicitly. Four-dimensional fermions carry three different indices: spacetime spinor indices, gauge-group representation indices and global/ flavor indices. Fermions obey reality conditions of the form

$$\bar{\lambda} = \lambda^t C V, \quad (2.9)$$

where C is a matrix acting on spinor and flavor indices and V is a matrix acting on the gauge-group representation indices which is taken to be unitary and antisymmetric. The representation matrices for the fermions obey

$$V T_R^{\hat{a}} V^\dagger = -(T_R^{\hat{a}})^*, \quad (2.10)$$

¹For notational clarity, we will hat all gauge-group indices throughout the paper.

D	$n_F(D, P, \dot{P})$	conditions	flavor group
5	P	R	$SO(P)$
6	$P + \dot{P}$	RW	$SO(P) \times SO(\dot{P})$
7	$2P$	R	$SO(P)$
8	$4P$	R/W	$U(P)$
9	$8P$	PR	$USp(2P)$
10	$8P + 8\dot{P}$	PRW	$USp(2P) \times USp(2\dot{P})$
11	$16P$	PR	$USp(2P)$
12	$16P$	R/W	$U(P)$
$k + 8$	$16 n_F(k, P, \dot{P})$	as for k	as for k

Table 1. Parameters in the construction of $\mathcal{N} = 2$ homogeneous Maxwell-Einstein supergravities as double copies. The second column gives the number n_F of irreducible spinors once the non-supersymmetric gauge-theory factor is reduced to $4D$. Reality (R), pseudo-reality (PR) or Weyl (W) conditions are used to obtain irreducible spinors in D dimensions.

i.e. the representation R is pseudo-real. The spinor λ includes P flavors of irreducible $SO(D - 1, 1)$ spinors, which obey reality (R) or pseudo-reality (PR) conditions, according to the choice of C in equation (2.9). Specifically, we have

$$R: C = \mathcal{C}_D, \quad PR: C = \mathcal{C}_D \Omega, \quad (2.11)$$

where \mathcal{C}_D is the $SO(D - 1, 1)$ charge-conjugation matrix and Ω is an antisymmetric real matrix acting on the flavor indices. The dimensions in which R and PR conditions are appropriate are listed in table 1 together with the corresponding flavor symmetry. For $D = 6, 10 \pmod{8}$, Weyl conditions are compatible with R and PR conditions. Since representations with opposite chiralities are inequivalent, we need to introduce two distinct integers P and \dot{P} giving the number of irreducible D -dimensional spinors for each chirality. Results of this paper are more naturally expressed in terms of the number of four-dimensional fermions n_F , which is also listed in table 1.

The left gauge theory entering the double-copy construction is a $\mathcal{N} = 2$ sYM theory with a single half-hypermultiplet transforming in the same pseudo-real representation R which appears in the other gauge theory. This choice is motivated by the fact that a pseudo-real half-hypermultiplet is CPT self-conjugate and hence does not need to be augmented to a full hypermultiplet to have a sensible theory. It is convenient to consider a sYM theory in $D = 6$ spacetime dimensions with Lagrangian

$$\mathcal{L}_L = -\frac{1}{4} F_{\mu\nu}^{\hat{a}} F^{\hat{a}\mu\nu} + i\bar{\psi}\Gamma^\mu D_\mu\psi + D_\mu\bar{\varphi}D^\mu\varphi + \frac{i}{2}\bar{\chi}\Gamma^\mu D_\mu\chi + \sqrt{2}\bar{\psi}^{\hat{a}}\bar{\varphi}T_R^{\hat{a}}\chi + \sqrt{2}\bar{\chi}T_R^{\hat{a}}\varphi\psi^{\hat{a}}, \quad (2.12)$$

where the covariant derivatives are

$$(D_\mu\psi)^{\hat{a}} = \partial_\mu\psi^{\hat{a}} + gf^{\hat{a}\hat{b}\hat{c}}A_\mu^{\hat{b}}\psi^{\hat{c}}, \quad (2.13)$$

$$D_\mu\chi = \partial_\mu\chi - igT_R^{\hat{a}}A_\mu^{\hat{a}}\chi, \quad (2.14)$$

$$D_\mu\varphi = \partial_\mu\varphi - igT_R^{\hat{a}}A_\mu^{\hat{a}}\varphi. \quad (2.15)$$

The spin-1/2 field $\psi^{\hat{a}}$ transforms in the adjoint representation and is the supersymmetric partner of the gluon. It obeys a chirality condition of the form

$$\Gamma_7 \psi = \psi, \quad \Gamma_7 = \Gamma^0 \cdots \Gamma^5. \quad (2.16)$$

The complex scalar φ and the spinor χ are the components of the half-hypermultiplet. They obey the conditions,

$$\bar{\chi} = \chi^t C V, \quad \Gamma_7 \chi = \chi. \quad (2.17)$$

Amplitudes in this theory can be conveniently organized into superamplitudes with manifest $\mathcal{N} = 2$ supersymmetry. While the Lagrangian in this section can be used for Feynman-rule computations, the quickest way to obtain amplitudes for the left theory is to start from $\mathcal{N} = 4$ amplitudes and use an orbifold procedure, as explained in the next section.

Generic homogeneous supergravities are characterized by the property that their scalar manifolds admit a transitive group of isometries. In four-dimensions they have a U-duality algebra which can be decomposed into grade 0, 1, 2 generators, with the grade-zero part

$$\mathcal{G}_0 = \text{so}(1, 1) \oplus \text{so}(D - 4, 2) \oplus \mathcal{S}(P, \dot{P}), \quad (2.18)$$

where $\mathcal{S}(P, \dot{P})$ is the flavor algebra listed in table 1. A $\text{so}(D - 4) \oplus \mathcal{S}(P, \dot{P})$ subalgebra is already manifest at the level of the gauge theories entering the construction.

In particular cases, however, symmetry enhancement takes place so that the scalar manifold becomes a symmetric space. This results in the appearance of additional grade -1 and -2 generators in the U-duality algebra. There exists one infinite family of theories with symmetric scalar manifolds both in five and four dimensions, the so-called Generic Jordan family [67, 68]. It has two distinct double-copy realizations, one with $P = 0$ and arbitrary D and the other with $D = 6$ and arbitrary P . There also exist four exceptional cases of theories with symmetric scalar manifolds, the so-called *Magical Supergravities* [67, 69], that correspond to $D = 7, 8, 10, 14$ and $P = 1$. We will discuss more in detail the UV properties of these theories in the following sections.²

3 One-loop gauge amplitudes with hypermultiplets

Field-theory orbifolds [70] are constructed from a parent theory, which in this case is $\mathcal{N} = 4$ sYM, by projecting out states that are not invariant under the transformation

$$\Phi \rightarrow R_i g_i \Phi g_i^\dagger, \quad (R_i, g_i) \in \Gamma, \quad (3.1)$$

where Φ is a generic field of the theory, g_i is an element of the gauge group G and R_i is a matrix acting on the R-symmetry indices of the theory. Γ is taken to be a discrete abelian

²The literature has long discussed the relation between supergravities with symmetric scalar manifolds in five dimensions and Jordan algebras of degree three, as well as the connection between the Magical Supergravities and the four division algebras. However, discussing these connections lies outside the scope of the present paper.

subgroup of $G \times \text{SU}(4)$. Orbifold amplitudes can be conveniently obtained by inserting the projectors

$$\mathcal{P}_\Gamma \Phi = \frac{1}{|\Gamma|} \sum_{(R_i, g_i) \in \Gamma} R_i g_i \Phi g_i^\dagger, \tag{3.2}$$

in the propagators of the parent theory, where $|\Gamma|$ is the order of the orbifold group. As a consequence of the invariance of the Feynman-rule vertices under gauge and R-symmetry, projectors can be moved around the diagram past individual vertices. For tree-level amplitudes, this observation implies that orbifold amplitudes are identical to the ones of the parent theory, provided that the external states are chosen to be invariant under the orbifold group. One-loop amplitudes that preserve color/kinematics duality for general abelian orbifolds of $\mathcal{N} = 4$ sYM theory were obtained in ref. [66]. They are generally organized in a presentation based on cubic graphs in which the internal line labeled by the loop momentum is dressed with an extra phase factor, which is in turn expressed in terms of the entries of a diagonal matrix encoding the action of the R-symmetry part of the orbifold generators on fundamental $\text{SU}(4)$ indices.

Without repeating the derivation in ref. [66] and referring to appendix C for details, we focus on amplitudes with four external hypermultiplets. Hypermultiplet asymptotic states can be conveniently organized in an on-shell superfield \mathcal{Q} ,

$$\mathcal{Q} = \chi_+ + \eta^r \varphi_r + \eta^1 \eta^2 \tilde{\chi}_-, \quad r = 1, 2, \tag{3.3}$$

together with the CPT-conjugate superfield $\bar{\mathcal{Q}}$. When the representation is pseudo-real, we can identify $\mathcal{Q} = \bar{\mathcal{Q}}$ and $\chi_\pm = \tilde{\chi}_\pm$. The superamplitude between four hypermultiplets is then written as

$$\mathcal{F}_4^{\text{1-loop}}(1\mathcal{Q}, 2\mathcal{Q}, 3\bar{\mathcal{Q}}, 4\bar{\mathcal{Q}}) = -g^4 \sum_{\mathcal{G}_3} \frac{1}{S_i} \frac{n_i C_i}{D_i}. \tag{3.4}$$

Color and numerator factors are listed in table 2. While we have obtained these amplitudes with a particular choice of gauge group and representation matrices, amplitudes with any other choice of gauge group and complex representation have the same formal expression. In later sections, we will simply use the numerators without any explicit reference to the orbifold procedure we have employed in their derivation.

It is interesting to look at the color and numerator relations obeyed by the above amplitude. Some color relations stem from the representation-matrices commutation relations $[\tilde{T}^{\hat{a}}, \tilde{T}^{\hat{b}}] = \tilde{f}^{\hat{a}\hat{b}\hat{c}} \tilde{T}^{\hat{c}}$.³

$$\begin{aligned} C_1 - C_{2b} &= C_{5b} = C_{7b}, \\ C_3 - C_{2a} &= C_{8b} = -C_{9b}, \\ C_{11a} &= 2C_{8b} = -2C_{9b}, \\ C_{12a} &= 2C_{5b} = 2C_{7b}. \end{aligned} \tag{3.5}$$

The reader can verify that these relations are obeyed by the corresponding numerator factors.

³Throughout the paper we have $\tilde{f}^{\hat{a}\hat{b}\hat{c}} = i\sqrt{2}f^{\hat{a}\hat{b}\hat{c}}$ and $\tilde{T}^{\hat{a}} = \sqrt{2}T^{\hat{a}}$.

i	\mathcal{G}	C_i, n_i
1		$C_1 = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}}^{\hat{\delta}}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}}^{\hat{\gamma}}$ $n_1 = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.6)$
2a		$C_{2a} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}}^{\hat{\gamma}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}})_{\hat{\beta}}^{\hat{\delta}}$ $n_{2a} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.7)$
2b		$C_{2b} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}}^{\hat{\delta}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}})_{\hat{\beta}}^{\hat{\gamma}}$ $n_{2b} = -\frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.8)$
3		$C_3 = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}}^{\hat{\gamma}}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}}^{\hat{\delta}}$ $n_3 = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.9)$
5a		$C_{5a} = -\tilde{T}_{\hat{\beta}}^{\hat{\alpha}}{}^{\hat{\gamma}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}}^{\hat{\delta}} = (T(G) - 2C(R))\tilde{T}_{\hat{\alpha}}^{\hat{\delta}}\tilde{T}_{\hat{\beta}}^{\hat{\alpha}}{}^{\hat{\gamma}}$ $n_{5a} = \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.10)$
5b		$C_{5b} = \tilde{T}_{\hat{\beta}}^{\hat{\alpha}}{}^{\hat{\gamma}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\alpha}}^{\hat{\delta}} = -T(G)\tilde{T}_{\hat{\alpha}}^{\hat{\delta}}\tilde{T}_{\hat{\beta}}^{\hat{\alpha}}{}^{\hat{\gamma}}$ $n_{5b} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.11)$

<p>7a</p>		$C_{7a} = -\tilde{T}_{\hat{\alpha}}^{\hat{\delta}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}}^{\hat{\gamma}} = (T(G) - 2C(R))\tilde{T}_{\hat{\alpha}}^{\hat{\delta}}\tilde{T}_{\hat{\beta}}^{\hat{\gamma}}$ $n_{7a} = \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.12)$
<p>7b</p>		$C_{7b} = \tilde{T}_{\hat{\alpha}}^{\hat{\delta}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\beta}}^{\hat{\gamma}} = -T(G)\tilde{T}_{\hat{\alpha}}^{\hat{\delta}}\tilde{T}_{\hat{\beta}}^{\hat{\gamma}}$ $n_{7b} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.13)$
<p>8a</p>		$C_{8a} = -\tilde{T}_{\hat{\alpha}}^{\hat{\gamma}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}}^{\hat{\delta}} = (T(G) - 2C(R))\tilde{T}_{\hat{\alpha}}^{\hat{\gamma}}\tilde{T}_{\hat{\beta}}^{\hat{\delta}}$ $n_{8a} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.14)$
<p>8b</p>		$C_{8b} = \tilde{T}_{\hat{\alpha}}^{\hat{\gamma}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\beta}}^{\hat{\delta}} = -T(G)\tilde{T}_{\hat{\alpha}}^{\hat{\gamma}}\tilde{T}_{\hat{\beta}}^{\hat{\delta}}$ $n_{8b} = -\frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.15)$
<p>9a</p>		$C_{9a} = \tilde{T}_{\hat{\beta}}^{\hat{\delta}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}}^{\hat{\gamma}} = -(T(G) - 2C(R))\tilde{T}_{\hat{\alpha}}^{\hat{\gamma}}\tilde{T}_{\hat{\beta}}^{\hat{\delta}}$ $n_{9a} = -\frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.16)$
<p>9b</p>		$C_{9b} = -\tilde{T}_{\hat{\beta}}^{\hat{\delta}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\alpha}}^{\hat{\gamma}} = T(G)\tilde{T}_{\hat{\alpha}}^{\hat{\gamma}}\tilde{T}_{\hat{\beta}}^{\hat{\delta}}$ $n_{9b} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.17)$

11a		$C_{11a} = \tilde{T}_{\hat{\alpha}}^{\hat{\gamma}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} \tilde{f}^{\hat{b}\hat{c}\hat{d}} \tilde{T}_{\hat{\beta}}^{\hat{\delta}} = -2T(G) \tilde{T}_{\hat{\alpha}}^{\hat{\gamma}} \tilde{T}_{\hat{\beta}}^{\hat{\delta}}$ $n_{11a} = -2 \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.18)$
11b		$C_{11b} = -\tilde{T}_{\hat{\alpha}}^{\hat{\gamma}} \text{Tr}(\tilde{T}^{\hat{a}} \tilde{T}^{\hat{b}}) \tilde{T}_{\hat{\beta}}^{\hat{\delta}} = -2T(R) \tilde{T}_{\hat{\alpha}}^{\hat{\delta}} \tilde{T}_{\hat{\beta}}^{\hat{\delta}}$ $n_{11b} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.19)$
12a		$C_{12a} = \tilde{T}_{\hat{\alpha}}^{\hat{\delta}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} \tilde{f}^{\hat{b}\hat{c}\hat{d}} \tilde{T}_{\hat{\beta}}^{\hat{\gamma}} = -2T(G) \tilde{T}_{\hat{\alpha}}^{\hat{\delta}} \tilde{T}_{\hat{\beta}}^{\hat{\gamma}}$ $n_{12a} = -2 \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.20)$
12b		$C_{12b} = -\tilde{T}_{\hat{\alpha}}^{\hat{\delta}} \text{Tr}(\tilde{T}^{\hat{a}} \tilde{T}^{\hat{b}}) \tilde{T}_{\hat{\beta}}^{\hat{\gamma}} = -2T(R) \tilde{T}_{\hat{\alpha}}^{\hat{\delta}} \tilde{T}_{\hat{\beta}}^{\hat{\gamma}}$ $n_{12b} = \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.21)$

Table 2. Color and numerator factors for the 1-loop superamplitude with four external hypermultiplets in case of a complex representation. Curly lines denote adjoint fields, while solid lines denote hypermultiplet matter fields. Arrows are assigned according to the representation and we use the short-hand notation $\delta^4(Q) = \delta^4(\sum_i \eta_i^r |i\rangle)$. The numbering of the graphs follows the one in ref. [66]. Gauge-group representation indices $\hat{\alpha}, \hat{\beta}$ are written explicitly.

Other relations stem from the two-term identity $T_{\hat{\alpha}}^{\hat{\gamma}} T_{\hat{\beta}}^{\hat{\delta}} = T_{\hat{\alpha}}^{\hat{\delta}} T_{\hat{\beta}}^{\hat{\gamma}}$:

$$\begin{aligned}
 C_1 &= C_3, \\
 C_{2b} &= -C_{8a} = C_{9a}, \\
 C_{2a} &= -C_{5a} = -C_{7a}, \\
 C_{11b} &= C_{8a} = -C_{9a}, \\
 C_{12b} &= C_{7a} = C_{5a}.
 \end{aligned} \quad (3.22)$$

These relations are also obeyed by the corresponding numerator factors. If we chose a different complex representation with respect to the one from orbifolding, the color relations

would no longer hold, but numerator relations would be unaltered. For later convenience, color factors in table 2 are rewritten in terms of the representation's index $T(R)$ and quadratic Casimir $C(R)$,

$$\text{Tr}(\tilde{T}_R^{\hat{a}}\tilde{T}_R^{\hat{b}}) = 2T(R)\delta^{\hat{a}\hat{b}}, \quad \tilde{T}_R^{\hat{a}}\tilde{T}_R^{\hat{a}} = 2C(R) \mathbf{1}_R. \quad (3.23)$$

While the numerators obtained so far are appropriate for complex representations, the construction outlined in the previous section notably uses pseudo-real representations as a key ingredient. It is quite instructive to see how the construction is modified in this case. On general grounds, using a reality condition allows us to relate fermions with their Dirac conjugates. For four-fermion amplitudes, this implies that a (fermionic) permutation symmetry needs to be introduced on all external legs. This symmetry extends to superamplitudes with four-external half-hypermultiplets, taking into account that hypermultiplet on-shell superfields anticommute with each other (since their lowest component is Grassmann-odd). In short, duality-satisfying numerators can be written down for the pseudo-real case as the unique set of numerators with the following two properties:

1. Numerators are invariant under the permutation of all external legs up to a sign which is assigned according to the signature of the permutation.
2. Numerators corresponding to color factors which are non-zero in the case of a complex representation reproduce the ones listed in the table 2.

The complete list of numerators is given in table 3.

Numerators with different assignments of external legs are related by permutation symmetry, as it can be verified using the identities

$$\frac{s}{\langle 12 \rangle \langle 34 \rangle} = -\frac{u}{\langle 13 \rangle \langle 24 \rangle} = \frac{t}{\langle 14 \rangle \langle 23 \rangle}. \quad (3.24)$$

The identity

$$(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})^t = -\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}, \quad (3.25)$$






which relies on the antisymmetry of V , is also useful. An important difference between amplitudes in the complex and pseudo-real cases is that the symmetry factor for the bubble integrals with internal half-hypermultiplets needs to be changed to take into account that the corresponding graphs no longer carry arrows.

The additional nonzero color and numerator factors obey identities of the form







$$\begin{aligned} C_{1b} - C_{3b} &= -C_{6b} = -C_{4b}, \\ C_{4b} + C_{6b} &= C_{10a}, \end{aligned} \quad (3.26)$$

which stem from the gauge-group generators commutation relations. However, the extra identities in which gauge-group generators are contracted by an adjoint index require a more detailed discussion. In contrast to the complex case, color factors will no longer obey identities of the form $T_{\hat{\alpha}}^{\hat{a}} T_{\hat{\beta}}^{\hat{a}} = T_{\hat{\alpha}}^{\hat{a}} T_{\hat{\beta}}^{\hat{a}}$. While it would be natural to impose additional three-term identities for both color and numerator factors in the pseudo-real

i	\mathcal{G}	C_i, n_i
1a		$C_{1a} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\delta}}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}\hat{\gamma}}$ $n_{1a} = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.27)$
1b		$C_{1b} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\beta}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}})_{\hat{\gamma}\hat{\delta}}$ $n_{1b} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.28)$
2a		$C_{2a} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\gamma}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}})_{\hat{\beta}\hat{\delta}}$ $n_{2a} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.29)$
2b		$C_{2b} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\delta}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}})_{\hat{\beta}\hat{\gamma}}$ $n_{2b} = -\frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.30)$
3a		$C_{3a} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\gamma}}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}\hat{\delta}}$ $n_{3a} = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.31)$

3b		$C_{3b} = (\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\beta}}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\gamma}\hat{\delta}}$ $n_{3b} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.32)$
4a		$C_{4a} = -\tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\gamma}\hat{\delta}} = (T(G) - 2C(R))\tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}}\tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}$ $n_{4a} = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.33)$
4b		$C_{4b} = \tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\gamma}\hat{\delta}} = -T(G)\tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}}\tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}$ $n_{4b} = -\frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.34)$
5a		$C_{5a} = -\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\delta}} = (T(G) - 2C(R))\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}$ $n_{5a} = \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.35)$
5b		$C_{5b} = \tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\alpha}\hat{\delta}} = -T(G)\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}$ $n_{5b} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.36)$

6a		$C_{6a} = -\tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\beta}} = (T(G) - 2C(R))\tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}}\tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}$ $n_{6a} = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.37)$
6b		$C_{6b} = \tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\alpha}\hat{\beta}} = -T(G)\tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}}\tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}$ $n_{6b} = -\frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.38)$
7a		$C_{7a} = -\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}\hat{\gamma}} = (T(G) - 2C(R))\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}$ $n_{7a} = \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.39)$
7b		$C_{7b} = \tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}\tilde{f}^{\hat{a}\hat{b}\hat{c}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{c}})_{\hat{\beta}\hat{\gamma}} = -T(G)\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}$ $n_{7b} = -\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.40)$
8a		$C_{8a} = -\tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}}(\tilde{T}^{\hat{b}}\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})_{\hat{\beta}\hat{\delta}} = (T(G) - 2C(R))\tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}}\tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}}$ $n_{8a} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.41)$

<p>8b</p>		$C_{8b} = \tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} (\tilde{T}^{\hat{b}} \tilde{T}^{\hat{c}})_{\hat{\beta}\hat{\delta}} = -T(G) \tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}}$ $n_{8b} = -\frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.42)$
<p>9a</p>		$C_{9a} = \tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}} (\tilde{T}^{\hat{b}} \tilde{T}^{\hat{a}} \tilde{T}^{\hat{b}})_{\hat{\alpha}\hat{\gamma}} = -(T(G) - 2C(R)) \tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}}$ $n_{9a} = -\frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.43)$
<p>9b</p>		$C_{9b} = -\tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} (\tilde{T}^{\hat{b}} \tilde{T}^{\hat{c}})_{\hat{\alpha}\hat{\gamma}} = T(G) \tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}}$ $n_{9b} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.44)$
<p>10a</p>		$C_{10a} = \tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} \tilde{f}^{\hat{b}\hat{c}\hat{d}} \tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{d}} = -2T(G) \tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}} \tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}$ $n_{10a} = -2 \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.45)$
<p>10b</p>		$C_{10b} = -\tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}} \text{Tr}(\tilde{T}^{\hat{a}} \tilde{T}^{\hat{b}}) \tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{b}} = -2T(R) \tilde{T}_{\hat{\alpha}\hat{\beta}}^{\hat{a}} \tilde{T}_{\hat{\gamma}\hat{\delta}}^{\hat{a}}$ $n_{10b} = \frac{s^2}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.46)$
<p>11a</p>		$C_{11a} = \tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} \tilde{f}^{\hat{b}\hat{c}\hat{d}} \tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{d}} = -2T(G) \tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}}$ $n_{11a} = -2 \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.47)$

11b		$C_{11b} = -\tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}} \text{Tr}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})\tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{b}} = -2T(R)\tilde{T}_{\hat{\alpha}\hat{\gamma}}^{\hat{a}}\tilde{T}_{\hat{\beta}\hat{\delta}}^{\hat{a}}$ $n_{11b} = \frac{su}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.48)$
12a		$C_{12a} = \tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}} \tilde{f}^{\hat{a}\hat{b}\hat{c}} \tilde{f}^{\hat{b}\hat{c}\hat{d}} \tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{d}} = -2T(G)\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}$ $n_{12a} = -2\frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.49)$
12b		$C_{12b} = -\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}} \text{Tr}(\tilde{T}^{\hat{a}}\tilde{T}^{\hat{b}})\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{b}} = -2T(R)\tilde{T}_{\hat{\alpha}\hat{\delta}}^{\hat{a}}\tilde{T}_{\hat{\beta}\hat{\gamma}}^{\hat{a}}$ $n_{12b} = \frac{st}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q) \quad (3.50)$

Table 3. Color and numerator factors for the 1-loop superamplitude with four external half-hypermultiplets in case of a pseudo-real representation. Note that all bubbles have symmetry factor $S_i = 2$. Representation matrices with two low indices are defined as $T_{\hat{\alpha}\hat{\beta}}^{\hat{a}} = (T^{\hat{a}}V^{-1})_{\hat{\alpha}\hat{\beta}}$, where V is the antisymmetric matrix entering the pseudo-reality condition.

case, the numerators listed in table 3 still obey the same set of two-term identities as in the complex case. It should be noted that these identities are not necessary for the consistency of the theory from the double-copy, i.e. the numerator identities stemming from the gauge-group generator commutation relations are sufficient for ensuring that the double-copy amplitudes obey the relevant Ward identities. However, it might be possible to find different amplitude presentations which apply specifically to hypermultiplets in pseudo-real representations and obey additional three-term identities in place of the two-term identities. As a consequence of the numerator relations and permutation symmetry, all numerators can be obtained in terms of a single master box numerator.

Finally, we note that the amplitude presentation in table 3 can be extended to the case of an arbitrary number n_H of half-hypermultiplets by dressing the numerator factors with Kronecker deltas with indices running over the number of half-hypermultiplets. For example, the first numerator is modified as

$$n_{1a} = \frac{s^2 \delta^{IJ} \delta^{KL}}{\langle 12 \rangle \langle 34 \rangle} \delta^4(Q). \quad (3.51)$$

Additionally, bubble numerators with a matter loop acquire an extra factor of n_H . This procedure leaves the numerator relations corresponding to (3.5) unaltered, but the extra two-term relations are lost in the generic case.⁴

4 One-loop supergravity amplitudes from the double copy

We now focus on four-dimensional supergravity superamplitudes with four vector multiplets⁵

$$\mathcal{M}_4^{1\text{-loop}}(1\mathcal{V}_a, 2\mathcal{V}_b, 3\mathcal{V}_c, 4\mathcal{V}_d), \quad (4.1)$$

where a, b, c, d are global indices running over the number of supergravity vectors that are realized as the product of two spin-1/2 asymptotic states. Note that these vectors transform as $SO(D-d)$ spinors under the global symmetry. These amplitudes can be obtained taking the double-copy of the amplitudes from the previous section with a four-fermion amplitude in the non-supersymmetric theory discussed in section 2,

$$\mathcal{A}_4^{1\text{-loop}}(1\lambda_a, 2\lambda_b, 3\lambda_c, 4\lambda_d). \quad (4.2)$$

The choice of chiralities for the external fermions determines whether a given external leg in (4.1) is associated to a $\mathcal{N} = 2$ vector on-shell superfield or its conjugate. We now focus on divergent contributions to the amplitude (4.2). Since the non-supersymmetric theory is renormalizable, there is no UV-divergent box integral, and all the amplitude's divergences are linked to the divergences of three- and two-point Green functions. In particular, we write the one-loop corrections to vertices and inverse propagators as

$$\begin{aligned} i(\Pi_\phi^{1\text{-loop}})^{\mu\nu, \hat{a}\hat{b}}(k^2) &= -ig^2 \delta^{\hat{a}\hat{b}} (\eta^{\mu\nu} - \eta_d^{\mu\nu}) k^2 \tilde{\Pi}_\phi^{1\text{-loop}}(k^2), \\ i(\Pi_A^{1\text{-loop}})^{\mu\nu, \hat{a}\hat{b}}(k^2) &= -ig^2 \delta^{\hat{a}\hat{b}} (k^2 \eta_d^{\mu\nu} - k^\mu k^\nu) \tilde{\Pi}_A^{1\text{-loop}}(k^2), \\ i\mathcal{V}_{A\bar{\lambda}\lambda}^{1\text{-loop}}(k_1, k_2, k_3) &= g^2 \mathcal{V}_{A\bar{\lambda}\lambda}^{\text{tree}}(k_1, k_2, k_3) \tilde{\mathcal{V}}_{A\bar{\lambda}\lambda}^{1\text{-loop}}(k_1, k_2, k_3), \\ i\mathcal{V}_{\phi\bar{\lambda}\lambda}^{1\text{-loop}}(k_1, k_2, k_3) &= g^2 \mathcal{V}_{\phi\bar{\lambda}\lambda}^{\text{tree}}(k_1, k_2, k_3) \tilde{\mathcal{V}}_{\phi\bar{\lambda}\lambda}^{1\text{-loop}}(k_1, k_2, k_3). \end{aligned} \quad (4.3)$$

Here we have split the D -dimensional gluons into d -dimensional gluons and d -dimensional scalars according to the values of the spacetime indices μ, ν . This is done by introducing the d -dimensional metric η_d defined as

$$\eta_d = \text{diag}\left(\underbrace{1, -1, \dots, -1}_d, \underbrace{0, \dots, 0}_{D-d}\right). \quad (4.4)$$

η_d is generated by the integral reduction identities collected in appendix D, since the loop momentum is taken to be in $d = 4 - 2\epsilon$ dimensions.

⁴Double-copying the numerators for amplitudes with n_H external half-hypermultiplets with the ones in table 3, one can reproduce the $(1 + n_V/2)$ dependence of the divergence of the amplitudes of $\mathcal{N} = 4$ supergravity with n_V vector multiplets [15]. This is an easy check of the consistency of our construction (the number of supergravity vector multiplets is related to the number of half-hypermultiplets as $n_V = n_H + 2$).

⁵By abuse of notation, we will use \mathcal{M} to denote both superamplitudes and component amplitudes.

We further split the tree-level, four-point amplitude in contributions corresponding to the three channels,

$$\mathcal{A}_4^{\text{tree}} = \mathcal{A}_{s,A}^{\text{tree}} + \mathcal{A}_{s,\phi}^{\text{tree}} + \mathcal{A}_{t,A}^{\text{tree}} + \mathcal{A}_{t,\phi}^{\text{tree}} + \mathcal{A}_{u,A}^{\text{tree}} + \mathcal{A}_{u,\phi}^{\text{tree}}, \quad (4.5)$$

where the vector and scalar channels have been written separately. Taking into account the color factors in table 3, the integrated numerator factors of the non-supersymmetric theory are related to the part of the one-loop corrections to vertices and propagators which is proportional to the indices of the adjoint and matter representations:

$$\begin{aligned} \int \frac{\tilde{n}_{4a} - \tilde{n}_{4b}}{D_4} &= A_{s,\phi}^{\text{tree}} \tilde{\mathcal{V}}_{\phi\lambda\lambda}^{\text{1-loop}}(-k_1 - k_2, k_1, k_2) \Big|_{T(G)} + A_{s,A}^{\text{tree}} \tilde{\mathcal{V}}_{A\lambda\lambda}^{\text{1-loop}}(-k_1 - k_2, k_1, k_2) \Big|_{T(G)}, \\ \frac{1}{S_{10}} \int \frac{\tilde{n}_{10a}}{D_{10}} &= \frac{1}{2} A_{s,A}^{\text{tree}} \tilde{\Pi}_A^{\text{1-loop}}(s) \Big|_{T(G)} + \frac{1}{2} A_{s,\phi}^{\text{tree}} \tilde{\Pi}_\phi^{\text{1-loop}}(s) \Big|_{T(G)}, \\ \frac{1}{S_{10}} \int \frac{\tilde{n}_{10b}}{D_{10}} &= \frac{1}{2} A_{s,A}^{\text{tree}} \tilde{\Pi}_A^{\text{1-loop}}(s) \Big|_{T(R)} + \frac{1}{2} A_{s,\phi}^{\text{tree}} \tilde{\Pi}_\phi^{\text{1-loop}}(s) \Big|_{T(R)}, \end{aligned}$$

where analogous relations for the other diagrams can be obtained by permutation. Using the numerators in table 3, we have the following expression for the UV-divergent part of the supergravity amplitude,

$$\begin{aligned} \mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} &= -\frac{2s\delta^4(Q)}{\langle 12 \rangle \langle 34 \rangle} \left(\frac{\kappa}{2} \right)^4 \left\{ s A_{s,\phi}^{\text{tree}} \left(\tilde{\mathcal{V}}_{\phi\lambda\lambda}^{\text{1-loop}} \Big|_{T(G),\text{div}} - \frac{1}{2} \tilde{\Pi}_\phi^{\text{1-loop}} \Big|_{T(G),\text{div}} + \frac{1}{4} \tilde{\Pi}_\phi^{\text{1-loop}} \Big|_{T(R),\text{div}} \right) \right. \\ &\quad \left. + s A_{s,A}^{\text{tree}} \left(\tilde{\mathcal{V}}_{A\lambda\lambda}^{\text{1-loop}} \Big|_{T(G),\text{div}} - \frac{1}{2} \tilde{\Pi}_A^{\text{1-loop}} \Big|_{T(G),\text{div}} + \frac{1}{4} \tilde{\Pi}_A^{\text{1-loop}} \Big|_{T(R),\text{div}} \right) \right\} + \text{Perms.} \end{aligned} \quad (4.6)$$

We recognize that the combination of one-loop vertex and propagator corrections in (4.6) corresponds to a piece of the beta function of the non-supersymmetric theory. The beta function $\beta(T(R), T(G), C(R))$ will formally depend on the index and quadratic Casimir of the representation, which are independent as long as no particular assumption is made on R . We can then rewrite the result above as

$$\begin{aligned} \mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} &= -\frac{s\delta^4(Q)}{\langle 12 \rangle \langle 34 \rangle} \left(\frac{\kappa}{2} \right)^4 \left\{ s A_{s,\phi}^{\text{tree}} \left(\beta_\phi \Big|_{T(G)} - \frac{\beta_\phi}{2} \Big|_{T(R)} \right) + s A_{s,A}^{\text{tree}} \left(\beta_A \Big|_{T(G)} - \frac{\beta_A}{2} \Big|_{T(R)} \right) \right\} \frac{c_T}{\epsilon} \\ &\quad + \text{Perms,} \end{aligned} \quad (4.7)$$

where $\beta_\phi \Big|_{T(G)}$ denotes the part of the beta function for the Yukawa couplings which is proportional to $T(G)$. Contributions from the wave-function renormalization of the external fermions are proportional to $C(R)$ and hence do not appear in the above expression. The factor of $-1/2$ in front of the second piece of the beta function in equation (4.7) is to be expected on physical grounds. We know that $\mathcal{N} = 4$ sYM amplitudes need to be recovered from our numerators as a particular case in which the theory has a single hypermultiplet




Figure 1. Triangle Feynman diagrams used in the computation for the non-supersymmetric gauge-theory factor.

in the adjoint representation. It is well known that amplitudes in $\mathcal{N} = 4$ sYM only have contributions from box diagrams. Hence, the numerators corresponding to diagrams 10a and 10b need to come with opposite sign. Furthermore, the factor of 1/2 arises because the amplitudes we are considering only involve one half-hypermultiplet circulating in the loop.

At one loop, the corrections to the vertices for the right-hand YM theory are obtained from the Feynman diagrams in figure 1 (see appendix B for the Feynman rules employed in the calculation),

$$\begin{aligned}
 i\mathcal{V}^{1\text{-loop}} &= g^3(VT^bT^aT^b) \frac{\bar{u}_2\Gamma^\nu(l+p_1)\Gamma^\mu(l-p_2)\Gamma_\nu u_1}{l^2(l+p_1)^2(l-p_2)^2} - \\
 &ig^3 f^{abc}(VT^bT^c)(\bar{u}_2\Gamma_\nu l\Gamma_\rho u_1) \frac{(l+2p_1+p_2)^\rho\eta^{\mu\nu} + (l-p_1-2p_2)^\nu\eta^{\mu\rho} + (p_2-2l-p_1)^\mu\eta^{\rho\nu}}{l^2(l+p_1)^2(l-p_2)^2}.
 \end{aligned}
 \tag{4.8}$$

This expression can be further simplified by employing the integral reduction identities collected in appendix D. We obtain:

$$\begin{aligned}
 \tilde{\mathcal{V}}_{A\bar{\lambda}\lambda}^{1\text{-loop}}(p_1, p_2) &= -\frac{i}{2}C(R)\{4I_2(p_1) + 4I_2(p_2) + (D - 10)I_2(p_1 + p_2)\} - iT(G)I_2(p_1 + p_2), \\
 \tilde{\mathcal{V}}_{\phi\bar{\lambda}\lambda}^{1\text{-loop}}(p_1, p_2) &= -iC(R)\{2I_2(p_1) + 2I_2(p_2) + (D - 6)I_2(p_1 + p_2)\} + \\
 &\frac{i}{2}T(G)(D - 4)I_2(p_1 + p_2),
 \end{aligned}
 \tag{4.9}$$

where we have omitted a finite part proportional to the triangle integral $I_3(p_1, p_2)$. $I_2(k)$ denote bubble integrals. Bubble-on-external-leg integrals $I_2(p_1)$ and $I_2(p_2)$ have dropped out of the part of the vertex corrections which is proportional to $T(G)$. This implies that the $1/\epsilon$ divergence of the vertex corrections that we will use in (4.6) is interpreted as a genuine UV divergence without any infrared contamination.

The calculation for the propagator corrections is performed along similar lines (see figure 2 for relevant diagrams). The final result is:

$$\begin{aligned}
 \tilde{\Pi}_\phi^{1\text{-loop}}(k^2) &= i(2T(G) - n_F T(R))I_2(k), \\
 \tilde{\Pi}_A^{1\text{-loop}}(k^2) &= -\frac{i}{6}\{(D - 14)T(G) + 4n_F T(R)\}I_2(k),
 \end{aligned}
 \tag{4.10}$$



Figure 2. Bubble Feynman diagrams used in the computation for the non-supersymmetric gauge-theory factor.

where n_F is the number of four-dimensional fermions. Equation (4.6) then becomes

$$\mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} = \frac{2is\delta^4(Q)}{\langle 12 \rangle \langle 34 \rangle} \left(\frac{\kappa}{2} \right)^4 \left\{ sA_{s,\phi}^{\text{tree}} \left(3 - \frac{D}{2} + \frac{n_F}{4} \right) + sA_{s,A}^{\text{tree}} \left(\frac{13}{6} - \frac{D}{12} + \frac{n_F}{6} \right) \right\} \frac{c_\Gamma}{\epsilon} + \text{Perms}, \quad (4.11)$$

where $c_\Gamma = i/(4\pi)^2$. By setting $D = 4 + n_S$, where n_S is the number of four-dimensional scalars, we obtain our master formula which expresses the value of the superamplitude's one-loop divergence in terms of the parameters of the construction for homogeneous supergravities:

$$\mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} = \frac{2is\delta^4(Q)}{\langle 12 \rangle \langle 34 \rangle} \left(\frac{\kappa}{2} \right)^4 \left\{ sA_{s,\phi}^{\text{tree}} \left(1 + \frac{n_F}{4} - \frac{n_S}{2} \right) + sA_{s,A}^{\text{tree}} \left(\frac{11}{6} + \frac{n_F}{6} - \frac{n_S}{12} \right) \right\} \frac{c_\Gamma}{\epsilon} + \text{Perms}. \quad (4.12)$$

4.1 Examples

To simplify the expression (4.12) further we specialize on amplitudes between four supergravity vectors,

$$\mathcal{M}^{\text{1-loop}}(1A_{a-}, 2A_{b-}, 3A_+^c, 4A_+^d) = \mathcal{A}_{\mathcal{N}=2}^{\text{1-loop}}(1\chi_-, 2\chi_-, 3\chi_+, 4\chi_+) \otimes \mathcal{A}_{\mathcal{N}=0}^{\text{1-loop}}(1\lambda_{a-}, 2\lambda_{b-}, 3\lambda_+^c, 4\lambda_+^d), \quad (4.13)$$

where the global indices of the last two vectors are raised with the inverse charge-conjugation matrix. With this assignment of external polarizations and after using spinor-helicity identities which are collected in appendix A, we get the following expressions for tree amplitudes in the three channels,

$$sA_{s,\phi}^{\text{tree}} = \frac{i}{2} \langle 12 \rangle [34] (\tilde{C}\tilde{\Gamma}^I)_{ab} (\tilde{\Gamma}_I \tilde{C}^{-1})^{cd}, \quad (4.14)$$

$$tA_{t,A}^{\text{tree}} = i \langle 12 \rangle [34] \tilde{\delta}_a^d \delta_b^c, \quad (4.15)$$

$$uA_{u,A}^{\text{tree}} = i \langle 12 \rangle [34] \tilde{\delta}_a^c \delta_b^d, \quad (4.16)$$

where we have written the higher-dimensional Dirac and charge-conjugation matrices as $\Gamma^I = \gamma_5 \otimes \tilde{\Gamma}^I$ and $C = \mathcal{C}_4 \otimes \tilde{C}$. In this case, the expression (4.12) can be simplified as

$$\mathcal{M}^{\text{1-loop}} \Big|_{\text{div}} (1A_{a-}, 2A_{-b}, 3A_+^c, 4A_+^d) = 2 \langle 12 \rangle^2 [34]^2 \left(\frac{\kappa}{2} \right)^4 \left\{ \frac{1}{2} \left(1 + \frac{n_F}{4} - \frac{n_S}{2} \right) (\tilde{C}\tilde{\Gamma}^I)_{ab} (\tilde{\Gamma}_I \tilde{C}^{-1})^{cd} + \left(\frac{11}{6} + \frac{n_F}{6} - \frac{n_S}{12} \right) (\delta_a^d \delta_b^c + \delta_a^c \delta_b^d) \right\} \frac{c_\Gamma}{\epsilon}. \quad (4.17)$$

We now consider some interesting particular cases. $n_S = 0$ ($D = 4$) corresponds to the so-called $\mathbb{CP}(n)$ or Luciani model [71]. Supergravities in this family do not uplift to dimension higher than four and have symmetric scalar manifold

$$M_{4D} = \frac{U(1, n)}{U(1) \times U(n)}, \tag{4.18}$$

where n is the number of vector multiplets. The corresponding matter amplitudes between four vectors have one-loop divergence

$$\mathcal{M}^{1\text{-loop}} \Big|_{\text{div}} (1A_{a-}, 2A_{-b}, 3A_{+}^c, 4A_{+}^d) = 2\langle 12 \rangle^2 [34]^2 \left(\frac{\kappa}{2} \right)^4 \left\{ \left(\frac{5}{3} + \frac{n}{6} \right) (\delta_a^d \delta_b^c + \delta_a^c \delta_b^d) \right\} \frac{c_\Gamma}{\epsilon}. \tag{4.19}$$

Another important example is the Generic Jordan Family [67, 68], which can be obtained by setting $D = 6$ and keeping the number of 4D fermions arbitrary (with the relation $n = n_F + 3$).⁶ This theory has symmetric target space

$$M_{4D} = \frac{SO(2, n-1)}{SO(n-1) \times SO(2)} \times \frac{SU(1, 1)}{U(1)}. \tag{4.20}$$

Focusing on amplitudes between two identical vectors and their CPT-conjugate states, we get the expression

$$\mathcal{M}^{1\text{-loop}} \Big|_{\text{div}} = 2\langle 12 \rangle^2 [34]^2 \left(\frac{\kappa}{2} \right)^4 \left\{ \left(\frac{7}{3} + \frac{n}{3} \right) \right\} \frac{c_\Gamma}{\epsilon}, \tag{4.21}$$

which reproduces the earlier result in ref. [15].

By inspecting the two terms contributing to (4.12), we see that the contribution corresponding to the vector exchange never vanishes.⁷ However, the contribution linked to the scalar exchange vanishes for

$$\begin{aligned} D = 7, \quad P = 1 & \quad (n_S = 3, \quad n_F = 2), \\ D = 8, \quad P = 1 & \quad (n_S = 4, \quad n_F = 4), \\ D = 10, \quad P = 1 & \quad (n_S = 6, \quad n_F = 8), \\ D = 14, \quad P = 1 & \quad (n_S = 10, \quad n_F = 16). \end{aligned} \tag{4.22}$$

These are precisely the four Magical supergravities [67, 69]. Additionally, the contribution from the remaining channel matches the earlier computation in ref. [17].

5 Discussion

In this paper, we have calculated the one-loop divergence for selected amplitudes between four vector multiplets in $\mathcal{N} = 2$ homogeneous Maxwell-Einstein supergravities with the

⁶As previously discussed, the Generic Jordan Family has two distinct double-copy realizations. The other realization involves a non-supersymmetric gauge theory with no matter fermions and an arbitrary number of scalars.

⁷One can also verify that the two terms cannot combine to give a vanishing divergence by contracting (4.19) with $\delta_a^c \delta_b^d$. The resulting expression never vanishes for the values of the parameters given in table 1.

double-copy construction, focusing on amplitudes between vector constructed as the double copy of two spin-1/2 fields. Supergravity amplitudes are constructed using as building blocks gauge-theory amplitudes between four hypermultiplets in a presentation that obeys color/kinematics duality. Such amplitudes were first obtained in ref. [66] in terms of kinematical numerators which do not possess any explicit dependence on the loop momentum. Because of this property, the supergravity divergence is directly linked to the beta function of the non-supersymmetric gauge theory. In a sense, our calculation presents analogies with the one in ref. [7], where the absence of some one- and two-loops divergences in half-maximal supergravity was linked to the renormalizability of the non-supersymmetric gauge theory entering the construction thanks to the absence of loop-momenta dependence in $\mathcal{N} = 4$ sYM numerators at one and two loops.

Our calculation employs an “asymmetric” version of the double-copy construction, taking advantage of the fact that only one of the two sets of numerators needs to manifestly obey color/kinematics duality. This allowed us to obtain the second set of numerators from a Feynman rule calculation, avoiding the cumbersome task of finding duality-satisfying numerators, much along the lines of the computation for $\mathcal{N} = 4$ and $\mathcal{N} = 5$ supergravity at three and four loops [6–9]. Choosing a double copy in which all the $\mathcal{N} = 2$ supersymmetry is realized by one of the two gauge-theory factors further enabled us to establish a direct link between the physical properties of the supergravity and the ones of the non-supersymmetric gauge theory. Ultimately, our calculation would have been more opaque had we chosen to focus on a $(\mathcal{N} = 1) \times (\mathcal{N} = 1)$ realization of the same supergravity theories.

Among the homogeneous supergravities, we do not find any matter amplitude which remains finite at one loop. An open question is how robust is this finding with respect to modifications of the construction. For example, we can generalize the construction by including n_ϕ complex matter scalars in the non-supersymmetric gauge theory. If the scalars are in the same pseudo-real representation as the half-hypermultiplets, the resulting supergravity theory will contain hypermultiplets in addition to the vector multiplets which are already present in the basic construction. The contribution to the supergravity divergence is modified by hypermultiplet loops and eq. (4.12) becomes

$$\mathcal{M}|_{\text{div}} = \frac{2is\delta^4(Q)}{(12)\langle 34 \rangle} \left(\frac{\kappa}{2}\right)^4 \left\{ sA_{s,\phi}^{\text{tree}} \left(1 + \frac{n_F}{4} - \frac{n_S}{2}\right) + sA_{s,A}^{\text{tree}} \left(\frac{11}{6} + \frac{n_F}{6} - \frac{n_S}{12} + \frac{n_\phi}{12}\right) \right\} \frac{c_\Gamma}{\epsilon} + \text{Perms.} \tag{5.1}$$

Hence, the additional contribution increases the divergence with respect to the Maxwell-Einstein case. Another possible modification is to add adjoint fermions. Since adjoint and matter representation contributions to (4.6) come with opposite sign, adjoint fermions alleviate the UV divergence. However, they can be introduced in a way that is consistent with color/kinematics duality only if the gauge theory becomes supersymmetric [66].

Our results can be understood in terms of counterterm analysis [72–84], which relies on the fact that the appearance of a divergence in gravity can be directly related to the existence of a local operator of the appropriate mass dimension that satisfies all non-anomalous symmetries of the theory. Hence, the challenging task of understanding loop-level divergences can be simplified by a combination of computing tree-level amplitudes and symmetry

consideration. A delicate part of the process is to analyze the constraint imposed by non-linearly-realized symmetries on candidate counterterms. This can be done by a careful inspection of single- and double-soft scalar limits [72, 75, 76], by studying solutions of Laplace equation on the relevant scalar manifolds [73, 74], or by employing a superspace formulation in which U-duality is manifest [77, 79]. Such considerations have allowed, for example, to rule out all candidate counterterms for $\mathcal{N} = 8$ supergravity up to and including six loops [75]. In contrast, counterterms in theories with matter and reduced supersymmetry are much less understood. In this respect, it is interesting to look at the counterterms corresponding to the divergences observed in our calculation. Counterterms probed by a four-vector amplitude will involve four powers of the field strength. Specializing to four dimensions, it is advantageous to write separately self-dual and anti-self-dual components of the vector field strengths by using the two-component spinor notation $F_{\alpha\beta} = \frac{1}{2}F_{\mu\nu}\sigma_{\alpha\beta}^{\mu\nu}$, $F_{\dot{\alpha}\dot{\beta}} = \frac{1}{2}F_{\mu\nu}\bar{\sigma}_{\dot{\alpha}\dot{\beta}}^{\mu\nu}$. Hence, we need to consider counterterms of the form:

$$M_{ab,cd}^{(1)}(F_{\alpha\beta}^a F^{b\alpha\beta})(F_{\dot{\alpha}\dot{\beta}}^c F^{d\dot{\alpha}\dot{\beta}}), \tag{5.2}$$

$$M_{ab,cd}^{(2)}(F_{\alpha\beta}^a F^{b\alpha\beta})(F_{\gamma\delta}^c F^{d\gamma\delta}), \tag{5.3}$$

$$M_{abcd}^{(3)}(F_{\alpha\beta}^a F^{b\beta\gamma} F_{\gamma\delta}^c F^{d\delta\alpha}), \tag{5.4}$$

where a, b, c, d are spinor indices of $\text{SO}(D - 4)$ and $M_{ab,cd}^{(1)}, M_{ab,cd}^{(2)}, M_{abcd}^{(3)}$ are arbitrary matrices to be constrained by the symmetries of the theory. These candidate counterterms need to be considered together with their conjugates and in combination with their supersymmetric completions. Given the choice of polarization in our calculation, we are able to probe only counterterms of the kind (5.2). The matrix $M_{ab,cd}^{(1)}$ needs to obey manifestly the linearly realized $\text{SO}(D - 4)$ symmetry which is present in our amplitudes. Additionally, it needs to be symmetric under the exchange of the indices a, b and under exchange of the indices c, d . It is convenient to find an expression in terms of Clifford algebra elements. Putting this all together, the observed divergences correspond to the appearance of the linearized counterterms

$$\mathcal{O}_1 = \tilde{C}_{ac}\tilde{C}_{bd}(F_{\alpha\beta}^a F^{b\alpha\beta})(F_{\dot{\alpha}\dot{\beta}}^c F^{d\dot{\alpha}\dot{\beta}}), \quad \mathcal{O}_2 = ((\tilde{C}\tilde{\Gamma}^I)_{ab}F_{\alpha\beta}^a F^{b\alpha\beta})((\tilde{C}\tilde{\Gamma}^I)_{cd}F_{\dot{\alpha}\dot{\beta}}^c F^{d\dot{\alpha}\dot{\beta}}), \tag{5.5}$$

together with their supersymmetric completions. The second counterterm does not appear in the case of the Magical supergravities, which signals symmetry enhancement corresponding to the enlarged U-duality groups of these theories. Note that, for D sufficiently large, additional $\text{SO}(D - 4)$ -invariant candidate counterterms built out of higher-rank elements of the Clifford algebra become possible which are associated to zero divergence from our calculations. A systematic study of the effect of non-linearly realized symmetries on candidate counterterms for the theories we are considering is left to future work.

Finally, it would be interesting to see if the relation between supergravity divergences and physical quantities of the non-supersymmetric theory entering the construction (i.e. beta functions) can carry over to other matter amplitudes or higher loops. When supergravity vectors constructed as vector times scalar are taken into account, the numerators for the supersymmetric gauge theory contain explicit dependence on the loop momentum,

which makes it difficult to observe a relation between integrated quantities. As for extending the computation to higher loops, amplitudes which manifestly satisfy color/kinematics duality at two loops have recently become available [85] and are likely to trigger further progress.

Acknowledgments

M.C. would like to thank Murat Günaydin, Radu Roiban and Henrik Johansson for many insightful conversations and collaboration on related topics. We thank Henrik Johansson and Murat Günaydin for valuable comments on an earlier draft of this paper. We also thank Gustav Mogull, Gregor Kälin and Alex Ochirov for sharing a draft of their paper. The research of M.C. is supported in part by the Knut and Alice Wallenberg Foundation under grant KAW 2013.0235 and the Swedish Research Council under grant 621-2014-5722.

A Conventions

In this appendix we collect the conventions employed throughout this paper. Our notation can be obtained from the one of Elvang and Huang [86] by replacing $\eta_{\mu\nu} \rightarrow -\eta_{\mu\nu}$. Our metric has mostly-minus signature and the Clifford algebra relation is

$$\{\Gamma^\mu, \Gamma^\nu\} = 2\eta^{\mu\nu}. \tag{A.1}$$

Γ^0 is hermitian while the other gamma matrices are antihermitian. The four-dimensional gamma matrices γ^μ are

$$\begin{aligned} \gamma^0 &= \sigma^1 \otimes 1, \\ \gamma^1 &= i\sigma^2 \otimes \sigma^1, \\ \gamma^2 &= i\sigma^2 \otimes \sigma^2, \\ \gamma^3 &= i\sigma^2 \otimes \sigma^3, \\ \gamma_5 &= \sigma^3 \otimes 1. \end{aligned} \tag{A.2}$$

Four-dimensional charge-conjugation and B matrix are taken to be

$$\mathcal{C}_4 = \begin{pmatrix} \epsilon^{\alpha\beta} & 0 \\ 0 & t_1 \epsilon_{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \quad B = \begin{pmatrix} 0 & t_1 \epsilon_{\dot{\alpha}\dot{\beta}} \\ \epsilon^{\alpha\beta} & 0 \end{pmatrix}, \tag{A.3}$$

with $\epsilon^{12} = -\epsilon_{12} = +1$. t_1 is a sign to be assigned according to the value of the parameter D in the construction. The charge conjugation matrix obeys the conditions $\mathcal{C}_4^t = -\mathcal{C}_4$ and $(\gamma^\mu)^t = -t_1 \mathcal{C}_4^{-1} \gamma^\mu \mathcal{C}_4$.

Higher dimensional gamma matrices are written as

$$\begin{aligned} \Gamma^\mu &= \gamma^\mu \otimes \mathbf{1}, & \mu < 4, \\ \Gamma^\mu &= \gamma_5 \otimes \tilde{\Gamma}^\mu, & \mu \geq 4. \end{aligned} \tag{A.4}$$

We introduce indices I, J running over the internal dimensions. \tilde{C} and $\tilde{\Gamma}^I$ denote the components of the charge-conjugation matrix acting on the spinor indices corresponding to the internal $(D - d)$ dimensions. The sign t_1 is fixed by the requirement that $\tilde{C}\tilde{\Gamma}^I$ is always symmetric or, alternatively, that $C\Gamma^\mu$ is always antisymmetric.

Introducing the spinor-helicity variables as

$$\lambda(p) = \begin{pmatrix} |p]_\alpha \\ |p\rangle^{\dot{\alpha}} \end{pmatrix}, \tag{A.5}$$

the Majorana condition is rewritten as

$$\lambda^* = B\lambda \quad \rightarrow \quad \begin{cases} (|p]_\alpha)^* = \epsilon_{\dot{\alpha}\beta} |p\rangle^{\dot{\beta}} = |p\rangle_{\dot{\alpha}} \\ (|p\rangle^{\dot{\alpha}})^* = \epsilon^{\alpha\beta} |p]_\beta = |p]^\alpha \end{cases}. \tag{A.6}$$

With this condition we have

$$([pq])^* = \langle qp \rangle \tag{A.7}$$

for real momenta. We can expand null momenta using spinor-helicity variables as

$$\not{p} = -|p\rangle[p] - |p]\langle p|. \tag{A.8}$$


We also have the identities


$$\langle pq \rangle [qp] = 2p \cdot q, \tag{A.9}$$


$$\langle p|\gamma^\mu|q\rangle \langle r|\gamma_\mu|s\rangle = 2\langle pr \rangle [sq]. \tag{A.10}$$


B Feynman rules


In this appendix, we collect the Feynman rules for the non-supersymmetric gauge theory entering the double-copy construction, which are obtained from the Lagrangian (2.5). All momenta are taken as in-going.

Fermion propagator:  = $\frac{ip_\mu \Gamma^\mu}{p^2} C^{-1} V^{-1}$. (B.1)


Gluon propagator:  = $\frac{-i\eta_{\mu\nu}}{p^2}$. (B.2)

Ghost propagator:  = $\frac{-i\delta^{ab}}{p^2}$. (B.3)

Fermion vertex:  = $igVt^a C\Gamma^\mu$. (B.4)

Gluon vertex:  $= g f^{abc} (\eta^{\mu\nu} (k-p)^\rho + \eta^{\nu\rho} (p-q)^\mu + \eta^{\rho\mu} (q-k)^\nu) .$

(B.5)

Ghost vertex:  $= g f^{abc} p^\mu .$

(B.6)

C Details on the orbifold numerators

To obtain an explicit presentation of the one-loop (super)amplitude with four external hypermultiplets in a complex representation, it is convenient to start from the amplitude specified in eq. (5.46) of ref. [66] and perform the summation over the orbifold group elements. We consider a \mathbb{Z}_3 orbifold taking $SU(3N)$ as the gauge group for the parent theory. The orbifold action is given by

$$r_n = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & e^{in\theta} & 0 \\ 0 & 0 & 0 & e^{-in\theta} \end{pmatrix}, \quad g_n = \begin{pmatrix} I_N & 0 & 0 \\ 0 & e^{in\theta} I_N & 0 \\ 0 & 0 & e^{2in\theta} I_N \end{pmatrix}, \quad \theta = \frac{2\pi}{3}. \quad (\text{C.1})$$

This choice breaks the gauge group as $SU(3N) \rightarrow SU(N)^3 \times U(1)^2$. We then split accordingly the $SU(3N)$ adjoint indices as $\hat{A} = (\hat{a}, \hat{\alpha}, \hat{\bar{\alpha}})$, where the indices run over the following representations of the (product) gauge group,

$$\begin{aligned} \hat{a} : & \quad (\mathbf{N}^2 - 1, 1, 1) \oplus (1, \mathbf{N}^2 - 1, 1) \oplus (1, 1, \mathbf{N}^2 - 1) \oplus 2(1, 1, 1), \\ \hat{\alpha} : & \quad (\mathbf{N}, \bar{\mathbf{N}}, 1) \oplus (1, \mathbf{N}, \bar{\mathbf{N}}) \oplus (\bar{\mathbf{N}}, 1, \mathbf{N}), \\ \hat{\bar{\alpha}} : & \quad (\bar{\mathbf{N}}, \mathbf{N}, 1) \oplus (1, \bar{\mathbf{N}}, \mathbf{N}) \oplus (\mathbf{N}, 1, \bar{\mathbf{N}}). \end{aligned} \quad (\text{C.2})$$

Projectors into the three sets of representations are written as

$$\begin{aligned} (\mathcal{P}_G \Phi)^{\hat{A}} &= \sum_{\Gamma} g^{\hat{A}\hat{B}} \Phi^{\hat{B}} = (\Phi^{\hat{a}}, 0, 0), \\ (\mathcal{P}_R \Phi)^{\hat{A}} &= \sum_{\Gamma} r_3^3 g^{\hat{A}\hat{B}} \Phi^{\hat{B}} = (0, \Phi_{\hat{\alpha}}, 0), \\ (\mathcal{P}_{\bar{R}} \Phi)^{\hat{A}} &= \sum_{\Gamma} r_4^4 g^{\hat{A}\hat{B}} \Phi^{\hat{B}} = (0, 0, \Phi^{\hat{\alpha}}), \end{aligned} \quad (\text{C.3})$$

where Φ is a generic field of the parent theory. Representation matrices of the R representation are then given by

$$\tilde{T}_{\hat{\alpha}}^{\hat{a}}{}^{\hat{\beta}} = -(\mathcal{P}_R)_{\hat{\alpha}}^{\hat{A}} (\mathcal{P}_{\bar{R}})^{\hat{B}\hat{B}} (\mathcal{P}_G)^{\hat{a}\hat{C}} \tilde{f}^{\hat{A}\hat{B}\hat{C}}. \quad (\text{C.4})$$




Figure 3. Orientation for external and loop momenta used in the integral reduction identities.

At four points, supersymmetry implies that amplitudes with four external hypermultiplet fields can be organized in superamplitudes which can be directly obtained from the amplitudes between two identical scalars and their conjugates given in ref. [66],

$$\mathcal{F}_4^{1\text{-loop}}(1\mathcal{Q}, 2\mathcal{Q}, 3\bar{\mathcal{Q}}, 4\bar{\mathcal{Q}}) = \frac{\delta^4(\sum_i \eta_i^\alpha |i\rangle)}{\langle 12 \rangle \langle 34 \rangle} \mathcal{A}_4^{1\text{-loop}}(1\varphi, 2\varphi, 3\bar{\varphi}, 4\bar{\varphi}). \quad (\text{C.5})$$

Component amplitudes can be easily extracted by acting with derivatives with respect to the Grassmann variables η .

It should be noted that the amplitudes in [66] were obtained with a procedure that is not sensitive to bubble-on-external-leg graphs. In principle, it is possible to add back these graphs in a way that preserves color/kinematics duality by adding to all numerators terms proportional to the squares of the external momenta p_i^2 . When the external momenta are put on-shell, i.e. the limit $p_i^2 \rightarrow 0$ is taken, the additional contributions drop out of the final expression in all graphs except the ones with bubbles on one external leg, which have a $1/p_i^2$ factor in the propagators. However, these graphs can be safely ignored in the present context as they do not contribute to the gravity amplitudes because each of the two numerators entering the double-copy formula is proportional to p_i^2 . At the level of gauge-theory amplitudes, bubbles-on-external-legs integrals vanish in dimensional regularization. However, they can lead to non-vanishing contributions if particular kinematical limits (UV or infrared) are inspected.

D Integral reduction

Orientation of loop and external momenta are taken as shown in figure 3. The integral-reduction identities for bubble diagrams are:

$$I_2(l^\mu; p) = -\frac{1}{2} I_2(p) p^\mu, \quad (\text{D.1})$$

$$I_2(l^\mu l^\nu; p) = \frac{d}{4d-4} I_2(p) p^\mu p^\nu - \frac{p^2}{4d-4} I_2(p) \eta_d^{\mu\nu}. \quad (\text{D.2})$$

For triangle diagrams, integral reduction identities are as follows:

$$I_3(l^\mu; p_1, p_2) = \frac{p_1^\mu (I_2(p_1 + p_2) - I_2(p_1))}{2p_1 \cdot p_2} + \frac{p_2^\mu (I_2(p_1, p_2) - I_2(p_1 + p_2))}{2p_1 \cdot p_2}, \quad (\text{D.3})$$

$$\begin{aligned}
 I_3(l^\mu l^\nu; p_1, p_2) &= \frac{\eta_d^{\mu\nu} I_2(p_1 + p_2)}{2(d-2)} + p_1^\mu p_1^\nu \frac{I_2(p_1) - I_2(p_1 + p_2)}{4p_1 \cdot p_2} + \\
 &\quad p_2^\mu p_2^\nu \frac{I_2(p_2) - I_2(p_1 + p_2)}{4p_1 \cdot p_2} + \\
 &\quad p_1^{(\mu} p_2^{\nu)} \frac{(d-4)I_2(p_1 + p_2)}{4(d-2)p_1 \cdot p_2}
 \end{aligned} \tag{D.4}$$

Note that the above identities depend on the d -dimensional metric $\eta_d^{\mu\nu}$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Z. Bern, J.J.M. Carrasco and H. Johansson, *New Relations for Gauge-Theory Amplitudes*, *Phys. Rev. D* **78** (2008) 085011 [[arXiv:0805.3993](https://arxiv.org/abs/0805.3993)] [[INSPIRE](#)].
- [2] Z. Bern, J.J.M. Carrasco and H. Johansson, *Perturbative Quantum Gravity as a Double Copy of Gauge Theory*, *Phys. Rev. Lett.* **105** (2010) 061602 [[arXiv:1004.0476](https://arxiv.org/abs/1004.0476)] [[INSPIRE](#)].
- [3] Z. Bern, J.J. Carrasco, W.-M. Chen, H. Johansson and R. Roiban, *Gravity Amplitudes as Generalized Double Copies of Gauge-Theory Amplitudes*, *Phys. Rev. Lett.* **118** (2017) 181602 [[arXiv:1701.02519](https://arxiv.org/abs/1701.02519)] [[INSPIRE](#)].
- [4] Z. Bern, J.J.M. Carrasco, W.-M. Chen, H. Johansson, R. Roiban and M. Zeng, *Five-loop four-point integrand of $N = 8$ supergravity as a generalized double copy*, *Phys. Rev. D* **96** (2017) 126012 [[arXiv:1708.06807](https://arxiv.org/abs/1708.06807)] [[INSPIRE](#)].
- [5] Z. Bern et al., *Ultraviolet Properties of $\mathcal{N} = 8$ Supergravity at Five Loops*, *Phys. Rev. D* **98** (2018) 086021 [[arXiv:1804.09311](https://arxiv.org/abs/1804.09311)] [[INSPIRE](#)].
- [6] Z. Bern, S. Davies, T. Dennen and Y.-t. Huang, *Absence of Three-Loop Four-Point Divergences in $N = 4$ Supergravity*, *Phys. Rev. Lett.* **108** (2012) 201301 [[arXiv:1202.3423](https://arxiv.org/abs/1202.3423)] [[INSPIRE](#)].
- [7] Z. Bern, S. Davies, T. Dennen and Y.-t. Huang, *Ultraviolet Cancellations in Half-Maximal Supergravity as a Consequence of the Double-Copy Structure*, *Phys. Rev. D* **86** (2012) 105014 [[arXiv:1209.2472](https://arxiv.org/abs/1209.2472)] [[INSPIRE](#)].
- [8] Z. Bern, S. Davies, T. Dennen, A.V. Smirnov and V.A. Smirnov, *Ultraviolet Properties of $N = 4$ Supergravity at Four Loops*, *Phys. Rev. Lett.* **111** (2013) 231302 [[arXiv:1309.2498](https://arxiv.org/abs/1309.2498)] [[INSPIRE](#)].
- [9] Z. Bern, S. Davies and T. Dennen, *Enhanced ultraviolet cancellations in $\mathcal{N} = 5$ supergravity at four loops*, *Phys. Rev. D* **90** (2014) 105011 [[arXiv:1409.3089](https://arxiv.org/abs/1409.3089)] [[INSPIRE](#)].
- [10] Z. Bern, M. Enciso, J. Parra-Martinez and M. Zeng, *Manifesting enhanced cancellations in supergravity: integrands versus integrals*, *JHEP* **05** (2017) 137 [[arXiv:1703.08927](https://arxiv.org/abs/1703.08927)] [[INSPIRE](#)].
- [11] J.J.M. Carrasco, R. Kallosh, R. Roiban and A.A. Tseytlin, *On the $U(1)$ duality anomaly and the S -matrix of $N = 4$ supergravity*, *JHEP* **07** (2013) 029 [[arXiv:1303.6219](https://arxiv.org/abs/1303.6219)] [[INSPIRE](#)].

- [12] Z. Bern, J. Parra-Martinez and R. Roiban, *Canceling the U(1) Anomaly in the S Matrix of $N = 4$ Supergravity*, *Phys. Rev. Lett.* **121** (2018) 101604 [[arXiv:1712.03928](#)] [[INSPIRE](#)].
- [13] Z. Bern, C. Boucher-Veronneau and H. Johansson, *$N \geq 4$ Supergravity Amplitudes from Gauge Theory at One Loop*, *Phys. Rev. D* **84** (2011) 105035 [[arXiv:1107.1935](#)] [[INSPIRE](#)].
- [14] H. Johansson and A. Ochirov, *Pure Gravities via Color-Kinematics Duality for Fundamental Matter*, *JHEP* **11** (2015) 046 [[arXiv:1407.4772](#)] [[INSPIRE](#)].
- [15] J.J.M. Carrasco, M. Chiodaroli, M. Günaydin and R. Roiban, *One-loop four-point amplitudes in pure and matter-coupled $\mathcal{N} \leq 4$ supergravity*, *JHEP* **03** (2013) 056 [[arXiv:1212.1146](#)] [[INSPIRE](#)].
- [16] Z. Bern, S. Davies, T. Dennen, Y.-t. Huang and J. Nohle, *Color-Kinematics Duality for Pure Yang-Mills and Gravity at One and Two Loops*, *Phys. Rev. D* **92** (2015) 045041 [[arXiv:1303.6605](#)] [[INSPIRE](#)].
- [17] M. Chiodaroli, M. Günaydin, H. Johansson and R. Roiban, *Complete construction of magical, symmetric and homogeneous $\mathcal{N} = 2$ supergravities as double copies of gauge theories*, *Phys. Rev. Lett.* **117** (2016) 011603 [[arXiv:1512.09130](#)] [[INSPIRE](#)].
- [18] A. Anastasiou et al., *Twin supergravities from Yang-Mills theory squared*, *Phys. Rev. D* **96** (2017) 026013 [[arXiv:1610.07192](#)] [[INSPIRE](#)].
- [19] A. Anastasiou, L. Borsten, M.J. Duff, A. Marrani, S. Nagy and M. Zoccali, *Are all supergravity theories Yang-Mills squared?*, *Nucl. Phys. B* **934** (2018) 606 [[arXiv:1707.03234](#)] [[INSPIRE](#)].
- [20] H. Johansson, G. Kälin and G. Mogull, *Two-loop supersymmetric QCD and half-maximal supergravity amplitudes*, *JHEP* **09** (2017) 019 [[arXiv:1706.09381](#)] [[INSPIRE](#)].
- [21] M. Chiodaroli, M. Günaydin, H. Johansson and R. Roiban, *Scattering amplitudes in $\mathcal{N} = 2$ Maxwell-Einstein and Yang-Mills/Einstein supergravity*, *JHEP* **01** (2015) 081 [[arXiv:1408.0764](#)] [[INSPIRE](#)].
- [22] M. Chiodaroli, *Simplifying amplitudes in Maxwell-Einstein and Yang-Mills-Einstein supergravities*, in *Space — Time — Matter: Analytic and Geometric Structures*, J. Brüning and M. Staudacher eds., *De Gruyter* (2018), pp. 266–287 [[arXiv:1607.04129](#)] [[INSPIRE](#)].
- [23] M. Chiodaroli, M. Günaydin, H. Johansson and R. Roiban, *Spontaneously Broken Yang-Mills-Einstein Supergravities as Double Copies*, *JHEP* **06** (2017) 064 [[arXiv:1511.01740](#)] [[INSPIRE](#)].
- [24] M. Chiodaroli, M. Günaydin, H. Johansson and R. Roiban, *Gauged Supergravities and Spontaneous Supersymmetry Breaking from the Double Copy Construction*, *Phys. Rev. Lett.* **120** (2018) 171601 [[arXiv:1710.08796](#)] [[INSPIRE](#)].
- [25] H. Johansson and J. Nohle, *Conformal Gravity from Gauge Theory*, [arXiv:1707.02965](#) [[INSPIRE](#)].
- [26] H. Johansson, G. Mogull and F. Teng, *Unraveling conformal gravity amplitudes*, *JHEP* **09** (2018) 080 [[arXiv:1806.05124](#)] [[INSPIRE](#)].
- [27] H. Johansson and A. Ochirov, *Color-Kinematics Duality for QCD Amplitudes*, *JHEP* **01** (2016) 170 [[arXiv:1507.00332](#)] [[INSPIRE](#)].
- [28] L. de la Cruz, A. Kniss and S. Weinzierl, *Proof of the fundamental BCJ relations for QCD amplitudes*, *JHEP* **09** (2015) 197 [[arXiv:1508.01432](#)] [[INSPIRE](#)].

- [29] G. Chen and Y.-J. Du, *Amplitude Relations in Non-linear σ -model*, *JHEP* **01** (2014) 061 [[arXiv:1311.1133](#)] [[INSPIRE](#)].
- [30] G. Chen, Y.-J. Du, S. Li and H. Liu, *Note on off-shell relations in nonlinear σ -model*, *JHEP* **03** (2015) 156 [[arXiv:1412.3722](#)] [[INSPIRE](#)].
- [31] C. Cheung and C.-H. Shen, *Symmetry for Flavor-Kinematics Duality from an Action*, *Phys. Rev. Lett.* **118** (2017) 121601 [[arXiv:1612.00868](#)] [[INSPIRE](#)].
- [32] Y.-J. Du and C.-H. Fu, *Explicit BCJ numerators of nonlinear sigma model*, *JHEP* **09** (2016) 174 [[arXiv:1606.05846](#)] [[INSPIRE](#)].
- [33] G. Chen, S. Li and H. Liu, *Off-shell BCJ Relation in Nonlinear σ -model*, [arXiv:1609.01832](#) [[INSPIRE](#)].
- [34] M. Chiodaroli, M. Günaydin, H. Johansson and R. Roiban, *Explicit Formulae for Yang-Mills-Einstein Amplitudes from the Double Copy*, *JHEP* **07** (2017) 002 [[arXiv:1703.00421](#)] [[INSPIRE](#)].
- [35] S. Stieberger and T.R. Taylor, *New relations for Einstein-Yang-Mills amplitudes*, *Nucl. Phys. B* **913** (2016) 151 [[arXiv:1606.09616](#)] [[INSPIRE](#)].
- [36] S. Stieberger and T.R. Taylor, *Graviton as a Pair of Collinear Gauge Bosons*, *Phys. Lett. B* **739** (2014) 457 [[arXiv:1409.4771](#)] [[INSPIRE](#)].
- [37] S. Stieberger and T.R. Taylor, *Graviton Amplitudes from Collinear Limits of Gauge Amplitudes*, *Phys. Lett. B* **744** (2015) 160 [[arXiv:1502.00655](#)] [[INSPIRE](#)].
- [38] D. Nandan, J. Plefka, O. Schlotterer and C. Wen, *Einstein-Yang-Mills from pure Yang-Mills amplitudes*, *JHEP* **10** (2016) 070 [[arXiv:1607.05701](#)] [[INSPIRE](#)].
- [39] L. de la Cruz, A. Kniss and S. Weinzierl, *Relations for Einstein-Yang-Mills amplitudes from the CHY representation*, *Phys. Lett. B* **767** (2017) 86 [[arXiv:1607.06036](#)] [[INSPIRE](#)].
- [40] F. Teng and B. Feng, *Expanding Einstein-Yang-Mills by Yang-Mills in CHY frame*, *JHEP* **05** (2017) 075 [[arXiv:1703.01269](#)] [[INSPIRE](#)].
- [41] Y.-J. Du, B. Feng and F. Teng, *Expansion of All Multitrace Tree Level EYM Amplitudes*, *JHEP* **12** (2017) 038 [[arXiv:1708.04514](#)] [[INSPIRE](#)].
- [42] D. Nandan, J. Plefka and G. Travaglini, *All rational one-loop Einstein-Yang-Mills amplitudes at four points*, *JHEP* **09** (2018) 011 [[arXiv:1803.08497](#)] [[INSPIRE](#)].
- [43] S. Stieberger, *Open & Closed vs. Pure Open String Disk Amplitudes*, [arXiv:0907.2211](#) [[INSPIRE](#)].
- [44] N.E.J. Bjerrum-Bohr, P.H. Damgaard and P. Vanhove, *Minimal Basis for Gauge Theory Amplitudes*, *Phys. Rev. Lett.* **103** (2009) 161602 [[arXiv:0907.1425](#)] [[INSPIRE](#)].
- [45] N.E.J. Bjerrum-Bohr, P.H. Damgaard, T. Sondergaard and P. Vanhove, *Monodromy and Jacobi-like Relations for Color-Ordered Amplitudes*, *JHEP* **06** (2010) 003 [[arXiv:1003.2403](#)] [[INSPIRE](#)].
- [46] P. Tourkine and P. Vanhove, *Higher-loop amplitude monodromy relations in string and gauge theory*, *Phys. Rev. Lett.* **117** (2016) 211601 [[arXiv:1608.01665](#)] [[INSPIRE](#)].
- [47] S. Hohenegger and S. Stieberger, *Monodromy Relations in Higher-Loop String Amplitudes*, *Nucl. Phys. B* **925** (2017) 63 [[arXiv:1702.04963](#)] [[INSPIRE](#)].

- [48] C.-H. Fu, P. Vanhove and Y. Wang, *A Vertex Operator Algebra Construction of the Colour-Kinematics Dual numerator*, *JHEP* **09** (2018) 141 [[arXiv:1806.09584](#)] [[INSPIRE](#)].
- [49] C.R. Mafra, O. Schlotterer and S. Stieberger, *Explicit BCJ Numerators from Pure Spinors*, *JHEP* **07** (2011) 092 [[arXiv:1104.5224](#)] [[INSPIRE](#)].
- [50] C.R. Mafra and O. Schlotterer, *Towards one-loop SYM amplitudes from the pure spinor BRST cohomology*, *Fortsch. Phys.* **63** (2015) 105 [[arXiv:1410.0668](#)] [[INSPIRE](#)].
- [51] S. He, R. Monteiro and O. Schlotterer, *String-inspired BCJ numerators for one-loop MHV amplitudes*, *JHEP* **01** (2016) 171 [[arXiv:1507.06288](#)] [[INSPIRE](#)].
- [52] C.R. Mafra and O. Schlotterer, *Two-loop five-point amplitudes of super Yang-Mills and supergravity in pure spinor superspace*, *JHEP* **10** (2015) 124 [[arXiv:1505.02746](#)] [[INSPIRE](#)].
- [53] S. Stieberger, *Closed superstring amplitudes, single-valued multiple zeta values and the Deligne associator*, *J. Phys. A* **47** (2014) 155401 [[arXiv:1310.3259](#)] [[INSPIRE](#)].
- [54] C.R. Mafra and O. Schlotterer, *Double-Copy Structure of One-Loop Open-String Amplitudes*, *Phys. Rev. Lett.* **121** (2018) 011601 [[arXiv:1711.09104](#)] [[INSPIRE](#)].
- [55] Y.-t. Huang, O. Schlotterer and C. Wen, *Universality in string interactions*, *JHEP* **09** (2016) 155 [[arXiv:1602.01674](#)] [[INSPIRE](#)].
- [56] T. Azevedo, M. Chiodaroli, H. Johansson and O. Schlotterer, *Heterotic and bosonic string amplitudes via field theory*, *JHEP* **10** (2018) 012 [[arXiv:1803.05452](#)] [[INSPIRE](#)].
- [57] F. Cachazo, S. He and E.Y. Yuan, *Scattering of Massless Particles in Arbitrary Dimensions*, *Phys. Rev. Lett.* **113** (2014) 171601 [[arXiv:1307.2199](#)] [[INSPIRE](#)].
- [58] F. Cachazo, S. He and E.Y. Yuan, *Scattering of Massless Particles: Scalars, Gluons and Gravitons*, *JHEP* **07** (2014) 033 [[arXiv:1309.0885](#)] [[INSPIRE](#)].
- [59] F. Cachazo, S. He and E.Y. Yuan, *Scattering equations and Kawai-Lewellen-Tye orthogonality*, *Phys. Rev. D* **90** (2014) 065001 [[arXiv:1306.6575](#)] [[INSPIRE](#)].
- [60] F. Cachazo, S. He and E.Y. Yuan, *Einstein-Yang-Mills Scattering Amplitudes From Scattering Equations*, *JHEP* **01** (2015) 121 [[arXiv:1409.8256](#)] [[INSPIRE](#)].
- [61] F. Cachazo, S. He and E.Y. Yuan, *Scattering Equations and Matrices: From Einstein To Yang-Mills, DBI and NLSM*, *JHEP* **07** (2015) 149 [[arXiv:1412.3479](#)] [[INSPIRE](#)].
- [62] N.E.J. Bjerrum-Bohr, J.L. Bourjaily, P.H. Damgaard and B. Feng, *Manifesting Color-Kinematics Duality in the Scattering Equation Formalism*, *JHEP* **09** (2016) 094 [[arXiv:1608.00006](#)] [[INSPIRE](#)].
- [63] L. Mason and D. Skinner, *Ambitwistor strings and the scattering equations*, *JHEP* **07** (2014) 048 [[arXiv:1311.2564](#)] [[INSPIRE](#)].
- [64] E. Casali, Y. Geyer, L. Mason, R. Monteiro and K.A. Roehrig, *New Ambitwistor String Theories*, *JHEP* **11** (2015) 038 [[arXiv:1506.08771](#)] [[INSPIRE](#)].
- [65] B. de Wit and A. Van Proeyen, *Special geometry, cubic polynomials and homogeneous quaternionic spaces*, *Commun. Math. Phys.* **149** (1992) 307 [[hep-th/9112027](#)] [[INSPIRE](#)].
- [66] M. Chiodaroli, Q. Jin and R. Roiban, *Color/kinematics duality for general abelian orbifolds of $N = 4$ super Yang-Mills theory*, *JHEP* **01** (2014) 152 [[arXiv:1311.3600](#)] [[INSPIRE](#)].
- [67] M. Günaydin, G. Sierra and P.K. Townsend, *The Geometry of $N = 2$ Maxwell-Einstein Supergravity and Jordan Algebras*, *Nucl. Phys. B* **242** (1984) 244 [[INSPIRE](#)].

- [68] M. Günaydin, G. Sierra and P.K. Townsend, *Gauging the $d = 5$ Maxwell-Einstein Supergravity Theories: More on Jordan Algebras*, *Nucl. Phys. B* **253** (1985) 573 [INSPIRE].
- [69] M. Günaydin, G. Sierra and P.K. Townsend, *Exceptional Supergravity Theories and the MAGIC Square*, *Phys. Lett. B* **133** (1983) 72 [INSPIRE].
- [70] M. Bershadsky and A. Johansen, *Large N limit of orbifold field theories*, *Nucl. Phys. B* **536** (1998) 141 [hep-th/9803249] [INSPIRE].
- [71] J.F. Luciani, *Coupling of $O(2)$ Supergravity with Several Vector Multiplets*, *Nucl. Phys. B* **132** (1978) 325 [INSPIRE].
- [72] H. Elvang, D.Z. Freedman and M. Kiermaier, *A simple approach to counterterms in $N = 8$ supergravity*, *JHEP* **11** (2010) 016 [arXiv:1003.5018] [INSPIRE].
- [73] G. Bossard, P.S. Howe and K.S. Stelle, *On duality symmetries of supergravity invariants*, *JHEP* **01** (2011) 020 [arXiv:1009.0743] [INSPIRE].
- [74] G. Bossard, P.S. Howe, K.S. Stelle and P. Vanhove, *The vanishing volume of $D = 4$ superspace*, *Class. Quant. Grav.* **28** (2011) 215005 [arXiv:1105.6087] [INSPIRE].
- [75] N. Beisert, H. Elvang, D.Z. Freedman, M. Kiermaier, A. Morales and S. Stieberger, *$E_{7(7)}$ constraints on counterterms in $\mathcal{N} = 8$ supergravity*, *Phys. Lett. B* **694** (2011) 265 [arXiv:1009.1643] [INSPIRE].
- [76] H. Elvang, D.Z. Freedman and M. Kiermaier, *SUSY Ward identities, Superamplitudes and Counterterms*, *J. Phys. A* **44** (2011) 454009 [arXiv:1012.3401] [INSPIRE].
- [77] G. Bossard, P.S. Howe and K.S. Stelle, *Anomalies and divergences in $N = 4$ supergravity*, *Phys. Lett. B* **719** (2013) 424 [arXiv:1212.0841] [INSPIRE].
- [78] D.Z. Freedman and E. Tonni, *The $D^{2k} R^4$ Invariants of $\mathcal{N} = 8$ Supergravity*, *JHEP* **04** (2011) 006 [arXiv:1101.1672] [INSPIRE].
- [79] G. Bossard, P.S. Howe and K.S. Stelle, *Invariants and divergences in half-maximal supergravity theories*, *JHEP* **07** (2013) 117 [arXiv:1304.7753] [INSPIRE].
- [80] J. Broedel and L.J. Dixon, *\mathcal{R}^4 counterterm and $E_{7(7)}$ symmetry in maximal supergravity*, *JHEP* **05** (2010) 003 [arXiv:0911.5704] [INSPIRE].
- [81] R. Kallosh, *$E_{7(7)}$ Symmetry and Finiteness of $N = 8$ Supergravity*, *JHEP* **03** (2012) 083 [arXiv:1103.4115] [INSPIRE].
- [82] R. Kallosh, *$N = 8$ Counterterms and $E_{7(7)}$ Current Conservation*, *JHEP* **06** (2011) 073 [arXiv:1104.5480] [INSPIRE].
- [83] R. Kallosh, *On Absence of 3-loop Divergence in $N = 4$ Supergravity*, *Phys. Rev. D* **85** (2012) 081702 [arXiv:1202.4690] [INSPIRE].
- [84] R. Kallosh and T. Ortín, *New $E_{7(7)}$ invariants and amplitudes*, *JHEP* **09** (2012) 137 [arXiv:1205.4437] [INSPIRE].
- [85] G. Kälin, G. Mogull and A. Ochirov, *Two-loop $\mathcal{N} = 2$ SQCD amplitudes with external matter from iterated cuts*, arXiv:1811.09604 [INSPIRE].
- [86] H. Elvang and Y.-t. Huang, *Scattering Amplitudes in Gauge Theory and Gravity*, Cambridge University Press, Cambridge U.K. (2015).