

Scaling black holes and modularity

Aradhita Chattopadhyaya, Jan Manschot and Swapnamay Mondal

*School of Mathematics, Trinity College,
Dublin 2, Ireland*

*Hamilton Mathematical Institute, Trinity College,
Dublin 2, Ireland*

E-mail: aradhita@maths.tcd.ie, manschot@maths.tcd.ie,
swapno@maths.tcd.ie

ABSTRACT: Scaling black holes are solutions of supergravity with multiple black hole singularities, which can be adiabatically connected to a single center black hole solution. We develop techniques to determine partition functions for such scaling black holes, if each constituent carries a non-vanishing magnetic charge corresponding to a D4-brane in string theory, or equivalently M5-brane in M-theory. For three constituents, we demonstrate that the partition function is a mock modular form of depth two, and we determine the appropriate non-holomorphic completion using generalized error functions. From the four-dimensional perspective, the modular parameter is the axion-dilaton, and our results show that S -duality leaves this subset of the spectrum invariant. From the five-dimensional perspective, the modular parameter is the complex structure of a torus T^2 , and the scaling black holes are dual to states in the dimensional reduction of the M5-brane worldvolume theory to T^2 . As a case study, we specialize the compactification manifold to a K3 fibration, and explicitly evaluate holomorphic parts of partition functions.

KEYWORDS: Black Holes in String Theory, AdS-CFT Correspondence, D-Branes, M-Theory

ARXIV EPRINT: [2110.05504](https://arxiv.org/abs/2110.05504)

Contents

1	Introduction	1
2	Black hole solutions in $\mathcal{N} = 2$ supergravity	5
2.1	Black hole bound states	5
2.2	Bound state indices	8
3	Review of D4-brane black holes	13
3.1	Supergravity partition function	13
3.2	Attractor partition function	16
4	Partition functions for scaling black holes	17
4.1	Bound states and lattices	19
4.2	Partition functions	26
4.3	Convergence	28
4.4	Modular completion of Φ_μ	29
4.5	Modular completion of Ψ_μ	35
5	Relation to M5-branes and AdS₃/CFT₂	37
6	Case studies	39
6.1	Charge configuration 1: $P_1 = P_2 = (0, 1)$, $P_3 = (1, 1)$	40
6.2	Charge configuration 2: $P_1 = (1, 0)$, $P_2 = P_3 = (0, 1)$	45

1 Introduction

Solutions of supergravity with multiple black hole singularities provide interesting insights on the spectrum of quantum gravity and the dependence on the compactification moduli [1–10]. Such multi-center solutions are particularly intriguing if their solution space contains a region, where the distances between the centers becomes arbitrarily small. In other words, these solutions, known as “scaling” solutions [6, 11] in four dimensions, are adiabatically connected to a supergravity solution with a single black hole singularity.

We will study such scaling black holes formed from M5-branes in M-theory, or equivalently D4-branes in Type IIA string theory [12]. Such black holes are well studied using the AdS₃/CFT₂ correspondence [12–15], as well as the hypermultiplet geometry of the IIB string theory [16–18]. The D4/M5-branes wrap a four-cycle (or divisor) $P \in H_4(X, \mathbb{Z})$ of the Calabi-Yau threefold X , which can also be studied within algebraic geometry [19–23].

Multi-center solutions exist with each center having a positive D4-brane charge but vanishing D6-brane charge. Their walls of marginal stability extend to the large volume

regime of the Kähler moduli space [24–27], which corresponds to $\text{Im}(t) = J = \lambda \underline{J}$, where \underline{J} is the normalized, dimensionless Kähler modulus, $\underline{J}^3 = 1$, and $\lambda \gg 1$. While attractor points lie typically in the interior of the Kähler moduli space, there is an analogue of the attractor point at large volume t_γ^λ defined in eq. (3.14) [18, 25, 27]. Similarly, single center black holes, whose internal degrees of freedom are independent of the asymptotic moduli, have an analogue at large volume, where families of black hole solutions appear effectively as single center black holes. Much like nucleons can be considered as elementary particles at sufficiently low energies.

To deal with these effectively single center solutions, we introduce the notion of “ λ -core”, or “core” for short. The defining property of a “ λ -core” is that the distances between constituents in a λ -core are bounded by $C\lambda^{-3}$ for sufficiently large λ , and for some fixed length C . Examples of such bound black holes states are of course proper single center black holes, as well as bound states of a D6-brane and anti-D6-brane, for which the wall of marginal stability lies in the interior of the Kähler moduli space.

The λ -core solutions are also distinguished in the uplift to five dimensions, and decoupling limit. Recall that the four dimensional Newton’s constant equals $G_4 = \ell_5^3/R$, with ℓ_5 the five dimensional Planck length and R the radius of the M-theory circle. Then, λ is related to these variables by $\lambda = R/\ell_5$ [27]. When uplifted to five dimensions and in the decoupling limit $\ell_5 \rightarrow 0$, states with D4-D2-D0 charges and within a λ -core will develop an asymptotically AdS_3 throat, whereas bound states with larger separation will decouple from the spectrum since the energy of their fluctuations diverges [27].

In a series of works [18, 28–30], the modular properties of partition functions enumerating D4-D2-D0 black holes are studied from a complementary perspective, namely by mapping this D-brane system to D3-D1-D(-1) instantons. These D-instantons correct the hypermultiplet geometry [17, 31–33], which is constrained by IIB $\text{SL}(2, \mathbb{Z})$ S-duality group. This duality group acts on the axion-dilation field $\chi := C^{(0)} + ie^{-\phi}$ by linear fractional transformations. Here $C^{(0)}$ is the RR scalar and ϕ is the dilaton. Alternatively, the duality can be identified with the modular symmetry of the worldvolume reduction of M5-branes. These have proven to be fruitful connections to determine the modular and analytic properties of the partition functions for D4-D2-D0 black holes. In this way, non-holomorphic contributions to the partition function are determined, which imply that the partition functions involve mock modular forms [34, 35], and mock modular forms of higher depth [36–39]. This implies potentially interesting arithmetic of the BPS indices, while the non-holomorphic contribution is also interesting independently, and a generalization of similar non-holomorphic terms in partition functions of $\mathcal{N} = 4$ Yang-Mills on four-manifolds [40, 41]. The origins and explicit expressions of these terms have been understood better recently [30, 42–48].

We will study in this paper the partition function of scaling black holes with three constituents. Each constituent is a λ -core carrying a positive D4-brane charge, but vanishing D6-brane charge. Intriguingly, each core gives rise to an associated $\text{AdS}_3/\text{CFT}_2$ after uplifting to M-theory, while the near coincident region gives rise to an asymptotic AdS_3 solution for the total magnetic charge, and should thus be captured by the $\text{AdS}_3/\text{CFT}_2$ correspondence for the total charge. In other words, they are examples of AdS fragmentation,

and attributed to the Coulomb branch of the CFT [1]. The existence of scaling black holes with these charges puts constraints on the topology of the Calabi-Yau threefold. In particular, these only exist if the second Betti number of the Calabi-Yau manifold $b_2 \geq 2$.

Our explicit analysis of scaling solutions gives rise to a decomposition for the partition function of attractor black holes \mathcal{Z}_P^λ with fixed magnetic charge P in terms of partition functions of single core and n -core scaling black holes \mathcal{Z}_P^{nT} . This reads explicitly (See also eq. (4.63).),

$$\mathcal{Z}_P^\lambda = \mathcal{Z}_P^T + \mathcal{Z}_P^{3T} + \dots, \tag{1.1}$$

where \mathcal{Z}_P^T is the partition function of single core black holes, \mathcal{Z}_P^{3T} the partition function of scaling black holes consisting of three cores, and the dots stand for scaling black holes with $n > 3$ cores. We determine the holomorphic part of the partition function using formula's for the degeneracies of black hole bound states such as studied in [6, 11, 49–53], which gives rise to a holomorphic indefinite theta series Ψ_μ of signature $(2, 2b_2 - 2)$. Since its coefficients grow polynomially, this demonstrates that the entropy arising from these solutions is exponentially smaller than the entropy of a single center black hole. This is expected since we have not included pure Higgs degeneracies [11, 49] or single center [51]. We have worked out two explicit case studies, where we specialize the CY three-fold to a K3 fibration and determine explicit q -series for the partition function. We do observe that the exponent of the leading term in the q -expansions is rather large.

We have moreover demonstrated that the partition function \mathcal{Z}_P^{3T} admits a theta function decomposition as a consequence of a spectral flow symmetry familiar from MSW CFT. This in contrast to generic bound states with non-vanishing D4-brane charge for which this symmetry is not present [25]. On the other hand for modular transformations, \mathcal{Z}_P^{3T} needs to be complemented with additional non-holomorphic terms, a phenomena which is also familiar for the attractor partition function \mathcal{Z}_P^λ as mentioned above. We distinguish the completed functions from the original functions by a hat, thus $\widehat{\mathcal{Z}}_P^\lambda$ for \mathcal{Z}_P^λ and $\widehat{\mathcal{Z}}_P^{3T}$ for \mathcal{Z}_P^{3T} and similarly for other functions. We then establish that $\widehat{\mathcal{Z}}_P^\lambda$ and $\widehat{\mathcal{Z}}_P^{3T}$ transform identically. Alexandrov and Pioline have derived in ref. [29] the non-holomorphic terms for the r.h.s., $\widehat{\mathcal{Z}}_P^\lambda$. It would be interesting to combine this with the non-holomorphic terms derived in the present paper for $\widehat{\mathcal{Z}}_P^{3T}$ to deduce the non-holomorphic terms of $\widehat{\mathcal{Z}}_P^T$.

The identical transformation properties of $\widehat{\mathcal{Z}}_P^\lambda$ and $\widehat{\mathcal{Z}}_P^{nT}$ raises the question, which term(s) correspond to the partition function of the conformal field theory. Since upon taking the decoupling limit to vanishing 5d Planck length $\ell_5 \rightarrow 0$ [27], the supergravity solutions contributing to $\widehat{\mathcal{Z}}_P^{3T}$ decouple from the AdS₃ geometry. This term is therefore not expected to correspond to states within the MSW conformal field theory, and it seems therefore plausible to us to expect that $\widehat{\mathcal{Z}}_P^T$ is to be identified with the MSW CFT partition function. It would be interesting to understand whether the finite difference between $\widehat{\mathcal{Z}}_P^T$ and $\widehat{\mathcal{Z}}_P^\lambda$ can arise by turning on the irrelevant perturbation in the CFT, which corresponds to moving away from the near horizon geometry and up the attractor flow [54].

Determination of the completion $\widehat{\mathcal{Z}}_P^{3T}$ amounts to determining the completion of the indefinite theta series Ψ_μ . We determine furthermore the non-holomorphic completion of a closely related function Φ_μ (4.58), which enumerates the number of “scaling” charge

configurations. Although still involved, determination of the non-holomorphic completion of this function is simpler than for generating function of the BPS indices. The analysis of Φ_μ and Ψ_μ demonstrates that these are mock modular forms of depth 2. There are different representations of the completion:

1. As a non-holomorphic kernel of the theta series. This involves (generalized) error functions [36]. The modular properties of the completion follow from application of results by Vignéras [55]. This is applied to the partition functions for scaling black holes in section 4.4. For the partition function $\widehat{\Phi}_\mu$ see equation (4.87), and for $\widehat{\Psi}_\mu$, see equation (4.116).
2. Another useful representation, well-known for mock modular forms [34, 35], is as a holomorphic q -series plus an (iterated) integral of modular forms [36, 43, 56, 57]. This form is determined for three-center scaling black holes in equation (4.90) and (4.105). The modular transformations of the q -series follow directly from those of the iterated integral. This representation is also relevant physically, where the non-holomorphic part is contributed to continuum of states [58], or the Coulomb branch [44, 48].
3. The third representation we mention here is as an integral over a domain in a symmetric space, studied by Funke, Kudla and Millson [38, 39, 59], also known as Narain moduli space in the context of conformal field theory. In the case of a 3-center scaling solution, this would be a union of triangles in $\text{SO}(2, 2b_2 - 2; \mathbb{Z}) \backslash \text{SO}(2, 2b_2 - 2; \mathbb{R}) / \text{SO}(2; \mathbb{R}) \times \text{SO}(2b_2 - 2; \mathbb{R})$. While we have not explored this representation in detail, it is interesting to mention in light of recent discussion on averaging over Narain moduli space and the $\text{AdS}_3/\text{CFT}_2$ correspondence [60, 61].

Though technically involved, we believe that our results can be extended to scaling solutions with $n > 3$ cores, and will give rise to mock modular forms of depth $n - 1$. Any Calabi-Yau manifold with $b_2 \geq 2$ gives rise to such mock modular forms, which thus provides a large resource of holomorphic higher depth mock modular forms. Further, it will be interesting to include single center degeneracies. Moreover, we hope that our results could be used for the study of AdS_3 fragmentation, and the interpretation of these solutions in the dual CFT. In this way, it may be possible to derive the non-holomorphic terms within gravity or the worldvolume theory of intersecting D-branes.

The outline of this paper is as follows. Section 2 reviews aspects of multi-center black holes, and in particular the index of scaling solutions. Section 3 reviews partition functions of D4-brane black holes. Section 4 discusses charge lattices for D4-brane bound states, and defines the partition functions of scaling black holes. We define here also the partition functions Φ_μ and Ψ_μ , which enumerate scaling configurations and their BPS indices, and determine its modular completion. Section 5 discusses the relation to M-theory and the decoupling limit AdS_3 . Section 6 considers two case studies for a specific Calabi-Yau 3-fold and charges, and presents the holomorphic q -series which are mock modular forms of depth 2.

2 Black hole solutions in $\mathcal{N} = 2$ supergravity

We briefly review in this section supersymmetric black holes in $\mathcal{N} = 2$ supergravity and partition functions.

2.1 Black hole bound states

Let X be a simply connected Calabi-Yau threefold, with triple intersection product d_{abc} , $a, b, c = 1, \dots, b_2$. The intersection product d_{abc} is symmetric in its indices. The classical central charge of a BPS state is given by

$$Z(\gamma, t) = - \int_X e^{-t} \wedge \gamma, \quad (2.1)$$

where γ on the r.h.s. is the Poincaré dual differential form of the homology class of the cycle which supports the D-branes, and is in 1-to-1 correspondence with the electric-magnetic charge of the BPS state. Moreover, t is the Kähler modulus of the Calabi-Yau three-fold.

The scalar fields X^I , $I = 0, \dots, b_2$ of the vector multiplets are related to the Calabi-Yau moduli, $t^a = B^a + iJ^a$, as $t^a = X^a/X^0$ for $a = 1, \dots, b_2$. Near the horizon, their values are determined by the attractor equations at the horizon in terms of the electric-magnetic charges of the black hole [62–64]. On the other hand, their asymptotic values for $|\vec{r}| \rightarrow \infty$ are boundary conditions for the equations of motion.

Besides the single center black hole, the equations of motion of $\mathcal{N} = 2$ supergravity give rise to intricate multi-center black hole solutions [3, 6, 65, 66]. Upon varying the asymptotic values of the scalar fields, multi-center solutions can cease to exist as proper solutions to the supergravity equations of motions, or reversely new solutions can appear. If the asymptotic values are chosen equal to the attractor values only few multi-center solutions exist. That is to say, only multi-center solutions which can be continuously connected to a single center black hole exist. These are the scaling solutions mentioned above, and are the main focus of this paper.

To understand this more explicitly, recall that an n -center solution, is required to satisfy the following $n - 1$ Denef equations,

$$\sum_{i \neq j} \frac{\langle \gamma_i, \gamma_j \rangle}{r_{ij}} = 2 \operatorname{Im} \left(e^{-i\alpha} Z(\gamma_i, t) \right) \Big|_{r=\infty}, \quad (2.2)$$

where \langle, \rangle is the symplectic innerproduct between the charges,

$$\langle \gamma_1, \gamma_2 \rangle = -P_1^0 Q_{0,2} + P_1 \cdot Q_2 - P_2 \cdot Q_1 + P_2^0 Q_{0,1}. \quad (2.3)$$

Moreover, $r_{ij} = |\vec{r}_i - \vec{r}_j|$ is the distance between the centers i and j , and α is the phase of the central charge $Z(\gamma, t)$ for the total charge $\gamma = \sum_{i=1}^n \gamma_i$. We set

$$c_j = 2 \operatorname{Im} \left(e^{-i\alpha} Z(\gamma_j, t) \right) \Big|_{r=\infty}. \quad (2.4)$$

We fix \vec{r}_1 at the origin \mathbb{R}^3 and let \mathcal{M}_n be the solution space for $\vec{r}_j \in \mathbb{R}^3$, $j = 2, \dots, n$ to (2.2). Then \mathcal{M}_n has dimension $2n - 2$. The low energy degrees of freedom of the supersymmetric

multi-center black hole give rise to $\mathcal{N} = 4$ quiver quantum mechanics [65]. The quiver for an n -center bound state with charges $\{\gamma_j\}$ consists of n nodes, and $\gamma_{ij} > 0$ arrows from node i to node j .

Note that the equations (2.2) are necessary but not sufficient for the multi-center solution to exist, to this end one needs to verify that the full supergravity solution is regular away from the black hole singularities, and without time-like curves [3]. Since we restrict to the large volume limit of the Calabi-Yau moduli space, we assume that this is the case in the following, and that we can determine the existence of bound states from (2.2).

The gravity perspective has led to the following form for the index of an n -center bound state [6, 9]. To express this, we first introduce the rational index $\bar{\Omega}(\gamma)$ associated to the integer index $\Omega(\gamma)$,

$$\bar{\Omega}(\gamma) = \sum_{m|\gamma} \frac{\Omega(\gamma/m)}{m^2}. \tag{2.5}$$

The single center invariants $\Omega_S(\gamma)$ are the internal degeneracies of a supersymmetric particle or black hole with charge γ . It is expected to be a positive integer for a black hole. To analyze the spectrum of bound states it is convenient to introduce a fugacity y for angular momentum. The rational variant of the refined index is defined as

$$\bar{\Omega}(\gamma, y) = \sum_{m|\gamma} \frac{1}{m} \frac{y - y^{-1}}{y^m - y^{-m}} \Omega(\gamma/m, y^m). \tag{2.6}$$

This reproduces (2.5) in the limit $y \rightarrow 1$. A few variants of BPS indices will be important for us. We mention,

- The BPS index $\Omega(\gamma; t)$, which enumerates BPS states for a given value t of the asymptotic moduli. This includes single-center BPS states as well as bound states.
- The attractor index $\Omega(\gamma; t_\gamma)$ with the moduli t fixed at the attractor point determined by γ . These indices are the input for determining $\Omega(\gamma; t)$ using attractor flow trees [3, 4, 6, 26, 70].
- The single-center index $\Omega_S(\gamma)$, which is the internal degeneracy of a BPS particle or black hole center. This invariant is independent of the moduli t . These invariants are the input determining $\Omega(\gamma; t)$ using the Coulomb branch formula [67, 68].
- The total invariant $\Omega_T(\gamma)$, which is a composite of $\Omega_S(\gamma)$ and independent of the moduli t . We give the expression below in (2.9).

The rational refined BPS index $\bar{\Omega}(\gamma, y; t)$ can be expressed as a sum over partitions of γ , each representing a BPS bound state. It takes the form [9, 67]

$$\bar{\Omega}(\gamma, y; t) = \sum_{\gamma = \sum_{i=1}^n \gamma_i} \frac{g_C(\{\gamma_j\}, \{c_j\}, y)}{|\text{Aut}(\{\gamma_j\})|} \prod_{j=1}^n \bar{\Omega}_T(\gamma_j, y), \tag{2.7}$$

with,

- $|\text{Aut}(\{\gamma_j\})|$ is the order of the subgroup of the permutation group, which preserves the ordered set $\{\gamma_1, \dots, \gamma_n\}$.
- The index g_C can be determined using localization of the black hole solution with respect to rotation around a fixed axis generated by J_3 , say the z -axis [68]. A fixed point $p \in \mathcal{M}_n$ corresponds to a collinear solution with all centers placed on the z -axis. If the associated bound state quiver has no oriented loop, g_C is the refined index of the $\mathcal{N} = 4$ quiver quantum mechanics describing the bound state [9],

$$g_C(\{\gamma_j\}, \{c_j\}, y) = \text{Tr}'_{\mathcal{H}_{\text{qm}}}(-y)^{2J_3}, \tag{2.8}$$

where the trace is over the BPS Hilbert space \mathcal{H}_{qm} of the quiver quantum mechanics, and J_3 is one of the generators of $\text{SU}(2)$.

- Ω_T the *total invariant* defined in terms of single centered indices Ω_S by

$$\Omega_T(\gamma, y) = \Omega_S(\gamma, y) + \sum_{\sum_{j=1}^n m_j \gamma_j = \gamma} H(\{\gamma_i, m_i\}, y) \prod_{i=1}^n \Omega_S(\gamma_i, y^{m_i}), \tag{2.9}$$

where $m_j \in \mathbb{N}$ are multiplicities of the charges in the partition of γ . For bound states whose associated quiver has no closed loops, the $H(\{\gamma_i, m_i\}, y)$ vanish. Otherwise they are determined by the “minimal modification hypothesis”. This has the effect that if we express (2.7) as

$$\bar{\Omega}(\gamma, y; t) = \sum_{\gamma = \sum_{i=1}^n \gamma_i} \frac{\bar{g}_C(\{\gamma_j\}, \{c_j\}, y)}{|\text{Aut}(\{\gamma_j\})|} \prod_{j=1}^n \bar{\Omega}_S(\gamma_j, y), \tag{2.10}$$

then the \bar{g}_C are $\text{SU}(2)$ characters.

To determine g_C using localization, it is convenient to introduce the refined index

$$g_C(\{\gamma_j\}, \{c_j\}, y) = \text{Tr}'_{\mathcal{H}_{\text{qm}}}(-y)^{2J_3}.$$

Let z_j be the position of the center with charge γ_j . The localization technique then gives the following sum over collinear fixed points with respect to rotation around this axis,

$$g_C(\{\gamma_j\}, \{c_j\}; y) = (-1)^{n-1} (y - y^{-1})^{-n+1} \sum_{p \in \mathcal{M}_n} s(p) (-y)^{\sum_{i < j} \gamma_{ij} \text{sgn}(z_j - z_i)}, \tag{2.11}$$

where $s(p) \in \pm 1$ is a sign depending on the details of the fixed point, and z_j is the z -coordinate of center j . If the associated quiver does not contain loops, this is the complete index and the $y \rightarrow 1$ limit is well-defined. However, if the quiver contains loops, the distances between the black hole centers may be arbitrarily small [6]. Such solutions are known as scaling solutions, and additional fixed points need to be included in (2.11). An (n -center) *scaling black hole* is a multi-center solution of n black holes, whose phase space

\mathcal{M}_n contains a region where the centers can approach each other arbitrarily close. Thus, while the centers are spatially separated for generic points of \mathcal{M}_n , they are adiabatically connected to the black hole solution with a single center.

While many BPS bound states decay if we tune the moduli to their attractor values, scaling solutions remain part of the BPS spectrum. Since the index is evaluated at the attractor point c_j^* , each term on the r.h.s. of (2.7) with $n \geq 3$ corresponds to a scaling solution.

Various quantities may diverge in the limit $y \rightarrow 1$, such as Ω_T and g_C . In order to arrive at numerical counterparts for these quantities, we propose to regularize a rational function of the form

$$\frac{f(y)}{(y - y^{-1})^\ell}, \quad \text{with} \quad \lim_{y \rightarrow 1} f(y) \neq 0, \quad (2.12)$$

as follows

$$\frac{f(y)}{(y - y^{-1})^\ell} \longrightarrow \frac{1}{2^\ell \ell!} \left(y \frac{d}{dy} \right)^\ell f(y) \Big|_{y=1}. \quad (2.13)$$

2.2 Bound state indices

Let us consider the equations (2.2) for small values of n . For $n = 2$, there is a single equation,

$$\frac{\gamma_{12}}{r_{12}} = c_1. \quad (2.14)$$

We deduce that the two-center solution only exists as a physical solution if $\gamma_{12} c_1 > 0$. This depends on the moduli t . If t approaches a value where c_1 vanishes, r_{12} diverges and the solution disappears as a solution to the supergravity equations of motion. In particular for the attractor point, the two-center solution never exists.

The quantum states of the two-center solution correspond to the product of the internal degeneracies of the centers times the states of a spin $(|\gamma_{12}| - 1)/2$ multiplet, which arises due to the electric-magnetic fields sourced by the charges of the two-centers [65]. We express it here as the product

$$\Omega_2(\gamma_1 + \gamma_2; t) = g_C(\{\gamma_1, \gamma_2\}, \{c_1, c_2\}) \Omega(\gamma_1) \Omega(\gamma_2), \quad (2.15)$$

with

$$g_C(\{\gamma_1, \gamma_2\}, \{c_1, c_2\}) = \frac{1}{2} (\text{sgn}(\gamma_{12}) + \text{sgn}(c_1)) (-1)^{\gamma_{12}-1} \gamma_{12}, \quad (2.16)$$

and $\Omega(\gamma_j)$ are degeneracies of the individual centers. For sgn , we use the definition

$$\text{sgn}(x) = \begin{cases} 1, & x > 0, \\ 0, & x = 0, \\ -1, & x < 0. \end{cases} \quad (2.17)$$

The function $\frac{1}{2} (\text{sgn}(\gamma_{12}) + \text{sgn}(c_1))$ equals 1 if the solution to (2.14) is physical, i.e. $r_{12} > 0$, and it vanishes if the sign of r_{12} is unphysical, $r_{12} < 0$. The factor $(-1)^{\gamma_{12}-1} \gamma_{12}$ is the number of states of the bound state, assuming that it exists. The case that $c_1 = 0$ is a special case, we aim to avoid. At the attractor point for the total charge, the c_j (2.4) are equal to c_j^* , given by [3, 4]

$$c_j^* = |Z(\gamma, t_\gamma^*)| \langle \gamma, \gamma_j \rangle. \quad (2.18)$$

As a result, two-center solutions do not exist at the attractor point, because substituting c_1^* in (2.14) gives a negative value for r_{12} which is unphysical.

For $n = 3$ distinct charges, (2.2) gives two independent equations,

$$\begin{aligned} \frac{\gamma_{12}}{r_{12}} + \frac{\gamma_{13}}{r_{13}} &= c_1, \\ \frac{\gamma_{21}}{r_{12}} + \frac{\gamma_{23}}{r_{23}} &= c_2. \end{aligned} \tag{2.19}$$

An intriguing aspect of these equations is that for appropriate values of γ_{ij} , they can be satisfied with positive r_{ij} for all pairs $i \neq j$, for t at the attractor point. Then, there is a one-parameter family of solutions [6], with

$$r_{ij} = \varepsilon \gamma_{ij} + O(\varepsilon^2), \quad \varepsilon \rightarrow 0, \tag{2.20}$$

for ij equal to 12, 23 and 31.

We can in fact do better, and give an all order solution in ε . The parameter ε together with three angular variables form the 4-dimensional solution space to (2.19). We set

$$\gamma_{12} = a, \quad \gamma_{23} = b, \quad \gamma_{31} = c, \tag{2.21}$$

in the following, and assume $a, b, c > 0$.¹ We can then verify that the following 1-parameter family of distances r_{ij} satisfy Denef's equations (2.2),

$$\frac{1}{r_{12}} = \frac{1}{a\varepsilon} - \rho_a, \quad \frac{1}{r_{23}} = \frac{1}{b\varepsilon} - \rho_b, \quad \frac{1}{r_{31}} = \frac{1}{c\varepsilon} - \rho_c, \tag{2.22}$$

where ρ_a, ρ_b, ρ_c satisfy $(\rho_c c - \rho_a a) = c_1$, $(\rho_a a - \rho_b b) = c_2$, $(\rho_b b - \rho_c c) = c_3$. The range of ε is determined by triangle inequalities and positivity of r_{12}, r_{23}, r_{31} . The equations for $\rho_{a,b,c}$ allow for a shift, which modifies the range of ε accordingly.

We discuss this now in detail for the attractor point, where we can use (2.18) such that $|Z(\gamma, t_\gamma^*)| = M$ and $\rho_a = \rho_b = \rho_c = M$. We then have,

$$\frac{1}{r_{12}} = \frac{1}{a\varepsilon} - M, \quad \frac{1}{r_{23}} = \frac{1}{b\varepsilon} - M, \quad \frac{1}{r_{31}} = \frac{1}{c\varepsilon} - M. \tag{2.23}$$

The free parameter ε is bounded from below by 0. In $\varepsilon \ll 1/M$ regime, this solution reduces to eq. (2.20); thus existence of the scaling solution requires a, b, c to obey triangle inequalities [6]. As ε is increased, the shape of the triangle changes, and we need to determine the correct maximum of ε domain. First, positivity of r_{12}, r_{23}, r_{31} imposes the upper bound $\varepsilon \leq \frac{1}{M \max(a,b,c)}$. However, we also need to impose that r_{ij} satisfy the triangle inequalities, which imposes an even stronger upper bound. E.g. the condition $r_{12} + r_{23} \geq r_{31}$ requires

$$(a + b - c) - 2ab\varepsilon M + abc\varepsilon^2 M^2 \geq 0. \tag{2.24}$$

If $(c - a)(c - b) < 0$, this condition is always satisfied since $a + b - c > 0$ and the l.h.s. does not have real roots in this case. Moreover if $(c - a)(c - b) \geq 0$, the condition

¹We apologize for the multiple use of a, b and c .

is saturated for $\varepsilon_c^\pm = \frac{1}{Mc} \left[1 \pm \sqrt{\frac{(c-a)(c-b)}{ab}} \right]$ and violated for $\varepsilon_c^- < \varepsilon < \varepsilon_c^+$. Both roots are non-negative provided (a, b, c) obeys the triangle inequality $a + b \geq c$. Noting that $\varepsilon_c^+ > \frac{1}{Mc}$, we see $\varepsilon \geq \varepsilon_c^+$ makes r_{31} negative. Thus we must have $\varepsilon \leq \varepsilon_c^-$. Using two other triangle inequalities, we have $\varepsilon \in (0, \min(\varepsilon_a^-, \varepsilon_b^-, \varepsilon_c^-)_{\mathbb{R}}]$ where $\varepsilon_a^- = \frac{1}{Ma} \left[1 - \sqrt{\frac{(a-b)(a-c)}{bc}} \right]$, $\varepsilon_b^- = \frac{1}{Mb} \left[1 - \sqrt{\frac{(b-a)(b-c)}{ac}} \right]$, and $\min(\varepsilon_a^-, \varepsilon_b^-, \varepsilon_c^-)_{\mathbb{R}}$ means the minimum among the $\varepsilon_{a,b,c}^- \in \mathbb{R}$. One of $\varepsilon_{a,b,c}^-$ may be complex. At the maximal value of ε the configuration is collinear. With the three additional angular variables, one can align the solution along the z -axis, thus giving a fixed point for rotations around this axis.

For three centers, the sum over fixed points reads

$$g_C(\{\gamma_j\}; \{c_j\}; y) = \frac{(-1)^{a+b+c}}{(y-y^{-1})^2} \left[F(123)y^{a+b-c} + F(321)y^{-a-b+c} + F(213)y^{-a+b-c} \right. \\ \left. + F(312)y^{a-b+c} + F(132)y^{a-b-c} + F(231)y^{-a+b+c} \right], \quad (2.25)$$

where

$$F(ijk) := F(ijk; \{c_j\}) = \begin{cases} s(p), & \exists \text{ a fixed point } p \in \mathcal{M}_n \text{ with } z_i < z_j < z_k, \\ 0, & \nexists \text{ a fixed point } p \in \mathcal{M}_n \text{ with } z_i < z_j < z_k. \end{cases} \quad (2.26)$$

Since $a, b, c \in \mathbb{Z}$, the signs $(-1)^{a \pm b \pm c}$ equal $(-1)^{a+b+c}$, and we can factor this out from the sum over permutations. The dependence on the r.h.s. on $\{c_j\}$ is through \mathcal{M}_n .

Since $F(123; \{c_j\}) = F(321; \{c_j\})$, we can shorten g_C to

$$g_C(\{\gamma_j\}; \{c_j\}; y) = \frac{(-1)^{a+b+c}}{(y-y^{-1})^2} \left[F(123; \{c_j\}) (y^{a+b-c} + y^{-a-b+c}) \right. \\ \left. + F(213; \{c_j\}) (y^{-a+b-c} + y^{a-b+c}) + F(132; \{c_j\}) (y^{a-b-c} + y^{-a+b+c}) \right]. \quad (2.27)$$

This expression does not have a smooth $y \rightarrow 1$ limit if \mathcal{M}_n is non-compact and contains a scaling region. In that case, only one of $F(ijk; \{c_j\})$'s is non-vanishing. Turning on a fugacity is indeed known to be subtle for non-compact phase spaces [69].

In the present context, the minimal modification hypothesis is put forward in [68] to correct this. It adds a term with minimal angular momentum corresponding to the fixed point with coincident centers. The effect of the minimal modification hypothesis for three distinct charges is that the refined index $g_C(\{\gamma_j\}; \{c_j\})$ is completed to

$$\bar{g}_C(\{\gamma_j\}; \{c_j\}, y) = g_C(\{\gamma_j\}; \{c_j\}, y) + H(\{\gamma_j\}, y), \quad (2.28)$$

with

$$H(\{\gamma_j\}, y) = \begin{cases} -\frac{2}{(y-y^{-1})^2}, & \text{if } a + b + c \in 2\mathbb{Z}, \\ \frac{y+y^{-1}}{(y-y^{-1})^2}, & \text{if } a + b + c \in 2\mathbb{Z} + 1, \end{cases} \quad (2.29)$$

\bar{g}_C has a well-defined $y \rightarrow 1$ limit, which reads,

$$\begin{aligned} \bar{g}_C(\{\gamma_j\}; \{c_j\}) &= \lim_{y \rightarrow 1} \bar{g}_C(\{\gamma_j\}; \{c_j\}; y) \\ &= (-1)^{a+b-c} [F(123) s(a, b, c) + F(213) s(a, c, b) + F(132) s(c, b, a)], \end{aligned} \tag{2.30}$$

with

$$s(a, b, c) = \begin{cases} \frac{(a+b-c)^2}{4}, & \text{if } a + b + c \in 2\mathbb{Z}, \\ \frac{(a+b-c)^2-1}{4}, & \text{if } a + b + c \in 2\mathbb{Z} + 1. \end{cases} \tag{2.31}$$

We note that for degenerate cases, where one or more among a, b and c vanish, g_C can be a multiple of $1/2$ or $1/4$ rather than in $\{-1, 0, 1\}$.

Using the regularization (2.13), we obtain for the numerical version (2.29) of H ,

$$H(\{\gamma_j\}) = \begin{cases} 0, & \text{if } a + b + c \in 2\mathbb{Z}, \\ \frac{1}{4}, & \text{if } a + b + c \in 2\mathbb{Z} + 1, \end{cases} \tag{2.32}$$

which in turn gives the numerical counterpart for Ω_T ,

$$\Omega_T(\gamma) = \Omega_S(\gamma) + \begin{cases} 0, & \text{if } a + b + c \in 2\mathbb{Z}, \\ \frac{1}{4} \prod_{j=1}^3 \Omega_S(\gamma_j), & \text{if } a + b + c \in 2\mathbb{Z} + 1. \end{cases} \tag{2.33}$$

We obtain similarly using (2.13) for the numerical version of g_C ,

$$\begin{aligned} g_C(\{\gamma_j\}; \{c_j\}) &= \frac{(-1)^{a+b+c}}{4} \times [F(123; \{c_j\}) (a + b - c)^2 \\ &\quad + F(213, \{c_j\}) (a - b + c)^2 + F(132, \{c_j\}) (a - b - c)^2]. \end{aligned} \tag{2.34}$$

Thus the numerical g_C essentially corresponds to the $y \rightarrow 1$ limit of the equivariant volume of the solution space \mathcal{M}_3 [68].

The term $F(123, \{c_j\})$ is determined in [51, equation (2.57)]. For our purposes we rewrite this in terms of sgn rather than the step function. This reads,

$$F(123, \{c_j\}) = F_1(a, b, c, \{c_j\}) + F_2(a, b, c), \tag{2.35}$$

with

$$\begin{aligned} F_1(a, b, c, \{c_j\}) &= \frac{1}{4} (\text{sgn}(a) + \text{sgn}(c_1)) (\text{sgn}(b) - \text{sgn}(c_3)), \\ F_2(a, b, c) &= \frac{1}{4} (\text{sgn}(a) + \text{sgn}(b)) (\text{sgn}(a + b - c) - \text{sgn}(a + b)). \end{aligned} \tag{2.36}$$

At special charge configurations, where one or more arguments of the signs vanish, this may differ from [51, equation (2.57)]. In such cases, the F_j can be a fraction rather than an integer. To deal properly with such cases, we include additional terms below in eqs. (2.42) and (2.45).

Our interest in this paper is the index at the attractor point, thus

$$g_C(\{\gamma_j\}; \{c_j^*\}), \tag{2.37}$$

where c_j^* (2.18) is c_j evaluated at the attractor point t_γ^* . Then F_1 reads

$$F_1(a, b, c) = \frac{1}{4} (\text{sgn}(a) + \text{sgn}(c - a)) (\text{sgn}(b) + \text{sgn}(c - b)). \quad (2.38)$$

Assuming non-vanishing arguments of the sign functions, we can simplify the sum $F^*(123) = F_1 + F_2$ using the identity [70, eq. (A.1)]²

$$(\text{sgn}(x_1) + \text{sgn}(x_2)) \text{sgn}(x_1 + x_2) - \text{sgn}(x_1) \text{sgn}(x_2) = 1, \quad (x_1, x_2) \neq (0, 0), \quad (2.39)$$

to $F^*(123) = F_1(a, b, c) + F_2(a, b, c)$ [70, eq. (4.10)],

$$F^*(123) = \frac{1}{4} (1 + \text{sgn}(a - c) \text{sgn}(b - c) + \text{sgn}(b - c) \text{sgn}(c - a - b) + \text{sgn}(c - a - b) \text{sgn}(a - c)). \quad (2.40)$$

We stress that this expression may differ from (2.35) if the arguments of some products of sgn 's vanish. For example for $a = 0, b = c = 1$, $F(123) = 0$, while $F^*(123)$ equals $\frac{1}{4}$.

For the other permutations, we also define

$$\begin{aligned} F_3(a, b, c) &= \frac{1}{4} (\text{sgn}(a) + \text{sgn}(b - a)) (\text{sgn}(c) + \text{sgn}(b - c)), \\ F_4(a, b, c) &= \frac{1}{4} (\text{sgn}(a) + \text{sgn}(c)) (\text{sgn}(a + c - b) - \text{sgn}(a + c)), \\ F_5(a, b, c) &= \frac{1}{4} (\text{sgn}(c) + \text{sgn}(a - c)) (\text{sgn}(b) + \text{sgn}(a - b)), \\ F_6(a, b, c) &= \frac{1}{4} (\text{sgn}(c) + \text{sgn}(b)) (\text{sgn}(c + b - a) - \text{sgn}(c + b)), \end{aligned} \quad (2.41)$$

and

$$\begin{aligned} F^*(213) &= \frac{1}{4} (1 + \text{sgn}(a - b) \text{sgn}(c - b) + \text{sgn}(c - b) \text{sgn}(b - a - c) + \text{sgn}(b - a - c) \text{sgn}(a - b)), \\ F^*(132) &= \frac{1}{4} (1 + \text{sgn}(c - a) \text{sgn}(b - a) + \text{sgn}(b - a) \text{sgn}(a - b - c) + \text{sgn}(a - b - c) \text{sgn}(c - a)). \end{aligned}$$

Having defined the $F^*(ijk)$, we turn to $g_C(\{\gamma_j\}; \{c_j^*\})$ and take care of the special cases where both arguments of a products of sgn 's vanish. Let us first determine for which values of a, b and c , $g_C(\{\gamma_j\}; \{c_j^*\})$ is affected. When the last two products of sgn 's of $F^*(123)$ (2.40) vanish, the angular momentum factor $(a + b - c)^2/4$ also vanishes. Thus replacing these products of sgn 's with a non-vanishing value will not affect g_C . For the remaining product, $\text{sgn}(a - c) \text{sgn}(b - c)$'s, the arguments vanish in the equilateral case $a = b = c$ for which the angular momentum factor is $a^2/4$. This is the same as for the other permutations, $F^*(213)$ and $F^*(132)$, such that we can take all three into account by adding a single additional term with yet undetermined coefficient A . We obtain thus for g_C at the attractor point,

$$\begin{aligned} g_C(\{\gamma_j\}; \{c_j^*\}) &= \frac{(-1)^{a+b+c}}{4} \left[F^*(123) (a + b - c)^2 + F^*(213) (a - b + c)^2 \right. \\ &\quad \left. + F^*(132) (-a + b + c)^2 + \frac{1}{4} A \delta_{a,c} \delta_{b,c} a^2 \right]. \end{aligned} \quad (2.42)$$

²We thank Sergey Alexandrov for stressing the simplifying power of this identity.

At this point, we can make a “guess” of the preferred “physical” value for A . From the gravity perspective, the equilateral configurations with $a = b = c$ are perfectly fine multi-center solutions, such that it is most natural that these contribute with $1 \times (-1)^a a^2/4$ to g_C . For (2.42), we have instead $(3 + A) \times (-1)^a a^2/16$, thus suggesting $A = 1$. We will demonstrate in section 4.5 that modularity of the q -series leads to exactly the same value.

With this we arrive at our final expression for the numerical attractor index $\Omega(\gamma; t_\gamma)$ with γ partitioned into at most three constituent charges is given by,

$$\bar{\Omega}(\gamma; t_\gamma) = \Omega_T(\gamma) + \sum_{\gamma = \sum_{i=1}^3 \gamma_i} g_C(\{\gamma_j\}; \{c_j^*\}) \prod_{j=1}^3 \bar{\Omega}_T(\gamma_j), \tag{2.43}$$

with $g_C(\{\gamma_j\}; \{c_j^*\})$ given by (2.42), and $\Omega_T(\gamma)$ given in terms of Ω_S by (2.33).

Besides the bound state index g_C , we are also interested to enumerate the number of charge configurations giving rise to scaling black holes. Up to vanishing arguments, the sum $F_{\text{total}}(a, b, c) = \sum_{j=1}^6 F_j(a, b, c)$ can be further simplified to

$$F_{\text{total}}(a, b, c) = \frac{1}{4} \left[1 + \text{sgn}(a + b - c) \text{sgn}(a + c - b) + \text{sgn}(a + c - b) \text{sgn}(b + c - a) + \text{sgn}(b + c - a) \text{sgn}(a + b - c) \right]. \tag{2.44}$$

This expression has the advantage that for $a + b + c$ odd, the arguments of the sgn 's never vanish. With inclusion of additional terms to deal with vanishing arguments, we define number f_C ,

$$f_C(\{\gamma_j\}, \{c_j^*\}) = F_{\text{total}}(a, b, c) (-1)^{a+b+c} + \frac{1}{4} A_1 \delta_{a,0} \delta_{b,c} + \frac{1}{4} A_2 \delta_{c,0} \delta_{a,b} + \frac{1}{4} A_3 \delta_{b,0} \delta_{a,c}, \tag{2.45}$$

where the constants A_j are yet to be determined. We will determine these from the modular completion. To our surprise, these numbers are typically *irrational* for $a + b + c \in 2\mathbb{Z}$, while they do not contribute for $a + b + c \in 2\mathbb{Z} + 1$. We find the irrationality quite peculiar. On the other hand, f_C is not a BPS index, such that it is not really violating any physical principles.

3 Review of D4-brane black holes

We review in this section aspects of partition functions of D4-brane black holes. We start in subsection 5 by reviewing the uplift of D4-brane black holes to M-theory. In subsection 3.1, we discuss the “supergravity” partition function, which enumerates D4-brane BPS indices $\Omega(\gamma; t)$ for fixed Kähler modulus t . In subsection 3.2, we discuss the “attractor” partition function, which is a generating function of BPS indices $\Omega(\gamma; t_\gamma^*)$ evaluated at the attractor point t_γ^* of the corresponding charge γ .

3.1 Supergravity partition function

From the perspective of asymptotically flat \mathbb{R}^4 , the most natural BPS partition function enumerates the BPS indices for a fixed, asymptotic value of the (Kähler) moduli t , and in

the mixed ensemble where the magnetic charge P is fixed [6, 13, 14, 71] and the electric charge Q is varied. The electric charge takes value in the lattice $\Lambda \simeq \mathbb{Z}^{b_2}$ with bi-linear form $D_{ab} = d_{abc}P^c$. For a positive magnetic charge P^a , $d_{abc}P^c$ provides a non-degenerate quadratic form with signature $(1, b_2 - 1)$ for the lattice Λ of magnetic charges. The electric charges Q_a take values in dual lattice Λ^* , with quadratic form $D^{ab} = (d_{abc}P^c)^{-1}$. We abbreviate the pairing between an element $Q \in \Lambda^*$ and $k \in \Lambda$ as

$$\sum_{a=1}^{b_2} Q_a P^a = Q \cdot P \tag{3.1}$$

and extend this by linearity in each argument to $\Lambda^* \otimes \mathbb{R}$ and $\Lambda \otimes \mathbb{R}$. For later use, we also introduce the notation $\mu \in \Lambda^*/\Lambda$,

$$\Lambda_\mu^* = \{Q \in \Lambda + \mu + P/2\}. \tag{3.2}$$

We stress that the elements of Λ_μ^* do not necessarily lie in Λ^* due to the shift by $P/2$. Using this notation, the partition function for D4-D2-D0 black holes reads schematically

$$\mathcal{Z}_{SG}(\tau, C, t) = \sum_{Q_0, Q_a} \bar{\Omega}(\gamma, t) (-1)^{P \cdot Q} e^{-2\pi\tau_2 |Z(\gamma, t)| + 2\pi i \tau_1 (Q_0 - Q \cdot B + B^2/2) + 2\pi i C \cdot (Q - B/2)}, \tag{3.3}$$

where $\bar{\Omega}(\gamma, t)$ is the rational index (2.5), $\tau_1 \in \mathbb{R}$ is the RR 1-form C_1 , and $\tau_2 = e^{-\phi} \in \mathbb{R}_+$ with ϕ being the dilaton, C the RR 3-form and B the B -field.

Here $Z(\gamma, t)$ is the central charge of the $\mathcal{N} = 2$ algebra. Ignoring the non-perturbative terms in the strict large volume limit, Z reads

$$Z(\gamma, t) = \frac{1}{2} P \cdot (J^2 - B^2) + Q \cdot B - Q_0 + i(Q - BP) \cdot J. \tag{3.4}$$

The BPS mass becomes in this limit,

$$|Z(\gamma, t)| = \frac{1}{2} P \cdot (J^2 - B^2) + Q \cdot B - Q_0 + \frac{((Q - BP) \cdot J)^2}{P \cdot J^2}, \tag{3.5}$$

up to inverse powers of J . By assumption $J^2 > 0$, hence J^a lies in the positive cone of Λ . Thus $\frac{k \cdot J J}{J^2}$ is the component of the vector k along J .

In the large volume limit, supergravity has an $\text{Sp}(2b_2 + 2, \mathbb{Z})$ symmetry, generated by matrices

$$\mathbb{K}(k) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ k^a & \mathbb{I}_{b_2} & 0 & 0 \\ \frac{1}{2} d_{abc} k^b k^c & d_{abc} k^c & \mathbb{I}_{b_2} & 0 \\ \frac{1}{6} d_{abc} k^a k^b k^c & \frac{1}{2} d_{abc} k^b k^c & k^a & 1 \end{pmatrix}, \quad k \in \mathbb{Z}^{b_2}, \tag{3.6}$$

and acts linearly on $\gamma = (P^0, P^a, Q_a, Q_0)$. For $P^0 = 0$, these transformations preserve the magnetic charge, and act on remaining charges and moduli as follows:

$$\begin{aligned} Q_a &\rightarrow Q_a + d_{abc} k^b P^c, \\ Q_0 &\rightarrow Q_0 + k^a Q_a + \frac{1}{2} d_{abc} k^a k^b P^c, \\ t^a &\rightarrow t^a + k^a. \end{aligned} \tag{3.7}$$

We introduce the abbreviations:

$$\begin{aligned}
 \hat{Q}_0 &:= -Q_0 + \frac{1}{2}Q^2, \\
 \hat{Q} &:= Q - B, \\
 \hat{Q}_-^2 &:= \hat{Q}^2 - \hat{Q}_+^2, \\
 (Q - B)_+^2 &:= \frac{((Q - BJ).P)^2}{PJ^2}.
 \end{aligned}
 \tag{3.8}$$

Note that the combinations \hat{Q}, \hat{Q}_0 and the conjugacy class μ of electric charge vector are invariant under the transformations (3.7). Invariance of the conjugacy class is seen by noting that spectral flow shifts Q by integer lattice vectors, when mapped to magnetic lattice Λ and such shifts do not change the conjugacy class.

The holomorphic and anti-holomorphic dependence on τ can be made more manifest, if we rewrite \mathcal{Z}_{SG} as

$$\mathcal{Z}_{SG}(\tau, C, t) = e^{-\pi\tau_2 J^2} \sum_{Q_0, Q} \bar{\Omega}(P, Q, Q_0; t) (-1)^{P.Q} q^{\hat{Q}_0 - \hat{Q}_-^2/2} q^{\hat{Q}_+^2/2} e^{2i\pi C.(Q-B/2)}, \tag{3.9}$$

where $q = e^{2\pi i\tau}$. The transformations (3.7) act on \mathcal{Z}_{SG} as

$$\mathcal{Z}_{SG}(\tau, C, t) \rightarrow (-1)^{P.k} e^{\pi i C.k} \mathcal{Z}_{SG}(\tau, C, t), \tag{3.10}$$

under the action of $\mathbb{K}(k)$.

We can map the system of D4-D2-D0 branes to IIB string theory, by a T-duality along time circle. The RR 1-form C_1 is mapped to the RR 0-form C_0 , while the D4, D2 and D0-branes respectively become D3, D1 branes and D(-1)-instantons. Moreover, the action of the IIB strong-weak duality on the instantonic branes carries over to the spectrum of D4-D2-D0 branes in IIA string theory. The duality acts as follows. The modular parameter is $\tau = C_1 + ie^{-\phi}$, with ϕ being the dilaton and C_1 the Ramond-Ramond 1-form flux along S_t^1 . For an element $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z})$. This duality acts as

$$\begin{aligned}
 \tau &\rightarrow \frac{a\tau + b}{c\tau + d}, \\
 \begin{pmatrix} C \\ B \end{pmatrix} &\rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} C \\ B \end{pmatrix}, \\
 J &\rightarrow |c\tau + d|J,
 \end{aligned}
 \tag{3.11}$$

In a series of papers [6, 13, 14, 18, 25, 26, 29], the supergravity partition function (3.3) has been analyzed. We summarize the main properties:

- Quasi-periodicity in the two-form fields B and C .

For $k \in \Lambda$, we have

$$\begin{aligned}
 \mathcal{Z}_{SG}(\tau, C, t + k) &= (-1)^{P.k} e^{\pi i C.k} \mathcal{Z}_{SG}(\tau, C, t), \\
 \mathcal{Z}_{SG}(\tau, C + k, t) &= (-1)^{P.k} e^{-\pi i B.k} \mathcal{Z}_{SG}(\tau, C, t).
 \end{aligned}
 \tag{3.12}$$

These can be seen as large gauge transformations.

- $SL(2, \mathbb{Z})$ S-duality:

$$\begin{aligned} S : \quad & \mathcal{Z}_{SG}(-1/\tau, -B, C + |\tau|J) = \tau^{1/2} \bar{\tau}^{-3/2} \varepsilon(S) \mathcal{Z}_{SG}(\tau, C, t), \\ T : \quad & \mathcal{Z}_{SG}(\tau + 1, C + B, t) = \varepsilon(T) \mathcal{Z}_{SG}(\tau, C, t), \end{aligned} \tag{3.13}$$

where $\varepsilon(S) = \varepsilon(T)^{-3}$, $\varepsilon(T) = e^{-i\pi \frac{c_2(X).P}{12}}$ are phases. The factor $\bar{\tau}^{-3/2}$ can be understood from the non-compact bosons in the CFT due to the center of mass in \mathbb{R}^3 .

- The partition function involves an intricate dependence on the moduli t through wall-crossing. The proper modular invariant partition function differs from (3.9) by additional non-holomorphic, subleading terms which are non-holomorphic in τ .

3.2 Attractor partition function

The supergravity partition function is a rather complicated function. It has become clear that an alternative partition function is a more amenable object to study [18]. This is the attractor partition function \mathcal{Z}_P^λ , which is the generating function of indices $\bar{\Omega}(\gamma; t_\gamma^\lambda)$, where each index is evaluated at its (large volume) attractor point t_γ^λ . The indices $\bar{\Omega}(\gamma; t_\gamma^\lambda)$ were also referred to Ω_{CFT} in [25], and MSW invariants Ω^{MSW} in [18]. For irreducible magnetic charge P and magnetic charges which can be written as a sum of at most 2 magnetic charges, these indices are conjectured to coincide with the CFT indices. However as discussed in section 5, our findings in this paper suggest that there is a difference for generic magnetic charges due to scaling black holes.

This partition function is obtained by replacing $\Omega(\gamma, t)$ in the definition of \mathcal{Z}_{SG} by its attractor value $\Omega(\gamma, t_\gamma^\lambda)$. For a D4-D2-D0 black hole with charge $\gamma = (0, P^a, Q_a, Q_0)$, we have for the “large volume” attractor value $(t_\gamma^*)^a$,

$$(t_\gamma^\lambda)^a = D^{ab} Q_b + i\lambda P^a, \tag{3.14}$$

with λ sufficiently large, such that subleading terms in a λ expansion can be ignored. Eq. (3.14) is equivalent to

$$B_\gamma^a = D^{ab} Q_b, \quad (J_\gamma^\lambda)^a = \lambda P^a. \tag{3.15}$$

The precise proportionality constant λ between J^a and P^a will not be important for us, since we will restrict to the large volume limit, $|J| \gg 1$. On the other hand, we do not take the limit $\lambda \rightarrow \infty$, since physical quantities such as the BPS mass diverge in this limit.

The attractor or MSW partition function then reads,

$$\mathcal{Z}_P^\lambda(\tau, C, t) = \sum_{Q_0, Q_a} \bar{\Omega}(\gamma, t_\gamma^\lambda) (-1)^{P \cdot Q} e^{-2\pi\tau_2 |Z(\gamma, t)| + 2\pi i \tau_1 (Q_0 - Q \cdot B + B^2/2) + 2\pi i C \cdot (Q - B/2)}. \tag{3.16}$$

Note that although the degeneracy is evaluated at the attractor point t_γ^λ , the mass in the exponent is the ADM mass evaluated for moduli at infinity being t . The other moduli dependence in the exponent is also similar to that of $\mathcal{Z}_{SG}(\tau, C, t)$.

Since black holes states contributing to the attractor index exist everywhere in moduli space, their quantum degeneracies should be moduli independent. Moreover, one can show that $\Omega(\gamma; t_\gamma^\lambda)$ are invariant under spectral flow transformations (3.7). This entails that $\Omega(\gamma; t_\gamma^\lambda)$ depends on the charges only through the invariant combination \hat{Q}_0 and the conjugacy class μ . These imply the sum

$$h_{P,\mu}(\tau) := \sum_{Q_0} \bar{\Omega}(\gamma; t_\gamma^\lambda) q^{\hat{Q}_0}, \quad (3.17)$$

for fixed Q is also invariant under spectral flow transformations and apart from the magnetic charge, depends solely on the conjugacy class μ . It has been understood from wall-crossing and the perspective of D-instantons, that $h_{P,\mu}$ receives additional non-holomorphic contributions $h_{P,\mu}$ [18, 29]. The resultant is the non-holomorphic function $\hat{h}_{P,\mu}(\tau, \bar{\tau})$.

Hence one is led to the following theta function decomposition of the attractor partition function,

$$\mathcal{Z}_P^\lambda(\tau, C, t) = e^{-\pi\tau_2 J^2} \sum_{\mu \in \Lambda^*/\Lambda} \overline{h_{P,\mu}(\tau)} \Theta_\mu(\tau, \bar{\tau}, C, B), \quad (3.18)$$

$$\text{where } \Theta_\mu(\tau, \bar{\tau}, C, B) = \sum_{Q \in \Lambda_\mu^*} (-1)^{P \cdot Q} q^{\hat{Q}_+^2/2} \bar{q}^{-\hat{Q}_-^2/2} e^{2\pi i C \cdot (Q-B/2)}, \quad (3.19)$$

with the set Λ_μ^* is given in (3.2). We define analogously the completion $\hat{\mathcal{Z}}_P^\lambda$ by replacing $h_{P,\mu}$ by $\hat{h}_{P,\mu}$ in (3.18).

Various factors in the right hand side of (3.18) also have definite modular transformation properties. The prefactor $e^{-\pi\tau_2 J^2}$ is modular invariant. Moreover, the Siegel-Narain theta function $\Theta_\mu(\tau, C, B)$ transforms as

$$S: \quad \Theta_\mu(-1/\tau, -1/\bar{\tau}, -B, C) = \frac{1}{\sqrt{|\Lambda^*/\Lambda|}} (-i\tau)^{b_2^+/2} (i\bar{\tau})^{b_2^-/2} e^{-i\pi P^2/2} \\ \times \sum_{\nu \in \Lambda^*/\Lambda} e^{-2\pi i \mu \cdot \nu} \Theta_\nu(\tau, \bar{\tau}, C, B), \quad (3.20)$$

$$T: \quad \Theta_\mu(\tau + 1, \bar{\tau} + 1, C + B, B) = e^{i\pi(\mu+P/2)^2} \Theta_\mu(\tau, \bar{\tau}, B, C).$$

This implies that the appropriate completion $\hat{h}_{P,\mu}$ transforms as a vector valued modular form,

$$S: \quad \hat{h}_{P,\mu}(-1/\tau, -1/\bar{\tau}) = -\frac{1}{\sqrt{|\Lambda^*/\Lambda|}} (-i\tau)^{-b_2/2-1} \varepsilon(S)^* e^{-i\pi P^2/2} \sum_{\delta \in \Lambda^*/\Lambda} e^{-2\pi i \delta \cdot \mu} \hat{h}_{P,\delta}(\tau, \bar{\tau}), \\ T: \quad \hat{h}_{P,\mu}(\tau + 1, \bar{\tau} + 1) = \varepsilon(T)^* e^{i\pi(\mu+P/2)^2} \hat{h}_{P,\mu}(\tau, \bar{\tau}), \quad (3.21)$$

where $\varepsilon(S)$ and $\varepsilon(T)$ are as below (3.13).

4 Partition functions for scaling black holes

Our aim in this section is determine and analyze the first multi-center contribution to the attractor partition function (3.17) with the moduli at the large volume attractor points

t_γ^λ (3.14). The large volume attractor points are in many ways analogous to the proper attractor points t_γ . For this value of the moduli and assuming that γ and γ_j both carry D4-brane charge, the c_j (2.4) evaluated at t_γ^λ are to leading order equal to c_j^λ ,

$$c_j^\lambda = 2\lambda \langle \gamma, \gamma_j \rangle. \quad (4.1)$$

Since the factor 2λ on the r.h.s. is positive, the large volume attractor point is similar in nature to the c_j^* (2.18) for the proper attractor point.

On the other hand, specific bound state solutions may exist at t_γ^λ which are unstable at t_γ . Examples of such bound states are pairs of D6 and anti-D6 branes. The spatial separation of these such bounds state is of the order ℓ_5^3/R^2 , which decreases as λ^{-3} in the large λ limit. On the other hand, the separation of centers that carry D4-brane charge, but vanishing D6 charge scales as (4.1), scales as λ^{-1} except for the near coincident region. From the perspective of the large volume limit, the (D6, anti-D6) bound states are effectively single centers. To emphasize that they are bound states with a small separation, we introduce the notion of “core” for a black hole solution with separation of the order ℓ_5^3/R^2 between the centers. The discussion of section 2 is easily adapted to the large volume regime by reading “single core” for “single center”. One naturally defines a single core index Ω_{SC} and total core invariant Ω_{TC} analogous to the single center index Ω_{SC} and total invariant Ω_{TC} .

The first multi-center, or multi-core, contribution to the attractor partition function (3.17) consists of bound states with three cores, which each carry a positive D4-brane charge and vanishing D6-brane charge. The symplectic innerproducts a, b and c (2.21) read in this case,

$$\begin{aligned} a &= P_1 Q_2 - P_2 Q_1, \\ b &= P_2 Q_3 - P_3 Q_2, \\ c &= P_3 Q_1 - P_1 Q_3. \end{aligned} \quad (4.2)$$

This gives for the c_j^λ , $c_1^\lambda = 2\lambda(c - a)$, $c_2^\lambda = 2\lambda(a - b)$. Since the c_j^* (2.18) and c_j^λ (4.1) are simply related by the substitution $|Z(\gamma; t_\gamma^*)| \rightarrow 2\lambda$, the discussion on the solution to Denef’s equations below (2.23) is applicable, and in particular gives the separation of collinear solutions.

At the attractor point t_γ^λ many multi-center and multi-core configurations do not exist as physical solutions. However we have seen that some multi-center solutions do exist. These solutions are distinguished from generic solutions, since the centers can approach each other arbitrarily close. The scaling solutions do respect the spectral flow symmetry, since the symplectic innerproducts are invariant under (3.7).

We note that the existence scaling solutions poses constraints on the magnetic charges P_j . For example since for a scaling solution a, b, c must have the same sign, we deduce that there are no scaling solutions for $P_1 = P_2 = P_3 = P/3$, since this gives rise to $\gamma_{12} + \gamma_{23} + \gamma_{31} = a + b + c = 0$. Therefore, there must be an asymmetry in the magnetic charge of the centers for such scaling solutions to exist. To see that F vanishes in the case of three equal magnetic charges, note that the second factors in F_2, F_4, F_6 vanish immediately as $\text{sgn}(2x) - \text{sgn}(x) = 0$. Moreover, in F_1 we can replace $c = -a - b$ and when $a > 0$ the

first factor imposes the constraint $-2a - b \geq 0$ so $b < -2a$ and b must be negative. Hence the second factor in F_1 vanishes. The argument goes similarly for F_3, F_5 . Similarly one may show that such scaling black holes only exist for $b_2 > 1$.

4.1 Bound states and lattices

We discuss in this subsection various aspects of charge lattices of bound states, and the projection of the charge lattice to a sublattice with fixed total charge. The discussion in this subsection does not rely on the existence of scaling black holes for these charges and lattices.

Lattice decomposition. We apply some techniques of decompositions and gluings of integral lattices to charge lattices of bound states. See for example [72, 73]. We will consider a black hole bound state of n centers, with non-vanishing, positive magnetic charge $P_j \in \Lambda_j$, $j = 1, 2, \dots, n$, where Λ_j is the b_2 -dimensional lattice associated to the j 'th center with innerproduct D_j ,

$$D_{jab} = d_{abc} P_j^c, \quad a, b, c = 1, \dots, b_2. \quad (4.3)$$

The total magnetic charge is $P = \sum_{j=1}^n P_j$ with quadratic form D and b_2 -dimensional lattice Λ . The signature Λ_j and Λ is $(1, b_2 - 1)$.

We use boldface notation for the lattices for boundstates. For n centers, we introduce $(n b_2)$ -dimensional lattices and vectors,

$$\begin{aligned} \vec{k} &= (k_1, k_2, \dots, k_n) \in \mathbf{\Lambda} := \Lambda_1 \oplus \Lambda_2 \oplus \dots \oplus \Lambda_n, \\ \vec{x} &= (x_1, x_2, \dots, x_n) \in \mathbf{\Lambda}^* := \Lambda_1^* \oplus \Lambda_2^* \oplus \dots \oplus \Lambda_n^*, \\ \vec{Q} &= (Q_1, Q_2, \dots, Q_n) \in \mathbf{\Lambda}^* + \vec{P}/2 = \Lambda_1^* \oplus \Lambda_2^* \oplus \dots \oplus \Lambda_n^* + (P_1, P_2, \dots, P_n)/2. \end{aligned} \quad (4.4)$$

We denote the quadratic form for $\mathbf{\Lambda}$ by $\vec{D} = \text{diag}(D_1, D_2, \dots, D_n)$.

Since we typically sum over bound states with fixed total electric charge, we aim to decompose the lattice $\mathbf{\Lambda}$ in a b_2 -dimensional sublattice $\bar{\mathbf{\Lambda}} \subset \mathbf{\Lambda}$ representing the total charge, and its orthogonal complement $\underline{\mathbf{\Lambda}}$ with dimension $(n - 1)b_2$ representing the relative charge distribution over the constituents. To introduce this properly, let $\bar{\mathbf{\Lambda}}$ be the sublattice $\bar{\mathbf{\Lambda}} \subset \mathbf{\Lambda}$,

$$\bar{\mathbf{\Lambda}} = \{\vec{k} = (k, k, \dots, k) \in \mathbf{\Lambda} \mid k \in \mathbb{Z}^{b_2}\}. \quad (4.5)$$

The lattice $\mathbf{\Lambda}$ induces a quadratic form on $\bar{\mathbf{\Lambda}}$. Namely for $\vec{k} \in \mathbf{\Lambda}$, $\vec{D}(\vec{k}) = \sum_{j=1}^n D_j(k) = D(k)$. For $k \in \mathbb{Z}^{b_2}$, this is the quadratic form for the total charge P as desired. As a result, we have the group isomorphism $\mathbf{\Lambda}^*/\mathbf{\Lambda} = \bar{\mathbf{\Lambda}}^*/\bar{\mathbf{\Lambda}}$.

Let $\bar{\pi}$ be the orthogonal projection,

$$\bar{\pi} : \mathbf{\Lambda} \rightarrow \bar{\mathbf{\Lambda}} \otimes \mathbb{Q}. \quad (4.6)$$

For $\vec{k} \in \mathbf{\Lambda}$ and $\vec{m} \in \bar{\mathbf{\Lambda}}$, we have

$$\vec{D}(\bar{\pi}(\vec{k}), \vec{m}) = \vec{D}(\vec{k}, \bar{\pi}(\vec{m})) = \vec{D}(\vec{k}, \vec{m}) \in \mathbb{Z}. \quad (4.7)$$

Therefore, $\bar{\pi}$ gives an injection of $\mathbf{\Lambda}$ to the dual lattice $\bar{\mathbf{\Lambda}}^*$,

$$\bar{\pi} : \mathbf{\Lambda} \rightarrow \bar{\mathbf{\Lambda}}^*. \quad (4.8)$$

Moreover, if we extend $\bar{\pi}$ by linearity to Λ^* , a similar argument to (4.7) shows that $\bar{\pi} : \Lambda^* \rightarrow \bar{\Lambda}^*$. The projection $\bar{\pi}$ is explicitly given for $\vec{k} = (k_1, k_2, \dots, k_n) \in \Lambda$ by,

$$\bar{\pi}(\vec{k}) = (k, k, \dots, k) \in (\bar{\Lambda})^* \quad \text{with} \quad k = D^{-1} \sum_{j=1}^n D_j k_j. \quad (4.9)$$

We define furthermore the kernel $\underline{\Lambda} = \text{Ker}(\bar{\pi})$,

$$\underline{\Lambda} := \left\{ \vec{k} \in \Lambda \left| \sum_{j=1}^n D_j k_j = 0 \right. \right\}. \quad (4.10)$$

Elements in $\underline{\Lambda}$ have vanishing innerproduct with $\bar{\Lambda}$, such that they are indeed each others orthogonal complement in Λ .

Since $\bar{\pi}(\vec{k}) \in \bar{\Lambda}$ for $\vec{k} \in \Lambda$, we have that $\vec{k} - \bar{\pi}(\vec{k}) \in \underline{\Lambda}$. Thus $\bar{\Lambda} \oplus \underline{\Lambda}$ is in the kernel of the homomorphism $\bar{h} : \Lambda \rightarrow \bar{\Lambda}^*/\bar{\Lambda}$. We call

$$G = \Lambda / (\bar{\Lambda} \oplus \underline{\Lambda}), \quad (4.11)$$

the glue group for the decomposition of Λ , and the image of G under \bar{h} , $\bar{h}(G) \subset \bar{\Lambda}^*/\bar{\Lambda}$, the glue group for $\bar{\Lambda}$ [72]. The homomorphism \bar{h} gives an injection of G to the subgroup $\bar{h}(G) \subset \bar{\Lambda}^*/\bar{\Lambda}$. Therefore, the number of glue vectors, $N_g = |\Lambda / (\bar{\Lambda} \oplus \underline{\Lambda})|$ is a factor in $|\bar{\Lambda}^*/\bar{\Lambda}| = \det(\bar{D})$. In the special case that $\det(\bar{D})$ is prime, or more generally co-prime with $\det(D_j)$ for all j , $N_g = \det(\bar{D})$. By the same arguments, there is a projection $\underline{\pi} : \Lambda \rightarrow \underline{\Lambda}^*$, and a homomorphism $\underline{h} : G \rightarrow \underline{\Lambda}^*/\underline{\Lambda}$. Therefore, N_g is also a factor in $\det(\underline{D})$. This gives for the number of glue vectors N_g in general

$$N_g = \sqrt{\frac{\det(\bar{D}) \det(\underline{D})}{\prod_{j=1}^n \det(D_j)}}. \quad (4.12)$$

The order of the quotient group $(\underline{\Lambda}^*/\underline{\Lambda})/\underline{h}(G)$ is

$$N_q = \frac{\det(\underline{D})}{N_g}. \quad (4.13)$$

This will be useful for us in the following way. If we consider the class of vectors $\vec{k} \in \Lambda$ with fixed projection to $\bar{\Lambda}^*$, for example $\bar{\pi}(\vec{k}) = 0$, this fixes an element of the glue group G , and thus also of the images of G in $\bar{\Lambda}^*/\bar{\Lambda}$ and $\underline{\Lambda}^*/\underline{\Lambda}$. As a result, the number of possible conjugacy classes of $\underline{\pi}(\vec{k}) \in \underline{\Lambda}^*/\underline{\Lambda}$ of $\bar{\pi}(\vec{k})$ is N_q (4.13). This is also the case for a vector $\vec{x} \in \Lambda^*$ with fixed projection $\bar{\pi}(\vec{x}) \in \bar{\Lambda}^*$.

Similarly to Λ_μ^* (3.2), we introduce the notation $\underline{\Lambda}_\mu^*$,

$$\underline{\Lambda}_\mu^* := \left\{ \vec{Q} = (Q_1, Q_2, \dots, Q_n) \in \Lambda + (\mu_1, \mu_2, \dots, \mu_n) + \vec{P}/2 \left| \sum_{j=1}^n Q_j = \mu + P/2 \right. \right\}, \quad (4.14)$$

where the subscript $\mu \in \underline{\Lambda}^*/\underline{\Lambda}$, $\mu = \sum_{j=1}^n \mu_j \in \Lambda^*$ and $P = \sum_{j=1}^n P_j \in \Lambda$. We define the quadratic form Q^2 for $\underline{\Lambda}_\mu^*$,

$$\begin{aligned} Q^2 &= -Q^2 + \sum_{j=1}^3 (Q_j)_j^2 \\ &= -(\mu + P/2)^2 + \sum_{j=1}^3 (Q_j)_j^2. \end{aligned} \tag{4.15}$$

This form of the quadratic form appears naturally, when we determine the partition function h_μ^s (4.53) of scaling solutions in the next subsection. On the other hand, it should match with the inverse of the quadratic form for $\underline{\Lambda}$. In the cases considered below, we find that this is indeed the case.

Note that in components, P in (4.14) has a lower index and equals $P_a = d_{abc} P^b P^c = d_{abc} (\sum_j P_j)^b (\sum_j P_j)^c$. To understand constraint $\sum_j Q_j = \mu + P/2$ better, we change variables from $Q_j \in \Lambda_j^*$ to $k_j \in \Lambda_j$ using

$$\begin{aligned} Q_{1,a} &= \mu_{1,a} + D_{1ab} (k_1^b + P_2^b + P_1^b/2), \\ Q_{2,a} &= \mu_{2,a} + D_{2ab} (k_2^b + P_3^b + P_2^b/2), \\ &\vdots \\ Q_{n-1,a} &= \mu_{n-1,a} + D_{(n-1)ab} (k_n^b + P_n^b + P_{n-1}^b/2), \\ Q_{n,a} &= \mu_{n,a} + D_{nab} (k_n^b + P_1^b + P_n^b/2), \end{aligned} \tag{4.16}$$

with $k_j^b \in \mathbb{Z}^{b_2}$ for $j = 1, \dots, n$. The shifts of k_j^b by P_j^b are included such that the required identity for the k_j takes a compact form,

$$\sum_{j=1}^n D_{jab} k_j^b = 0, \quad k_j \in \mathbb{Z}^{b_2}, \tag{4.17}$$

which is indeed identical to the defining condition for the lattice $\underline{\Lambda}$ in (4.10). Solving this relation over the integers, $k_j \in \mathbb{Z}^{b_2}$, is in general a complicated problem depending on the D_j . Solving say for k_n , we have

$$k_n = -D_n^{-1} \sum_{j=1}^{n-1} D_j k_j. \tag{4.18}$$

Thus we find that if $D_n^{-1} D_j$, $j = 1, \dots, n-1$ are not integral matrices, not all $k_j \in \mathbb{Z}^{b_2}$ can correspond to bound state charges for these conjugacy classes. In some cases, this problem can be avoid by solving for another k_j instead of k_n , but not in general. We restrict to special cases in the following.

2- and 3-center bound states for $b_2 = 1$. For $b_2 = 1$, the D_j are simply positive numbers. For $n = 2$, the relation (4.17) becomes

$$D_1 k_1 + D_2 k_2 = 0. \tag{4.19}$$

The solutions with $k_{1,2} \in \mathbb{Z}$ are

$$k_1 = \frac{D_2}{\gcd(D_1, D_2)} m, \quad k_2 = -\frac{D_1}{\gcd(D_1, D_2)} m, \quad m \in \mathbb{Z}, \quad (4.20)$$

where \gcd stands for the greatest common divisor. Substituting (4.20) in the quadratic form $\sum_{j=1,2} D_j k_j^2$, gives for the quadratic form \underline{D} on $\underline{\Lambda}$,

$$\underline{D} = \frac{D_1 D_2 (D_1 + D_2)}{\gcd(D_1, D_2)^2}. \quad (4.21)$$

Thus the number of glue vectors (4.12) and order N_q (4.13) are in this case,

$$n = 2 : \quad N_g = \frac{D_1 + D_2}{\gcd(D_1, D_2)}, \quad N_q = \frac{D_1 D_2}{\gcd(D_1, D_2)}. \quad (4.22)$$

Moving on to $n = 3$, we need to find the integral solutions to

$$D_1 k_1 + D_2 k_2 + D_3 k_3 = 0. \quad (4.23)$$

To this end, we first consider $k_3 = 0$. Then, the solutions are obviously given by (4.20). Next for a fixed non-vanishing k_3 , there are only solutions if $\gcd(D_1, D_2)$ divides $D_3 k_3$ by Bézout's identity. As a result, if we choose $\gcd(D_1, D_2) / \gcd(D_1, D_2, D_3)$ for k_3 , Bézout's identity asserts that there is an integral solution (ℓ_1, ℓ_2) for (k_1, k_2) ,

$$D_1 \ell_1 + D_2 \ell_2 + D_3 \frac{\gcd(D_1, D_2)}{\gcd(D_1, D_2, D_3)} = 0. \quad (4.24)$$

Since this choice for k_3 has the smallest non-vanishing magnitude with integral solutions, the other choices for k_3 follow by multiplication by an integer m_1 . Including also the solutions (4.20), we find for the general solution,

$$\begin{aligned} k_1 &= \ell_1 m_1 + \frac{D_2}{\gcd(D_1, D_2)} m_2, \\ k_2 &= \ell_2 m_1 - \frac{D_1}{\gcd(D_1, D_2)} m_2, \\ k_3 &= \frac{\gcd(D_1, D_2)}{\gcd(D_1, D_2, D_3)} m_1. \end{aligned} \quad (4.25)$$

Substitution of (4.25) in $\sum_{j=1}^3 D_j k_j^2$, one finds for the 2-dimensional quadratic form \underline{D} for (m_1, m_2) ,

$$\underline{D} = \begin{pmatrix} D_1 \ell_1^2 + D_2 \ell_2^2 + D_3 \frac{\gcd(D_1, D_2)^2}{\gcd(D_1, D_2, D_3)^2} & \frac{D_1 D_2 (\ell_1 - \ell_2)}{\gcd(D_1, D_2)} \\ \frac{D_1 D_2 (\ell_1 - \ell_2)}{\gcd(D_1, D_2)} & \frac{D_1 D_2 (D_1 + D_2)}{\gcd(D_1, D_2)^2} \end{pmatrix}. \quad (4.26)$$

Using (4.24), we find for its determinant,

$$\det(\underline{D}) = \frac{D_1 D_2 D_3 (D_1 + D_2 + D_3)}{\gcd(D_1, D_2, D_3)^2}, \quad (4.27)$$

which is symmetric in the D_j . For the number of glue vectors N_g and order N_q , we now obtain,

$$n = 3 : \quad N_g = \frac{D_1 + D_2 + D_3}{\gcd(D_1, D_2, D_3)}, \quad N_q = \frac{D_1 D_2 D_3}{\gcd(D_1, D_2, D_3)}. \quad (4.28)$$

Together with the discussion below, this is very suggestive that the generalization to $n > 3$ is

$$\det(\underline{D}) = \frac{(\sum_{j=1}^n D_j) \prod_{j=1}^n D_j}{\gcd(D_1, D_2, \dots, D_n)^2}. \quad (4.29)$$

2- and 3-center bound states for $b_2 \geq 1$ with simplifications. We continue to discuss the general case $b_2 \geq 1$. Let us first consider $n = 2$. We make the assumption that $D_2^{-1}D_1$ is an integral matrix. The equation (4.17) can be solved by setting $k_2 = -D_2^{-1}D_1 k_1$. Substituting this in the quadratic form \vec{D} , gives us for the b_2 -dimensional quadratic form on $\underline{\Lambda}$

$$\underline{D} = D_1 + D_1 D_2^{-1} D_1. \quad (4.30)$$

This agrees with (4.21) upon specialization to $b_2 = 1$. We find then for N_g and N_q ,

$$n = 2 : \quad N_g = \det(D_2^{-1}D_1), \quad N_q = \det(D_1). \quad (4.31)$$

To deal with our main case, $n = 3$, we make two technical simplifications:

1. We assume that $D_3^{-1}D_1$ and $D_3^{-1}D_2$ are integral matrices, such that $k_3 \in \mathbb{Z}^{b_2}$ in (4.18) if k_1 and $k_2 \in \mathbb{Z}^2$.
2. We assume that $d_{abc} \in 2\mathbb{Z}$ for all a, b, c . Then $\Lambda^* + P/2 = \Lambda^*$ for any P such that shifts by P (and P_j) in (4.14), (4.15) and (4.16) are unnecessary.

These assumptions are satisfied in the examples in section 6.

If we substitute now (4.18) in \mathcal{Q}^2 , we arrive at

$$\begin{aligned} \mathcal{Q}^2 = & -\mu^2 + (\mu_1)_1^2 + (\mu_2)_2^2 + (\mu_3)_3^2 + 2(\mu_1 - D_3^{-1}D_1\mu_3) \cdot k_1 + 2(\mu_2 - D_3^{-1}D_2\mu_3) \cdot k_2 \\ & + (k_1, k_2) \underline{D} (k_1, k_2)^T, \end{aligned} \quad (4.32)$$

where \underline{D} is the quadratic form of the lattice $\underline{\Lambda}$,

$$\underline{D} = \begin{pmatrix} D_1 + D_1 D_3^{-1} D_1 & D_1 D_3^{-1} D_2 \\ D_2 D_3^{-1} D_1 & D_2 + D_2 D_3^{-1} D_2 \end{pmatrix}. \quad (4.33)$$

The determinant of \underline{D} is given by³

$$\begin{aligned} \det(\underline{D}) &= \det(D_1) \det(D_2) \det(D_3^{-1}) \det(D_1 + D_2 + D_3) \\ &= \det(D_1) \det(D_2) \det(D_3^{-1}D). \end{aligned} \quad (4.34)$$

We have for N_g and N_q in this case,

$$N_g = \det(D_3^{-1}D), \quad N_q = \det(D_1) \det(D_2). \quad (4.35)$$

³We use that for an invertible $n \times n$ matrix A and $n \times m$ matrices U and V , we have $\det(A + UV^T) = \det(A) \det(1_m + V^T A^{-1}U)$, https://en.wikipedia.org/wiki/Matrix_determinant_lemma.

These formulas are in agreement with (4.27) for $b_2 = 1$, and $\gcd(D_1, D_2, D_3) = D_3$. Moreover for generic b_2 , we can consider cases where we can solve (4.17) in terms of k_2 as well as k_3 , and we expect a symmetry in $D_2 \leftrightarrow D_3$. Indeed, then $D_2^{-1}D_1$ and $D_2^{-1}D_3$ are integral matrices too. The only way for $D_2^{-1}D_3$ and $D_3^{-1}D_2$ both to be integral is to satisfy $|\det(D_2)| = |\det(D_3)|$, in which case (4.34) reduces to $\det \underline{D} = \pm \det(D_1) \det(D_1 + D_2 + D_3)$, which is symmetric under the exchange of 2 and 3. Similar comments hold of course as well for k_1 .

Using general formula's for inverses of block matrices [74], we derive that the inverse of \underline{D} reads

$$\begin{aligned} \underline{D}^{-1} &= \begin{pmatrix} D^{-1}(D_2 + D_3)D_1^{-1} & -D^{-1} \\ -D^{-1} & D^{-1}(D_1 + D_3)D_2^{-1} \end{pmatrix} \\ &= \begin{pmatrix} D_1^{-1} - D^{-1} & -D^{-1} \\ -D^{-1} & D_2^{-1} - D^{-1} \end{pmatrix}, \end{aligned} \tag{4.36}$$

One can verify that the determinant of \underline{D}^{-1} is indeed the inverse of (4.34). Indeed, if we introduce the two components $\boldsymbol{\mu}_1$ and $\boldsymbol{\mu}_2$,

$$\begin{aligned} \boldsymbol{\mu}_1 &= \mu_1 - D_1 D_3^{-1} \mu_3, \\ \boldsymbol{\mu}_2 &= \mu_2 - D_2 D_3^{-1} \mu_3, \end{aligned} \tag{4.37}$$

the quadratic form \mathbf{Q}^2 (4.32) for $k_1 = k_2 = 0$, $\mathbf{Q}^2 = \boldsymbol{\mu}^2$, can be written as

$$\boldsymbol{\mu}^2 = -(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2)^2 + (\boldsymbol{\mu}_1)_1^2 + (\boldsymbol{\mu}_2)_2^2 = (\boldsymbol{\mu}_1, \boldsymbol{\mu}_2) \underline{D}^{-1} (\boldsymbol{\mu}_1, \boldsymbol{\mu}_2)^T. \tag{4.38}$$

For k_1, k_2 non-zero, we have $\mathbf{Q} = (\mathbf{Q}_1, \mathbf{Q}_2)$ with

$$\begin{aligned} \mathbf{Q}_1 &= \mu_1 - D_1 D_3^{-1} \mu_3 + (D_1 + D_1 D_3^{-1} D_1) k_1 + D_1 D_3^{-1} D_2 k_2, \\ \mathbf{Q}_2 &= \mu_2 - D_2 D_3^{-1} \mu_3 + D_2 D_3^{-1} D_1 k_1 + (D_2 + D_2 D_3^{-1} D_2) k_2, \end{aligned} \tag{4.39}$$

or equivalently

$$\begin{aligned} \mathbf{Q}_1 &= Q_1 - D_1 D_3^{-1} Q_3, \\ \mathbf{Q}_2 &= Q_2 - D_2 D_3^{-1} Q_3, \end{aligned} \tag{4.40}$$

which we can write more compactly as $\mathbf{Q} = \boldsymbol{\mu} + \underline{D} \mathbf{k}$ with $\mathbf{k} = (\mathbf{k}_1, \mathbf{k}_2)^T$. Thus an element $\boldsymbol{\mu} \in \underline{\Lambda}^*/\underline{\Lambda}$ is completely determined by

$$\boldsymbol{\mu} = \{(\mu_1, \mu_2, \mu_3, \mu) \mid \mu_j \in \Lambda_j^*, \quad \mu_1 + \mu_2 + \mu_3 = \mu \in \Lambda^*\}. \tag{4.41}$$

Generic number of constituents (with simplifications). The analysis for a generic number n of constituents (including $n = 2$) follows analogously. The quadratic form reads as in (4.15), but with 3 replaced by n in the summation, and with constraint $\sum_{j=1}^n Q_j = \mu + P/2$. Similarly to (4.17), the constraint can be expressed as

$$\sum_{j=1}^n D_{jab} k_j^b = 0. \tag{4.42}$$

With the *assumption* that $D_n^{-1}D_j$ is an integer matrix for all $j = 1, \dots, n$, this can be solved and the quadratic form becomes

$$\underline{D} = \text{diag}(D_1, \dots, D_{n-1}) + \begin{pmatrix} D_1 \\ \dots \\ D_{n-1} \end{pmatrix} (D_n^{-1}D_1, \dots, D_n^{-1}D_{n-1}), \quad (4.43)$$

with inverse

$$\underline{D}^{-1} = \text{diag}(D_1^{-1}, \dots, D_{n-1}^{-1}) - \begin{pmatrix} D^{-1} \\ \dots \\ D^{-1} \end{pmatrix} (1, \dots, 1), \quad (4.44)$$

where $D = \sum_{j=1}^n D_j$. Moreover, the determinant of \underline{D} is

$$\begin{aligned} \det(\underline{D}) &= \det(D_n^{-1}) \det(D) \prod_{j=1}^{n-1} \det(D_j) \\ &= \det(D_n^{-1}D) \prod_{j=1}^{n-1} \det(D_j), \end{aligned} \quad (4.45)$$

from which N_g and N_q are easily determined.

Characteristic vectors. We briefly discuss here a characteristic vector for the lattice $\underline{\Lambda}$ with $n = 3$, which is important for the theta series of the scaling solutions. A sign which frequently occurred in section 2 is $(-1)^{a+b+c}$. Such signs in a theta series are typically written in terms of a characteristic vector. We therefore express $a + b + c$ as

$$a + b + c = \mathbf{K} \cdot \mathbf{Q}, \quad (4.46)$$

with

$$\mathbf{K} = (P_3 - P_2, P_1 - P_3). \quad (4.47)$$

This is a characteristic vector of $\underline{\Lambda}$. Indeed, we have with $\vec{P} = (P_1, P_2, P_3)$,

$$\vec{k} \cdot \vec{P} + \vec{k}^2 \in 2\mathbb{Z}, \quad (4.48)$$

since P_j is a characteristic vector for Λ_j , $j = 1, 2, 3$. We can decompose with respect to the lattice decomposition $\underline{\Lambda} \oplus \overline{\Lambda}$,

$$\mathbf{k} \cdot \mathbf{K} + \mathbf{k}^2 + k \cdot P + k^2. \quad (4.49)$$

Since P is a characteristic vector for Λ , this shows that \mathbf{K} is a characteristic vector for $\underline{\Lambda}$.

However, if we express $a + b + c + P \cdot Q$ in terms of vectors in $\underline{\Lambda} = \sum_j \Lambda_j$, such that $a + b + c = \vec{K} \cdot \vec{Q} \pmod{2}$, then $\vec{K} = (P_3 - P_2, P_1 - P_3, P_2 - P_1)$. Then $(\vec{K} + \vec{P}) \cdot \vec{Q} = \mathbf{K} \cdot \mathbf{Q} + \vec{P} \cdot \vec{Q} = P \cdot Q \pmod{2}$. Moreover, \vec{P} is a characteristic vector of $\underline{\Lambda}$, and

$$\vec{P}^2 = P^3 + \mathbf{K}^3 \pmod{4}. \quad (4.50)$$

4.2 Partition functions

We consider black hole bound states with three cores. The j^{th} core carries electric and magnetic charges Q_j and P_j respectively. It is natural to work with a mixed ensemble with total magnetic charge P held fixed. For the present purpose, we shall fix the total electric D2-brane charge $Q = \sum_j Q_j \in \Lambda^* + P/2$ with $\mu \in \Lambda^*/\Lambda$ as well. We work at the attractor value of the moduli, corresponding to total charge vector (P, Q) . At this point apart from single core black holes, the only other black holes to survive are the scaling black holes. A natural question is — for a fixed total charge, how many scaling black holes are there and what is their contribution to the index?

In analogy to the attractor indices $\bar{\Omega}(\gamma, t_\lambda^\lambda)$, we define the generating function $h_\mu^T(\tau)$ of numerical total core invariants $\bar{\Omega}_{TC}$ introduced earlier in this section,

$$h_{P,\mu}^T(\tau) = \sum_{Q_0} \bar{\Omega}_{TC}(\gamma) q^{\bar{Q}_0}. \quad (4.51)$$

We can similarly define the partition function of single core indices $h_{P,\mu}^S(\tau)$, with Ω_T replaced by Ω_S . The $\Omega_T(\gamma)$ are determined from the refined ones (2.9) using the regularization (2.13). For the 3-core case, this gives (2.33). If P is irreducible, i.e. it can not be written as a sum of more than 1 positive magnetic charge, and the three partition functions agree.

Based on (2.7), we can express the attractor partition function $h_{P,\mu}$ in terms of the partition function $h_{P,\mu}^T$. We have schematically

$$h_{P,\mu}(\tau) = h_{P,\mu}^T(\tau) + \sum_{n>1} \sum_{\substack{\sum_{j=1}^n P_j = P \\ \sum_{j=1}^n Q_j = Q}} \frac{g_C(\{\gamma_j\}, \{c_j^\lambda\})}{|\text{Aut}(\{\gamma_j\})|} q^{\mu^2/2 - \sum_j (Q_j)^2/2} \prod_{j=1}^n h_{P_j,\mu_j}^T(\tau). \quad (4.52)$$

Recall that there are no 2-center/core scaling black holes, such that there is no contribution from $n = 2$ on the r.h.s.

We will proceed by considering the term in (4.52) with $n = 3$. Using the notation introduced in section 4.1, we can enumerate the number of three-core scaling black holes as

$$h_{\{P_j\},\mu}^{3T}(\tau) = \sum_{\substack{\mu_j \in \Lambda_j^*/\Lambda_j, j=1,2,3, \\ \mu_1 + \mu_2 + \mu_3 = \mu}} h_{P_1,\mu_1}^T(\tau) h_{P_2,\mu_2}^T(\tau) h_{P_3,\mu_3}^T(\tau) \Psi_\mu(\tau), \quad (4.53)$$

with μ as in (4.41), and where Ψ_μ is the indefinite theta series

$$\Psi_\mu(\tau) = \sum_{Q \in \Lambda_\mu^*} g_C(\{\gamma_j\}, \{c_j^\lambda\}) q^{-Q^2/2}. \quad (4.54)$$

With $y = e^{2\pi iz}$, we define the refined series as

$$\Psi_\mu(\tau, z) = (y - y^{-1})^2 \sum_{Q \in \Lambda_\mu^*} g_C(\{\gamma_j\}, \{c_j^\lambda\}; y) q^{-Q^2/2}, \quad (4.55)$$

with $g_C(\{\gamma_j\}, \{c_j^\lambda\})$ as in (2.42) and $g_C(\{\gamma_j\}, \{c_j^\lambda\}; y)$ as in (2.27). Note that Ψ_μ is symmetric as function of z ,

$$\Psi_\mu(\tau, -z) = \Psi_\mu(\tau, z). \quad (4.56)$$

The two functions are related by (2.13),

$$\Psi_{\mu}(\tau) = \left(\frac{1}{4\pi i} \frac{\partial}{\partial z} \right)^2 \Psi_{\mu}(\tau, z)|_{z=0}. \quad (4.57)$$

The kernel $g_C(\{\gamma_j\}, \{c_j^\lambda\})$ and therefore Ψ_{μ} is unchanged under a symplectic transformation (3.7), such that $h_{\{P_j\}, \mu}^T$ is invariant under spectral flow as required. The number of terms in the sum over μ_j in (4.53) is given by N_q (4.13).

To determine the modular properties of Ψ_{μ} , we consider first the generating function of f_C (2.45), which enumerates the number of scaling charge configurations for a given total charge. We define this function Φ_{μ} as the following theta series,

$$\Phi_{\mu}(\tau) = \sum_{Q \in \Lambda_{\mu}^*} f_C(\{\gamma_j\}, \{c_j^\lambda\}) q^{-Q^2/2}, \quad (4.58)$$

with $f_C(\{\gamma_j\}, \{c_j^\lambda\})$ as in (2.45).

The following subsections will demonstrate that $\Phi_{\mu}(\tau)$ is a convergent q -series, which can be completed to a function $\widehat{\Phi}_{\mu}$ which transform as a vector-valued modular form. The transformation properties under the S and T transformations are

$$\begin{aligned} \widehat{\Phi}_{\mu}(-1/\tau, -1/\bar{\tau}) &= -\frac{(-i\tau)^{b_2}}{\sqrt{|\Lambda^*/\Lambda|}} e^{\pi i K^2/2} \sum_{\nu \in \Lambda^*/\Lambda} e^{2\pi i \mu \cdot \nu} \widehat{\Phi}_{\nu}(\tau, \bar{\tau}), \\ \widehat{\Phi}_{\mu}(\tau + 1, \bar{\tau} + 1) &= e^{\pi i (\mu + K/2)^2} \widehat{\Phi}_{\mu}(\tau, \bar{\tau}). \end{aligned} \quad (4.59)$$

The partition function $\widehat{\Psi}_{\mu}(\tau)$ can be obtained by introducing a suitable elliptic variable in $\widehat{\Phi}_{\mu}$ and subsequently differentiating twice to this variable. As a result, the modular transformations of the completed function $\widehat{\Psi}_{\mu}$ equal those of the completed $\widehat{\Phi}_{\mu}$ except that the weight of $\widehat{\Psi}_{\mu}$ is increased by two compared to $\widehat{\Phi}_{\mu}$. The weight of $\widehat{\Psi}_{\mu}$ is thus $b_2 + 2$. The non-holomorphic terms are determined in this way in section 4.5, specifically eq. (4.116). The end result is that $\widehat{\Psi}_{\mu}$ transforms as

$$\begin{aligned} \widehat{\Psi}_{\mu}(-1/\tau, -1/\bar{\tau}) &= \frac{(-i\tau)^{b_2+2}}{\sqrt{|\Lambda^*/\Lambda|}} e^{\pi i K^2/2} \sum_{\nu \in \Lambda^*/\Lambda} e^{2\pi i \mu \cdot \nu} \widehat{\Psi}_{\nu}(\tau, \bar{\tau}), \\ \widehat{\Psi}_{\mu}(\tau + 1, \bar{\tau} + 1) &= e^{\pi i (\mu + K/2)^2} \widehat{\Psi}_{\mu}(\tau, \bar{\tau}). \end{aligned} \quad (4.60)$$

Therefore, the completion of $h_{\{P_j\}, \mu}^{3T}$ (4.53),

$$\widehat{h}_{\{P_j\}, \mu}^{3T}(\tau, \bar{\tau}) = \sum_{\substack{\mu_j \in \Lambda_j^*/\Lambda_j, \ j=1,2,3, \\ \mu_1 + \mu_2 + \mu_3 = \mu}} \widehat{h}_{P_1, \mu_1}^T(\tau) \widehat{h}_{P_2, \mu_2}^T(\tau) \widehat{h}_{P_3, \mu_3}^T(\tau) \widehat{\Psi}_{\mu}(\tau), \quad (4.61)$$

transforms as $\widehat{h}_{P, \mu}$ (3.21) as we aimed to show. We can furthermore combine $\widehat{h}_{P, \mu}$ with the theta series Θ_{μ} ,

$$\widehat{\mathcal{Z}}_P^{3T}(\tau, C, t) = \sum_{\sum_{j=1}^3 P_j = P} \sum_{\mu \in \Lambda^*/\Lambda} \widehat{h}_{\{P_j\}, \mu}^{3T}(\tau, \bar{\tau}) \Theta_{\mu}(\tau, \bar{\tau}, C, B). \quad (4.62)$$

We can then decompose the attractor partition $\widehat{\mathcal{Z}}_P^\lambda$ in terms of the multi-core partition functions $\widehat{\mathcal{Z}}_P^{nT}$,

$$\widehat{\mathcal{Z}}_P^\lambda(\tau, C, t) = \widehat{\mathcal{Z}}_P^T(\tau, C, t) + \widehat{\mathcal{Z}}_P^{3T}(\tau, C, t) + \dots \tag{4.63}$$

Since the partition functions transform the same way, this raises the question which terms are captured by the MSW conformal field theory. As mentioned in the introduction, it will also be interesting to deduce the non-holomorphic terms of $\widehat{\mathcal{Z}}_P^T$ using those determined for $\widehat{\mathcal{Z}}_P^{3T}$ in this paper, and those for $\widehat{\mathcal{Z}}_P^\lambda$ in [29].

4.3 Convergence

A crucial aspect of Φ_μ (and Ψ_μ) is whether the sum on the r.h.s. of (4.58) (and (4.54)) is convergent. If Q^2 would be negative definite, convergence of these series would be guaranteed. However this is not the case since the electric charge lattice has signature $(2, 2b_2 - 2)$, i.e. has 2 positive directions.

To prove the convergence, we first introduce a theta series $\Theta_\mu[\mathcal{K}](\tau)$ with kernel \mathcal{K} for a generic indefinite theta lattice L and $\mu \in L^*$,

$$\Theta_\mu[\mathcal{K}](\tau; L) = \sum_{x \in L + \mu} \mathcal{K}(x) q^{-B(x)/2}, \tag{4.64}$$

with integral quadratic form B . If L is negative definite, we also use

$$\theta_\mu(\tau; L) = \Theta_\mu[1](\tau; L). \tag{4.65}$$

For an indefinite lattice the kernel $\mathcal{K}(x) = \mathcal{K}(x, \mathcal{V})$ depends on a collection

$$\mathcal{V} = \{V_1, V_2, \dots, V_N\},$$

of positive vectors. For signature $(2, 2b_2 - 2)$, $\mathcal{K}(x, \mathcal{V})$ can be expressed as [28, 36, 75]

$$\mathcal{K}(x, \mathcal{V}) = \frac{1}{4} \left(w(\mathcal{V}) + \sum_{j=1}^N \text{sgn}(B(x, V_j)) \text{sgn}(B(x, V_{j+1})) \right), \tag{4.66}$$

where for any strictly positive vector $v \in L$, $v^2 > 0$,

$$w(\mathcal{V}) = - \sum_{j=1}^N \text{sgn}(B(v, V_j)) \text{sgn}(B(v, V_{j+1})), \tag{4.67}$$

which is independent of the choice of positive vector v [75]. There are various sufficient conditions for convergence put forward in the literature [28, 36, 38, 39, 75]. We will consider here the following N -gon conditions put forward in [28, 36, 75], which read

$$\begin{aligned} B(V_j, V_j) &> 0, \\ B(V_j, V_j) B(V_{j+1}, V_{j+1}) - B(V_j, V_{j+1})^2 &> 0, \\ B(V_j, V_j) B(V_{j-1}, V_{j+1}) - B(V_j, V_{j-1}) B(V_j, V_{j+1}) &< 0. \end{aligned} \tag{4.68}$$

Now let us return to the sum (4.58) at hand. It comprises of six individual sums, which are each of the form

$$\sum_{\substack{Q_i \in \mu_i + \Lambda_i + P_i/2 \\ Q_1 + Q_2 + Q_3 = \mu + P/2}} F_\ell(a, b, c) (-1)^{a+b+c} q^{Q^2/2 - \sum_i (Q_i)^2/2} = \sum_{Q \in \underline{\Lambda}_\mu^*} F_\ell(a, b, c) (-1)^{a+b+c} q^{-Q^2/2}, \quad (4.69)$$

with $\ell = 1, \dots, 6$.

The simplification $F_{\text{total}}(a, b, c)$ for $\sum_{j=1}^6 F_j(a, b, c)$ put forward in (2.44) is precisely of the form (4.66) with $N = 3$. To present the vectors, we define

$$\begin{aligned} C_a &= (-P_2, P_1, 0), \\ C_b &= (0, -P_3, P_2), \\ C_c &= (P_3, 0, -P_1), \end{aligned} \quad (4.70)$$

such that $C_a \cdot Q = a$, $C_b \cdot Q = b$ and $C_c \cdot Q = c$, with a, b and c as in (4.2). The V_j are then identified with $C_j \in \underline{\Lambda}$ with the C_j given by,

$$\begin{aligned} C_1 &= C_a + C_b - C_c = (-P_2 - P_3, P_1 - P_3, P_1 + P_2), \\ C_2 &= C_a - C_b + C_c = (-P_2 + P_3, P_1 + P_3, -P_1 - P_2), \\ C_3 &= -C_a + C_b + C_c = (P_2 + P_3, -P_1 - P_3, -P_1 + P_2). \end{aligned} \quad (4.71)$$

If we assume that P_j is an ample divisor for each $j \in 1, 2, 3$, triple intersections $P_i P_j P_k > 0$ for all $i, j, k \in \{1, 2, 3\}$. The conditions for convergence (4.68) are then satisfied.

It is also useful to consider the convergence for the kernel due to a single permutation separately, since the different permutations are weighted by a different factor in g_C . The vectors V_j for the kernel $F^*(123)$ (2.40) can be chosen as $C_j^{(123)}$,

$$\begin{aligned} C_1^{(123)} &= C_a - C_c = (C_1 - C_2)/2, \\ C_2^{(123)} &= C_b - C_c = (C_1 - C_3)/2, \\ C_3^{(123)} &= C_c - C_a - C_b = -C_1. \end{aligned} \quad (4.72)$$

Again one may verify that with the assumption $P_i P_j P_k > 0$ for all $i, j, k \in \{1, 2, 3\}$, these vectors satisfy the conditions for convergence (4.68).

4.4 Modular completion of Φ_μ

Having discussed the convergence of Φ_μ , we proceed in this section to discuss its modularity. Since Φ_μ is a sum over a subset of an indefinite lattice, the function is not modular in the classical sense. Our task is to determine a modular completion $\widehat{\Phi}_\mu$, which differs from Φ_μ by subleading non-holomorphic terms, and which does transform as a modular form. Essentially, products of sgn-functions are replaced by a generalized error function [36]. Following this approach, we will demonstrate that the difference between $\widehat{\Phi}_\mu$ and Φ_μ , is given by iterated integrals of modular forms.

Such non-holomorphic contributions have appeared in similar contexts. In specific cases, the non-holomorphic contributions are derived from different physical points of view, for example the continuum of multi-particle states in \mathbb{R}^4 [58], or the quantum field theory on the world volume of the D-branes [44, 47, 48], or the perspective of D3-instantons in the hypermultiplet moduli space [18].

To this end let us consider $F_{\text{total}}(a, b, c)$ in (2.44). Under modular completion, one adds certain extra terms to $\text{sgn}(V_1, x) \text{sgn}(V_2, x)$, therefore replacing $\text{sgn}(V_1, x) \text{sgn}(V_2, x)$ with the double error function $E_2(\alpha, u_1, u_2)$:

$$\text{sgn}(V_1, x) \text{sgn}(V_2, x) \rightarrow E_2(\alpha, \sqrt{2\tau_2} \mathbf{u}), \tag{4.73}$$

with [36]

$$E_2(\alpha; \mathbf{u}) = \int_{\mathbb{R}^2} e^{-\pi(u_1-u'_1)^2 - \pi(u_2-u'_2)^2} \text{sgn}(u'_2) \text{sgn}(u'_1 + \alpha u'_2) du'_1 du'_2, \tag{4.74}$$

whose arguments are given in terms of V_1, V_2 and x by

$$\alpha = \alpha(V_1, V_2) = \frac{(V_1, V_2)}{\sqrt{V_1^2 V_2^2 - (V_1, V_2)^2}}, \tag{4.75}$$

$$\mathbf{u} = \mathbf{u}(V_1, V_2; x) = (u_1(V_1, V_2; x), u_2(V_1, V_2; x)),$$

with

$$u_1(V_1, V_2; x) = \frac{(V_{1\perp 2}, x)}{\sqrt{(V_{1\perp 2}, V_{1\perp 2})}}, \tag{4.76}$$

$$u_2(V_1, V_2; x) = \frac{(V_2, x)}{\sqrt{(V_2, V_2)}} \tag{4.77}$$

and $V_{1\perp 2}$ the component of V_1 orthogonal to V_2 ,

$$V_{1\perp 2} = V_1 - \frac{(V_1, V_2)}{(V_2, V_2)} V_2. \tag{4.78}$$

To stress the dependance of E_2 on the vectors V_1, V_2 and x , we will also use E_2 with alternative arguments,

$$E_2(\alpha, \mathbf{u}) \equiv E_2(V_1, V_2; x), \tag{4.79}$$

with the identifications as in (4.75).

In addition to (4.74), another (equivalent) expression for E_2 is in terms of Eichler integrals. To this end, we first define the Eichler integrals M_1 ,

$$M_1(u) = \begin{cases} \frac{i u}{\sqrt{2\tau_2}} q^{\frac{u^2}{4\tau_2}} \int_{-\bar{\tau}}^{i\infty} \frac{e^{-\frac{i\pi u^2 w}{2\tau_2}}}{\sqrt{-i(w+\tau)}} dw, & u \neq 0, \\ 0, & u = 0. \end{cases} \tag{4.80}$$

and the (iterated) Eichler integral M_2 and m_2 , for $u_1 \neq 0$, and $u_2 - \alpha u_1 \neq 0$,

$$\begin{aligned}
 m_2(u_1, u_2) &= \begin{cases} \frac{u_1 u_2}{2\tau_2} q^{\frac{u_1^2}{4\tau_2} + \frac{u_2^2}{4\tau_2}} \int_{-\bar{\tau}}^{i\infty} dw_2 \int_{w_2}^{i\infty} dw_1 \frac{e^{\frac{\pi i u_1^2 w_1}{2\tau_2} + \frac{\pi i u_2^2 w_2}{2\tau_2}}}{\sqrt{-(w_1 + \tau)(w_2 + \tau)}}, & u_1 \neq 0 \\ 0, & u_1 = 0. \end{cases} \\
 M_2(\alpha; u_1, u_2) &= \begin{cases} -m_2(u_1, u_2) - m_2\left(\frac{u_2 - \alpha u_1}{\sqrt{1 + \alpha^2}}, \frac{u_1 + \alpha u_2}{\sqrt{1 + \alpha^2}}\right) & u_1 \neq 0, u_2 - \alpha u_1 \neq 0, \\ -m_2\left(\frac{u_2 - \alpha u_1}{\sqrt{1 + \alpha^2}}, \frac{u_1 + \alpha u_2}{\sqrt{1 + \alpha^2}}\right) & u_1 = 0, u_2 \neq 0, \\ -m_2(u_1, u_2) & u_1 \neq 0, u_2 - \alpha u_1 = 0, \\ \frac{2}{\pi} \arctan \alpha & u_1 = u_2 = 0. \end{cases} \quad (4.81)
 \end{aligned}$$

With $\mathbf{u} = (u_1, u_2)$ as before, the double error function E_2 is then defined as a linear combination of M_1 and M_2 [36, 57]

$$\begin{aligned}
 E_2(\alpha; \mathbf{u}) &= \text{sgn}(u_2) \text{sgn}(u_1 + \alpha u_2) + \text{sgn}(u_1) M_1(u_2) \\
 &\quad + \text{sgn}(u_2 - \alpha u_1) M_1\left(\frac{u_1 + \alpha u_2}{\sqrt{1 + \alpha^2}}\right) + M_2(\alpha; u_1, u_2). \quad (4.82)
 \end{aligned}$$

See [36] for other representations of E_2 .

Thus E_2 consists of the original $\text{sgn}(V_1, x) \text{sgn}(V_2, x)$ plus four more terms. Noting that

$$\begin{aligned}
 \frac{u_2(V_2, x) - \alpha u_1(V_1, V_2, x)}{\sqrt{1 + \alpha^2}} &= u_1(V_2, V_1, x), \\
 \frac{u_1(V_1, V_2, x) + \alpha u_2(V_2, x)}{\sqrt{1 + \alpha^2}} &= u_2(V_1, x), \quad (4.83)
 \end{aligned}$$

we can write

$$\begin{aligned}
 E_2(\alpha; \mathbf{u}) &= \text{sgn}(V_1.x) \text{sgn}(V_2.x) + \text{sgn}(u_1(V_1, V_2, x)) M_1(u_2(V_2, x)) \\
 &\quad + \text{sgn}(u_1(V_2, V_1, x)) M_1(u_2(V_1, x)) \\
 &\quad - m_2(u_1(V_1, V_2, x), u_2(V_2, x)) - m_2(u_1(V_2, V_1, x), u_2(V_1, x)). \quad (4.84)
 \end{aligned}$$

E_2 satisfies an identity similar to (2.39). This reads

$$E_2(V_1, V_1 + V_2; x) + E_2(V_2, V_1 + V_2; x) - E_2(V_1, V_2; x) = 1 \quad (4.85)$$

and is valid for any choice of the arguments such that the corresponding α , u_1 and u_2 's are in \mathbb{R} .

The sum at hand (2.44) has the form

$$\text{sgn}(C_1.x) \text{sgn}(C_2.x) + \text{sgn}(C_2.x) \text{sgn}(C_3.x) + \text{sgn}(C_3.x) \text{sgn}(C_1.x), \quad (4.86)$$

with C_1, C_2, C_3 given in (4.71). The modular completion $\widehat{\Phi}_\mu$ of Φ_μ (4.58) follows by the adding to the coefficient f_C (2.45) the following terms

$$\frac{1}{4} \sum_{\ell=1,2,3} \left[E_2(C_\ell, C_{\ell+1}; \sqrt{2\tau_2}x) - \text{sgn}(C_\ell.x) \text{sgn}(C_{\ell+1}.x) - A_\ell \delta_{(C_\ell.x)} \delta_{(C_{\ell+1}.x)} \right]. \quad (4.87)$$

This essentially amounts to replacing f_C by a linear combination of E_2 's. Since the latter satisfies Vignéras equation, modular transformation properties are ensured [55].

Our first aim is to determine the value of A_ℓ such that the completion is subleading, i.e. it vanishes in the limit that $\text{Im}(\tau) = \tau_2 \rightarrow \infty$. This follows from realizing that the difference $E_2(C_\ell, C_{\ell+1}; \sqrt{2\tau_2}x) - \text{sgn}(C_\ell.x) \text{sgn}(C_{\ell+1}.x)$ vanishes in this limit except if $(C_\ell.x) = (C_{\ell+1}.x) = 0$, when it equals $\frac{2}{\pi} \arctan(\alpha_\ell)$, $\alpha_\ell = \alpha(C_\ell, C_{\ell+1})$ (4.75). Requiring that (4.87) vanish, we thus arrive at

$$A_\ell = \frac{2}{\pi} \arctan(\alpha_\ell). \quad (4.88)$$

Surprisingly this implies that A_ℓ can be irrational, as we will see in the explicit case studies in section 6. This of course obstructs an interpretation as a ‘‘counting’’ function for the coefficients to which A_ℓ contribute. On the other hand, since the function Φ_μ is not a proper physical partition function summing over a Hilbert space, we are not very concerned about this.

To proceed with determining the completion, we rearrange the terms and write (4.87) as

$$\begin{aligned} & \frac{1}{4} \sum_{\ell=1,2,3} [\text{sgn}(u_1(C_{\ell+1}, C_\ell, x)) + \text{sgn}(u_1(C_{\ell-1}, C_\ell, x))] M_1(\sqrt{2\tau_2} u_2(C_\ell, x)) \\ & - m_2(\sqrt{2\tau_2} u_1(C_{\ell+1}, C_\ell, x), \sqrt{2\tau_2} u_2(C_\ell, x)) - m_2(\sqrt{2\tau_2} u_1(C_{\ell-1}, C_\ell, x), \sqrt{2\tau_2} u_2(C_\ell, x)). \end{aligned} \quad (4.89)$$

We determine now the A_ℓ by requiring that this expression vanishes in the limit $y \rightarrow \infty$.

Using these expressions, we thus naturally separate the holomorphic part Φ_μ from the completion $\widehat{\Phi}_\mu$,

$$\widehat{\Phi}_\mu(\tau, \bar{\tau}) = \Phi_\mu(\tau) + R_\mu^\Phi(\tau, \bar{\tau}), \quad (4.90)$$

with the non-holomorphic completion R_μ^Φ defined by

$$\begin{aligned} R_\mu^\Phi(\tau, \bar{\tau}) &= \sum_{Q \in \Delta_\mu^*} \sum_{\ell=1,2,3} \\ & \left[[\text{sgn}(u_1(C_{\ell+1}, C_\ell, x)) + \text{sgn}(u_1(C_{\ell-1}, C_\ell, x))] M_1(\sqrt{2\tau_2} u_2(C_\ell, x)) \right. \\ & - m_2(\sqrt{2\tau_2} u_1(C_{\ell+1}, C_\ell, x), \sqrt{2\tau_2} u_2(C_\ell, x)) \\ & \left. - m_2(\sqrt{2\tau_2} u_1(C_{\ell-1}, C_\ell, x), \sqrt{2\tau_2} u_2(C_\ell, x)) \right] \\ & \times (-1)^{\mathbf{K} \cdot \mathbf{Q}} q^{-Q^2/2}. \end{aligned} \quad (4.91)$$

Our next aim is to write the non-holomorphic part as an (iterated) integral over modular forms. This makes the modular properties of the holomorphic q -series manifest, since the modular properties of integrals of modular forms are readily determined. Moreover, it is straightforward to determine the non-holomorphic anomaly.

To determine this form of the non-holomorphic part, we write R_μ^Φ as

$$R_\mu^\Phi(\tau, \bar{\tau}) = \sum_{\ell=1,2,3} \left[R_{\mu,1,\ell}(\tau, \bar{\tau}) + R_{\mu,2,(\ell-1,\ell)}(\tau, \bar{\tau}) + R_{\mu,2,(\ell+1,\ell)}(\tau, \bar{\tau}) \right] \quad (4.92)$$

term	form	dual lattice	signature
\mathbf{Q}^2	\underline{D}	$\underline{\Lambda}_\mu^*$	$(2, 2b_2 - 2)$
$u_{2,\ell}^2$	$\underline{D}_\ell = C_\ell ^2$	$(L_\ell)_\mu^* \subset \underline{\Lambda}_\mu^*$	$(1,0)$
$\mathbf{Q}^2 - u_{2,\ell}^2$	$\underline{D}_{\perp\ell}$	$(L_\ell^\perp)_\mu^* \subset \underline{\Lambda}_\mu^*$	$(1, 2b_2 - 2)$
$u_{1,(k,l)}^2$	$\underline{D}_{(k,l)} = C_{k\perp\ell} ^2$	$(L_{k\ell})_\mu^* \subset (L_\ell^\perp)_\mu^*$	$(1,0)$
$\mathbf{Q}^2 - u_{1,(k,l)}^2 - u_{2,\ell}^2$	$\underline{D}_{(k\ell)}^\perp$	$(L_{k\ell}^\perp)_\mu^* \subset (L_\ell^\perp)_\mu^*$	$(0, 2b_2 - 2)$

Table 1. Lattices and quadratic forms associated to the splitting in equation (4.94).

and $R_{\mu,1,\ell}, R_{\mu,2,(k,\ell)}$ are defined as

$$\begin{aligned}
 R_{\mu,1,\ell}(\tau, \bar{\tau}) &= \sum_{\mathbf{Q} \in \underline{\Lambda}_\mu^*} \left[\text{sgn}(u_{1,(\ell+1,\ell)} + \text{sgn}(u_{1,(\ell-1,\ell)}) \right] M_1(\sqrt{2y}u_{2,\ell})(-1)^{\mathbf{K} \cdot \mathbf{Q}} q^{-\mathbf{Q}^2/2}, \\
 R_{\mu,2,(k,\ell)}(\tau, \bar{\tau}) &= - \sum_{\mathbf{Q} \in \underline{\Lambda}_\mu^*} m_2(\sqrt{2\tau_2}u_{1,(k,l)}, \sqrt{2\tau_2}u_{2,\ell})(-1)^{\mathbf{K} \cdot \mathbf{Q}} q^{-\mathbf{Q}^2/2},
 \end{aligned} \tag{4.93}$$

where we have used the abbreviations $u_{1,(k,\ell)} = u_1(C_k, C_\ell, x)$, $u_{2,\ell} = u_2(C_\ell, x)$.

To evaluate the sums (4.93), it is useful to split \mathbf{Q}^2 as follows

$$\mathbf{Q}^2 = u_{1,(k,\ell)}^2 + u_{2,\ell}^2 + (\mathbf{Q}^2 - u_{1,(k,\ell)}^2 - u_{2,\ell}^2), \tag{4.94}$$

$u_{1,(k,\ell)}^2$, $u_{2,\ell}^2$ and $(\mathbf{Q}^2 - u_{1,(k,\ell)}^2 - u_{2,\ell}^2)$ are naturally associated with quadratic forms and lattices as described in table 1.

We let L_ℓ be the 1-dimensional lattice spanned by C_ℓ . The quadratic form is a number in this case, $D_\ell = |C_\ell|^2$. We denote the dual lattice with quadratic form $|C_\ell|^{-2}$ by $(L_\ell)^*$. The projection of μ to $(L_\ell)^*$ is $\mu_\ell = (\mu \cdot C_\ell) C_\ell \in \mathbb{Z}^{2b_2}$. To express a generic vector in $\underline{\Lambda}_\mu^*$ as an element of $(L_\ell)^* \oplus (L_\ell^\perp)^*$, we introduce glue vectors ρ . We denote by $(L_\ell)_\mu^*$ the set of vectors $\mu_\ell \bmod L_\ell \in (L_\ell)^*$, and by $(L_\ell)_{\mu+\rho}^*$ the set of vectors $\mu_\ell + \rho \bmod L_\ell \in (L_\ell)^*$. We introduce similar notation for L_ℓ^\perp . Using this notation, the direct sum $(L_\ell)_\mu^* \oplus (L_\ell^\perp)_\mu^*$ is a subset of the lattice $\underline{\Lambda}_\mu^*$, $(L_\ell)_\mu^* \oplus (L_\ell^\perp)_\mu^* \subset \underline{\Lambda}_\mu^*$.

A generic element \mathbf{k} of $\underline{\Lambda}_\mu^*$ can be written as a sum, $\mathbf{k} = \mathbf{l}_\ell + \mathbf{l}_\ell^\perp \in (L_\ell)_{\mu,\rho}^* \oplus (L_\ell^\perp)_{\mu,\rho}^*$ for some ρ , where the projection \mathbf{l}_ℓ to $(L_\ell^\perp)^*$ vanishes, and similarly for the projection of \mathbf{l}_ℓ^\perp to $(L_\ell)^*$. We have a similar decomposition of vectors in $(L_\ell)^\perp$ with respect to the decomposition $(L_{k\ell})^* \oplus (L_{k\ell}^\perp)^*$. The representatives of minimal length in $(L_\ell)_\mu^* \oplus (L_\ell^\perp)_\mu^*$ appearing in such splits are called ‘‘glue vectors’’. We use glue vectors ρ for the splitting $(L_\ell)_{\mu,\rho}^* \oplus (L_\ell^\perp)_{\mu,\rho}^*$ and ν for the splitting $(L_{k\ell})^* \oplus (L_{k\ell}^\perp)^*$. We then have,

$$\begin{aligned}
 \underline{\Lambda}_\mu^* &= \sum_{\rho} (L_\ell)_{\mu+\rho}^* \oplus (L_\ell^\perp)_{\mu+\rho}^*, \\
 (L_\ell^\perp)_\rho^* &= \sum_{\nu} (L_k)_{\rho+\nu}^* \oplus (L_{k\ell}^\perp)_{\rho+\nu}^*.
 \end{aligned} \tag{4.95}$$

Now let us evaluate the sum $R_{\mu,1}$ and $R_{\mu,2}$ (4.93). The embedding of L_ℓ in $\underline{\Lambda}_\mu \otimes \mathbb{Q}$ is spanned by $k C_\ell$ with $k \in \mathbb{Z} + \rho$ where $\rho \in \mathbb{Q}$ is (the projection of) the glue vector. Then,

$u_{2,\ell} = |C_\ell|k$, such that $R_{\mu,1,\ell}$ reads

$$R_{\mu,1,\ell}(\tau, \bar{\tau}) = \sum_{\rho} \sum_{\substack{l_\ell^\perp \in (L_\ell^\perp)^* \\ k \in \mathbb{Z} + \mu + \rho}} (\text{sgn}(u_{1,(\ell-1,\ell)}) + \text{sgn}(u_{1,(\ell+1,\ell)})) \times (-1)^{\mathbf{K} \cdot l_\ell^\perp + \mathbf{K} \cdot C_\ell k} M_1(\sqrt{2\tau_2} |C_\ell| k) q^{-(l_\ell^\perp)^2/2 - C_\ell^2 k^2/2}. \quad (4.96)$$

To write this more compactly, we define for a generic lattice L of signature $(1, \dim(L) - 1)$ and characteristic vector K ,

$$\Theta_\alpha(\tau; L, \{V, V'\}) := \sum_{x \in L + \alpha} (\text{sgn}(V, x) - \text{sgn}(V', x)) (-1)^{K \cdot x} q^{-x^2/2}. \quad (4.97)$$

Convergence of Θ_α requires [34]

$$V^2 > 0, (V')^2 > 0, (V, V') > 0. \quad (4.98)$$

Moreover, we define the unary theta series Υ_α ,

$$\Upsilon_\alpha(\tau, M, N) = \sum_{x \in \mathbb{Z} + \alpha} x (-1)^{Nx} e^{\pi i \tau M x^2}. \quad (4.99)$$

For $\sigma \in \mathbb{H}$, we introduce the period integral

$$R_\alpha(\tau, \sigma; M, N) := i \sum_{x \in \mathbb{Z} + \alpha} (-1)^{Nx} x \int_{-\sigma}^{i\infty} dw \frac{e^{\pi i w M x^2}}{\sqrt{-i(w + \tau)}} = i \int_{-\sigma}^{i\infty} dw \frac{\Upsilon_\alpha(w, M, N)}{\sqrt{-i(w + \tau)}}. \quad (4.100)$$

The non-holomorphic modular completion $\widehat{\Theta}_\alpha$ of Θ_α is expressed in terms of R_α as

$$\begin{aligned} \widehat{\Theta}_\alpha(\tau, \bar{\tau}; L, \{V, V'\}) &= \Theta_\alpha(\tau; L, V, V') \\ &+ \sum_{\nu} \theta_{\alpha+\nu}(\tau; L_V^\perp) R_{\alpha+\nu}(\tau, \bar{\tau}; V^2, K_L \cdot V) \\ &- \sum_{\nu'} \theta_{\alpha+\nu'}(\tau; L_{V'}^\perp) R_{\alpha+\nu'}(\tau, \bar{\tau}; (V')^2, K_L \cdot V'), \end{aligned} \quad (4.101)$$

with θ_α as in (4.65).

Returning to $R_{\mu,1,\ell}$, we can now write this as

$$R_{\mu,1,\ell}(\tau, \bar{\tau}) = \sum_{\rho} \Theta_{\mu+\rho}(\tau, L_\ell^\perp, \{C_{\ell-1}, C_{\ell+1}\}) R_{\mu+\rho}(\tau, \bar{\tau}; C_\ell^2, \mathbf{K} \cdot C_\ell). \quad (4.102)$$

The relations for convergence of $\Theta_{\mu+\rho}$ are indeed satisfied for the vectors given in (4.71). We will continue to demonstrate that these are captured by the term $R_{\mu,2}$ in (4.92). We have

$$\begin{aligned} R_{\mu,2,(k,\ell)}(\tau, \bar{\tau}) &= - \sum_{\mathbf{Q} \in \underline{\Lambda}_\mu^*} m_2(\sqrt{2\tau_2} u_1(C_k, C_\ell, \mathbf{Q}), \sqrt{2\tau_2} u_2(C_\ell, \mathbf{Q})) (-1)^{\mathbf{K} \cdot \mathbf{Q}} q^{-\mathbf{Q}^2/2} \\ &= - \sum_{\mathbf{Q} \in \underline{\Lambda}_\mu^*} u_1(C_k, C_\ell, \mathbf{Q}) u_2(C_\ell, \mathbf{Q}) (-1)^{\mathbf{K} \cdot \mathbf{Q}} q^{(u_1(C_k, C_\ell, \mathbf{Q})^2 + u_2(C_\ell, \mathbf{Q})^2 - \mathbf{Q}^2)/2} \\ &\quad \times \int_{-\bar{\tau}}^{i\infty} dw_2 \int_{w_2}^{i\infty} dw_1 \frac{e^{\pi i (w_1 u_1(C_k, C_\ell, \mathbf{Q})^2 + w_2 u_2(C_\ell, \mathbf{Q})^2)}}{\sqrt{-(w_1 + \tau)(w_2 + \tau)}}. \end{aligned} \quad (4.103)$$

Using the splits of the lattices and Υ_α (4.99), we can express this as

$$R_{\mu,2,(k,\ell)}(\tau, \bar{\tau}) = - \sum_{\rho} \theta_{\mu+\rho+\nu}(\tau; L_{k\ell}^\perp) \times \int_{-\bar{\tau}}^{i\infty} dw_2 \int_{w_2}^{i\infty} dw_1 \frac{\Upsilon_{\mu+\rho}(w_2; C_\ell^2, \mathbf{K} \cdot C_\ell) \Upsilon_{\mu+\rho+\nu}(w_1; C_{k\perp\ell}^2, \mathbf{K} \cdot C_{k\perp\ell})}{\sqrt{-(w_1 + \tau)(w_2 + \tau)}}. \quad (4.104)$$

Next, we aim to combine the sums in (4.91). We recognize the two Υ 's in the integrand in (4.104). The first, $\Upsilon_{\mu+\rho}$, appears in the integrand of $R_{\mu+\rho}$ on the r.h.s. of (4.102), whereas the second $\Upsilon_{\mu+\rho+\nu}$ matches with one of the non-holomorphic terms on the r.h.s. of (4.101). More precisely and concisely, we arrive at

$$R_\mu^\Phi(\tau, \bar{\tau}) = \sum_{\ell=1,2,3} i \int_{-\bar{\tau}}^{i\infty} dw \frac{\sum_{\rho} \hat{\Theta}_{\mu+\rho}(\tau, -w; L_\ell^\perp, \{C_{\ell-1}C_{\ell+1}\}) \Upsilon_{\mu+\rho}(w; C_\ell^2, \mathbf{K} \cdot C_\ell)}{\sqrt{-i(w + \bar{\tau})}}. \quad (4.105)$$

We thus have succeeded to express R_μ^Φ in (4.90) as an iterated integral of modular forms.

Note that for $\ell = 1$, there is a symmetry exchanging $P_1 \leftrightarrow P_2$, and there are similar symmetries for $\ell = 2$ and $\ell = 3$. The form (4.105) makes the determination of the anti-holomorphic derivative of $\hat{\Phi}_\mu$ immediate. We have,

$$\frac{\partial \hat{\Phi}_\mu(\tau, \bar{\tau})}{\partial \bar{\tau}} = \sum_{\ell=1,2,3} \sum_{\rho} \frac{\hat{\Theta}_{\mu+\rho}(\tau, \bar{\tau}; L_\ell^\perp, \{C_{\ell-1}, C_{\ell+1}\}) \Upsilon_{\mu+\rho}(-\bar{\tau}; C_\ell^2, \mathbf{K} \cdot C_\ell)}{\sqrt{2\tau_2}}. \quad (4.106)$$

4.5 Modular completion of Ψ_μ

We will treat in this subsection the completion of Ψ_μ . Eq. (4.57) related the holomorphic q -series $\Psi_\mu(\tau)$ to that the second derivative of the function $\Psi_\mu(\tau, z)$. To determine the non-holomorphic completion $\hat{\Psi}_\mu(\tau, \bar{\tau})$, we consider the non-holomorphic differential operator

$$- \frac{1}{4\pi^2} \left(\frac{\partial^2}{\partial z^2} + \frac{2\pi m}{\tau_2} \right), \quad (4.107)$$

on the completion $\hat{\Psi}_\mu(\tau, \bar{\tau}, z, \bar{z})$ of $\Psi_\mu(\tau, z)$ (4.55). The differential operator maps a Jacobi form of weight k and index m to a Jacobi form of weight $k + 2$ and index m . The second non-holomorphic term in (4.107) is required for modularity but does is not relevant for the holomorphic part.

To determine the completion $\hat{\Psi}_\mu(\tau, \bar{\tau}, z, \bar{z})$, we set $z = \beta\tau + \delta$ with $\beta, \delta \in \mathbb{R}$, so that $\beta = \text{Im}(z)/\tau_2$. We first note that completing the square gives for a generic vector $V \in \underline{\Lambda}$,

$$y^{V \cdot \mathbf{Q}} q^{-\frac{1}{2}\mathbf{Q}^2} = q^{\frac{\beta^2}{2}V^2 - \frac{1}{2}(\mathbf{Q} - \beta V)^2} e^{2\pi i(V \cdot \mathbf{Q})\delta}, \quad (4.108)$$

for an arbitrary vector V . To write the modular completion, we can treat the three permutations separately. With C_1 as in (4.71), and $C_\ell^{(123)}$ as in (4.72), we find that the

kernel $(-1)^{a+b-c} F^*(123) (y^{a+b-c} + y^{-a-b+c})$ is to be completed to

$$\widehat{F}^*(123, y) = \frac{(-1)^{a+b-c}}{4} \sum_{\pm} \left[1 + \sum_{\ell=1}^3 E_2(C_\ell^{(123)}, C_{\ell+1}^{(123)}; \sqrt{2\tau_2} (\mathbf{Q} \mp \beta C_1)) \right] y^{\pm(a+b-c)}. \quad (4.109)$$

Then the completion $\widehat{\Psi}_\mu(\tau, \bar{\tau}, z, \bar{z})$ reads,

$$\widehat{\Psi}_\mu(\tau, \bar{\tau}, z, \bar{z}) = \sum_{\mathbf{Q} \in \underline{\Lambda}_\mu^*} [\widehat{F}^*(123, y) + \widehat{F}^*(213, y) + \widehat{F}^*(132, y)] q^{-\mathbf{Q}^2/2}. \quad (4.110)$$

The completion $\widehat{\Psi}_\mu(\tau, \bar{\tau}, z, \bar{z})$ transforms as a Jacobi form of weight $b_2/2$ and index m_P given by,

$$m_P = -\frac{1}{2} C_1^2 = -P_1 P_2 P_3 - \frac{1}{2} \sum_{\ell=1}^3 P_\ell (P_{\ell+1}^2 + P_{\ell+2}^2). \quad (4.111)$$

This is symmetric under permutations, and thus also equal to $-C_2^2/2 = -C_3^2/2$.

To write the modular completion, we define the function

$$\begin{aligned} G(V_1, V_2, V_3; \mathbf{Q}, \tau_2) &= -\frac{1}{4\pi^2} \partial_z^2 \left(E_2(V_1, V_2; \sqrt{2\tau_2} (\mathbf{Q} - \beta V_3)) y^{V_3 \cdot \mathbf{Q}} \right) |_{z=0} \\ &= (V_3 \cdot \mathbf{Q})^2 E_2(V_1, V_2; \sqrt{2\tau_2} \mathbf{Q}) \\ &\quad - \frac{1}{4\pi^2} \partial_z^2 E_2(V_1, V_2; \sqrt{2\tau_2} (\mathbf{Q} - \beta V_3)) |_{z=0} \\ &\quad + \frac{1}{2\pi i} (V_3 \cdot \mathbf{Q}) \partial_z E_2(V_1, V_2; \sqrt{2\tau_2} (\mathbf{Q} - \beta V_3)) |_{z=0}. \end{aligned} \quad (4.112)$$

The limit $\tau_2 \rightarrow \infty$ is determined by the first term,

$$\lim_{\tau_2 \rightarrow \infty} G(V_1, V_2, V_3; \mathbf{Q}, \tau_2) = \begin{cases} (V_3 \cdot \mathbf{Q})^2 \arctan(\alpha), & \text{if } V_1 \cdot \mathbf{Q} = V_2 \cdot \mathbf{Q} = 0, \\ (V_3 \cdot \mathbf{Q})^2 \operatorname{sgn}(V_1 \cdot \mathbf{Q}) \operatorname{sgn}(V_2 \cdot \mathbf{Q}), & \text{otherwise,} \end{cases} \quad (4.113)$$

with α as in (4.75).

To write the modular completion, we can treat the three permutations separately. We set

$$\widehat{G}^*(123) = \frac{(-1)^{a+b-c}}{4} \left((a+b-c)^2 + \sum_{\ell=1}^3 G(C_\ell^{(123)}, C_{\ell+1}^{(123)}, C_1; \mathbf{Q}, \tau_2) \right), \quad (4.114)$$

with C_1 as in (4.71), and $C_\ell^{(123)}$ as in (4.72). The other permutations (213) and (132) give similarly rise to $\widehat{F}^*(213)$ and $\widehat{F}^*(132)$. We define,

$$\widehat{g}_C(\{\gamma_j\}; \{c_j^*\}) = \frac{1}{4} \left[\widehat{G}^*(123) + \widehat{G}^*(213) + \widehat{G}^*(132) \right]. \quad (4.115)$$

The modular completion $\widehat{\Psi}_\mu$ now reads

$$\widehat{\Psi}_\mu(\tau, \bar{\tau}) = \sum_{\mathbf{Q} \in \underline{\Lambda}_\mu^*} \widehat{g}_C(\{\gamma_j\}; \{c_j^*\}) q^{-\mathbf{Q}^2/2} - \frac{m_P}{2\pi\tau_2} \widehat{\Phi}_\mu(\tau, \bar{\tau}), \quad (4.116)$$

where the last term is due to the m -dependent term in (4.107). We let this expression be our final form for $\widehat{\Psi}_\mu(\tau, \bar{\tau})$.

Our last task is to determine the constant A in $g_C(\{\gamma_j\}; \{c_j^*\})$ (2.42). To this end, we consider the $\tau_2 \rightarrow \infty$ limit of $\widehat{g}_C(\{\gamma_j\}; \{c_j^*\})$ and require that it reduces to $g_C(\{\gamma_j\}; \{c_j^*\})$. If we consider the term $F^*(123)(a+b-c)^2$, we see that only the completion of the term $\text{sgn}(a-c)\text{sgn}(b-c)$ in g_C can contribute a non-vanishing difference. Indeed, if we include the other permutations, the only remaining terms in the difference are the equilateral cases with $a = b = c$,⁴

$$\begin{aligned} \lim_{\tau_2 \rightarrow \infty} \widehat{g}_C(\{\gamma_j\}; \{c_j^*\}) - g_C(\{\gamma_j\}; \{c_j^*\}) &= \frac{(-1)^a}{4} a^2 \delta_{a,b} \delta_{b,c} \\ &\times \lim_{\tau_2 \rightarrow \infty} \left(E_2(C_a - C_c, C_b - C_c; \sqrt{2\tau_2} \mathbf{Q}) + E_2(C_a - C_b, C_c - C_b; \sqrt{2\tau_2} \mathbf{Q}) \right. \\ &\quad \left. + E_2(C_c - C_a, C_b - C_a; \sqrt{2\tau_2} \mathbf{Q}) - A \right). \end{aligned} \quad (4.117)$$

Now the sum of E_2 's is precisely of the form (4.85), and thus equals 1. As a result, we find that with $A = 1$ the limit vanishes for any choice of Calabi-Yau and charge configurations. This matches perfectly with the physical expectation. Note that the individual values for E_2 are given by an arctan, and are generically irrational, but that this combination adds up to 1.

It is quite striking that the values of E_2 's precisely confirm the physical expectation. Also in other cases [30, 43], the values of the generalized error functions for vanishing arguments has matched with the expectations based on BPS invariants.

5 Relation to M5-branes and AdS₃/CFT₂

In this brief section, we discuss our findings from the point of view of M-theory and the MSW CFT. We discuss how the partition functions \widehat{Z}_P^{nT} , $n > 2$, may arise from the 2-dimensional perspective.

The uplift of the D4-branes to M5-branes in M-theory is useful to understand the modular properties of the partition functions [6, 12–14]. The spatial dimensions of the M5-brane are $P \times S_M^1$, where $P \in H_4(X, \mathbb{Z})$ is a four-cycle in the Calabi-Yau threefold and S_M^1 is the M-theory circle. The D2 branes of IIA string theory are realized as excitations of the self-dual 2-form field on M5 brane world volume, while the D0-branes are realized as momenta of the brane system along S_M^1 with radius $R = g_s \ell_s = \ell_{11}^3 / \ell_s^2$ in terms of the string coupling g_s , string length ℓ_s , and eleven dimensional Planck length ℓ_{11} . The world volume theory of the M5 brane gives a low energy description of the system, provided gravitational effects can be ignored. This can be ensured by taking the volume of the CY₃ to be large, namely V_X / ℓ_{11}^6 large but fixed. Supersymmetry of the M5 world volume theory implies that the MSW CFT has (0,4) supersymmetry, and is dual to the near horizon AdS₃ geometry [12, 76]. Bosons of this CFT include, moduli of the divisor P inside X , translations along flat \mathbb{R}^3 and (anti)-chiral scalars coming from reduction of self-dual two-form field in

⁴We stress that the equilateral condition $a = b = c$, is the condition on \mathbf{Q} to satisfy $C_a \cdot \mathbf{Q} = C_b \cdot \mathbf{Q} = C_c \cdot \mathbf{Q}$, and does *not* imply equalities among C_a , C_b and C_c .

the M5 brane world volume. The number of fields can be determined using geometric data of X , when P is a very ample divisor. Consequently the central charges can be determined. The central charge of the left-moving, non-supersymmetric sector is

$$c_L = P^3 + c_2(X) \cdot P, \tag{5.1}$$

where $c_2(X)$ is the second Chern class of the Calabi-Yau three-fold X . Using Cardy's formula, Reference [12] demonstrated that microscopic entropy is in agreement with one loop corrected Bekenstein-Hawking entropy.

As for any CFT, a key feature of the MSW CFT is the modular symmetry. When time direction is compactified, the CFT lives on a 2-torus $T^2 = S_M^1 \times S_t^1$. The linear fractional transformation of the complex structure modulus τ of T^2 by an element of $SL(2, \mathbb{Z})$ corresponds to the same torus T^2 . Thus the $SL(2, \mathbb{Z})$ symmetry, or modular symmetry, appears as a symmetry of the CFT. This is a strong constraint on the degeneracies. Another property of the CFT is the spectral flow of the $U(1)^{b_2}$ current algebra. This leads to a symmetry of degeneracies as function of the charges, such that the partition function can be decomposed as a finite sum of theta series \times modular functions, much as is the case for the attractor partition function in section 3.2.

We proceed by considering the AdS_3 dual to the MSW CFT following [27]. The relation of λ in (3.14) to the five-dimensional quantities can be understood from four-dimensional Newton's constant. From the IIA perspective, we have

$$G_4 = \frac{g_s^2 \ell_s^8}{V_X} = \frac{g_s^2 \ell_s^2}{\lambda^3}, \tag{5.2}$$

while in terms of the five-dimensional quantities,

$$G_4 = \frac{\ell_5^3}{R} = g_s^2 \ell_s^2 \frac{\ell_5^3}{R^3}, \tag{5.3}$$

such that one has,

$$\lambda = \frac{R}{\ell_5}. \tag{5.4}$$

Starting from five dimensional asymptotically flat geometries with appropriate charges, one takes the decoupling limit as follows: size of M-theory circle R is kept fixed (in absolute units), Calabi-Yau volume is kept finite in units of five dimensional Planck length ℓ_5 . In the decoupling limit to AdS_3 , $\ell_5 \rightarrow 0$, hence $\lambda \rightarrow \infty$ by (5.4). Multi-centered geometries where centers have mutual distance of the order of $\ell_5^3 \sim \lambda^{-3}$ or less, survive this limit and go over to asymptotically $AdS_3 \times S^2$ geometries, with asymptotic moduli fixed to their attractor values. These are the λ -core geometries mentioned in the introduction, and include centers with non-vanishing individual D6 brane charges, which add up to zero. On the other hand, if the distances between the centers is larger than $\sim \ell_5^3/R^2$ for $\ell_5 \rightarrow 0$, the energy of their fluctuations diverges and such bound states decouple from the spectrum. As we can see from the Denef equations (2.2), this is for example the case if the centers carry a non-vanishing D4-brane charge with vanishing D6-brane charge. Then the distances between the centers scale as ℓ_5 for non-scaling bound states, and the centers give rise to disconnected AdS_3

throats in the decoupling limit [27]. On the other hand for scaling solutions, the distance between the centers contains a regime where the centers can come arbitrarily close.

As explained in section 2, the BPS index can be determined using localization with respect to rotation around the z -axis, and we can therefore concentrate on collinear solutions to Denef’s equations [68]. These collinear solutions admit two branches, one corresponding to centers at finite distances, and the other one is when the centers are nearly coincident. The second branch, sometimes referred to as “deep scaling regime” reproduces pure Higgs states and goes to a single throat in $\ell_5 \rightarrow 0$ limit [53]. This is in accordance with the expectation that MSW CFT captures the near coincident regime of scaling solutions since these are smoothly connected to the single center black hole. Since $c_j \sim \lambda$ (4.1), The separation between the centers at the collinear fixed point is of order $\lambda^{-1} \sim \ell_5$, and these therefore decouple from the AdS_3 geometry. As a result, these do not appear to be captured by the CFT. This leads us to speculate that the first term on the r.h.s. of (1.1), $\widehat{\mathcal{Z}}_P^T$, corresponds to the $\text{AdS}_3/\text{CFT}_2$ partition function while the other terms on the r.h.s. do not.

It is intriguing that the terms due to scaling solutions $\widehat{\mathcal{Z}}_P^{nT}$ with $n \geq 3$ do satisfy the restrictive modular transformations as well as the spectral flow symmetry, and it is desirable to understand the origin. We think that these terms can appear after turning on an irrelevant deformation in the $(0, 4)$ CFT, away from the conformal fixed point and reversing the attractor flow and $\ell_5 \rightarrow 0$ limit. While such a deformation does not lead to a finite change in the partition function for $(4, 4)$ CFT [54], it seems plausible to us that this can happen with reduced supersymmetry. This deformation is distinguished among other deformations spanning the space of attractor moduli, since this deformation does preserve the spectral flow symmetry which is in general not the case for variations of the Kähler moduli orthogonal to P [25]. We leave a further exploration of these interesting aspects for future work.

6 Case studies

To make a suitable choice of a Calabi-Yau threefold, we note that the lattices involved have dimensions linear in b_2 , the second Betti number. Thus the computations are less involved for smaller b_2 . However for the simplest case $b_2 = 1$, there are no cyclic quivers and hence no scaling solutions.⁵ Also, the lattices concerned are positive definite, and therefore do not give rise to mock modular forms. So we settle for the next simplest case $b_2 = 2$.

In order to define quadratic forms on the lattices, one needs the intersection numbers. For a Calabi Yau threefold with $b_2 = 2$, there are $2^3 = 8$ intersection numbers of which only 4 are independent. For the sake of simplicity we shall choose a Calabi Yau threefold X , for which all intersection numbers are $\in 2\mathbb{Z}$ such that Simplification 2 of section 4.1 holds, and for which most of these numbers vanish. To be specific, we choose X to be a $K3$ fibration over \mathbb{P}^1 . This Calabi-Yau manifold corresponds to Polytope ID # =14 in the online database [77]. X can be realized as a divisor in an ambient toric variety $\mathbb{P}^3 \times \mathbb{P}^1$,

⁵Consider triangular quiver to start with. For $b_2 = 1$, electric and magnetic charges are numbers and satisfy the identity $aP_3 + bP_1 + cP_2 = 0$. Since $P_i > 0$ this implies a, b, c can not all have the same sign hence the quiver can not be cyclic. This is easily generalized to any cyclic quiver.

specified by the weight matrix

$$W = \begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{pmatrix}. \quad (6.1)$$

W defines the toric variety through the identification on $\mathbb{C}^6/\{0\}$

$$(z_1, \dots, z_6) \sim (\lambda^{W_{i1}} z_1, \dots, \lambda^{W_{i6}} z_6), \quad \lambda \in \mathbb{C}^*, \quad i = 1, 2, \quad (6.2)$$

where z_1, \dots, z_6 are coordinates of \mathbb{C}^6 . Anti-canonical divisor of this toric variety defines the Calabi Yau threefold under consideration. It has the following Hodge numbers

$$h^{1,1} = 2, \quad h^{2,1} = 86 \quad (6.3)$$

and consequently the Euler number is $\chi = 2(h^{1,1} - h^{2,1}) = -168$. Intersection polynomial and second Chern class of X are given respectively by

$$\begin{aligned} \text{intersection polynomial} &= 4J_1 J_2^2 + 2J_2^3, \\ \text{second Chern class} &= 8J_1 J_2 + 6J_2^2, \end{aligned} \quad (6.4)$$

where J_i 's are a basis of 2-forms on X and generators of the Kähler cone. X is favorable, meaning that Kähler forms on X descend from the ambient toric variety. In our notation, the intersection numbers d_{abc} and second Chern class $c_{2,a}$ read

$$\begin{aligned} d_{111} = d_{112} = 0, \quad d_{122} = 4, \quad d_{222} = 2, \\ c_{2,1} = 24, \quad c_{2,2} = 44 \end{aligned} \quad (6.5)$$

and the d_{abc} for other indices follow by permutation.

6.1 Charge configuration 1: $P_1 = P_2 = (0, 1)$, $P_3 = (1, 1)$

In this example we choose the total magnetic charge $P = (1, 3)$ magnetic charges split into 3 centers as

$$P_1 = P_2 = (0, 1), \quad P_3 = (1, 1). \quad (6.6)$$

We note that $P_{1,2}$ are irreducible while P_3 is reducible. Therefore $h_{P_{1,2},\mu}$ is a weakly holomorphic modular form, while $h_{P_3,\mu}$ is a mock modular form. The corresponding central charges of the left-moving sector of the CFT (5.1) are,

$$c_L(P_{1,2}) = 46, \quad c_L(P_3) = 92, \quad c_L(P) = 318. \quad (6.7)$$

For the j -th center, we have for the innerproduct $D_j = d_{abc} P_j^c$,

$$D_1 = D_2 = \begin{pmatrix} 0 & 4 \\ 4 & 2 \end{pmatrix}, \quad D_3 = \begin{pmatrix} 0 & 4 \\ 4 & 6 \end{pmatrix}. \quad (6.8)$$

For this choice, $D_3^{-1} D_1 = D_3^{-1} D_2$ is an integral matrix, such that Simplification 1. of section 4.1 is satisfied, and we can use the results of that section. We have, $D_1 D_3^{-1} D_1 =$

$\begin{pmatrix} 0 & 4 \\ 4 & -2 \end{pmatrix}$ which produces the quadratic form of $\underline{\Lambda}$ (4.33),

$$\underline{D} = \begin{pmatrix} 0 & 8 & 0 & 4 \\ 8 & 0 & 4 & -2 \\ 0 & 4 & 0 & 8 \\ 4 & -2 & 8 & 0 \end{pmatrix}, \quad \det \underline{D} = 2304. \quad (6.9)$$

The quadratic form on the lattice of electric charges \underline{D}^{-1} is given by,

$$\underline{D}^{-1} = \begin{pmatrix} -\frac{1}{18} & \frac{1}{6} & \frac{5}{72} & -\frac{1}{12} \\ \frac{1}{6} & 0 & -\frac{1}{12} & 0 \\ \frac{5}{72} & -\frac{1}{12} & -\frac{1}{18} & \frac{1}{6} \\ -\frac{1}{12} & 0 & \frac{1}{6} & 0 \end{pmatrix}. \quad (6.10)$$

Electric charge vectors $Q_j \in \Lambda_j^*$ have the following form

$$Q_j = \begin{pmatrix} q_{j1} \\ q_{j2} \end{pmatrix}.$$

There are $\det(D_1) \det(D_2) = 256$ conjugacy classes of the form

$$\begin{aligned} q_{11} &= 4k_{12} + \mu_{11}, & q_{12} &= 4k_{11} + 2k_{12} + \mu_{12}, \\ q_{21} &= 4k_{22} + \mu_{21}, & q_{22} &= 4k_{21} + 2k_{22} + \mu_{22}, \end{aligned} \quad (6.11)$$

where $\mu_{ij} \in \{0, 1, 2, 3\}$, $k_{ij} \in \mathbb{Z}$. These conjugacy classes have the exchange symmetry between μ_1, μ_2 . Since D_3 divides D_1 and D_2 , $N_q = \det(D_1) \det(D_2)$ (4.35) is the number of conjugacy classes of $\underline{\Lambda}^*/\underline{\Lambda}$ for fixed $\mu_1 + \mu_2 + \mu_3 = \mu \in \Lambda^*/\Lambda$, since the class $\mu_3 \in \Lambda_3^*/\Lambda_3$ by the simplification. Using the equations in (4.39), we can write

$$\begin{aligned} \mathcal{Q}_1 &= \begin{pmatrix} 8k_{12} + 4k_{22} + \mu_{11} - \mu_{31} \\ 8k_{11} + 4k_{21} - 2k_{22} + \mu_{12} + \mu_{31} - \mu_{32} \end{pmatrix}, \\ \mathcal{Q}_2 &= \begin{pmatrix} 8k_{22} + 4k_{12} + \mu_{21} - \mu_{31} \\ 8k_{21} + 4k_{11} - 2k_{12} + \mu_{22} + \mu_{31} - \mu_{32} \end{pmatrix}. \end{aligned} \quad (6.12)$$

We restrict to $\mu = 0 = \mu_1 + \mu_2 + \mu_3$, such that

$$\begin{aligned} \mathcal{Q}_1 &= \begin{pmatrix} 8k_{12} + 4k_{22} + 2\mu_{11} + \mu_{21} \\ 8k_{11} + 4k_{21} - 2k_{22} + 2\mu_{12} + \mu_{22} - \mu_{11} - \mu_{21} \end{pmatrix}, & \boldsymbol{\mu}_1 &= \begin{pmatrix} 2\mu_{11} + \mu_{21} \\ 2\mu_{12} + \mu_{22} - \mu_{11} - \mu_{21} \end{pmatrix}, \\ \mathcal{Q}_2 &= \begin{pmatrix} 8k_{22} + 4k_{12} + 2\mu_{21} + \mu_{11} \\ 8k_{21} + 4k_{11} - 2k_{12} + 2\mu_{22} + \mu_{12} - \mu_{11} - \mu_{21} \end{pmatrix}, & \boldsymbol{\mu}_2 &= \begin{pmatrix} 2\mu_{21} + \mu_{11} \\ 2\mu_{22} + \mu_{12} - \mu_{11} - \mu_{21} \end{pmatrix}. \end{aligned}$$

List of Φ_μ . We next present explicit q -expansions of Φ_μ for a various choices of μ . All the convergence conditions of the types $C_i^2 C_{i+1}^2 - (C_i \cdot C_{i+1})^2 > 0$ and $C_i^2 (C_{i-1} \cdot C_{i+1}) - (C_i \cdot C_{i-1}) \cdot (C_i \cdot C_{i+1}) < 0$ are satisfied in this example along with $C_j^2 > 0$. As in explained in section 4.1, μ is fully determined by specifying (μ_1, μ_2, μ_3) and $\mu = \mu_1 + \mu_2 + \mu_3$. In the

following example, we fix $\mu = \mu_1 = 0$ and $\mu_3 = -\mu_2$, and list Φ_μ for 16 different values of μ_2 . Out of this set, 10 give rise to different q -series. The overall sign of the q -expansion is given by $(-1)^{(a+b+c)} = (-1)^{\mu_{11}-\mu_{21}} = (-1)^{\mu_{21}}$. Furthermore, the growth of the coefficients is moderate, $|c(n)| \leq C n^\alpha$ for some positive constants α and C , which is expected for a holomorphic (mock) modular form of weight 2.

It can be shown that in most of the sectors we study ($\mu_1 = (0, 0)$) there is no contribution to the holomorphic part when the arguments of E_2 are both zero. When $\mu_2 = (1, j)$ or $\mu_2 = (3, j)$ b is odd while a, c are both odd or both even. Under these conditions at least two of $(a + b - c)$, $(a - b + c)$, $(c + b - a)$ cannot vanish. Only in the conjugacy classes $\mu_2 = (0, 0), (0, 2), (2, 3)$ we find integral $k_{11}, k_{12}, k_{21}, k_{22}$ to satisfy at least two of $(a + b - c)$, $(a - b + c)$, $(c + b - a)$ to vanish. The contributions A_ℓ are given by

$$\begin{aligned} A_1 &= \frac{2}{\pi} \arctan(-5/\sqrt{11}), \\ A_2 = A_3 &= \frac{2}{\pi} \arctan(-1/\sqrt{8}), \end{aligned} \tag{6.13}$$

which are all irrational numbers.

For $\mu_2 = (0, 0)$, we have:

$$\begin{aligned} \Phi_\mu &= 2q^6 + 4q^{20} + 6q^{24} + 4q^{30} + 4q^{44} + 4q^{50} + 8q^{52} + 2q^{54} + 12q^{56} + 4q^{60} \\ &\quad + 4q^{64} + 4q^{68} + 4q^{70} + 12q^{80} + 2q^{88} + 8q^{90} + 8q^{92} + 8q^{94} + 14q^{96} + 16q^{100} \\ &\quad + \frac{(A_1 + A_2 + A_3 + 1)}{2} + (A_1 + A_2 + A_3 + 3)(q^8 + q^{32} + q^{72}) + \dots \end{aligned}$$

For $\mu_2 = (0, 1)$, we have:

$$\begin{aligned} \Phi_\mu &= q^5 \left(1 + q + 2q^{12} + 2q^{15} + 2q^{17} + 2q^{20} + q^{22} + 2q^{23} + q^{27} + 3q^{28} + q^{30} + 2q^{35} + 2q^{38} \right. \\ &\quad + 3q^{40} + 2q^{42} + 4q^{45} + 3q^{46} + 3q^{49} + 4q^{51} + 2q^{52} + 2q^{53} + 3q^{54} + 2q^{55} + 2q^{57} + 3q^{60} \\ &\quad + 2q^{61} + 2q^{63} + q^{64} + 2q^{65} + 5q^{69} + 6q^{70} + 2q^{71} + 2q^{73} + 3q^{75} + 2q^{78} + 2q^{79} + 2q^{81} \\ &\quad \left. + q^{82} + 2q^{83} + 6q^{84} + 2q^{86} + 3q^{87} + 4q^{88} + 5q^{90} + 4q^{91} + 2q^{92} + 6q^{93} + 2q^{94} + 2q^{95} \right) + \dots \end{aligned}$$

For $\mu_2 = (0, 2)$, we have:

$$\begin{aligned} \Phi_\mu &= 2q^{14} + 2q^{20} + 2q^{24} + 2q^{26} + 2q^{32} + 6q^{36} + 4q^{40} + 4q^{42} + 2q^{44} + 4q^{48} \\ &\quad + 10q^{56} + 6q^{60} + 4q^{62} + 4q^{68} + 4q^{70} + 8q^{72} + 8q^{76} + 10q^{80} + 4q^{82} + 4q^{86} \\ &\quad + 8q^{88} + 4q^{92} + 4q^{96} + 6q^{98} + 8q^{100} + (A_1 + 1)(q^{50} + q^{54} + q^{62} + q^{74} + q^{90}) + \dots \end{aligned}$$

For $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Phi_\mu &= -2q^{29/4} (1 + q^4 + q^{12} + q^{17} + q^{18} + q^{19} + 2q^{21} + q^{25} + q^{31} + q^{37} + 2q^{40} + q^{42} + q^{43} \\ &\quad + 2q^{45} + q^{48} + 2q^{49} + 3q^{50} + q^{52} + q^{53} + 2q^{54} + q^{55} + q^{56} + q^{57} + 2q^{60} \\ &\quad + q^{66} + 2q^{71} + 2q^{74} + q^{75} + q^{76} + 3q^{79} + q^{80} + 2q^{81} + q^{82} + 3q^{84} + q^{85} + q^{86} \\ &\quad + 3q^{87} + 2q^{90} + 3q^{91} + 3q^{92}) + \dots \end{aligned}$$

For $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{83/4} \left(1 + q^3 + q^5 + q^8 + q^9 + q^{10} + q^{13} + q^{16} + q^{22} + q^{23} + q^{24} + q^{26} + 3q^{31} \right. \\ & + 2q^{33} + q^{34} + q^{35} + q^{36} + 2q^{38} + q^{39} + q^{40} + q^{41} + 2q^{42} + q^{44} + q^{45} + q^{46} + q^{49} + 2q^{51} \\ & + 2q^{54} + q^{57} + q^{58} + 2q^{59} + 2q^{60} + 2q^{61} + q^{68} + q^{69} + q^{70} + 2q^{71} + 4q^{72} + 3q^{72} + q^{74} \\ & \left. + 2q^{75} + q^{76} + q^{77} + 3q^{78} + q^{79} \right) + \dots \end{aligned}$$

For $\mu_2 = (1, 2)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{45/4} \left(1 + 2q^8 + q^{15} + q^{18} + q^{21} + q^{23} + 2q^{27} + q^{28} + q^{29} + q^{32} + 2q^{35} + q^{39} + q^{41} \right. \\ & + 2q^{43} + q^{44} + q^{46} + 2q^{48} + q^{50} + q^{52} + q^{54} + q^{56} + 3q^{57} + q^{58} + 2q^{62} + 3q^{63} + 3q^{65} \\ & \left. + q^{66} + q^{68} + q^{70} + 2q^{72} + q^{73} + 2q^{74} + q^{75} + 3q^{80} + q^{82} + 2q^{85} + q^{86} + q^{87} + q^{88} \right) + \dots \end{aligned}$$

For $\mu_2 = (1, 3)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{15/4} \left(1 + q + q^{10} + q^{13} + q^{17} + q^{18} + q^{19} + q^{21} + q^{22} + 2q^{32} + q^{35} + q^{39} \right. \\ & + 2q^{40} + q^{42} + q^{45} + q^{47} + q^{50} + 2q^{51} + q^{53} + 3q^{54} + q^{56} + 2q^{59} + q^{63} + q^{65} + 2q^{66} \\ & + 2q^{68} + q^{71} + q^{72} + q^{74} + q^{75} + q^{76} + 3q^{77} + q^{79} + 2q^{80} + 2q^{83} + q^{84} + 2q^{85} + q^{86} \\ & \left. + q^{87} + q^{88} + 2q^{90} + 2q^{92} + 3q^{93} + 2q^{95} \right) + \dots \end{aligned}$$

For $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Phi_\mu = & 2 \left(q^9 + q^{15} + q^{19} + 2q^{27} + 2q^{29} + q^{33} + q^{35} + q^{43} + 4q^{45} + 4q^{53} + 2q^{55} + q^{57} + q^{59} \right. \\ & + 2q^{61} + 2q^{63} + 3q^{65} + q^{67} + 2q^{71} + q^{75} + 4q^{77} + 2q^{81} + 3q^{83} + 2q^{87} + q^{89} + 5q^{91} \\ & \left. + q^{93} + 6q^{95} + q^{97} + 2q^{99} \right) + \dots \end{aligned}$$

For $\mu_2 = (2, 1)$, we have:

$$\begin{aligned} \Phi_\mu = & 2q^{25} \left(2 + 2q + q^2 + q^4 + 2q^{15} + q^{16} + 2q^{18} + q^{22} + 2q^{25} + 2q^{29} + 4q^{32} + 3q^{34} + 4q^{37} \right. \\ & + 2q^{39} + q^{40} + 2q^{41} + q^{44} + q^{46} + 2q^{51} + 2q^{55} + q^{56} + 2q^{59} + 4q^{61} + 3q^{62} + 2q^{65} + 2q^{66} \\ & \left. + 2q^{68} + 4q^{70} + 2q^{73} + 2q^{74} + 2q^{75} \right) + \dots \end{aligned}$$

For $\mu_2 = (2, 3)$, we have:

$$\begin{aligned} \Phi_\mu = & 2q^5 \left(1 + q^6 + q^9 + q^{15} + q^{19} + 2q^{20} + q^{27} + q^{28} + q^{33} + 2q^{34} + q^{37} + q^{40} + q^{42} + 2q^{47} + 3q^{51} \right. \\ & + q^{54} + 2q^{55} + 2q^{56} + q^{60} + q^{63} + 2q^{64} + q^{66} + q^{69} + 2q^{70} + 4q^{71} + 2q^{73} + q^{75} + 2q^{78} + 3q^{81} \\ & \left. + 2q^{84} + 2q^{85} + 2q^{88} + 4q^{90} + 3q^{91} + 2q^{92} + 3q^{94} \right) + (A_3 + 1) q^2 (1 + q^{16} + q^{48} + q^{96}) + \dots \end{aligned}$$

List of Ψ_μ . Similarly to the Φ_μ , we list the q -expansion of Ψ_μ for a various choices of μ . In these examples, we keep $\mu = \mu_1 = 0$ and $\mu_3 = -\mu_2$, and list Ψ_μ for 16 different values of μ_2 . Out of these 16 only 10 give rise to different q -series. The factor $(-1)^{(a+b+c)} = (-1)^{\mu_{11}-\mu_{21}} = (-1)^{\mu_{21}}$ fixes the overall sign of the whole series. Comparison of the Φ_μ and Ψ_μ demonstrates that for small powers of q , scaling solutions can be present kinematically, but that the number of bound states vanishes quantum mechanically. We again observe a moderate, polynomial growth of the coefficients, as expected for a (mock) modular form of weight 4.

For $\mu_2 = (0, 0)$, we have:

$$\Psi_\mu = 16q^{30} \left(1 + 2q^{22} + 4q^{34} + q^{40} + 2q^{60} + 8q^{62} + 2q^{64} + 4q^{70} \right) + \dots$$

For $\mu_2 = (0, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{17} \left(1 + q^5 + q^8 + q^{11} + q^{16} + q^{23} + q^{26} + q^{28} + 4q^{30} + q^{33} + q^{34} + 9q^{37} + 5q^{39} + q^{40} \right. \\ & + q^{41} + q^{42} + 4q^{43} + 4q^{45} + 4q^{48} + q^{49} + 9q^{51} + q^{53} + q^{57} + 6q^{58} + q^{59} + q^{61} + q^{63} + 9q^{66} \\ & \left. + 4q^{67} + 16q^{69} + q^{71} + 5q^{72} + 16q^{74} + q^{75} + 13q^{76} + 26q^{78} + q^{79} + q^{80} + 13q^{81} + q^{82} + q^{83} \right) \\ & + \dots \end{aligned}$$

For $\mu_2 = (0, 2)$, we have:

$$\Psi_\mu = 16q^{36} \left(1 + q^4 + q^{12} + 2q^{20} + q^{24} + q^{32} + q^{34} + 2q^{36} + 2q^{40} + q^{44} + q^{46} + 2q^{52} + 4q^{56} + 4q^{62} + 2q^{64} \right) + \dots$$

For $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{29/4}}{2} \left(1 + q^4 + q^{12} + q^{17} + q^{18} + q^{19} + 10q^{21} + 9q^{25} + q^{31} + q^{37} + 10q^{40} + q^{42} + q^{43} \right. \\ & + 34q^{45} + q^{48} + 2q^{49} + 27q^{50} + q^{52} + q^{53} + 50q^{54} + q^{55} + q^{56} + q^{57} + 58q^{60} + 9q^{66} + 2q^{71} \\ & + q^{74} + 9q^{75} + q^{78} + 51q^{79} + 25q^{80} + 10q^{81} + q^{82} + 107q^{84} + 25q^{85} + 49q^{86} + 43q^{87} + 26q^{90} \\ & \left. + 107q^{91} + 51q^{92} \right) + \dots \end{aligned}$$

For $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{83/4}}{2} \left(9 + 9q^3 + q^5 + q^8 + q^9 + q^{10} + 9q^{13} + q^{16} + 9q^{22} + q^{23} + 9q^{24} + q^{26} + 19q^{31} \right. \\ & + 34q^{33} + 9q^{34} + 25q^{35} + 9q^{36} + 26q^{38} + 9q^{39} + 25q^{40} + q^{41} + 10q^{42} + 25q^{44} + 9q^{45} \\ & + 25q^{46} + q^{49} + 2q^{51} + 10q^{54} + 9q^{57} + 9q^{58} + 2q^{59} + 18q^{60} + 34q^{61} + q^{68} + 9q^{69} + 49q^{70} \\ & \left. + 74q^{71} + 140q^{72} + 131q^{73} + q^{74} + 18q^{75} + q^{76} + 49q^{77} + 43q^{78} + 25q^{79} \right) + \dots \end{aligned}$$

For $\mu_2 = (1, 2)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{45/4}}{2} \left(1 + 2q^8 + q^{15} + q^{18} + 9q^{21} + 9q^{23} + 2q^{27} + q^{28} + 9q^{29} + q^{32} + 18q^{35} + 9q^{39} \right. \\ & + 9q^{41} + 26q^{43} + 9q^{44} + 25q^{46} + 10q^{47} + 9q^{50} + q^{52} + 9q^{54} + 9q^{56} + 19q^{57} + 25q^{58} + 10q^{62} \\ & + 75q^{64} + 43q^{65} + 9q^{66} + 9q^{68} + q^{70} + 10q^{72} + q^{73} + 18q^{74} + q^{75} + 131q^{80} + 49q^{82} \\ & \left. + 50q^{85} + 44q^{86} + 49q^{87} + 25q^{88} \right) + \dots \end{aligned}$$

For $\mu_2 = (1, 3)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{15/4}}{2} \left(1 + q + q^{10} + q^{13} + q^{17} + q^{18} + q^{19} + 9q^{21} + q^{22} + 10q^{32} + q^{35} + 25q^{39} + 2q^{40} \right. \\ & + q^{42} + q^{45} + q^{47} + 9q^{50} + 50q^{51} + q^{53} + 27q^{54} + 9q^{56} + 10q^{59} + q^{63} + 9q^{65} + 10q^{66} \\ & + 50q^{68} + q^{71} + 25q^{72} + q^{74} + 9q^{75} + q^{76} + 83q^{77} + q^{79} + 10q^{80} + 10q^{83} + 25q^{84} \\ & \left. + 26q^{85} + 49q^{86} + q^{87} + q^{88} + 26q^{90} + 10q^{92} + 139q^{93} + 34q^{95} \right) + \dots \end{aligned}$$

For $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^9(1 + q^6 + q^{10} + 2q^{18} + 2q^{20} + q^{24} + q^{26} + q^{34} + 4q^{36} + 12q^{44} + 2q^{46} + q^{48} + 9q^{50} + 2q^{52} \\ & + 10q^{54} + 3q^{56} + 2q^{58} + 10q^{62} + q^{66} + 4q^{68} + 18q^{72} + 11q^{74} + 2q^{78} + q^{80} + 29q^{82} + q^{84} \\ & + 46q^{86} + q^{88} + 10q^{90}) + \dots \end{aligned}$$

For $\mu_2 = (2, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & 4q^{25} \left(1 + q + q^{15} + q^{18} + 4q^{25} + 4q^{29} + 5q^{32} + 4q^{34} + 8q^{37} + q^{39} + q^{41} + q^{51} + 4q^{55} \right. \\ & \left. + 4q^{59} + 5q^{61} + 4q^{62} + 4q^{65} + q^{66} + 16q^{68} + 16q^{70} + 9q^{73} + 9q^{74} + 4q^{75} \right) + \dots \end{aligned}$$

For $\mu_2 = (2, 3)$, we have:

$$\begin{aligned} \Psi_\mu = & 4 \left(q^{25} + q^{39} + q^{52} + 9q^{56} + q^{61} + 4q^{69} + q^{75} + 17q^{76} + 4q^{83} + 9q^{89} + q^{90} + q^{93} \right. \\ & \left. + 5q^{95} + q^{96} + 4q^{97} + 25q^{99} \right) + \dots \end{aligned}$$

6.2 Charge configuration 2: $P_1 = (1, 0)$, $P_2 = P_3 = (0, 1)$

In this example the total magnetic charge is $P = (1, 2)$ split into three different centers as $P_1 = (1, 0)$, $P_2 = P_3 = (0, 1)$. The charge P_1 corresponds to the K3 fiber. The CFT central charge (5.1) of $P_{2,3}$ is given above, while we have for P_1 and P ,

$$c_L(P_1) = 24, \quad c_L(P) = 176. \tag{6.14}$$

In the j -th center, the quadratic form $D_j = d_{abc}P_j^c$ is given by

$$D_2 = D_3 = \begin{pmatrix} 0 & 4 \\ 4 & 2 \end{pmatrix}, \quad D_1 = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}. \tag{6.15}$$

Note that D_1 is a singular matrix, and we only keep the direction with the non-vanishing eigenvalue. The partition functions for P_1 can be determined using Noether-Lefschetz theory [19, 21, 78]. On the other hand, the modular properties are in this case very restrictive, and determine the partition function with a little further input. First, the lowest exponent of $h_{P,\mu}$ is $-c_L/24 = -1$. Therefore, each of $h_{P,\mu}$ has a term with a negative coefficient, and the vector space of weakly holomorphic vector-valued modular forms is 3-dimensional. We can fix the coefficients following [13, 79]: the coefficient for $\mu = 0$ is the Euler number of the linear system of the divisor, which is $c_2 \cdot P_1/12 = 2$. Thus the

linear system is a \mathbb{P}^1 . For $\mu = 1$ and 2 , we need to require that the divisor contains a given curve class $\beta \in H^2(X, \mathbb{Z})$, and turn on a world volume flux through this curve. This gives two complex constraints, however since the linear system is a \mathbb{P}^1 , there are generically no solutions. As a result, the coefficients of the potential polar terms in $h_{P_1,1}$ and $h_{P_1,2}$ vanish. This fixes $h_{P_1,\mu}$ to be the weight $-3/2$ modular form,

$$h_{P_1,\mu}(\tau) = \frac{2 f_{P_1,\mu}(\tau)}{\eta(\tau)^{24}}, \tag{6.16}$$

with

$$\begin{aligned} f_{P_1,\mu}(\tau) &= \frac{11}{18} E_4(\tau) E_6(\tau) \Theta_\mu(\tau) \\ &\quad - 6 E_4(\tau)^2 D_{1/2} \Theta_\mu(\tau) \\ &\quad + 16 E_6(\tau) D_{5/2} D_{1/2} \Theta_\mu(\tau), \end{aligned} \tag{6.17}$$

with $\Theta_\mu(\tau) = \sum_{n \in 4\mathbb{Z} + \mu} q^{n^2/8}$, D_k the modular derivative acting on weight k modular forms, $D_k = \frac{1}{2\pi i} \frac{d}{d\tau} - \frac{k}{12} E_2(\tau)$, and E_ℓ the Eisenstein series of weight ℓ . The $f_{P_1,\mu}$ also appeared in the context of Gromov-Witten theory [78, 80]. Finally since $P_1^3 = 0$, $h_{P_1,\mu}$ is also expected to be a weakly holomorphic modular form (rather than a mock modular form), and given in terms of Hecke transformations of (6.16) [21, 40].

Returning to our original problem of Φ_μ and Ψ_μ , note that we have for the products of quadratic forms $D_1 D_3^{-1} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $D_1 D_3^{-1} D_1 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$, $D_2 D_3^{-1} D_1 = D_1 D_3^{-1} D_2 = D_1$ which produces the quadratic form (4.33),

$$\underline{D} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 4 \\ 0 & 0 & 0 & 8 \\ 0 & 4 & 8 & 4 \end{pmatrix}. \tag{6.18}$$

Excluding the null direction the determinant of $\underline{D} = -256$. The quadratic form on the lattice of electric charge lattice is given by the pseudo-inverse of \underline{D} as follows:

$$\underline{D}^{-1} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4} & -\frac{1}{8} & 0 \\ 0 & -\frac{1}{8} & 0 & \frac{1}{8} \\ 0 & 0 & \frac{1}{8} & 0 \end{pmatrix}. \tag{6.19}$$

Electric charge vectors $Q_j \in \Lambda_j^*$ have the following form

$$Q_1 = \begin{pmatrix} 0 \\ q_{12} \end{pmatrix}, \quad Q_2 = \begin{pmatrix} q_{21} \\ q_{22} \end{pmatrix}.$$

There are 64 conjugacy classes of the form

$$\begin{aligned} q_{12} &= 4k_{12} + \mu_{12}, \\ q_{21} &= 4k_{22} + \mu_{21}, \quad q_{22} = 4k_{21} + 2k_{22} + \mu_{22}, \end{aligned} \tag{6.20}$$

where $\mu_{ij} \in \{0, 1, 2, 3\}$, $k_{ij} \in \mathbb{Z}$. These are $N_q = 64$ (4.35) conjugacy classes for fixed μ .

Using equations in (4.39), we get

$$\begin{aligned} \mathcal{Q}_1 &= \begin{pmatrix} 0 \\ 4k_{12} + 4k_{22} + \mu_{12} - \mu_{31} \end{pmatrix}, \\ \mathcal{Q}_2 &= \begin{pmatrix} 8k_{22} + \mu_{21} - \mu_{31} \\ 8k_{21} + 4k_{22} + 4k_{12} + \mu_{22} - \mu_{32} \end{pmatrix}. \end{aligned} \tag{6.21}$$

When $\mu_1 + \mu_2 + \mu_3 = 0$, we have,

$$\begin{aligned} \mathcal{Q}_1 &= \begin{pmatrix} 0 \\ 4k_{12} + 4k_{22} + \mu_{12} + \mu_{21} \end{pmatrix}, & \mu_1 &= \begin{pmatrix} 0 \\ \mu_{12} + \mu_{21} \end{pmatrix}, \\ \mathcal{Q}_2 &= \begin{pmatrix} 8k_{22} + 2\mu_{21} \\ 8k_{21} + 4k_{22} + 4k_{12} + 2\mu_{22} + \mu_{12} \end{pmatrix}, & \mu_2 &= \begin{pmatrix} 2\mu_{21} \\ 2\mu_{22} + \mu_{12} \end{pmatrix}. \end{aligned}$$

List of Φ_μ . We list Φ_μ for a various choices of μ . For this choice of charges, we find that some of the inequalities for convergence are sometimes saturated. This however do not give rise to divergences in the q series. As in explained in section 4.1, μ is fully determined by specifying (μ_1, μ_2, μ_3) and $\mu = \mu_1 + \mu_2 + \mu_3$. The overall sign of the q -expansion is given by $(-1)^{a+b+c} = (-1)^{\mu_{12}}$. The contributions A_ℓ are for this example

$$A_1 = A_2 = A_3 = \frac{2}{\pi} \arctan(-3/4), \tag{6.22}$$

which is an irrational number.

For $\mu_1 = \mu_2 = 0$, we have:

$$\begin{aligned} \Phi_\mu &= 4q^6 + 4q^{24} + 12q^{30} + 6q^{40} + 4q^{54} + 12q^{64} + 12q^{70} + 12q^{78} + 12q^{88} + 4q^{96} \\ &\quad + \frac{(A_1 + A_2 + A_3 + 1)}{2} + (A_1 + A_2 + A_3 + 3)(q^8 + q^{32} + q^{72}) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Phi_\mu &= 2q^{33/8} \left(1 + q^4 + q^7 + q^{16} + 2q^{21} + q^{22} + q^{24} + q^{30} + 3q^{31} + q^{34} + q^{41} + q^{42} + q^{44} \right. \\ &\quad + 2q^{53} + q^{54} + q^{56} + 2q^{58} + q^{60} + 2q^{66} + 4q^{67} + q^{72} + 2q^{74} + q^{76} + 2q^{77} + 2q^{81} \\ &\quad \left. + q^{82} + q^{84} + q^{91} + q^{92} + 2q^{93} \right) + \frac{(A_3 + 1)}{2} q^{1/8} (1 + q^6 + q^{10} + q^{28} + q^{36} + q^{66} + q^{78}) \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Phi_\mu &= 2 \left(\frac{1}{2} q^{5/2} + q^{21/2} + q^{29/2} + q^{33/2} + q^{41/2} + \frac{3}{2} q^{45/2} + q^{61/2} + q^{65/2} + q^{77/2} + 2q^{81/2} + q^{85/2} \right. \\ &\quad + q^{89/2} + 3q^{101/2} + 2q^{105/2} + q^{109/2} + q^{117/2} + q^{121/2} + \frac{1}{2} q^{125/2} + 2q^{129/2} + q^{133/2} + q^{145/2} \\ &\quad + q^{149/2} + q^{153/2} + 2q^{157/2} + q^{161/2} + q^{165/2} + 2q^{173/2} + q^{181/2} + 4q^{185/2} + 2q^{189/2} \left. \right) \\ &\quad + q^{1/2} \frac{(A_3 + 1)}{2} (1 + q^4 + q^{12} + q^{24} + q^{40}) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (3, 0)$, we have:

$$\begin{aligned} \Phi_{\mu} = & 2q^{105/8} \left(2+q^5+q^6+q^{13}+q^{14}+2q^{24}+q^{30}+2q^{31}+2q^{33}+q^{34}+q^{38}+q^{40}+q^{43}+2q^{45} \right. \\ & \left. +q^{46}+q^{52}+3q^{55}+2q^{60}+q^{70}+2q^{71}+2q^{73}+q^{74}+q^{76}+2q^{78}+2q^{82}+q^{84}+q^{86} \right) \\ & + \frac{(A_3+1)}{2} q^{9/8} (1+q^2+q^{14}+q^{20}+q^{44}+q^{54}+q^{90}) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (0, 1)$, we have:

$$\begin{aligned} \Phi_{\mu} = & 2 \left(q^4+q^8+q^{12}+q^{20}+q^{24}+2q^{26}+q^{28}+2q^{34}+2q^{36}+q^{40}+2q^{46}+q^{48}+q^{56}+2q^{58} \right. \\ & \left. +2q^{60}+2q^{64}+4q^{70}+2q^{72}+2q^{76}+q^{80}+2q^{82}+2q^{84}+2q^{86}+q^{88}+2q^{96}+2q^{98}+q^{100} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (0, 2)$, we have:

$$\begin{aligned} \Phi_{\mu} = & 2 \left(q^{10}+2q^{16}+2q^{22}+2q^{32}+2q^{38}+2q^{42}+2q^{48}+4q^{52}+2q^{58}+2q^{62}+2q^{66}+4q^{76} \right. \\ & \left. +2q^{80}+2q^{82}+3q^{90}+4q^{94}+2q^{96} \right) + q^2 (A_3+1)(1+q^{16}+q^{48}+q^{96}) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$, $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Phi_{\mu} = & -2q^{19/4} \left(1+q^2+q^6+q^{17}+2q^{21}+q^{23}+q^{24}+q^{26}+q^{27}+q^{29}+q^{33}+q^{34}+q^{38}+q^{39} \right. \\ & \left. +q^{46}+q^{52}+q^{54}+q^{56}+q^{57}+q^{58}+q^{61}+2q^{62}+2q^{64}+q^{66}+2q^{68}+q^{69}+q^{71}+q^{72} \right. \\ & \left. +q^{75}+q^{76}+q^{77}+q^{78}+q^{81}+q^{82}+q^{86}+q^{87}+q^{88}+q^{89}+q^{90}+q^{95} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$, $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Phi_{\mu} = & -2q^{71/8} \left(1+q^5+q^9+q^{12}+q^{16}+q^{21}+q^{25}+q^{26}+q^{28}+q^{30}+q^{35}+q^{36}+q^{39}+q^{41}+q^{43} \right. \\ & \left. +q^{45}+q^{47}+q^{48}+q^{50}+2q^{54}+q^{58}+q^{60}+q^{63}+q^{65}+q^{67}+q^{68}+q^{69}+q^{71}+q^{72}+q^{75} \right. \\ & \left. +q^{76}+q^{78}+q^{80}+q^{81}+q^{82}+q^{86}+q^{88}+q^{89}+q^{90} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Phi_{\mu} = & q^{45/8} \left(3+2q^4+q^{10}+4q^{18}+2q^{22}+2q^{23}+4q^{25}+2q^{26}+2q^{30}+4q^{35}+2q^{36}+2q^{43}+4q^{48} \right. \\ & \left. +2q^{56}+4q^{57}+4q^{59}+4q^{60}+2q^{62}+4q^{66}+2q^{68}+q^{70}+4q^{71}+2q^{72}+4q^{73}+2q^{74} \right. \\ & \left. +4q^{82}+8q^{85}+4q^{90}+2q^{92} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (1, 3)$, we have:

$$\begin{aligned} \Phi_{\mu} = & q^{5/8} \left(1+2q^2+2q^{10}+2q^{13}+2q^{14}+2q^{16}+2q^{20}+4q^{21}+q^{30}+2q^{32}+4q^{39}+2q^{40}+4q^{42} \right. \\ & \left. +2q^{44}+3q^{50}+8q^{51}+2q^{54}+4q^{56}+4q^{63}+2q^{64}+2q^{66}+2q^{73}+4q^{77}+2q^{78} \right. \\ & \left. +2q^{80}+4q^{84}+2q^{86}+2q^{87}+2q^{90}+4q^{92}+8q^{93} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (2, 1)$, we have:

$$\begin{aligned} \Phi_{\mu} = & 4q^{15/2} \left(1+q^{12}+q^{20}+q^{24}+2q^{28}+q^{40}+2q^{48}+q^{52}+q^{60}+3q^{64}+2q^{72}+2q^{76} \right. \\ & \left. +q^{80}+2q^{92} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$, $\mu_2 = (2, 3)$, we have:

$$\Phi_\mu = 4q^{3/2} \left(1 + q^{12} + 3q^{16} + q^{36} + 3q^{44} + 3q^{48} + 3q^{56} + q^{72} + 3q^{84} + 3q^{92} + 3q^{96} \right) + \dots$$

For $\mu_1 = (0, 1)$, $\mu_2 = (0, 0)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{39/8} \left(1 + q^2 + q^3 + q^{17} + q^{21} + 2q^{22} + q^{24} + q^{26} + q^{27} + 2q^{28} + 2q^{33} + q^{39} + q^{46} + q^{52} \right. \\ & + 2q^{55} + q^{57} + 2q^{60} + q^{61} + 2q^{63} + q^{64} + q^{66} + q^{67} + q^{68} + 4q^{70} + 2q^{76} + q^{78} + 2q^{79} + 2q^{81} \\ & \left. + q^{87} + 2q^{89} + q^{95} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$, $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{45/4} \left(1 + q^3 + q^6 + q^{10} + q^{14} + q^{18} + q^{23} + 2q^{28} + q^{31} + q^{33} + q^{34} + q^{36} + 2q^{39} + q^{42} \right. \\ & + q^{45} + q^{46} + q^{49} + q^{51} + q^{53} + q^{55} + q^{58} + q^{59} + q^{62} + q^{65} + q^{67} + q^{70} + 2q^{72} + q^{75} + q^{77} \\ & \left. + q^{78} + q^{80} + q^{81} + q^{82} + q^{83} + 2q^{85} + 2q^{88} \right) \dots \end{aligned}$$

For $\mu_1 = (0, 1)$, $\mu_2 = (1, 2)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{47/4} \left(1 + q^3 + q^5 + q^9 + q^{13} + q^{17} + q^{23} + q^{28} + q^{29} + q^{31} + q^{32} + q^{35} + q^{36} + q^{37} + q^{38} \right. \\ & + q^{42} + 2q^{44} + q^{49} + q^{50} + q^{52} + q^{53} + 2q^{58} + q^{60} + q^{65} + q^{68} + q^{70} + q^{71} + q^{74} + q^{76} \\ & \left. + q^{77} + q^{78} + q^{79} + q^{80} + q^{83} + q^{84} + 3q^{85} + q^{87} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$, $\mu_2 = (1, 3)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{13/4} \left(1 + q^6 + q^{10} + q^{15} + q^{19} + q^{20} + q^{21} + q^{27} + q^{30} + q^{33} + q^{34} + q^{35} + q^{39} + q^{42} + q^{45} \right. \\ & + q^{46} + q^{50} + q^{51} + q^{52} + q^{54} + q^{56} + q^{58} + q^{60} + q^{63} + q^{64} + q^{65} + q^{70} + q^{71} + q^{72} + q^{74} \\ & \left. + q^{75} + q^{76} + q^{78} + q^{80} + q^{81} + q^{84} + q^{85} + q^{86} + q^{87} + q^{93} + q^{94} + q^{95} + q^{96} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$, $\mu_2 = (2, 2)$, we have:

$$\begin{aligned} \Phi_\mu = & -2q^{15/8} \left(1 + q^{10} + q^{13} + q^{15} + q^{16} + 2q^{18} + q^{27} + q^{33} + q^{38} + 2q^{40} + q^{42} + q^{45} + 3q^{47} \right. \\ & + q^{48} + 2q^{52} + 2q^{53} + 2q^{58} + q^{60} + q^{68} + 2q^{69} + q^{75} + 2q^{79} + q^{81} + 3q^{86} + 2q^{88} + q^{90} \\ & \left. + q^{93} + q^{94} + q^{95} + 3q^{96} + 2q^{97} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 2)$, $\mu_2 = (0, 1)$, we have:

$$\begin{aligned} \Phi_\mu = & 2q^{11/2} \left(1 + q^4 + q^{10} + 2q^{18} + q^{22} + q^{24} + 2q^{26} + q^{30} + q^{34} + 2q^{36} + q^{42} + q^{44} \right. \\ & + 2q^{48} + 2q^{56} + 2q^{58} + 3q^{60} + q^{64} + 2q^{68} + q^{70} + 3q^{72} + 2q^{74} + q^{80} + 3q^{84} + 2q^{86} \\ & \left. + 3q^{90} + q^{94} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 2)$, $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Phi_\mu = & 2q^{41/8} \left(1 + 2q^5 + q^{10} + 2q^{11} + 4q^{18} + q^{21} + q^{25} + 2q^{26} + 2q^{27} + 2q^{32} + 2q^{35} + 2q^{36} \right. \\ & + 2q^{37} + q^{41} + 2q^{44} + 3q^{45} + 4q^{48} + 2q^{55} + 2q^{56} + 2q^{57} + q^{58} + 2q^{59} + 2q^{60} + 2q^{61} \\ & + q^{66} + 2q^{68} + 3q^{70} + 2q^{71} + 2q^{73} + 2q^{74} + 4q^{75} + 2q^{81} + 2q^{83} + 4q^{86} + 4q^{87} + q^{88} \\ & \left. + 4q^{90} + 2q^{93} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 2)$, $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Phi_\mu = & q^{29/8} \left(1 + q^4 + 3q^9 + q^{15} + 2q^{16} + q^{19} + q^{22} + 2q^{23} + 2q^{24} + q^{29} + q^{30} + 4q^{33} + 2q^{38} \right. \\ & + q^{39} + q^{40} + 2q^{42} + 2q^{43} + 2q^{44} + q^{49} + 2q^{51} + q^{52} + 3q^{54} + 2q^{55} + 2q^{56} + q^{60} + 2q^{62} \\ & + 3q^{65} + 4q^{66} + 4q^{69} + q^{70} + q^{72} + 2q^{78} + 3q^{79} + 2q^{80} + 2q^{82} + 4q^{83} + 2q^{84} + q^{85} + q^{87} \\ & \left. + 2q^{91} + 2q^{93} + q^{94} + 4q^{95} + 2q^{96} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 2)$, $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Phi_\mu = & 4 \left(q^{13} + q^{19} + q^{27} + q^{37} + q^{43} + q^{45} + q^{47} + q^{53} + q^{57} + q^{59} + q^{67} + q^{69} + q^{73} + q^{83} \right. \\ & \left. + q^{85} + q^{87} + q^{89} + q^{93} + q^{97} + q^{99} \right) + \dots \end{aligned}$$

List of Ψ_μ . Similarly as the Φ_μ q -series we list the q -series of Ψ_μ for a various choices of μ . In this example $(-1)^{a+b+c} = (-1)^{\mu_{12}}$. For $\mu_1 = \mu_2 = (0, 0)$, we have:

$$\Psi_\mu = 16(q^6 + 4q^{24} + 3q^{30} + 9q^{54} + 12q^{64} + 3q^{70} + 3q^{78} + 16q^{96}) + \dots$$

For $\mu_1 = (0, 0)$ $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{33/8} \left(1 + 4q^4 + 9q^{16} + 2q^{21} + 4q^{22} + 16q^{24} + 4q^{30} + 8q^{31} + q^{42} + 25q^{44} + 18q^{53} + 16q^{54} \right. \\ & \left. + 36q^{56} + 2q^{58} + 4q^{60} + 17q^{66} + 40q^{67} + 4q^{72} + 8q^{74} + 8q^{81} + 9q^{82} + 49q^{84} + 2q^{93} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & 8q^{21/2} \left(1 + q^6 + q^{12} + 4q^{22} + q^{28} + 2q^{30} + 4q^{32} + 2q^{40} + 5q^{42} + q^{48} + 2q^{54} + 9q^{56} + \right. \\ & \left. + 4q^{66} + 8q^{68} + 9q^{70} + q^{72} + 2q^{76} + 10q^{82} + 13q^{84} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (3, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{105/8} \left(5 + 4q^6 + q^{14} + 25q^{24} + 4q^{30} + 2q^{31} + 8q^{33} + 16q^{34} + 4q^{40} + 8q^{45} + 9q^{46} + 2q^{55} \right. \\ & \left. + 61q^{60} + 16q^{70} + 18q^{71} + 32q^{73} + 36q^{74} + 4q^{76} + 2q^{78} + 8q^{82} + 16q^{84} + q^{86} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (0, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^4 \left(1 + q^4 + q^8 + 9q^{16} + q^{20} + 2q^{22} + 9q^{24} + 2q^{30} + 10q^{32} + q^{36} + 2q^{42} + 25q^{44} \right. \\ & + 9q^{52} + 18q^{54} + 26q^{56} + 2q^{60} + 20q^{66} + 34q^{68} + 2q^{72} + 9q^{76} + 2q^{78} + 2q^{80} + 18q^{82} \\ & \left. + 49q^{84} + 2q^{92} + 2q^{94} + 25q^{96} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (0, 2)$, we have:

$$\Psi_\mu = 16 \left(q^{16} + q^{32} + 4q^{42} + q^{48} + 2q^{52} + 4q^{66} + 2q^{76} + 9q^{80} + 4q^{90} + 8q^{94} + q^{96} \right) + \dots$$

For $\mu_1 = (0, 0)$ $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{45/8} \left(1 + q^4 + 13q^{18} + q^{22} + 2q^{25} + 9q^{26} + q^{30} + 2q^{32} + 4q^{36} + 41q^{48} + 9q^{56} + 8q^{57} \right. \\ & \left. + 18q^{59} + 26q^{60} + 2q^{66} + 9q^{68} + 2q^{71} + q^{72} + 18q^{73} + 16q^{74} + 2q^{82} + 10q^{85} + 85q^{90} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (1, 3)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{21/8} \left(1 + 4q^8 + q^{12} + 9q^{14} + q^{18} + 2q^{19} + 16q^{30} + 8q^{37} + 9q^{38} + 25q^{40} + q^{42} + 9q^{48} \right. \\ & + 20q^{49} + q^{52} + 2q^{54} + 2q^{61} + 4q^{62} + 36q^{64} + 32q^{75} + 25q^{76} + 49q^{78} + 8q^{82} + 9q^{84} + q^{88} \\ & \left. + 29q^{90} + 68q^{91} \right) \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (2, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & 4q^{15/2} \left(1 + q^{12} + 9q^{20} + q^{24} + 2q^{28} + 9q^{40} + 2q^{48} + 25q^{52} + 9q^{60} + 19q^{64} + 2q^{72} + 2q^{76} \right. \\ & \left. + 25q^{80} + 18q^{92} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 0)$ $\mu_2 = (2, 3)$, we have:

$$\Psi_\mu = 4q^{3/2} \left(1 + 9q^{12} + 3q^{16} + 25q^{36} + 27q^{44} + 3q^{48} + 3q^{56} + 49q^{72} + 75q^{84} + 27q^{92} + 3q^{96} \right) + \dots$$

For $\mu_1 = (0, 2)$ $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & 4 \left(q^{13} + q^{19} + q^{27} + 9q^{37} + q^{43} + q^{45} + 9q^{47} + q^{53} + q^{57} + 9q^{59} + q^{67} + q^{69} \right. \\ & \left. + 25q^{73} + 9q^{83} + 9q^{85} + 25q^{87} + q^{89} + q^{93} + 9q^{97} + q^{99} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{19/4}}{2} \left(1 + q^2 + q^6 + 25q^{17} + 26q^{21} + q^{23} + 9q^{24} + q^{26} + q^{27} + 25q^{29} + q^{33} + 9q^{34} + q^{38} \right. \\ & + 9q^{39} + 81q^{46} + 81q^{52} + 25q^{54} + 25q^{56} + 49q^{57} + q^{58} + 25q^{61} + 26q^{62} + 90q^{64} + q^{66} + 2q^{68} \\ & + 49q^{71} + 25q^{72} + q^{75} + q^{76} + 25q^{77} + 49q^{78} + 9q^{81} + 9q^{82} + q^{86} + 169q^{87} + 9q^{88} + q^{89} \\ & \left. + 9q^{90} + 169q^{95} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (2, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{71/8}}{2} \left(1 + q^5 + 9q^9 + q^{12} + 9q^{16} + 25q^{21} + 9q^{25} + q^{26} + q^{28} + 25q^{30} + q^{35} + 49q^{36} \right. \\ & + q^{39} + 25q^{41} + 9q^{43} + 9q^{45} + 49q^{47} + q^{48} + q^{50} + 90q^{54} + 9q^{58} + 49q^{60} + 25q^{63} \\ & + 25q^{65} + 81q^{67} + q^{68} + 9q^{69} + 9q^{71} + q^{72} + 121q^{75} + 25q^{76} + q^{78} + 25q^{80} \\ & \left. + q^{81} + 81q^{82} + 49q^{86} + 49q^{88} + q^{89} + 121q^{90} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (0, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{39/8}}{2} \left(1 + 9q^2 + q^3 + 25q^{17} + 49q^{21} + 2q^{22} + 25q^{24} + 9q^{26} + q^{27} + 18q^{28} + 2q^{33} + 9q^{39} \right. \\ & + 81q^{46} + 121q^{52} + 50q^{55} + 81q^{57} + 2q^{60} + 49q^{61} + 98q^{63} + 25q^{64} + 9q^{66} + q^{67} + q^{68} \\ & \left. + 68q^{70} + 18q^{76} + 49q^{78} + 2q^{79} + 2q^{81} + 169q^{87} + 18q^{89} + 225q^{95} \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{45/4}}{2} \left(1+q^3+q^6+9q^{10}+9q^{14}+9q^{18}+25q^{23}+26q^{28}+q^{31}+25q^{33}+q^{34}+q^{36} \right. \\ & + 50q^{39}+q^{42}+49q^{45}+9q^{46}+9q^{49}+49q^{51}+9q^{53}+9q^{55}+81q^{58}+9q^{59}+9q^{62}+81q^{65} \\ & + 25q^{67}+25q^{70}+82q^{72}+25q^{75}+25q^{77}+q^{78}+121q^{80}+q^{81}+25q^{82}+q^{83}+26q^{85}+122q^{88} \left. \right) \\ & + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (1, 2)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{47/4}}{2} \left(9+9q^3+q^5+9q^9+q^{13}+q^{17}+49q^{23}+49q^{28}+9q^{29}+9q^{31}+25q^{32}+9q^{35} \right. \\ & + 9q^{36}+q^{37}+49q^{38}+q^{42}+34q^{44}+9q^{49}+25q^{50}+q^{52}+q^{53}+122q^{58}+q^{60}+121q^{65}+49q^{68} \\ & + 49q^{70}+81q^{71}+9q^{74}+49q^{76}+49q^{77}+9q^{78}+121q^{79}+25q^{80}+9q^{83}+9q^{84}+25q^{85}+81q^{87} \left. \right) \\ & + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (1, 3)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{13/4}}{2} \left(1+9q^6+q^{10}+25q^{15}+q^{19}+q^{20}+9q^{21}+49q^{27}+q^{30}+9q^{33}+9q^{34}+25q^{35} \right. \\ & + q^{39}+81q^{42}+q^{45}+9q^{46}+25q^{50}+25q^{51}+49q^{52}+q^{54}+q^{56}+9q^{58}+121q^{60} \\ & + q^{63}+9q^{64}+25q^{65}+49q^{70}+49q^{71}+81q^{72}+q^{74}+q^{75}+9q^{76}+9q^{78}+25q^{80} \\ & + 169q^{81}+q^{84}+9q^{85}+25q^{86}+49q^{87}+81q^{93}+81q^{94}+121q^{95}+q^{96} \left. \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 1)$ $\mu_2 = (2, 2)$, we have:

$$\begin{aligned} \Psi_\mu = & -\frac{q^{15/8}}{2} \left(1+9q^{10}+25q^{13}+9q^{15}+q^{16}+2q^{18}+q^{27}+49q^{33}+81q^{38}+18q^{40}+49q^{42}+25q^{45} \right. \\ & + 59q^{47}+q^{48}+2q^{52}+18q^{53}+2q^{58}+25q^{60}+121q^{68}+2q^{69}+169q^{75}+98q^{79}+121q^{81} \\ & + 99q^{86}+162q^{88}+49q^{90}+25q^{93}+9q^{94}+9q^{95}+99q^{96}+50q^{97} \left. \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 2)$ $\mu_2 = (0, 1)$, we have:

$$\Psi_\mu = 16q^{47/2} \left(1+q^8+q^{18}+4q^{30}+q^{38}+q^{40}+4q^{42}+q^{50}+q^{54}+4q^{56}+q^{66}+q^{68}+9q^{72} \right) + \dots$$

For $\mu_1 = (0, 2)$ $\mu_2 = (1, 0)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{81/8} \left(1+q^6+5q^{13}+q^{21}+9q^{22}+q^{27}+q^{30}+4q^{31}+9q^{32}+q^{39}+q^{40}+25q^{43}+4q^{50} \right. \\ & + q^{51}+q^{52}+4q^{54}+9q^{55}+25q^{56}+q^{63}+9q^{65}+q^{66}+9q^{68}+16q^{69}+26q^{70}+q^{76} \\ & + 4q^{78}+10q^{81}+13q^{82}+61q^{85}+q^{88} \left. \right) + \dots \end{aligned}$$

For $\mu_1 = (0, 2)$ $\mu_2 = (1, 1)$, we have:

$$\begin{aligned} \Psi_\mu = & 2q^{101/8} \left(1+4q^7+q^{14}+4q^{15}+13q^{24}+q^{29}+q^{33}+4q^{34}+16q^{35}+4q^{42}+4q^{45}+9q^{46} \right. \\ & + 16q^{47}+q^{53}+4q^{56}+5q^{57}+41q^{60}+9q^{69}+4q^{70}+4q^{71}+9q^{73}+17q^{74} \\ & + 36q^{75}+q^{82}+4q^{84}+17q^{86}+4q^{87} \left. \right) + \dots \end{aligned}$$

Acknowledgments

We thank Sergey Alexandrov and Boris Pioline for useful discussions and comments on the draft. The authors are supported by Laureate Award 15175 “Modularity in Quantum Field Theory and Gravity” of the Irish Research Council.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.M. Maldacena, J. Michelson and A. Strominger, *Anti-de Sitter fragmentation*, *JHEP* **02** (1999) 011 [[hep-th/9812073](#)] [[INSPIRE](#)].
- [2] A. Maloney, M. Spradlin and A. Strominger, *Superconformal multiblack hole moduli spaces in four-dimensions*, *JHEP* **04** (2002) 003 [[hep-th/9911001](#)] [[INSPIRE](#)].
- [3] F. Denef, *Supergravity flows and D-brane stability*, *JHEP* **08** (2000) 050 [[hep-th/0005049](#)] [[INSPIRE](#)].
- [4] F. Denef, B.R. Greene and M. Raugas, *Split attractor flows and the spectrum of BPS D-branes on the quintic*, *JHEP* **05** (2001) 012 [[hep-th/0101135](#)] [[INSPIRE](#)].
- [5] I. Bena, C.-W. Wang and N.P. Warner, *Mergers and typical black hole microstates*, *JHEP* **11** (2006) 042 [[hep-th/0608217](#)] [[INSPIRE](#)].
- [6] F. Denef and G.W. Moore, *Split states, entropy enigmas, holes and halos*, *JHEP* **11** (2011) 129 [[hep-th/0702146](#)] [[INSPIRE](#)].
- [7] M.C.N. Cheng and E. Verlinde, *Dying Dyons Don't Count*, *JHEP* **09** (2007) 070 [[arXiv:0706.2363](#)] [[INSPIRE](#)].
- [8] J. de Boer, S. El-Showk, I. Messamah and D. Van den Bleeken, *Quantizing $N = 2$ Multicenter Solutions*, *JHEP* **05** (2009) 002 [[arXiv:0807.4556](#)] [[INSPIRE](#)].
- [9] J. Manschot, B. Pioline and A. Sen, *Wall Crossing from Boltzmann Black Hole Halos*, *JHEP* **07** (2011) 059 [[arXiv:1011.1258](#)] [[INSPIRE](#)].
- [10] A. Dabholkar, S. Murthy and D. Zagier, *Quantum Black Holes, Wall Crossing, and Mock Modular Forms*, [arXiv:1208.4074](#) [[INSPIRE](#)].
- [11] I. Bena, M. Berkooz, J. de Boer, S. El-Showk and D. Van den Bleeken, *Scaling BPS Solutions and pure-Higgs States*, *JHEP* **11** (2012) 171 [[arXiv:1205.5023](#)] [[INSPIRE](#)].
- [12] J.M. Maldacena, A. Strominger and E. Witten, *Black hole entropy in M-theory*, *JHEP* **12** (1997) 002 [[hep-th/9711053](#)] [[INSPIRE](#)].
- [13] D. Gaiotto, A. Strominger and X. Yin, *The M5-Brane Elliptic Genus: Modularity and BPS States*, *JHEP* **08** (2007) 070 [[hep-th/0607010](#)] [[INSPIRE](#)].
- [14] J. de Boer, M.C.N. Cheng, R. Dijkgraaf, J. Manschot and E. Verlinde, *A Farey Tail for Attractor Black Holes*, *JHEP* **11** (2006) 024 [[hep-th/0608059](#)] [[INSPIRE](#)].
- [15] P. Kraus and F. Larsen, *Partition functions and elliptic genera from supergravity*, *JHEP* **01** (2007) 002 [[hep-th/0607138](#)] [[INSPIRE](#)].
- [16] S. Alexandrov, J. Manschot and B. Pioline, *D3-instantons, Mock Theta Series and Twistors*, *JHEP* **04** (2013) 002 [[arXiv:1207.1109](#)] [[INSPIRE](#)].

- [17] S. Alexandrov, J. Manschot, D. Persson and B. Pioline, *Quantum hypermultiplet moduli spaces in $N = 2$ string vacua: a review*, *Proc. Symp. Pure Math.* **90** (2015) 181 [[arXiv:1304.0766](#)] [[INSPIRE](#)].
- [18] S. Alexandrov, S. Banerjee, J. Manschot and B. Pioline, *Multiple $D3$ -instantons and mock modular forms I*, *Commun. Math. Phys.* **353** (2017) 379 [[arXiv:1605.05945](#)] [[INSPIRE](#)].
- [19] A. Gholampour and A. Sheshmani, *Donaldson-Thomas invariants of 2-dimensional sheaves inside threefolds and modular forms*, *Adv. Math.* **326** (2018) 79 [[arXiv:1309.0050](#)] [[INSPIRE](#)].
- [20] A. Gholampour, A. Sheshmani and R. Thomas, *Counting curves on surfaces in Calabi-Yau 3-folds*, *Math. Ann.* **360** (2014) 67 [[arXiv:1309.0051](#)] [[INSPIRE](#)].
- [21] V. Bouchard, T. Creutzig, D.-E. Diaconescu, C. Doran, C. Quigley and A. Sheshmani, *Vertical $D4$ - $D2$ - $D0$ Bound States on $K3$ Fibrations and Modularity*, *Commun. Math. Phys.* **350** (2017) 1069 [[arXiv:1601.04030](#)] [[INSPIRE](#)].
- [22] S. Feyzbakhsh and R.P. Thomas, *Curve counting and S -duality*, [arXiv:2007.03037](#) [[INSPIRE](#)].
- [23] S. Feyzbakhsh and R.P. Thomas, *Rank r DT theory from rank 0*, [arXiv:2103.02915](#) [[INSPIRE](#)].
- [24] E. Andriyash and G.W. Moore, *Ample $D4$ - $D2$ - $D0$ Decay*, [arXiv:0806.4960](#) [[INSPIRE](#)].
- [25] J. Manschot, *Stability and duality in $N = 2$ supergravity*, *Commun. Math. Phys.* **299** (2010) 651 [[arXiv:0906.1767](#)] [[INSPIRE](#)].
- [26] J. Manschot, *Wall-crossing of $D4$ -branes using flow trees*, *Adv. Theor. Math. Phys.* **15** (2011) 1 [[arXiv:1003.1570](#)] [[INSPIRE](#)].
- [27] J. de Boer, F. Denef, S. El-Showk, I. Messamah and D. Van den Bleeken, *Black hole bound states in $AdS_3 \times S^2$* , *JHEP* **11** (2008) 050 [[arXiv:0802.2257](#)] [[INSPIRE](#)].
- [28] S. Alexandrov, S. Banerjee, J. Manschot and B. Pioline, *Multiple $D3$ -instantons and mock modular forms II*, *Commun. Math. Phys.* **359** (2018) 297 [[arXiv:1702.05497](#)] [[INSPIRE](#)].
- [29] S. Alexandrov and B. Pioline, *Black holes and higher depth mock modular forms*, *Commun. Math. Phys.* **374** (2019) 549 [[arXiv:1808.08479](#)] [[INSPIRE](#)].
- [30] S. Alexandrov, J. Manschot and B. Pioline, *S -duality and refined BPS indices*, *Commun. Math. Phys.* **380** (2020) 755 [[arXiv:1910.03098](#)] [[INSPIRE](#)].
- [31] D. Robles-Llana, M. Roček, F. Saueressig, U. Theis and S. Vandoren, *Nonperturbative corrections to 4D string theory effective actions from $SL(2, Z)$ duality and supersymmetry*, *Phys. Rev. Lett.* **98** (2007) 211602 [[hep-th/0612027](#)] [[INSPIRE](#)].
- [32] S. Alexandrov, B. Pioline, F. Saueressig and S. Vandoren, *D -instantons and twistors*, *JHEP* **03** (2009) 044 [[arXiv:0812.4219](#)] [[INSPIRE](#)].
- [33] S. Alexandrov, *D -instantons and twistors: Some exact results*, *J. Phys. A* **42** (2009) 335402 [[arXiv:0902.2761](#)] [[INSPIRE](#)].
- [34] S. Zwegers, *Mock Theta Functions*, Ph.D. Thesis, Utrecht University (2008). [[arXiv:0807.4834](#)] [[INSPIRE](#)].
- [35] D. Zagier, *Ramanujan's mock theta functions and their applications (after Zwegers and Ono-Bringmann)*, *Astérisque* **326** (2009) Exp. No. 986.
- [36] S. Alexandrov, S. Banerjee, J. Manschot and B. Pioline, *Indefinite theta series and generalized error functions*, *Selecta Math.* **24** (2018) 3927 [[arXiv:1606.05495](#)] [[INSPIRE](#)].
- [37] C. Nazaroglu, *r -Tuple Error Functions and Indefinite Theta Series of Higher-Depth*, *Commun. Num. Theor. Phys.* **12** (2018) 581 [[arXiv:1609.01224](#)] [[INSPIRE](#)].

- [38] S. Kudla, *Theta integrals and generalized error functions*, *Manuscripta Math.* **155** (2016) 303 [[arXiv:1608.03534](#)].
- [39] J. Funke and S. Kudla, *On some incomplete theta integrals*, *Compos. Math.* **155** (2019) 1711.
- [40] C. Vafa and E. Witten, *A Strong coupling test of S duality*, *Nucl. Phys. B* **431** (1994) 3 [[hep-th/9408074](#)] [[INSPIRE](#)].
- [41] J.A. Minahan, D. Nemeschansky, C. Vafa and N.P. Warner, *E strings and $N = 4$ topological Yang-Mills theories*, *Nucl. Phys. B* **527** (1998) 581 [[hep-th/9802168](#)] [[INSPIRE](#)].
- [42] J. Manschot, *BPS invariants of $\mathcal{N} = 4$ gauge theory on Hirzebruch surfaces*, *Commun. Num. Theor. Phys.* **6** (2012) 497 [[arXiv:1103.0012](#)] [[INSPIRE](#)].
- [43] J. Manschot, *Vafa-Witten Theory and Iterated Integrals of Modular Forms*, *Commun. Math. Phys.* **371** (2019) 787 [[arXiv:1709.10098](#)] [[INSPIRE](#)].
- [44] A. Dabholkar, P. Putrov and E. Witten, *Duality and Mock Modularity*, *SciPost Phys.* **9** (2020) 072 [[arXiv:2004.14387](#)] [[INSPIRE](#)].
- [45] S. Alexandrov, *Vafa-Witten invariants from modular anomaly*, *Commun. Num. Theor. Phys.* **15** (2021) 149 [[arXiv:2005.03680](#)] [[INSPIRE](#)].
- [46] S. Alexandrov, *Rank N Vafa-Witten invariants, modularity and blow-up*, [arXiv:2006.10074](#) [[INSPIRE](#)].
- [47] G. Bonelli, F. Fucito, J.F. Morales, M. Ronzani, E. Sysoeva and A. Tanzini, *Gauge theories on compact toric manifolds*, *Lett. Math. Phys.* **111** (2021) 77 [[arXiv:2007.15468](#)] [[INSPIRE](#)].
- [48] J. Manschot and G.W. Moore, *Topological correlators of $SU(2)$, $\mathcal{N} = 2^*$ SYM on four-manifolds*, [arXiv:2104.06492](#) [[INSPIRE](#)].
- [49] S.-J. Lee, Z.-L. Wang and P. Yi, *Quiver Invariants from Intrinsic Higgs States*, *JHEP* **07** (2012) 169 [[arXiv:1205.6511](#)] [[INSPIRE](#)].
- [50] D. Anninos, T. Anous, P. de Lange and G. Konstantinidis, *Conformal quivers and melting molecules*, *JHEP* **03** (2015) 066 [[arXiv:1310.7929](#)] [[INSPIRE](#)].
- [51] J. Manschot, B. Pioline and A. Sen, *On the Coulomb and Higgs branch formulae for multi-centered black holes and quiver invariants*, *JHEP* **05** (2013) 166 [[arXiv:1302.5498](#)] [[INSPIRE](#)].
- [52] D. Mirfendereski, J. Raeymaekers and D. Van Den Bleeken, *Superconformal mechanics of AdS_2 D-brane boundstates*, *JHEP* **12** (2020) 176 [[arXiv:2009.07107](#)] [[INSPIRE](#)].
- [53] G. Beaujard, S. Mondal and B. Pioline, *Multi-centered black holes, scaling solutions and pure-Higgs indices from localization*, *SciPost Phys.* **11** (2021) 023 [[arXiv:2103.03205](#)] [[INSPIRE](#)].
- [54] J. de Boer, J. Manschot, K. Papadodimas and E. Verlinde, *The Chiral ring of AdS_3/CFT_2 and the attractor mechanism*, *JHEP* **03** (2009) 030 [[arXiv:0809.0507](#)] [[INSPIRE](#)].
- [55] M.-F. Vignéras, *Séries thêta des formes quadratiques indéfinies*, in *Modular Functions of One Variable VI*, Lecture Notes in Mathematics, vol. 627, Springer, Berlin, Heidelberg (1977) [[DOI](#)].
- [56] J. Manschot and G.W. Moore, *A Modern Farey Tail*, *Commun. Num. Theor. Phys.* **4** (2010) 103 [[arXiv:0712.0573](#)] [[INSPIRE](#)].
- [57] K. Bringmann, J. Kaszian and A. Milas, *Higher depth quantum modular forms, multiple eichler integrals, and \mathfrak{sl}_3 false theta functions*, *Res. Math. Sci.* **6** (2019) 20 [[arXiv:1704.06891](#)].
- [58] B. Pioline, *Wall-crossing made smooth*, *JHEP* **04** (2015) 092 [[arXiv:1501.01643](#)] [[INSPIRE](#)].

- [59] S.S. Kudla and J.J. Millson, *The theta correspondence and harmonic forms. I*, *Math. Ann.* **274** (1986) 353.
- [60] N. Afkhami-Jeddi, H. Cohn, T. Hartman and A. Tajdini, *Free partition functions and an averaged holographic duality*, *JHEP* **01** (2021) 130 [[arXiv:2006.04839](#)] [[INSPIRE](#)].
- [61] A. Maloney and E. Witten, *Averaging over Narain moduli space*, *JHEP* **10** (2020) 187 [[arXiv:2006.04855](#)] [[INSPIRE](#)].
- [62] S. Ferrara, R. Kallosh and A. Strominger, *$N = 2$ extremal black holes*, *Phys. Rev. D* **52** (1995) R5412 [[hep-th/9508072](#)] [[INSPIRE](#)].
- [63] S. Ferrara and R. Kallosh, *Supersymmetry and attractors*, *Phys. Rev. D* **54** (1996) 1514 [[hep-th/9602136](#)] [[INSPIRE](#)].
- [64] M. Shmakova, *Calabi-Yau black holes*, *Phys. Rev. D* **56** (1997) 540 [[hep-th/9612076](#)] [[INSPIRE](#)].
- [65] F. Denef, *Quantum quivers and Hall/hole halos*, *JHEP* **10** (2002) 023 [[hep-th/0206072](#)] [[INSPIRE](#)].
- [66] B. Bates and F. Denef, *Exact solutions for supersymmetric stationary black hole composites*, *JHEP* **11** (2011) 127 [[hep-th/0304094](#)] [[INSPIRE](#)].
- [67] J. Manschot, B. Pioline and A. Sen, *The Coulomb Branch Formula for Quiver Moduli Spaces*, [arXiv:1404.7154](#) [[INSPIRE](#)].
- [68] J. Manschot, B. Pioline and A. Sen, *A Fixed point formula for the index of multi-centered $N = 2$ black holes*, *JHEP* **05** (2011) 057 [[arXiv:1103.1887](#)] [[INSPIRE](#)].
- [69] S.-J. Lee and P. Yi, *Witten Index for Noncompact Dynamics*, *JHEP* **06** (2016) 089 [[arXiv:1602.03530](#)] [[INSPIRE](#)].
- [70] S. Alexandrov and B. Pioline, *Attractor flow trees, BPS indices and quivers*, *Adv. Theor. Math. Phys.* **23** (2019) 627 [[arXiv:1804.06928](#)] [[INSPIRE](#)].
- [71] H. Ooguri, A. Strominger and C. Vafa, *Black hole attractors and the topological string*, *Phys. Rev. D* **70** (2004) 106007 [[hep-th/0405146](#)] [[INSPIRE](#)].
- [72] J. Conway and N. Sloane, *Sphere packings, lattices and groups*, Springer (1999) [[DOI](#)].
- [73] T. Gannon, *Lattices and Superstrings*, Ph.D. Thesis, McGill University (1989).
- [74] T.-T. Lu and S.-H. Shiou, *Inverses of 2×2 Block Matrices*, *Comput. Math. Appl.* **43** (2002) 119.
- [75] J. Funke and S. Kudla, *The case of an N -gon*, [arXiv:2109.10979](#).
- [76] R. Minasian, G.W. Moore and D. Tsimpis, *Calabi-Yau black holes and $(0, 4)$ sigma models*, *Commun. Math. Phys.* **209** (2000) 325 [[hep-th/9904217](#)] [[INSPIRE](#)].
- [77] *Toric calabi-yau database*, <http://rossealtman.com/toriccy/>.
- [78] D. Maulik and R. Pandharipande, *Gromov-Witten theory and Noether-Lefschetz theory*, [arXiv:0705.1653](#).
- [79] D. Gaiotto and X. Yin, *Examples of $M5$ -Brane Elliptic Genera*, *JHEP* **11** (2007) 004 [[hep-th/0702012](#)] [[INSPIRE](#)].
- [80] A. Klemm, M. Kreuzer, E. Riegler and E. Scheidegger, *Topological string amplitudes, complete intersection Calabi-Yau spaces and threshold corrections*, *JHEP* **05** (2005) 023 [[hep-th/0410018](#)] [[INSPIRE](#)].