Heterotic strings on $(K3 \times T^2)/\mathbb{Z}_3$ and their dual Calabi-Yau threefolds

Andreas Banlaki,^a Aradhita Chattopadhyaya,^b Abhiram Kidambi,^{a,d}
Thorsten Schimannek^c and Maria Schimpf^a

^a*Institute for Theoretical Physics, TU Wien,
A-1040 Vienna, Austria*

^b*Department of Mathematics, Trinity College Dublin,
Dublin 2, Ireland*

^c*Faculty of Physics, University of Vienna,
A-1090, Vienna, Austria*

^d*Stanford Institute for Theoretical Physics, Stanford University,
Palo Alto, U.S.A.*

E-mail: banlaki@hep.itp.tuwien.ac.at, aradhita@maths.tcd.ie,
abhiram.kidambi@tuwien.ac.at, thorsten.schimannek@univie.ac.at,
mschimpf@hep.itp.tuwien.ac.at

ABSTRACT: In this paper we study compactifications of the $\mathcal{N} = 2$ heterotic $E_8 \times E_8$ string on $(K3 \times T^2)/\mathbb{Z}_3$ with various gauge backgrounds and calculate the topological couplings in the effective supergravity action that arise from one-loop amplitudes. We then identify candidates for dual type IIA compactifications on Calabi-Yau threefolds and compare the heterotic results with the corresponding topological string amplitudes. We find that the dual Calabi-Yau geometries are $K3$ fibrations that are also genus one fibered with three-sections. Moreover, we show that the intersection form on the polarization lattice of the $K3$ fibration has to be three times the intersection form on the Narain lattice $\Gamma^{1,1}$.

KEYWORDS: String Duality, Superstrings and Heterotic Strings, Topological Strings

ARXIV EPRINT: [1911.09697](https://arxiv.org/abs/1911.09697)

Contents

1	Introduction	1
2	The new supersymmetric index for CHL orbifolds	3
2.1	The twisted elliptic genus of $K3$	4
2.2	Calculating the massless spectrum	8
2.3	The new supersymmetric index	11
3	Heterotic computation of Gopakumar-Vafa invariants	14
3.1	Extraction of Gopakumar-Vafa invariants from F_g	15
3.2	Conifold singularities	17
4	Dual type IIA compactifications on Calabi-Yau threefolds	18
4.1	A brief review of heterotic/type II duality	18
4.2	Constructing Calabi-Yau duals	20
4.3	All genus checks via the modular bootstrap	27
5	Conclusions	30
A	Conventions for modular forms	31
B	Higgsing of the gauge group	32

1 Introduction

The four-dimensional $\mathcal{N} = 2$ effective supergravity actions that arise from heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_N$ and type II compactifications on Calabi-Yau threefolds contain the well-known supersymmetry protected couplings

$$S = \int F_g(y, \bar{y}) \cdot F_+^{2g-2} R_+^2, \quad (1.1)$$

where F_+ and R_+ respectively denote the self-dual parts of the graviphoton field strength and the Riemann tensor. In particular, the coefficients $F_g(y, \bar{y})$ depend only on the vector moduli. On the heterotic side, one of the vector moduli corresponds to the axio-dilaton and all terms (1.1) that do not involve the dilaton arise already at one-loop level. The one-loop amplitudes have been evaluated for compactifications with various embeddings of the gauge connection on $K3 \times T^2$ by [1–3] and receive contributions only from BPS states that are encoded in the so-called new supersymmetric index [4]. Subsequently this analysis has been extended to compactifications on $(K3 \times T^2)/\mathbb{Z}_N$ [5] and again the result is essentially encoded in the new supersymmetric index. The indices in turn had been

calculated for various compactifications with standard embedding of the spin connection into the gauge connection on $(K3 \times T^2)/\mathbb{Z}_N$ and some non-standard embeddings when $N = 2$ [6]. Calabi-Yau duals for some of the $N = 2$ cases have been proposed in [7–9].

In this paper we extend the calculation of the new supersymmetric index to compactifications on $(K3 \times T^2)/\mathbb{Z}_3$ with non-standard embeddings and predict the geometric data of the dual Calabi-Yau manifold. To this end we take an explicit realization of the orbifold limit of a $K3$ as $T^4/\mathbb{Z}_3$. A list of the possible gauge embeddings is provided in table 3 on page 10. For two embeddings the effective theories are such that the gauge group on a generic point of the hypermultiplet moduli space is maximally broken to $U(1)^3$. For those models one can hope to find a dual Calabi-Yau compactification with only three Kähler moduli and without a larger non-higgsable gauge group. Indeed we will be able to identify many candidates. At certain points in the moduli space there might be singularities as additional vector bosons become massless. The strength of these singularities and the extraction of the Gopakumar-Vafa invariants are discussed in section 3.

On the type IIA side one can identify $\lim_{y \rightarrow \infty} F_g(y, \bar{y})$ with the complex conjugate of the topological string free energy at genus g in the holomorphic limit [2, 10–12]. Following arguments from [9] that we review in section 4.1, one expects that for heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_3$ the dual Calabi-Yau manifolds are genus one fibered with three sections. Moreover, based on the prepotential of the effective supergravity action, one can argue that the Calabi-Yau should also exhibit a $K3$ fibration [13]. We thus systematically construct all $K3$ fibered Calabi-Yau threefolds with $h^{1,1} = 3$ that exhibit a genus one fibration with three-sections and are realized as hypersurfaces in toric ambient spaces. We then apply the modular bootstrap that has been extended to genus one fibrations with multi-sections in [9] and obtain all-genus results for the topological string amplitudes. This provides all order checks of the duality. A list of the Calabi-Yau threefolds is provided in table 6 on page 23.

Let us note that on the heterotic side, the quotient acts as an order three symplectic automorphism on $K3$ together with a one-third shift along T^2 . Symplectic automorphisms of $K3$ manifolds have been classified and form subgroups of the Mathieu group M_{23} [14]. In particular, there is up to conjugacy only one automorphism of order three and this corresponds to the so-called 3A class. It turns out that the automorphism group of a non-linear sigma model into $K3$ is actually larger [15] and contains another element of order three. However, we will restrict to the geometric case.

The paper is structured as follows. In section 2 we study heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_3$ where the $K3$ is realized in the orbifold limit as $T^4/\mathbb{Z}_3$. We first calculate the twisted elliptic genus of $K3$ with order three twists by directly evaluating the corresponding trace. This confirms earlier results from the literature that were based on indirect arguments and determines the new supersymmetric index for the standard embedding. We then evaluate the new supersymmetric index for various non-standard embeddings.

In section 3 we describe how the new supersymmetric index encodes the topological couplings and extract predictions for the Gopakumar-Vafa invariants of the dual Calabi-Yau compactification spaces. We also discuss the singular behaviour of the generating functions at points where additional vector states become massless.

In section 4 we review the relevant aspects of heterotic/type II duality and then systematically construct candidate Calabi-Yau duals with $h^{1,1} = 3$ that in particular exhibit a $K3$ fibration and a compatible genus one fibration with three-sections. We find generic expressions for the Gopakumar-Vafa invariants at genus zero that only depend on the Euler characteristic of the Calabi-Yau and the intersection form on the polarization lattice of the $K3$ fibration. From the structure of the invariants we can see that for Calabi-Yau duals to heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_3$, the intersection form on the polarization lattice of the $K3$ fibration has to be three times the intersection form on the Narain lattice $\Gamma^{1,1}$. Physically this condition can be related to the presence of T-duality on the heterotic side. We will then apply the modular bootstrap to obtain all genus results for the Gopakumar-Vafa invariants of low degree with respect to the base of the $K3$ fiber. We find matching invariants for both of the compactifications with non-standard embedding on the heterotic side that lead to a maximally higgsable gauge group and for fifteen other geometries we expect that a heterotic dual exists.

The paper is supplemented by two appendices. In appendix A we summarize the definitions of the various modular and Jacobi forms that appear throughout the discussion. Finally, in appendix B, we discuss in some detail the higgsing of the gauge group for one of the models with non-standard embedding and give the necessary branching rules.

Acknowledgments

We thank Justin David, Rajesh Gopakumar, Timm Wrase, Harald Skarke, Cesar Fierro Cota and Albrecht Klemm for helpful discussions. The work of T.S. is supported by the Austrian Science Fund (FWF): P30904-N27. The work of A.B., A.K. and M.S. is supported by the Austrian Science Fund (FWF): P 28552. A.C. thanks the Centre for High Energy Physics at the Indian Institute of Science where a large portion of her work was done. A.C. is supported by the IRC Laurette fellowship. The work of A.B., A.C., A.K. and M.S. is supported by the OeAD ‘Scientific & Technological Cooperation with India’ grant Project IN 27/2018 titled ‘Mathieu moonshine and $\mathcal{N} = 2$ Heterotic - Type II string duality’. The work of A.K. is further supported the FWF project no. W1252-N27 and the Austrian Marshall Plan Fellowship.

2 The new supersymmetric index for CHL orbifolds

In this section we compute the massless spectrum as well as the new supersymmetric index for compactifications of the heterotic $E_8 \times E_8$ string on $(K3 \times T^2)/\mathbb{Z}_3$. The generator g' of $\mathbb{Z}_3$ acts via an order three shift along one of the circles of T^2 together with a symplectic automorphism of $K3$. The latter corresponds to the unique order three subgroup of $M_{23} < M_{24}$ that is in the conjugacy class $3A$ of M_{24} .

We first compute the twisted elliptic genus of order $N = 3$ orbifolds of $K3$. The twisted elliptic genus is defined as the trace

$$F^{(r,s)}(\tau, z) = \frac{1}{N} \text{Tr}_{g'^r}((-1)^{F_L + \bar{F}_R} g'^s e^{2\pi i z F_L} q^{L_0} \bar{q}^{\bar{L}_0}), \quad (2.1)$$

over the Ramond-Ramond sector of a non-linear sigma model into $K3$. For $N = 3$ the result can be expressed in terms of modular forms for $\Gamma_1(3)$ and reads

$$\begin{aligned} F^{(0,0)}(\tau, z) &= \frac{8}{3}A(\tau, z), & F^{(0,1)}(\tau, z) &= \frac{2}{3}A(\tau, z) - \frac{1}{2}B(\tau, z)\mathcal{E}_3(\tau), \\ F^{(r,rk)}(\tau, z) &= \frac{2}{3}A(\tau, z) + \frac{1}{6}B(\tau, z)\mathcal{E}_3\left(\frac{\tau+k}{3}\right), \end{aligned} \quad (2.2)$$

where

$$A(\tau, z) = \frac{\theta_2^2(\tau, z)}{\theta_2^2(\tau, 0)} + \frac{\theta_3^2(\tau, z)}{\theta_3^2(\tau, 0)} + \frac{\theta_4^2(\tau, z)}{\theta_4^2(\tau, 0)}, \quad B(\tau, z) = \frac{\theta_1^2(\tau, z)}{\eta^6(\tau)}, \quad (2.3)$$

while $r = 1, 2$ and $rk := rk \bmod 3$. This result was essentially bootstrapped in [16] and we reproduce it by directly evaluating the trace after taking a $T^4/\mathbb{Z}_3$ orbifold limit of the $K3$. In particular, it encodes the new supersymmetric index for compactifications on $(K3 \times T^2)/\mathbb{Z}_3$ with standard embedding, as predicted in [17]. The explicit calculation serves as a check that our construction of the orbifold is correct.

Next we move on to calculate the massless spectrum of the heterotic theory compactified on $(K3 \times T^2)/\mathbb{Z}_3$ for various gauge backgrounds. The realization of the $K3$ as $T^4/\mathbb{Z}_3$ allows for five inequivalent embeddings of $\mathbb{Z}_3$ into the gauge group which lead to an anomaly free theory [3, 18, 19]. They can be represented by *shift vectors* $V = (\gamma, \tilde{\gamma})$ where 3γ and $3\tilde{\gamma}$ take values in the two E_8 lattices. For the standard embedding, the difference between the number of hypermultiplets and vectormultiplets was computed in [5]. Among the four non-standard embeddings we find two models where the spectrum enables us to maximally higgs the gauge group.

Then we compute the new supersymmetric index of the heterotic theory compactified with standard embedding, where one of the E_8 groups remains unbroken. The new supersymmetric index is given by

$$\mathcal{Z}_{\text{new}}(\tau, \bar{\tau}) = \frac{1}{\eta^2(\tau)} \text{Tr}_R \left((-1)^F q^{L_0 - c/24} \bar{q}^{\bar{L}_0 - \bar{c}/24} \right), \quad (2.4)$$

where $q = e^{2\pi i \tau}$ and the trace is again performed over the Ramond sector of the internal CFT which has central charge $(c, \bar{c}) = (22, 9)$. The right moving worldsheet fermion number F is given by the sum of the fermion number on T^2 and that of $K3$. Our result matches that obtained by general arguments in [6] which acts as a further check to our orbifold construction.

We will then proceed to calculate the new supersymmetric index $\mathcal{Z}_{\text{new}}$ for the four non-standard embeddings. In each case of standard or non-standard embedding we can show that the difference between the number of hypermultiplets and vectormultiplets $N_h - N_v$ computed from the new supersymmetric index matches with the result obtained from explicitly counting the degrees of freedom.

2.1 The twisted elliptic genus of $K3$

We will now proceed to compute the twisted elliptic genus of the 3A orbifold of $K3$ where $K3$ is realized as $T^4/\mathbb{Z}_3$. Let us denote the coordinates on T^4 by $(z_1, z_2) \in \mathbb{C}^2/\mathbb{Z}^2$ with

periodicity

$$z_i \sim z_i + n_1 e_1 + n_2 e_2, \quad e_1 = e^{\frac{2\pi i}{3}}, e_2 = 1, \quad (n_1, n_2) \in \mathbb{Z}^2, \quad (2.5)$$

i.e. the A- and B-cycles of the torus e_1, e_2 generate the root lattice of $SU(3)$. Then the $\mathbb{Z}_3$ orbifold of T^4 which results in the orbifold limit of $K3$ can be implemented as

$$g^s : (z_1, z_2) \mapsto \left(e^{2\pi i s/3} z_1, e^{-2\pi i s/3} z_2 \right). \quad (2.6)$$

We also need to specify the action of the generator g' of the $\mathbb{Z}_3$ that is used in the CHL quotient $(K3 \times T^2)/\mathbb{Z}_3$. We consider the g' action as a one-third shift along the A-cycle and a two-third shift along the B-cycle,

$$g' : (z_1, z_2) \mapsto \left(z_1 + \frac{1}{3}e_1 + \frac{2}{3}e_2, z_2 \right), \quad (2.7)$$

with

$$g'^2 : (z_1, z_2) \mapsto \left(z_1 + \frac{2}{3}e_1 + \frac{1}{3}e_2, z_2 \right). \quad (2.8)$$

We first compute the twisted elliptic genus of the corresponding orbifold of $K3$ and show that it matches with the results obtained in [16, 20–25]. This serves as a check of our construction.

The twisted elliptic genus in the (r, s) -sector is given by the index

$$F^{(r,s)}(\tau, z) = \frac{1}{9} \sum_{a,b=0}^2 \text{Tr}_{g^a, g'^r} \left((-1)^{F_L + \bar{F}_R} g^b g'^s e^{2\pi i z F_L} q^{L_0} \bar{q}^{\bar{L}_0} \right). \quad (2.9)$$

The overall factor of $1/9$ comes from the projection of g, g' , both of which are $\mathbb{Z}_3$ orbifolds. The trace is taken over the Ramond-Ramond sector and F_L, F_R are the left- and right-moving fermion numbers. In defining the index we have suppressed the shifts $L_0 - c/24$ and $\bar{L}_0 - \bar{c}/24$, where $(c, \bar{c}) = (6, 6)$.

Let us now define the trace

$$\mathcal{F}(a, r; b, s) = \frac{1}{9} \text{Tr}_{g^a, g'^r} \left((-1)^{F_L + \bar{F}_R} g^b g'^s e^{2\pi i z F_L} q^{L_0} \bar{q}^{\bar{L}_0} \right). \quad (2.10)$$

To evaluate each of the above sectors in the twisted elliptic genus we will need the fixed points under the elements $g^a g'^r$, a, r being the twists in g and g' respectively. We also require to know which elements preserve these fixed points. We summarize the result in table 1.

Now we evaluate the different sectors of the twisted elliptic genus. The sector $(0, 0)$ comes with no twists in g' or insertions of g' and so the trace reduces to

$$F^{(0,0)}(\tau, z) = \frac{1}{3} Z_{K3}(\tau, z) = \frac{8}{3} A(\tau, z). \quad (2.11)$$

where Z_{K3} is the elliptic genus of $K3$.

	Fixed points	g'	g'^2	g	g^2	$g'g$	$g'g^2$	g'^2g	g'^2g^2
g	$(0,0), (2/3, 1/3), (1/3, 2/3)$	$\times$	$\times$	$\checkmark$	$\checkmark$	$\times$	$\times$	$\times$	$\times$
g^2	$(0,0), (2/3, 1/3), (1/3, 2/3)$	$\times$	$\times$	$\checkmark$	$\checkmark$	$\times$	$\times$	$\times$	$\times$
gg'	$(2/3,0), (0,2/3), (1/3,1/3)$	$\times$	$\times$	$\times$	$\times$	$\checkmark$	$\times$	$\times$	$\checkmark$
g^2g'	$(1/3,0), (0,1/3), (2/3,2/3)$	$\times$	$\times$	$\times$	$\times$	$\times$	$\checkmark$	$\checkmark$	$\times$
gg'^2	$(1/3,0), (0,1/3), (2/3,2/3)$	$\times$	$\times$	$\times$	$\times$	$\times$	$\checkmark$	$\checkmark$	$\times$
$g^2g'^2$	$(2/3,0), (0,2/3), (1/3,1/3)$	$\times$	$\times$	$\times$	$\times$	$\checkmark$	$\times$	$\times$	$\checkmark$

Table 1. Rows list the properties of points that are fixed under the action of the element $g^a g'^b$ in the first column. A $\times$ denotes that the point moves under the action of the corresponding element $g'^a g^b$ in the top row, while $\checkmark$ indicates that the point remains fixed. Positions are given as multiples of the A- and B-cycle e_1 and e_2 .

Let us now look into the untwisted sectors of the twisted elliptic genus. In the $(0,1)$ and $(0,2)$ sectors we obtain from table 1, that a single insertion of g' does not preserve any of the fixed points. Hence we get,

$$\mathcal{F}(a,0;b,1) = \mathcal{F}(a,0;b,2) = 0, \quad \text{for } a = 1, 2. \quad (2.12)$$

Evaluating the trace in the untwisted sector we see the contributions are

$$\begin{aligned} \mathcal{F}(0,0;0,s) &= 0, \\ \mathcal{F}(0,0;1,1) = \mathcal{F}(0,0;1,2) &= \frac{\theta_1(z + \frac{1}{3}, \tau) \theta_1(-z + \frac{1}{3})}{\theta_1^2(\frac{1}{3}, \tau)}, \\ \mathcal{F}(0,0;2,1) = \mathcal{F}(0,0;2,2) &= \frac{\theta_1(z + \frac{2}{3}, \tau) \theta_1(-z + \frac{2}{3})}{\theta_1^2(\frac{2}{3}, \tau)}. \end{aligned} \quad (2.13)$$

The coefficients (numerical) in the traces come from the contribution of fermionic zero modes. There are 9 right moving fermionic zero modes when g or g^2 are inserted in the trace. It can be easily shown by considering the q expansions that

$$\begin{aligned} \mathcal{F}(0,0;1,1) + \mathcal{F}(0,0;2,1) &= \mathcal{F}(0,0;1,2) + \mathcal{F}(0,0;2,2) \\ &= \frac{2}{3}A(\tau, z) - \frac{1}{2}B(\tau, z)\mathcal{E}_3(\tau) = F_{3A}^{(0,1)}(\tau, z) = F_{3A}^{(0,2)}(\tau, z), \end{aligned} \quad (2.14)$$

as predicted in [16, 20–24].

The other sectors can be obtained from $F^{(0,1)}$ through the modular transformation property of the twisted elliptic genus given by,

$$F^{(r,s)}\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) = \exp\left(2\pi i \frac{cz^2}{c\tau + d}\right) F^{(cs+ar, ds+br)}(\tau, z), \quad (2.15)$$

where $a, b, c, d \in \mathbb{Z}$, $ad - bc = 1$. However, with the knowledge of the fixed points we can directly compute the twisted sectors and re-evaluate the results of the twisted elliptic genus in each sector.

In the sectors twisted by g' without any insertion of g' we have the components of the twisted elliptic genus given by

$$F^{(1,0)} = \sum_{a,b=0}^2 \mathcal{F}(a, 1, b, 0).$$

Now $\mathcal{F}(0, 1, 1, 0) = 0 = \mathcal{F}(0, 1, 2, 0)$ and also $\mathcal{F}(1, 1, 1, 0) = 0 = \mathcal{F}(2, 1, 1, 0)$ due to the absence of any fixed point in the twisted sector (table 1). The same argument holds for $\mathcal{F}(1, 1, 2, 0) = 0 = \mathcal{F}(2, 1, 2, 0)$. Also $\mathcal{F}(0, 1, 0, 0) = 0$ due to the presence of zero modes of the right moving fermions. Therefore the only contributions are

$$F^{(1,0)} = \mathcal{F}(1, 1, 0, 0) + \mathcal{F}(2, 1, 0, 0).$$

Computing the trace we get

$$\begin{aligned} \mathcal{F}(1, 1, 1, 1) &= \frac{\theta_1(z + \frac{\tau}{3}, \tau) \theta_1(-z + \frac{\tau}{3})}{\theta_1^2(\frac{\tau}{3}, \tau)}, \\ \mathcal{F}(2, 1, 0, 0) &= \frac{\theta_1(z + \frac{2\tau}{3}, \tau) \theta_1(-z + \frac{2\tau}{3})}{\theta_1^2(\frac{2\tau}{3}, \tau)}, \\ F^{(1,0)} &= \frac{2}{3} A(\tau, z) + \frac{1}{6} B(\tau, z) \mathcal{E}_3\left(\frac{\tau}{3}\right). \end{aligned} \quad (2.16)$$

If we insert a g' in the trace we have the non-vanishing components in the twisted elliptic genus given by

$$\begin{aligned} \mathcal{F}(1, 1, 1, 1) &= \frac{\theta_1(z + \frac{\tau+1}{3}, \tau) \theta_1(-z + \frac{\tau+1}{3})}{\theta_1^2(\frac{\tau+1}{3}, \tau)}, \\ \mathcal{F}(2, 1, 2, 1) &= \frac{\theta_1(z + \frac{2\tau+2}{3}, \tau) \theta_1(-z + \frac{2\tau+2}{3})}{\theta_1^2(\frac{2\tau+2}{3}, \tau)}, \\ F^{(1,1)} &= \frac{2}{3} A(\tau, z) + \frac{1}{6} B(\tau, z) \mathcal{E}_3\left(\frac{\tau+1}{3}\right). \end{aligned} \quad (2.17)$$

The rest of the sectors vanish as all the fixed points change position. Similarly for the insertion of g'^2 we have

$$\begin{aligned} \mathcal{F}(1, 1, 1, 1) &= \frac{\theta_1(z + \frac{\tau+2}{3}, \tau) \theta_1(-z + \frac{\tau+2}{3})}{\theta_1^2(\frac{\tau+2}{3}, \tau)}, \\ \mathcal{F}(2, 1, 2, 1) &= \frac{\theta_1(z + \frac{2\tau+1}{3}, \tau) \theta_1(-z + \frac{2\tau+1}{3})}{\theta_1^2(\frac{2\tau+1}{3}, \tau)}, \\ F^{(1,2)} &= \frac{2}{3} A(\tau, z) + \frac{1}{6} B(\tau, z) \mathcal{E}_3\left(\frac{\tau+2}{3}\right). \end{aligned} \quad (2.18)$$

A very similar argument holds for the g'^2 twisted sector and the results can be summarized as:

$$\begin{aligned} F^{(2,0)} &= \frac{2}{3} A(\tau, z) + \frac{1}{6} B(\tau, z) \mathcal{E}_3\left(\frac{\tau}{3}\right), \\ F^{(2,1)} &= \frac{2}{3} A(\tau, z) + \frac{1}{6} B(\tau, z) \mathcal{E}_3\left(\frac{\tau+2}{3}\right), \\ F^{(2,2)} &= \frac{2}{3} A(\tau, z) + \frac{1}{6} B(\tau, z) \mathcal{E}_3\left(\frac{\tau+1}{3}\right). \end{aligned} \quad (2.19)$$

We note that these results were obtained previously in [16, 20–24], however, the authors are not aware of any paper that explicitly evaluates the traces.

2.2 Calculating the massless spectrum

In this section we derive the massless spectrum of the heterotic string theory compactified on $(K3 \times T^2)/\mathbb{Z}_N$. To this end we consider again an orbifold limit of the $K3$ such that the compactification space is $[(T^4/\mathbb{Z}_3) \times T^2]/\mathbb{Z}_3$. The orbifold action on the six toroidal coordinates can be given by

$$g : (z_1, z_2, z_3) \mapsto (e^{2\pi i/3} z_1, e^{-2\pi i/3} z_2, z_3), \quad (2.20)$$

where z_3 is a coordinate on T^2 and z_1, z_2 parametrize the T^4 . Again, the cycles of the T^4 form an $SU(3)$ lattice in $\mathbb{C}^2$ and we denote the A- and B-cycle by

$$e_1 = e^{\frac{2\pi i}{3}}, \quad e_2 = 1. \quad (2.21)$$

We can assume that the corresponding lattice for the T^2 is generated by 1 and i . The action of the CHL quotient is then

$$g' : (z_1, z_2, z_3) \mapsto \left(z_1 + \frac{1}{3}e_1 + \frac{2}{3}e_2, z_2, z_3 + \frac{1}{3} \right). \quad (2.22)$$

The spectrum of $K3$ realized as $T^4/\mathbb{Z}_3$ with the standard shift was first studied in [26] and we will follow the discussion of [26, 27]. Note that the orbifold action g' does not produce any fixed points and so it preserves $\mathcal{N} = 2$ supersymmetry.

The massless spectrum organizes into four-dimensional $\mathcal{N} = 2$ supergravity multiplets with N_v vectormultiplets and N_h hypermultiplets in the theory. The massless states in the g twisted sector are determined by setting left and right masses to zero

$$\begin{aligned} m_L^2 &= N_L + \frac{1}{2}(P + nV)^2 + E_n - 1 = 0, \\ m_R^2 &= N_R + \frac{1}{2}(r + nv)^2 + E_n - \frac{1}{2} = 0. \end{aligned} \quad (2.23)$$

Here P is an $E_8 \times E_8$ lattice vector and of the form

$$P = (P_{E_8}; P_{E'_8}), \quad (2.24)$$

while E_n is the shift in zero point energy on the ground state due to the twisting by g^n . It is given by

$$E_n = \frac{1}{9}n(\nu - n), \quad (2.25)$$

where $\nu = 3$ for the $T^4/\mathbb{Z}_3$ orbifold and $n = 0, 1, 2$. Moreover, r is an $SO(8)$ weight vector with

$$\sum_{i=1}^4 r_i = \text{odd}, \quad (2.26)$$

and the 4 dimensional vector v with

$$v = \frac{1}{3}(0, 0, 1, -1), \quad (2.27)$$

encodes the action that we used to define the $T^4/\mathbb{Z}_3$ orbifold. The value of V encodes the embedding of the $\mathbb{Z}_3$ into the gauge group. It takes values in the two E_8 lattices scaled by one third. For standard embedding we have the shift given by

$$V = \frac{1}{3}(1, -1, 0^6; 0^8). \quad (2.28)$$

The degeneracy of the massless states for $T^4/\mathbb{Z}_3$ model can be obtained from [27]

$$D(n) = \frac{1}{3} \sum_{m=0}^2 \chi(n, m) \Delta(n, m), \quad (2.29)$$

$$\Delta(n, m) = \exp \left\{ 2\pi i [(r + nv)mv - (P + nV)mV + \frac{1}{2}mn(V^2 - v^2) + m\rho] \right\}.$$

Here $\chi(n, m)$ are the number of fixed points in the g^n twisted sector invariant under the action of g^m and ρ is the phase by which oscillators are rotated under the $\mathbb{Z}_3$ action on the T^4 .

In the case of order three CHL orbifolds this formula is modified to

$$D(n; g') = \frac{1}{3} \sum_{m=0}^2 \frac{1}{3} \left[\chi(n, m) + \chi^{(g')}(n, m) + \chi^{(g'^2)}(n, m) \right] \Delta(n, m), \quad (2.30)$$

where $\chi^{(g'^s)}(n, m)$ is the number of fixed points in the g^n twisted sector invariant under the action of $g^m g'^s$. Essentially we implement the projection under the g' action. In the untwisted sector

$$\chi^{(g'^s)}(0, m) = \chi(0, m) = 1, \text{ for } s = 1, 2, \quad (2.31)$$

and the phases in (2.30) implement how the spectrum is projected under g^m . In the twisted sectors we can use again table 1 to obtain

$$\chi^{(g')}(1, m) = \chi^{(g'^2)}(1, m) = 0. \quad (2.32)$$

The sector $n = 2$ gives the anti-particles of the $n = 1$ sector. The untwisted sector, which counts the gravity multiplet and the number of vectors, remains the same as obtained in [27]. The result for the twisted sectors are just one third of that in [27].

In the following table 2 we summarize the untwisted and twisted hypermultiplet contents in the various embeddings of $K3 \times T^2$ where $K3$ is considered at the $T^4/\mathbb{Z}_3$ orbifold limit.

In the table 3 we show how the twisted sectors get modified under the order three orbifold action.

The sectors twisted by g' do not have massless modes as the winding numbers become one-third of integers. The untwisted sector contains the $\mathcal{N} = 2$ gravity multiplet and $\mathcal{N} = 2$ vector multiplets along with the hypers as obtained in [19]. The twisted sectors only

Group, shift	Untwisted sector	Twisted sector
$E_7 \times U(1) \times E_8$ $\frac{1}{3}(1, -1, 0^6; 0^8)$	$(56; 1) + 2(1; 1)$ $+(1; 1)$	$9(56; 1)$ $+ 45(1, 1)$ $18(1, 1)$
$SU(9) \times E_8$ $\frac{1}{3}(2, 1^4, 0^3; 0^8)$	$(84, 1) + 2(1, 1)$	$9(36, 1) + 18(9, 1)$
$SO(14)^2 \times U(1)^2$ $\frac{1}{3}(2, 0^7; 2, 0^7)$	$(14, 1) + (1, 14)$ $+(64, 1) + (1, 64)$ $+2(1, 1)$	$9(14, 1)$ $+9(1, 14)$ $+18(1, 1)$
$E_6 \times SU(3) \times E_7 \times U(1)$ $\frac{1}{3}(2, 1^2, 0^5; 1, -1, 0^6)$	$(27, 3, 1)$ $+(1, 1, 56)$ $+2(1) + (1)$	$9(27, 1, 1)$ $9(1, 3, 1)$ $+18(1, 3, 1)$
$SU(9) \times E_6 \times SU(3)$ $\frac{1}{3}(2, 1^2, 0^5; 2, 1^4, 0^3)$	$(84, 1, 1) + (1, 27, 3)$ $2(1)$	$9(9, 1, 3)$

Table 2. Hypermultiplet spectrum for different embeddings with $K3$ as $T^4/\mathbb{Z}_3$. We have not kept track of various $U(1)$ charges.

Group, shift	Untwisted sector	Twisted sector	$N_h - N_v$
$E_7 \times U(1) \times E_8$ $\frac{1}{3}(1, -1, 0^6; 0^8)$	$(56; 1) + 2(1; 1)$ $+(1; 1)$	$3(56; 1)$ $+15(1, 1)$ $+6(1, 1)$	-134
$SU(9) \times E_8$ $\frac{1}{3}(2, 1^4, 0^3; 0^8)$	$(84, 1) + 2(1, 1)$	$3(36, 1) + 6(9, 1)$	-80
$SO(14)^2 \times U(1)^2$ $\frac{1}{3}(2, 0^7; 2, 0^7)$	$(14, 1) + (1, 14)$ $+(64, 1) + (1, 64)$ $+2(1, 1)$	$3(14, 1)$ $+3(1, 14)$ $+6(1, 1)$	64
$E_6 \times SU(3) \times E_7 \times U(1)$ $\frac{1}{3}(2, 1^2, 0^5; 1, -1, 0^6)$	$(27, 3, 1)$ $+(1, 1, 56)$ $+2(1) + (1)$	$3(27, 1, 1)$ $+3(1, 3, 1)$ $+6(1, 3, 1)$	28
$SU(9) \times E_6 \times SU(3)$ $\frac{1}{3}(2, 1^2, 0^5; 2, 1^4, 0)$	$(84, 1, 1) + (1, 27, 3)$ $2(1)$	$3(9, 1, 3)$	82

Table 3. Hypermultiplet spectrum for different embeddings with 3A orbifold of $K3$ where $K3$ is realized as $T^4/\mathbb{Z}_3$. We have not kept track of various $U(1)$ charges.

contain the hypers. The hypermultiplet contents are summarized in table 2, 3. Evaluating the difference between N_h and N_v we have for the standard embedding

$$N_h - N_v = -134. \quad (2.33)$$

This was predicted in [6]. Note that when counting the number of vector multiplets N_v we only consider those that originate from the $E_8 \times E_8$ group and not the four generic

vectors that arise from the T^2 . When no orbifold acts on $K3$ the theory can be lifted to six dimensions. The value of $N_h - N_v$ then remains 244 for any realization of $K3$ and for any embeddings because of anomaly cancellation in six dimension [28]. This is different for $N = 2$ or $N = 3$ orbifold models.

By explicitly analysing the possible higgs chains for the gauge groups and spectra in table 3 we found that for two of the models the gauge group can be completely broken. These are the non-standard embeddings with $N_h - N_v = 64$ and $N_h - N_v = 82$. For the model with $N_h - N_v = 64$ we provide the details of the higgs chain in appendix 3. In section (4) we will construct Calabi-Yau geometries for which the enumerative invariants exactly match the predictions from the heterotic calculation of the gauge gravitational couplings for both of the models. We also find many more Calabi-Yau manifolds for which we expect that a heterotic dual compactification on $(K3 \times T^2)/\mathbb{Z}_3$ exists but where the embedding can not be realized with our construction.

2.3 The new supersymmetric index

In this section we shall evaluate the new supersymmetric index of the heterotic string compactified on $K3 \times T^2$ with an order 3 CHL action on $K3 \times T^2$ with various embeddings of $E_8 \times E_8$. We shall explicitly compute the results of standard embeddings and summarize the result of non-standard ones. We consider $K3$ at the orbifold limit of $T^4/\mathbb{Z}_3$ where the $\mathbb{Z}_3$ action on T^4 is given in (2.20). The CHL action is given by (2.22). The new supersymmetric index is defined as

$$\mathcal{Z}_{\text{new}}(\tau, \bar{\tau}) = \frac{1}{\eta^2(\tau)} \text{Tr}_R((-1)^F F q^{L_0 - c/24} \bar{q}^{\bar{L}_0 - \bar{c}/24}). \quad (2.34)$$

Under the standard embedding we have the shift action given by equation (2.28). The above trace for standard embedding splits into the following sectors [6]:

$$\mathcal{Z}_{\text{new}}(\tau, \bar{\tau}) = -\frac{1}{2\eta^{20}(\tau)} \sum_{a,b=0}^2 \sum_{r,s=0}^2 e^{-\frac{2\pi i ab}{9}} Z_{E_8}^{a,b}(\tau) \times E_4(\tau) \times \frac{1}{9} F(a, r, b, s, \tau) \Gamma_{2,2}^{(r,s)}(q, \bar{q}), \quad (2.35)$$

where only $\Gamma_{2,2}^{(r,s)}(q, \bar{q})$, defined in (2.44), has a non-holomorphic dependence which encodes the winding and momenta modes contribution of the T^2 bosons. The factor of E_4 is obtained from the untouched E_8 lattice. We can write the trace over the directions of T^4 as

$$F(a, r, b, s; \tau) = \text{Tr}_{g^a g^r} \left(g^b g^s e^{i\pi F T^4} q^{L_0} \bar{q}^{\bar{L}_0} \right), \quad (2.36)$$

and one can work out the explicit expression

$$F(a, r, b, s; \tau) = k^{a,r,b,s} \eta^2(\tau) q^{\frac{-a^2}{9}} \frac{1}{\theta_1^2(\frac{a\tau+b}{3}, \tau)}. \quad (2.37)$$

For standard embeddings the coefficients $k^{a,r,b,s}$ for different values of (r, s) are given by the following matrices:

$$\begin{aligned}
 k^{a,0,b,1} = k^{a,0,b,2} &= 9 \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & k^{a,1,b,0} = k^{a,2,b,0} &= 9 \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\
 k^{a,1,b,1} = k^{a,2,b,2} &= 9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & k^{a,2,b,1} = k^{a,1,b,2} &= 9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \\
 k^{a,0,b,0} &= 9 \begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}.
 \end{aligned} \tag{2.38}$$

Here the rows and columns are denoted by a and b respectively. The result for the coefficients $k^{a,r,b,s}$ comes from counting the fixed points as for the case of twisted elliptic genus in table 1.

However, if the embedding is non-standard the new supersymmetric index is given by [3, 6]

$$\mathcal{Z}_{\text{new}}(\tau, \bar{\tau}) = -\frac{1}{2\eta^{20}(\tau)} \sum_{a,b,r,s=0}^2 e^{-2\pi i \frac{ab}{\nu^2} \Gamma^2} Z_{E_8}^{a,b}(\tau) Z_{E'_8}^{a,b}(\tau) \frac{1}{\nu^2} F(a, r, b, s, \tau) \Gamma_{2,2}^{(r,s)}(\tau, \bar{\tau}), \tag{2.39}$$

where the partition functions over the shifted E_8 lattices are

$$\begin{aligned}
 Z_{E_8}^{a,b}(\tau) &= \sum_{\alpha, \beta=0}^1 e^{-i\pi\beta a \sum_{I=1}^8 \gamma^I} \prod_{I=1}^8 \theta \left[\begin{smallmatrix} \alpha+2a\gamma^I \\ \beta+2b\gamma^I \end{smallmatrix} \right], \\
 Z_{E'_8}^{a,b}(\tau) &= \sum_{\alpha, \beta=0}^1 e^{-i\pi\beta a \sum_{I=1}^8 \tilde{\gamma}^I} \prod_{I=1}^8 \theta \left[\begin{smallmatrix} \alpha+2a\tilde{\gamma}^I \\ \beta+2b\tilde{\gamma}^I \end{smallmatrix} \right],
 \end{aligned} \tag{2.40}$$

with $\gamma, \tilde{\gamma}$ being the shifts in the E_8, E'_8 lattices, while $\nu = 3$ is the order of the orbifold on T^4 . Moreover, $\Gamma^2 = \nu^2(\gamma^2 + \tilde{\gamma}^2)$ is the sum of the squares of all the shifts in the two E_8 lattices. Under non-standard embeddings we have

$$\begin{aligned}
 k^{a,0,b,1} = k^{a,0,b,2} &= 9 \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & k^{a,1,b,0} = k^{a,2,b,0} &= 9 \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\
 k^{a,1,b,1} = k^{a,2,b,1} &= 9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & e^{-\pi i(2-\Gamma^2)/9} & 0 \\ 0 & 0 & e^{-4\pi i(2-\Gamma^2)/9} \end{pmatrix} \\
 k^{a,1,b,2} = k^{a,2,b,1} &= 9 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & e^{-2\pi i(2-\Gamma^2)/9} \\ 0 & e^{-5\pi i(2-\Gamma^2)/9} & 0 \end{pmatrix}, \\
 k^{a,0,b,0} &= 9 \begin{pmatrix} 0 & 1 & 1 \\ 1 & e^{-\pi i(2-\Gamma^2)/9} & e^{-2\pi i(2-\Gamma^2)/9} \\ 1 & e^{-5\pi i(2-\Gamma^2)/9} & e^{-4\pi i(2-\Gamma^2)/9} \end{pmatrix},
 \end{aligned} \tag{2.41}$$

These matrices can be obtained as follows: when $r \cdot s = 0$, then $k^{a,r,b,s}$ is identical to the result that we obtained for standard embedding. Furthermore, modular invariance requires that the values in $k^{a,0,b,0}$ satisfy

$$k^{a,0,b,0} = e^{\pi i \frac{a^2}{9}(2-\Gamma^2)} k^{a,0,a+b,0}, \quad k^{a,0,b,0} = e^{-2\pi i \frac{ab}{9}(2-\Gamma^2)} k^{b,0,-a,0}, \quad (2.42)$$

where a, b are considered modulo three [29]. The relation

$$k^{a,0,b,0} = \sum_{\substack{r,s \\ rs > 0}} k^{a,r,b,s}, \quad (2.43)$$

together with the fact that the matrices differ from those for the standard embedding only by a phase, then fixes all of the other coefficients. The lattice sum involving the winding and momenta modes are given in each case by,

$$\Gamma_{2,2}^{(r,s)}(\tau, \bar{\tau}) = \sum_{\substack{m_1, m_2, n_2 \in \mathbb{Z} \\ n_1 = \mathbb{Z} + \frac{r}{3}}} q^{\frac{p_L^2}{2}} \bar{q}^{\frac{p_R^2}{2}} e^{2\pi i m_1 s/3}, \quad (2.44)$$

where the left- and right-moving momenta are given by

$$\begin{aligned} \frac{1}{2} p_R^2 &= \frac{1}{2T_2 U_2} | -m_1 U + m_2 + n_1 T + n_2 T U |^2, \\ \frac{1}{2} p_L^2 &= \frac{1}{2} p_R^2 + m_1 n_1 + m_2 n_2, \end{aligned} \quad (2.45)$$

with T, U being Kähler and complex structure of T^2 .

One can then define the different components of the new supersymmetric index $\mathcal{Z}_{\text{new}}^{r,s}$ via

$$\mathcal{Z}_{\text{new}}(\tau, \bar{\tau}) = -\frac{1}{\eta^{24}(\tau)} \sum_{r,s=0}^2 \mathcal{Z}_{\text{new}}^{r,s}(\tau) \Gamma_{2,2}^{(r,s)}(\tau, \bar{\tau}). \quad (2.46)$$

We note that $\mathcal{Z}_{\text{new}}^{r,s}$ are modular functions in congruent subgroups of $\text{SL}(2, \mathbb{Z})$ and for $N = 3$ orbifolds satisfy

$$\mathcal{Z}_{\text{new}}^{0,1} = \mathcal{Z}_{\text{new}}^{0,2}, \quad \mathcal{Z}_{\text{new}}^{1,1} = \mathcal{Z}_{\text{new}}^{2,2}, \quad \mathcal{Z}_{\text{new}}^{1,2} = \mathcal{Z}_{\text{new}}^{2,1}. \quad (2.47)$$

The value of $N_h - N_v$ can be extracted from the new supersymmetric index as

$$N_h - N_v = -\frac{1}{4} \frac{1}{\eta^{20}(\tau)} \sum_{0,s=0}^2 \mathcal{Z}_{\text{new}}^{0,s} \Big|_{q^{1/6}}, \quad (2.48)$$

where $\mathcal{Z}_{\text{new}}^{0,0} = \frac{4}{3} E_4 E_6$. The results for the $\mathcal{Z}_{\text{new}}^{r,s}$ sectors can be given in terms of modular forms for $\Gamma_1(3)$ and read

$$\begin{aligned} \mathcal{Z}_{\text{new}}^{0,1} &= \frac{4}{3} \left(\hat{a} E_4 E_6 + \hat{b} \mathcal{E}_3^2(\tau) E_6 + \hat{c} E_4 (E_6 + 3\mathcal{E}_3(\tau) E_4) + \hat{d} \mathcal{E}_3^2(\tau) (E_6 + 3\mathcal{E}_3(\tau) E_4) \right), \\ \mathcal{Z}_{\text{new}}^{r,rk} &= \frac{4}{3} \left(\hat{a} E_4 E_6 + \frac{\hat{b}}{9} \mathcal{E}_3'^2(\tau) E_6 + \hat{c} E_4 (E_6 - \mathcal{E}_3' E_4) + \frac{\hat{d}}{9} \mathcal{E}_3'^2(\tau) (E_6 - \mathcal{E}_3' E_4) \right), \end{aligned} \quad (2.49)$$

Shift	$N_h - N_v$	$\hat{a}$	$\hat{b}$	$\hat{c}$	$\hat{d}$
$\frac{1}{3}(1, -1, 0^6; 0^8)$	-134	0	0	$\frac{1}{4}$	0
$\frac{1}{3}(2, 1^4, 0^3; 0^8)$	-80	$\frac{1}{16}$	$-\frac{9}{16}$	$-\frac{3}{16}$	$\frac{9}{16}$
$\frac{1}{3}(2, 0^7; 2, 0^7)$	64	$\frac{-1}{48}$	$\frac{3}{16}$	$\frac{1}{48}$	$\frac{3}{16}$
$\frac{1}{3}(1, -1, 0^6; 2, 1^2, 0^5)$	28	0	0	$\frac{-1}{32}$	$\frac{9}{32}$
$\frac{1}{3}(2, 1^2, 0^5; 2, 1^4, 0^3)$	82	$\frac{-1}{32}$	$\frac{9}{32}$	$\frac{3}{64}$	$\frac{9}{64}$

Table 4. $\mathcal{Z}_{\text{new}}^{r,s}$ for different embeddings with $K3$ as $T^4/\mathbb{Z}_3$ and $N = 3$ CHL orbifold.

where we have introduced $\mathcal{E}'_3(\tau) = \mathcal{E}_3(\frac{\tau+k}{3})$ and the different values of $\hat{a}, \hat{b}, \hat{c}, \hat{d}$ are given in table 4. We note that the difference between the number of hypers and vectors in the standard embedding is as expected from earlier results in [6]. As we review in section 4.1, the difference $N_h - N_v$ directly encodes the Euler characteristic χ of a dual Calabi-Yau compactification space via

$$\chi = 2(N_v - N_h) + 8. \quad (2.50)$$

Let us note again that when counting the vector multiplets N_v we only consider those that originate from the $E_8 \times E_8$ and not the generic four vectors that arise from the T^2 . It turns out that the coefficients of the new supersymmetric indices for all of the four non-standard embeddings can be expressed as

$$\begin{aligned} \hat{a} &= \frac{1}{2^7 3^3} (48 + \chi), & \hat{b} &= -\frac{1}{2^7 3} 48 + \chi, \\ \hat{c} &= -\frac{1}{2^8 3^3} (456 + 5\chi), & \hat{d} &= \frac{1}{2^8 3} (264 + \chi). \end{aligned} \quad (2.51)$$

This would imply the values

$$(\hat{a}, \hat{b}, \hat{c}, \hat{d}) = \left(\frac{3}{32}, -\frac{27}{32}, -\frac{17}{64}, \frac{45}{64} \right), \quad (2.52)$$

for the standard embedding with $N_h - N_v = -134$, which is clearly different from what we found via an explicit calculation. Note that for compactifications on $K3 \times T^2$ and $(K3 \times T^2)/\mathbb{Z}_2$ it was found that the new supersymmetric index can be expressed in terms of the Euler characteristic for *all* of the standard and non-standard embeddings that had been considered [3, 6]. It would be interesting to understand what makes the standard embedding on $(K3 \times T^2)/\mathbb{Z}_3$ different and whether a non-standard embedding exists that realizes (2.52).

3 Heterotic computation of Gopakumar-Vafa invariants

In this section we compute the Gopakumar-Vafa invariants from the heterotic side for the non-standard embeddings discussed above. The Gopakumar-Vafa invariants n_β^g correspond to weighted sums of multiplicities of BPS states that arise from M2-branes wrapping curves of class $\beta \in H_2(M)$ in a Calabi-Yau threefold M [30, 31]. We will loosely refer to n_β^g as

Gopakumar-Vafa invariants of genus g because they only receive contributions from M2-branes that wrap curves with genus less than or equal to g . To obtain the invariants one needs to calculate the gravitational coupling in the low energy $\mathcal{N} = 2$ effective theory. The main ingredients of the calculation are different sectors of the new supersymmetric index.

At certain points in the moduli space one encounters singularities in the gravitational coupling. Towards the end of this section we shall discuss these singularities known as conifold singularities. The observations here are very similar to the ones obtained in [5]. The strength at these singular points are given by the Gopakumar-Vafa invariants at these points. It is also similar to the results for the standard embedding in [5] that the untwisted sector is devoid of these singular points.

3.1 Extraction of Gopakumar-Vafa invariants from F_g

The gravitational couplings F_g of the low energy effective $\mathcal{N} = 2$ theories appear as

$$S = \int F_g(y, \bar{y}) \cdot F_+^{2g-2} R_+^2, \quad (3.1)$$

where F_+ , R_+ are the self-dual parts of the graviphoton and Riemann tensor and $y, \bar{y}$ denote the dependence on vector moduli.

For $K3 \times T^2$ without any orbifolding, the couplings have been computed from the corresponding heterotic one-loop amplitude in [2] using the so-called Borchard lift [32]. The calculation without any graviphoton insertion, which leads to the prepotential of the supergravity theory, was done earlier in [1]. For compactifications on $(K3 \times T^2)/\mathbb{Z}_3$ with standard embeddings the calculation has been performed in [5].

Through the computation of this F_g we can obtain predictions for Gopakumar-Vafa invariants at genus g of a dual Calabi-Yau threefold. The essential ingredient in computing the F_g is given by the new supersymmetric index. Under standard embedding for any g' orbifold of $K3$ the results of this index were given in [6]. We can write the new supersymmetric index evaluated in the previous section for standard embeddings as a holomorphic part times the sum over winding and momenta. Re-writing the holomorphic component as $f^{(r,s)}$ for each sector of twist and insertion of g' (given by r, s respectively) action we have

$$\mathcal{Z}_{new}(q, \bar{q}) = -4 \sum_{r,s=0}^{N-1} \Gamma_{2,2}^{(r,s)} f^{(r,s)}(\tau), \quad (3.2)$$

where one introduces

$$f^{(r,s)}(\tau) = \frac{1}{2\eta^{24}(\tau)} E_4 \left[\frac{1}{4} \alpha_{g'}^{(r,s)} E_6 - \beta_{g'}^{(r,s)}(\tau) E_4 \right] = \sum_{l \in \frac{\mathbb{Z}}{N}} c^{(r,s)}(l) q^l. \quad (3.3)$$

The $\alpha_{g'}^{(r,s)}$ are numerical constants while $\beta_{g'}^{(r,s)}$ are weight 2 modular forms for $\Gamma_1(N)$, where N corresponds to the order of the orbifold group.

It has previously been shown [12, 33–35] that the one-loop integral F_g is

$$F_g = \frac{1}{2\pi^2(g!)^2} \int \frac{d^2\tau}{\tau_2} \left\{ \frac{1}{\tau_2^2 \eta^2(\tau)} \text{Tr}_R \left[(i\bar{\partial}X)^{(2g-2)} (-1)^F F q^{L_0 - \frac{c}{24}} \bar{q}^{\bar{L}_0 - \frac{\bar{c}}{24}} \right] \right. \\ \left. \times \left\langle \prod_{i=1}^g \int d^2x_i Z^1 \partial Z^2(x_i) \prod_{j=1}^g \int d^2y_j \bar{Z}^2 \partial \bar{Z}^1(y_j) \right\rangle \right\}. \quad (3.4)$$

Here X is the complex coordinate on the T^2 and Z^1, Z^2 are the complex coordinates of the transverse non-compact space time. The above result is obtained from the Wick contractions of the graviton and graviphoton vertex operators. Factorizing the non-compact directions, the internal CFT trace is given by

$$\begin{aligned} & \frac{1}{\eta^2(\tau)} \text{Tr}_R \left[(i\partial X)^{(2g-2)} (-1)^F F q^{L_0 - \frac{c}{24}} \bar{q}^{\tilde{L}_0 - \frac{\tilde{c}}{24}} \right] \\ &= 2 \frac{1}{\eta^{24}(\tau)} \sum_{r,s} \Gamma_{2,2}^{(r,s)} \left(\frac{(p_R^{(r,s)})}{\sqrt{2T_2 U_2}} \right)^{(2g-2)} q^{|p_L|^2/2} \bar{q}^{|p_R|^2/2} f^{r,s}(\tau). \end{aligned} \quad (3.5)$$

The only essential difference between this trace and the new supersymmetric index computation are the extra powers of $p_R^{(r,s)}$. It is evident that under non-standard embedding $f^{(r,s)}$ would have to be replaced by $\frac{1}{4\eta^{24}(\tau)} \mathcal{Z}_{\text{new}}^{r,s}$. The momenta $p_L^{(r,s)}, p_R^{(r,s)}$ are given in equation (2.45), assuming no other moduli dependence on the heterotic side.

In order to make contact with the enumerative geometry of a type II Calabi-Yau compactification that produces the same effective action we need to extract the holomorphic part of the integral F_g . The purely holomorphic contribution to the gravitational couplings are given by [5]

$$\begin{aligned} \bar{F}_g^{\text{hol}} = & \frac{(-1)^{g-1}}{\pi^2} \sum_{s=0}^{N-1} \left(\sum_{m>0} e^{-2\pi i n_2 s/N} c_{g-1}^{(r,s)}(n_1 n_2/N) \text{Li}_{3-2g}(e^{2\pi i m \cdot y}) \right. \\ & \left. + \frac{1}{2} c_{g-1}^{(0,s)}(0) \zeta(3-2g) \right). \end{aligned} \quad (3.6)$$

where $y = (T, U)$ with T, U being the complex and the complexified Kähler structure of T^2 . The coefficients $c_{g-1}^{(r,s)}(n_1 n_2/N)$ are obtained from

$$\frac{1}{4\eta^{24}} \mathcal{Z}_{\text{new}}^{r,s} \mathcal{P}_{2g}(G_2, G_4, \dots, G_{2g}) = \sum_{l \in \frac{\mathbb{Z}}{N}} c_{g-1}^{(r,s)}(l) q^l, \quad (3.7)$$

where $G_{2k} = 2\zeta(2k)E_{2k}$, ζ is the Riemann zeta function and E_{2k} denote the Eisenstein series of weight $2k$. $\mathcal{P}_{2g}$ is defined through the Schur polynomial of order g in the following manner

$$\mathcal{P}_{2g}(x_1, x_2, \dots, x_g) = -\mathcal{S} \left(x_1, \frac{1}{2}x_2, \dots, \frac{1}{g}x_g \right). \quad (3.8)$$

For an orbifold of order N the lattice sum $m > 0$ involves the points

$$\begin{aligned} m = (n_1, n_2) &\geq 0, \quad \text{but } (n_1, n_2) \neq (0, 0), \\ (r/N, -n_2), &\quad \text{with } n_2 > 0 \text{ and } rn_2 \leq N. \end{aligned} \quad (3.9)$$

The Gopakumar-Vafa invariants of the potential Calabi-Yau duals can now be extracted from the heterotic results by first writing down the Gopakumar-Vafa form of the genus g

topological amplitude. In terms of Gopakumar-Vafa invariants n_m^g and for $g > 1$ it can be written as [30, 31]

$$F_g^{\text{GV}} = \frac{(-1)^g |B_{2g} B_{2g-2}| \chi(X)}{4g(2g-2)(2g-2)!} + \sum_{\beta} \left[\frac{|B_{2g}| n_m^0}{2g(2g-2)!} + \frac{2(-1)^g n_m^2}{(2g-2)!} \pm \dots - \frac{g-2}{12} n_m^{g-1} + n_m^g \right] \text{Li}_{3-2g}(e^{2\pi i m \cdot y}), \quad (3.10)$$

where B_{2g} are the Bernoulli numbers and $\chi(X)$ is the Euler characteristic of the Calabi-Yau. The geometric interpretation of the curve class in the Calabi-Yau dual that is associated to $m = (n_1, n_2)$ is explained on page 25. For $g = 0$ we get

$$F_0^{\text{GV}} = \zeta(3) \frac{\chi(X)}{2} + \sum_{m>0} n_m^0 \text{Li}_3(e^{2\pi i m \cdot y}). \quad (3.11)$$

For $g = 1$ we have

$$F_1^{\text{GV}} = \sum_{m>0} \left(\frac{1}{12} n_m^0 + n_m^1 \right) \text{Li}_1(e^{2\pi i m \cdot y}). \quad (3.12)$$

Comparing the constant term of (3.6) and (3.10) gives the following relation between the topological amplitudes

$$F_g^{\text{GV}} = \frac{(-1)^{g+1}}{2(2\pi)^{2g-2}} \bar{F}_g^{\text{hol}}. \quad (3.13)$$

By proceeding recursively, one obtains expressions for the Gopakumar-Vafa invariants of lowest genus

$$\begin{aligned} n_{(n_1, n_2)}^0 &= 2 \sum_{s=0}^{N-1} e^{-\frac{2\pi i n_2 s}{N}} c_{-1}^{(r,s)}, \\ n_{(n_1, n_2)}^1 &= \frac{1}{2\pi^2} \sum_{s=0}^N e^{-\frac{2\pi i n_2 s}{N}} c_0^{(r,s)} (m^2/2) - \frac{1}{12} n_{(n_1, n_2)}^0, \\ n_{(n_1, n_2)}^2 &= \frac{1}{8\pi^4} \sum_{s=0}^{N-1} e^{-\frac{2\pi i n_2 s}{N}} c_1^{(r,s)} (m^2/2) - \frac{|B_4|}{8} n_{n_1, n_2}^0, \end{aligned} \quad (3.14)$$

where $r = n_1 N \bmod N$ and $m^2 = 2n_1 n_2$.

3.2 Conifold singularities

Conifold singularities are reflected in the gravitational thresholds as poles where vectors become massless. The polylogarithm $\text{Li}_a(z)$ is singular at $z = 1$ and the leading behaviour around the pole is for $a < 0$ given by

$$\text{Li}_a(z) = \left(z \frac{d}{dz} \right)^{|a|} \frac{1}{1-z} \Big|_{z \rightarrow 1} \sim a! \frac{1}{(1-z)^{|a|+1}}. \quad (3.15)$$

Hence the leading divergence for $g > 1$ in $\bar{F}_g^{\text{hol}}$ can be written as

$$\bar{F}_g^{\text{hol}}|_{m \cdot y \rightarrow 0} \sim \frac{(-1)^{g-1}}{\pi^2} \sum_{s=0}^{N-1} e^{-2\pi i n_2 s/N} c_{g-1}^{(r,s)} (n_1 n_2) (2g-3)! \frac{1}{\{1 - e^{2\pi i m \cdot y}\}^{2g-2}}, \quad (3.16)$$

where

$$m \cdot y = n_1 T_1 + n_2 U_1 = 0, \quad n_1 T_2 + n_2 U_2 = 0. \quad (3.17)$$

The Euler characteristic of a dual Calabi-Yau is then related to the coefficients via

$$\chi = 2 \sum_{s=0}^{N-1} c_{-1}^{(r,s)}(0), \quad (3.18)$$

and can be computed from the spectrum or $\mathcal{Z}_{\text{new}}$. Since $T_2, U_2 > 0$, being the imaginary parts of the Kähler and complex structure of the torus T^2 , and m is constrained by (3.1) we see that n_2 is negative at these singularities. Therefore in the untwisted sector a possible singularity lies at

$$(n_1, n_2) = (1, -1), \quad n_1 n_2 = -1, \quad (3.19)$$

However, a close inspection of the new supersymmetric index shows that for any standard or non-standard embedding for $N = 3$ CHL orbifolds there lies no conifold singularity in the untwisted sector. This was previously observed for the standard embedding results in [5].

4 Dual type IIA compactifications on Calabi-Yau threefolds

In the previous sections we have discussed in detail the calculation of the gravitational coupling

$$S = \int F_g(y, \bar{y}) F_+^{2g-2} R_+^2, \quad (4.1)$$

for heterotic $E_8 \times E_8$ compactifications on $(K3 \times T^2)/\mathbb{Z}_3$. Following the arguments that we review below, one expects that there exists a dual type IIA compactification on a genus one fibered Calabi-Yau threefold with three-sections that also exhibits a $K3$ fibration.

The calculations of the heterotic one-loop calculation in this paper extends earlier work [5, 6] in that we consider several non-standard embeddings of the spin connection into the gauge connection. In particular, two of the corresponding effective theories exhibit a spectrum such that the gauge group can be maximally higgsed to $U(1)^3$. We can therefore hope that dual type IIA compactifications on Calabi-Yau threefolds to these models exist such that the geometries are genus one and $K3$ -fibered but have the lowest possible number of Kähler moduli $h^{1,1} = 3$.

In this section we will systematically engineer such Calabi-Yau manifolds using tools from toric geometry and show that they are indeed dual to heterotic compactification on $(K3 \times T^2)/\mathbb{Z}_3$. To this end we employ the modular bootstrap and obtain all-genus topological string amplitudes on the corresponding geometries. We then match these with the results from the heterotic one-loop calculation.

4.1 A brief review of heterotic/type II duality

Let us briefly review the basics of heterotic/type II duality and why the Calabi-Yau manifolds require a particular fibration structure. A standard reference on this duality is still [36] and a more recent discussion can be found in [37].

In six dimensions, heterotic strings on T^4 are dual to type IIA strings on a $K3$ [38, 39]. The four dimensional duality between heterotic strings on $K3 \times T^2$ and type IIA strings on a Calabi-Yau threefold [40] can then be motivated by an adiabatic argument [41]. More precisely, one can fiber the geometries on both sides of the duality over a $\mathbb{P}^1$ such that the fibration of $K3$ over $\mathbb{P}^1$ is a Calabi-Yau threefold. Matching the amount of unbroken supersymmetries requires the fibration on the heterotic side to only involve a 2-torus inside T^4 such that the resulting geometry is $K3 \times T^2$. From the perspective of the type II compactification, the heterotic string arises from a 5-brane that wraps the generic $K3$ fiber. The adiabatic argument can fail at points where the $K3$ fiber degenerates and this can be related to the presence of 5-branes on the heterotic side of the duality [37].

The heterotic compactification on $K3 \times T^2$ exhibits a self-duality group which contains the $SL(2, \mathbb{Z})$ action on the complex structure of the torus. After taking a $\mathbb{Z}_N$ quotient which acts by an order N shift on the torus, this duality group is broken down to the congruence subgroup $\Gamma_1(N) \subset SL(2, \mathbb{Z})$ that preserves the generator of a $\mathbb{Z}_N$ subgroup of T^2 [9, 41]. The duality group of the heterotic string is realized on the type IIA side as monodromies in the stringy Kähler moduli space of the Calabi-Yau threefold [13, 42, 43]. This implies that for heterotic strings on $(K3 \times T^2)/\mathbb{Z}_N$ the dual Calabi-Yau compactification space should be a genus one fibration with N -sections [9, 44].

The moduli space of the four-dimensional $\mathcal{N} = 2$ effective theories factorizes into

$$\mathcal{M} = \mathcal{M}_{\text{vec.}} \times \mathcal{M}_{\text{hyp.}}, \quad (4.2)$$

where $\mathcal{M}_{\text{vec.}}$ is parametrized by the expectation values of scalar fields from vector multiplets and $\mathcal{M}_{\text{hyp.}}$ is correspondingly parametrized by scalars from hyper multiplets. The coefficients $F_g(y, \bar{y})$ in (4.1) only depend on the vector moduli.

On the heterotic side there are four massless vector fields from reducing the metric and the B-field along the T^2 of which one contributes to the single graviton multiplet. The remaining three vector fields combine with 6 real scalars that arise from the dual of the B-field in four dimensions, the dilaton and the Kaluza-Klein modes of the metric and the B-field along T^2 , as well as an appropriate number of fermions, into three vector multiplets. The dilaton and the dual of the B-field combine into a complex axio-dilaton while the Kaluza-Klein modes of metric and B-field can be interpreted as the complex structure and the complexified Kähler structure of T^2 . Additional massless vector multiplets might descend from the 10-dimensional $E_8 \times E_8$ gauge group but those can acquire a mass from the choice of gauge background.

On the type IIA side, the scalars in the vector multiplets arise as Kaluza-Klein modes of the Kähler form and the B-field, both of which can be expanded in harmonic 2-forms that in turn correspond to $H^{1,1}(M)$ classes of the Calabi-Yau M . The graviphoton on the other hand corresponds to the Ramond-Ramond one-form. This implies that for a heterotic compactification that leads to the minimum of four vector fields a dual Calabi-Yau compactification space needs to exhibit $h^{1,1} = 3$. The three minimal Kähler moduli of the Calabi-Yau can be related to the volume of the generic fiber, the volume of the base of the $K3$ fiber and the volume of the $\mathbb{P}^1$ base of the $K3$ fibration. Moreover, the

coefficients $F_g(y, \bar{y})$ are topological [12, 34] and can be interpreted as genus g amplitudes of the A-twisted topological string theory on M .

Under the duality, the complex structure and the complexified Kähler structure of the T^2 on the heterotic side can be identified with the complexified volumes of fiber and base of the generic $K3$ fiber. The heterotic axio-dilaton is identified with the complexified volume of the $\mathbb{P}^1$ base of the $K3$ fibration. On the heterotic side one can T-dualize along the cycle that is not involved in the $\mathbb{Z}_N$ quotient and this exchanges complex and complexified Kähler structure on T^2 . This duality is also realized as a monodromy in the complexified Kähler moduli space of the dual Calabi-Yau.

4.2 Constructing Calabi-Yau duals

We now want to construct Calabi-Yau threefolds that exhibit both a $K3$ fibration and a genus one fibration while $h^{1,1} = 3$. To this end let us first introduce some mathematical background about $K3$ surfaces [45, 46].

In contrast to the wealth of topologically distinct Calabi-Yau threefolds there is only one topological type of $K3$ surfaces. The non-vanishing Hodge numbers are always $h^{0,0} = h^{2,0} = h^{0,2} = h^{2,2} = 1$ and $h^{1,1} = 20$. However, for a given $K3$ surface S , the $H^2(S, \mathbb{Z})$ cohomology decomposes into the *Néron-Severi lattice*

$$\text{NS}_{K3} = H^2(S, \mathbb{Z}) \cap H^{1,1}(S), \quad (4.3)$$

and the orthogonal complement T which is called the *transcendental lattice*. The rank of NS_{K3} is called the *Picard number* $\rho(S)$. A $K3$ fibration is called Λ_S *polarized* if there is a set of divisors on the total space M of the fibration such that the restrictions of the divisors to a generic fiber generate a sublattice $\Lambda_S \subset \text{NS}_{K3}$. For $K3$ surfaces of Picard number two one can write the intersection form on NS_{K3} as

$$I = \begin{pmatrix} 2a & b \\ b & 2c \end{pmatrix}, \quad a, b, c \in \mathbb{Z}, \quad 4ac - b^2 < 0. \quad (4.4)$$

Moreover, the surface is genus one fibered with an N -section if and only if $b^2 - 4ac = N^2$.

Our goal in this section is to construct genus one fibered Calabi-Yau threefolds that also exhibit a $K3$ fibration and have $h^{1,1} = 3$. In particular, we want to construct genus one fibrations with 3-sections to find compactifications of Type IIA strings that are dual to the heterotic compactifications that we considered in the previous sections. Let us first note that every genus one fibration that exhibits 1-, 2- or 3-sections is birational to a hypersurface in a fibration of weighted projective spaces [45]. The respective fibers are $\mathbb{P}(1, 2, 3)$, $\mathbb{P}(1, 1, 2)$ and $\mathbb{P}^2$. Moreover, the base of a genus one fibration can only be $\mathbb{P}^2$, a Hirzebruch surface $\mathbb{F}_m$, the Enriques surface or blowups of these geometries [47]. A fibration over $\mathbb{P}^2$ will never exhibit a $K3$ fibration and, since we are interested in Calabi-Yau manifolds with $h^{1,1} = 3$, the only bases that we have to consider are Hirzebruch surfaces $\mathbb{F}_m$, $m = 0, 1, 2$. If we are only interested in genus one fibrations with N -sections for $N \leq 3$ we can therefore restrict ourselves to Calabi-Yau threefolds that are hypersurfaces in four-dimensional toric ambient spaces.

N -section	Polytope	Intersection form	Inequivalent n
$N = 1$	$\begin{pmatrix} 1 & 0 & 0 \\ -2 & -3 & 0 \\ -2 & -3 & 1 \\ -2 & -3 & -1 \end{pmatrix}$	$U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	
$N = 2$	$\begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ -1 & -1 & 0 \\ -2 & n & 1 \\ 0 & 0 & -1 \end{pmatrix}$	$I_n^{(2)} = \begin{pmatrix} 2n & 2 \\ 2 & 0 \end{pmatrix}$	$n \in \{0, 1, 2^*\}$
$N = 3$	$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & -1 & 0 \\ n & 0 & 1 \\ 0 & 0 & -1 \end{pmatrix}$	$I_n^{(3)} = \begin{pmatrix} 2(1+n) & 3 \\ 3 & 0 \end{pmatrix}$	$n \in \{-1, 0, 1, 2^*\}$

Table 5. The eight inequivalent $K3$ surfaces that are hypersurfaces in toric ambient spaces, genus one fibered and exhibit a toric Néron-Severi lattice of rank two. The values of n for which the toric N -section is actually a union of N sections are marked with an asterix.

Moreover, since the base of the $K3$ fibration has to be $\mathbb{P}^1$, the fibration has to arise from a compatible toric fibration of the ambient space. The generic $K3$ fiber will therefore be a hypersurface in a three-dimensional toric variety. Calabi-Yau hypersurfaces in d -dimensional toric ambient spaces correspond to reflexive polytopes of the same dimension [48]. Three dimensional reflexive polytopes have been classified [49] and the Picard number of the corresponding $K3$ hypersurface can easily be obtained from the polytope [48]. This allows us to perform an exhaustive classification.¹

An explicit scan over the 4319 three-dimensional reflexive polytopes shows that six $K3$ surfaces with Picard number two that exhibit a genus one fibration with N -sections are realized as generic hypersurfaces in three-dimensional toric ambient spaces. Two more cases have non-toric sections that are combined into a non-toric “pseudo” N -section. Those fibers can still lead to fibrations with $h^{1,1} = 3$ when the components of the pseudo N -section experience monodromy along the $\mathbb{P}^1$ base of the threefold. The data of the eight polytopes is respectively given in (4.8), (4.10) and (4.12). It is also summarized in table 5.

For each of the geometries we will determine the Néron-Severi lattice and the corresponding intersection form as follows: it turns out that each of the eight polytopes admits a unique triangulation that is compatible with a fibration of the toric ambient space over

¹Note that four-dimensional reflexive polytopes are also classified [50]. One can therefore in principle scan over the Calabi-Yau threefolds with $h^{1,1} = 3$ and in each case check if the appropriate fibration structures exist. However, in our case a direct construction based on the classification of $K3$ fibers turns out to be far more efficient and enlightening.

$\mathbb{P}^1$.² As reviewed above, the generic fiber will then be a hypersurface in $\mathbb{P}(1, 2, 3)$, $\mathbb{P}(1, 1, 2)$ or $\mathbb{P}^2$. Each of the fiber polytopes exhibits a ray that respectively corresponds to a 1-, 2- or 3-section. The N -sections, together with the class of the base $\mathbb{P}^1$, then generate the Néron-Severi lattice if the Picard number is two. For two cases we find that the toric N -section splits and there are additional non-toric divisors. We will later be interested in fibrations of $K3$ -surfaces that fuse these non-toric divisors into actual N -sections and will therefore consider only the toric part of the Néron-Severi lattice. The intersection form can then be determined using standard toric intersection calculus.

Given a three dimensional reflexive polytope that corresponds to a $K3$ we can construct a four-dimensional polytope by lifting Δ^{K3} into $\mathbb{Z}^3 \times \mathbb{Z}$ and adding the points

$$\nu'_1 = (\nu, 1), \quad \nu'_2 = (0, -1). \quad (4.5)$$

The convex hull is reflexive whenever $\nu \in 2\Delta^{K3}$ [45]. If the four dimensional polytope admits a fine regular star triangulation that is compatible with the fibration then the generic Calabi-Yau hypersurface in the corresponding toric ambient space will be a $K3$ fibration. In the case of genus one fibrations with 3-sections this will lead us to consider polytopes with the following points

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ n & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ \nu & & & -1 \end{pmatrix}, \quad (4.6)$$

where the admissible values for $n \in \mathbb{Z}$ and $\nu \in \mathbb{Z}^3$ are discussed below. As a result we obtain an exhaustive list of 14 genus one fibered Calabi-Yau threefolds that have 3-sections and also exhibit a compatible $K3$ fibration such that the polarization lattice is of rank two with anti-diagonal intersection form. The results are summarized in table 6 and discussed in more detail below. To facilitate the computations we have used the SageMath computer algebra system [52] as well as PALP [53, 54].

Elliptic fibrations There is only one case with a 1-section which is the degree twelve hypersurface in $\mathbb{P}(1, 1, 4, 6)$. The intersection form on the Néron-Severi lattice is

$$U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (4.7)$$

Let us write the points of the polytope $\Delta^{(1)}$ as

$$\begin{matrix} \nu^1 \\ \nu^2 \\ \nu^3 \\ \nu^4 \\ \nu^5 \end{matrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & -3 & 0 \\ -2 & -3 & 1 \\ -2 & -3 & -1 \end{pmatrix}, \quad (4.8)$$

²For a review of fibration structures in toric varieties see [51].

Polytope	n	ν	χ	Base B	Candidate shift
$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ n & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ & \nu & & -1 \end{pmatrix}$	2	(0, 0, -1)	-168	$\mathbb{F}_1$	
	2	(1, 0, 0)		$\mathbb{F}_0$	
	2	(0, 1, -1)	-156	$\mathbb{F}_1$	$\frac{1}{3}(2, 1^2, 0^5; 2, 1^4, 0^3)$
	-1	(0, 1, -1)		$\mathbb{F}_1$	
	-1	(0, 2, 0)		$\mathbb{F}_0$	
	-1	(-1, 0, 0)		$\mathbb{F}_0$	
	-1	(0, 0, -1)	-150	$\mathbb{F}_1$	
	2	(0, 0, 0)	-144	$\mathbb{F}_0$	
	-1	(0, 0, 0)		$\mathbb{F}_0$	
	-1	(1, 0, -1)	-138	$\mathbb{F}_1$	
	2	(0, 1, 0)	-132	$\mathbb{F}_0$	
	-1	(1, 0, 0)		$\mathbb{F}_0$	
	2	(-2, -2, 0)	-120	$\mathbb{F}_0$	$\frac{1}{3}(2, 0^7; 2, 0^7)$
	-1	(2, 0, 0)		$\mathbb{F}_0$	

Table 6. The 14 genus one fibered Calabi-Yau threefolds $M_{n,\nu}^{(3)}$ that have 3-sections and also exhibit a compatible $K3$ fibration such that the polarization lattice is of rank two with anti-diagonal intersection form. The table contains the polytope in terms of values $n \in \mathbb{Z}$, $\nu \in \mathbb{Z}^3$ that determines the toric ambient space, the Euler characteristic χ and the base B of the genus one fibration. For two of the Euler characteristics we also provide the gauge shifts for a heterotic compactification on $((T^4/\mathbb{Z}_3) \times T^2)/\mathbb{Z}_3$ that leads to the corresponding number of vector- and hypermultiplets that is expected via Heterotic-Type IIA duality. Some fiber Gopakumar-Vafa invariants of the Calabi-Yaus are listed for genus zero in table 7 on page 26 and higher genus invariants, obtained from the modular bootstrap, can be found in table 10 on page 30. They only depend on the Euler characteristic and match with the expectation from the heterotic one-loop computation. In particular, two geometries in this list that exhibit the same Euler characteristic and base are always homotopy equivalent [55, 56].

There are three more points in the interior of facets and they correspond to divisors that do not intersect the generic hypersurface. In general those points will not lift to points in the interior of facets of the four-dimensional polytope. The only inequivalent choices of points $\nu \in 2\Delta^{(1)}$ that lead to a Calabi-Yau threefold with $h^{1,1} = 3$ are

$$\nu = (-4, -6, -n, 1), \quad (4.9)$$

for $n \in \{0, 1, 2\}$. They lead to the well-known elliptic fibrations over $\mathbb{F}_n$. The Gopakumar-Vafa invariants of degree zero with respect to the $\mathbb{P}^1$ base of the $K3$ fibration are the same for all three geometries. Nevertheless, they differ already in the triple intersection numbers and correspond to heterotic compactifications on $K3 \times T^2$ with instantons distributed as $(12 + n, 12 - n)$ among the two E_8 's.

Genus one fibrations with two-sections. Let us also comment on the geometries with two-sections that have been discussed in [9]. Three $K3$ hypersurfaces can be obtained from

the polytope with points

$$\begin{matrix} \nu^1 \\ \nu^2 \\ \nu^3 \\ \nu^4 \\ \nu^5 \end{matrix} \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ -1 & -1 & 0 \\ -2 & n & 1 \\ 0 & 0 & -1 \end{pmatrix}, \quad (4.10)$$

for $n \in \{0, 1, 2\}$. The intersection forms on the corresponding Néron-Severi lattice are

$$I_n^{(2)} = \begin{pmatrix} 2n & 2 \\ 2 & 0 \end{pmatrix}. \quad (4.11)$$

The case $n = 2$ corresponds to a $K3$ with Picard rank three but again a part of NS_{K3} is generated by non-toric divisors that combine into a toric two-section. Note that the intersection forms $I_0^{(2)}$ and $I_2^{(2)}$ are related by a change of basis. In [8, 9] the Gopakumar-Vafa invariants of Calabi-Yau threefolds that exhibit a $K3$ fibration and a genus one fibration with two-sections have been compared against the heterotic one-loop calculation for compactifications on $(K3 \times T^2)/\mathbb{Z}_2$ from [5, 6]. We note that the matching cases are exactly those where the polarization lattice is $I_0^{(2)}$.

Genus one fibrations with three-sections. Three $K3$ surfaces that exhibit a genus one fibration with three-sections are realized as generic hypersurfaces in three-dimensional toric ambient spaces. They correspond to the reflexive polytopes that are the convex hull $\Delta_n^{(3)}$ of the points

$$\begin{matrix} \nu^1 \\ \nu^2 \\ \nu^3 \\ \nu^4 \\ \nu^5 \end{matrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & -1 & 0 \\ n & 0 & 1 \\ 0 & 0 & -1 \end{pmatrix}, \quad (4.12)$$

with $n \in \{-1, 0, 1\}$. The intersection forms on the Néron-Severi lattice are

$$I_n^{(3)} = \begin{pmatrix} 2(1+n) & 3 \\ 3 & 0 \end{pmatrix}. \quad (4.13)$$

For $n = 2$ the corresponding $K3$ surface has Picard rank four but only two generators of NS_{K3} descend from toric divisors on the ambient space. One of the toric divisors corresponds to a “pseudo” three-section with trivial monodromy acting on the intersections with the generic fiber. However, if this geometry is fibered over a $\mathbb{P}^1$, the pseudo three-section can turn into a real three-section. The fibration is then only polarized with respect to a sublattice of the generic fiber. The intersection form $I_2^{(3)}$ with respect to the toric divisors on the $K3$ is equivalent to $I_{-1}^{(3)}$.

Again we can lift $\Delta_n^{(3)}$ into $\mathbb{Z}^3 \times \mathbb{Z}$ and add the points (4.5),

$$\nu'_1 = (\nu, 1), \quad \nu'_2 = (0, -1). \quad (4.14)$$

The convex hull is reflexive whenever $\nu \in 2\Delta_n^{(3)}$. There is always a unique triangulation that is compatible with a fibration of the associated toric variety over a Hirzebruch surface that is induced by the lattice projection onto the last two coordinates. We denote the corresponding Calabi-Yau by $M_{n,\nu}^{(3)}$. Furthermore, we can use lattice automorphisms to impose $\nu_3 \leq 0$ and this does not affect the Calabi-Yau. However, it allows us to identify the divisor that corresponds to the point ν'_2 as the vertical divisor $\pi^{-1}(B)$ and the divisor corresponding to ν^5 as $\pi^{-1}(F)$, where B and F are the base and the fiber of the Hirzebruch surface that is the base of the genus one fibration and π is the projection from the Calabi-Yau onto that base.

The Hodge numbers can be calculated using the standard formula from [48], which is implemented e.g. in PALP [53, 54], and all of the corresponding Calabi-Yau threefolds will have $h^{1,1} = 3$ except if $n = 2$ and ν takes one of the three values

$$\nu = (0, 0, -2), \quad \nu = (1, 0, -1), \quad \nu = (2, 0, 0). \quad (4.15)$$

For all choices of n and ν we can express the Euler characteristic χ of the Calabi-Yau threefold in terms of the vector ν in (4.5),

$$\chi(M_{n,\nu}^{(3)}) = n(-12\nu_1 + 6\nu_2) + 6n^2\nu_3 - 144. \quad (4.16)$$

It turns out that it is exactly the fibrations constructed from the fibers with $n = -1$ and $n = 2$, i.e. those where the intersection form on the polarization lattice is

$$I = \begin{pmatrix} 0 & 3 \\ 3 & 0 \end{pmatrix}, \quad (4.17)$$

that can be matched with weakly coupled heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_3$. When $n \in \{0, 1\}$, the fiber-base duality of the $K3$ fiber and therefore the T-duality of any heterotic dual is broken. For hypersurfaces in toric ambient spaces, the Gopakumar-Vafa invariants at genus zero can be calculated using standard techniques [57]. We find that for $n \in \{-1, 2\}$ the invariants at genus zero and of degree zero with respect to the base of the $K3$ fibration only depend on the Euler characteristic and for low degrees we list the result in table 7. Here the degrees d_B, d_F respectively correspond to the $\mathbb{P}^1$ base and the genus one fiber of the generic $K3$ fiber. More precisely, for a given class $\beta \in H_2(M)$ one has

$$d_F = E_0 \cdot \beta, \quad d_B = \pi^{-1}(B) \cdot \beta, \quad (4.18)$$

where E_0 is the divisor associated to the three-section. This implies that $(d_B, d_F) = (3, 0)$ is the class of the generic genus one fiber while degrees $d_F \equiv 1, 2 \pmod{3}$ correspond to contributions from isolated I_2 singular fibers in special $K3$ fibers. We find a total of 17 inequivalent toric ambient spaces that lead to families of $K3$ fibered Calabi-Yau hypersurfaces with eight different Euler characteristics

$$\chi \in \{-192, -168, -156, -150, -144, -138, -132, -120\}. \quad (4.19)$$

Three models lead to $\chi = -192$ and are special in that the three-section is actually a union of three sections. The number of Kähler moduli is then $h^{1,1} = 5$.

$d_B \backslash d_F$	0	1	2	3	4
0	0	$\frac{\chi}{2} + 240$	$\frac{\chi}{2} + 240$	$-\chi$	$\frac{\chi}{2} + 240$
1	$\frac{\chi}{2} + 240$	$1962 - 3\chi$	$\frac{10\chi}{3} + 18016$	$\frac{15\chi}{2} + 95454$	$413280 - 30\chi$
2	0	$\frac{\chi}{2} + 240$	$\frac{10\chi}{3} + 18016$	$413280 - 30\chi$	$54\chi + 5694624$
3	0	0	$-\chi$	$\frac{15\chi}{2} + 95454$	$54\chi + 5694624$
4	0	0	0	$\frac{\chi}{2} + 240$	$413280 - 30\chi$
5	0	0	0	0	

Table 7. Genus zero Gopakumar-Vafa invariants of degree zero with respect to the base of the $K3$ fibration for the families of Calabi-Yau manifolds $M_{-1,\nu}^{(3)}$ and $M_{2,\nu}^{(3)}$. This matches the heterotic prediction from (2.49) with (2.51).

It turns out that the invariants exactly match those predicted from (3.6) with (2.49) and (2.51) when we identify $d_F = n_1$, $d_B = n_1 + n_2$. In other words, the complexified volume of the generic genus one fiber τ and the volume of the base of the $K3$ fiber t can be related to the complex structure and the complexified Kähler structure T, U on the heterotic side via

$$\tau = U, \quad t = T - U. \quad (4.20)$$

This identification is the same as for the original STU -model. In summary, we find that a $K3$ fibered Calabi-Yau threefold exhibits Gopakumar-Vafa invariants with respect to the $K3$ fiber that match the predictions from the corresponding one-loop amplitudes of heterotic strings on $(K3 \times T^2)/\mathbb{Z}_3$ if and only if the intersection form on the polarization lattice is (4.17).

Two of the heterotic compactifications with non-standard embedding on $(K3 \times T^2)/\mathbb{Z}_3$ led to a maximally higgsable gauge group and the corresponding predictions for the Euler characteristics were $\chi = -120$ and $\chi = -156$. The geometries with $(n; \nu)$ given by $(-1; 2, 0, 0)$ and $(2; -2, -2, 0)$ have Euler characteristic $\chi = -120$. However, both are genus one fibrations over Hirzebruch $\mathbb{F}_0$ and they turn out to have identical topological invariants, i.e. triple intersection numbers and intersections with the second Chern class. It follows from a theorem by Wall [55], see also [56], that these invariants classify the homotopy types of Calabi-Yau threefolds. In particular, all of the genus zero invariants for the two geometries appear to match and, assuming that they are also equivalent as complex manifolds, we therefore have a unique candidate Calabi-Yau dual.

On the other hand, the geometries with the data $(-1; 0, 2, 0)$, $(-1; -1, 0, 0)$, $(-1; 0, 1, -1)$ and $(2; 0, 1, -1)$ have Euler characteristic $\chi = -156$. The former two are again Wall equivalent and fibered over $\mathbb{F}_0$ while the latter two are also Wall equivalent but fibered over $\mathbb{F}_1$. Their intersection rings as well as the Gopakumar-Vafa invariants for non-zero degrees with respect to the base of the $K3$ fibration are different. We therefore find a situation that is similar to the STU models with instanton numbers $(12 - n, 12 + n)$ for $n = 0, 1, 2$. They also produce the same new supersymmetric index but are respectively dual to generic elliptic fibrations over $\mathbb{F}_n$, $n = 0, 1, 2$ [58].

$d_B \backslash d_F$	0	1	2	3	4	5
0	0	$\frac{\chi}{2} + 240$	$\frac{\chi}{2} + 240$	$-\chi$	$\frac{\chi}{2} + 240$	$\frac{\chi}{2} + 240$
1	$\frac{\chi}{2} + 84$	$318 - \chi$	$4776 - \frac{\chi}{2}$	$4\chi + 32262$	$156540 - \frac{11\chi}{2}$	$665484 - 2\chi$
2	0	$\frac{\chi}{2} + 84$	$4776 - \frac{\chi}{2}$	$156540 - \frac{11\chi}{2}$	$\frac{37\chi}{2} + 2492400$	$26767452 - \frac{17\chi}{2}$
3	0	-2	$\frac{\chi}{2} + 240$	$3\chi + 94806$	$5681624 - 27\chi$	$48\chi + 159180264$
4	0	0	0	$4776 - \frac{\chi}{2}$	$\frac{37\chi}{2} + 2492400$	$223507008 - 84\chi$
5	0	0	0	0	$156540 - \frac{11\chi}{2}$	$69\chi + 79435170$

Table 8. Genus zero Gopakumar-Vafa invariants of degree zero with respect to the base of the $K3$ fibration for the families of Calabi-Yau manifolds $M_{1,\nu}^{(3)}$.

$d_B \backslash d_F$	0	1	2	3	4	5
0	0	168	168	144	168	168
1	54	1080	9504	55080	258876	1045440
2	0	1080	55080	1045440	12531888	112746384
3	0	168	94248	5686200	159172380	2868991776
4	0	0	55080	12531888	828397800	29153182176
5	0	0	9504	12531888	2115255492	147357745992

Table 9. Genus zero Gopakumar-Vafa invariants of degree zero with respect to the base of the $K3$ fibration for the families of Calabi-Yau manifolds $M_{0,\nu}^{(3)}$.

Without the CHL orbifold, the instanton number of the non-standard embedding that leads to the predicted value $\chi = -156$ is $n = 3$ [3]. One might be tempted to divide this number by three and conjecture that the geometry with data $(-1; 0, 1, -1)/(2; 0, 1, 2)$ is the correct dual. However, determining the instanton number of the dual heterotic compactification directly from the Calabi-Yau geometry is difficult although a proposal was made in [37]. We leave a further investigation of this issue for future work. Note that the instanton number without CHL quotient of the non-standard embedding that predicts $\chi = -120$ is $n = 0$. This seems compatible with the fact that the unique Calabi-Yau hypersurface in our classification that reproduces this number is fibered over $\mathbb{F}_0$.

The Euler characteristic of all threefolds from $n = 0$ is $\chi(M_{0,\nu}^{(3)}) = -144$ and the Gopakumar-Vafa invariants at genus 0 with respect to the $K3$ fiber given in table 9 while for $n = 1$ the invariants are listed in table 8.

4.3 All genus checks via the modular bootstrap

We now want to use the modular bootstrap [9, 59] in order to obtain all-genus expressions for the topological string amplitudes. Before we apply it to the geometries constructed above, let us briefly review the general technique.

We assume that a Calabi-Yau threefold M is genus one fibered with N -sections over a rational base B such that $h^{1,1}(M) = 1 + h^{1,1}(B)$ and $N \leq 4$. More general cases are covered in [9] but will not be relevant for our discussion. A basis of divisors is given by the three section E_0 , as well as a set of vertical divisors $D_i = \pi^{-1}\tilde{D}_i$, $i = 1, \dots, b_2(B)$ for

a basis of divisors $\tilde{D}_i$ on B . There is a second set of vertical divisors D'_i , $i = 1, \dots, b_2(B)$ that is dual to the curves

$$C_i = \frac{1}{N} E_0 \cdot D_i. \quad (4.21)$$

Finally, one can find a vertical divisor D , such that $\tilde{E}_0 = E_0 + D$ is orthogonal to all of those curves. The complexified Kähler form ω on M can then be parametrized as

$$\omega = \tau \cdot \tilde{E}_0 + \sum_i \tilde{t}_i \cdot D'_i, \quad (4.22)$$

such that τ is the complexified volume of the generic fiber and $\tilde{t}_i$, $i = 1, \dots, h^{1,1}(B)$ are complexified volumes of the dual curves. To make the modular structure manifest one needs to introduce shifted Kähler parameters

$$t_i = \tilde{t}_i + \frac{\tilde{a}_i}{2N} \tau, \quad (4.23)$$

where the coefficients $\tilde{a}_i$ are defined as

$$\tilde{a}_i = \int_M \tilde{E}_0^2 \cdot D_i. \quad (4.24)$$

One can now expand the topological string partition function $Z_{\text{top.}}$ on M as

$$Z_{\text{top.}}(\tau, t, \lambda) = Z_0(\tau, \lambda) \left(1 + \sum_{\beta \in H_2(B)} Z_\beta(\tau, \lambda) Q^\beta \right), \quad (4.25)$$

where λ is the topological string coupling constant and $Q^\beta = \exp(2\pi i t \cdot \beta)$. The coefficients $Z_\beta(\tau, \lambda)$ are conjectured to be meromorphic Jacobi forms of weight 0 and index $\frac{1}{2}\beta \cdot (\beta - c_1(B))$ with respect to λ . More precisely, they take the form

$$Z_\beta(\tau, \lambda) = \frac{\Delta_{2N}(\tau)^{1 - \frac{r_\beta}{N}}}{\eta(N\tau)^{12\beta \cdot c_1(B)}} \frac{\phi_\beta(\tau, \lambda)}{\prod_{i=1}^{b_2(B)} \prod_{k=1}^{\beta_i} \phi_{-2,1}(N\tau, k\lambda)}, \quad (4.26)$$

where $\Delta_{2N}(\tau)$ is a particular modular form for $\Gamma_1(N)$, $\eta(\tau)$ is the Dedekind eta function and $\phi_{-2,1}(\tau, \lambda)$ is a weak Jacobi form. The exponent of Δ_{2N} is the smallest positive solution to the congruence condition

$$1 - \frac{r_\beta}{N} \equiv \frac{1}{2} \left[N c_1(B) - \frac{\tilde{a}}{N} \right] \cdot \beta \pmod{1}. \quad (4.27)$$

The numerator $\phi_\beta(\tau, \lambda)$ is some weak Jacobi form for $\Gamma_1(N)$ and for low degree β it can be fixed, e.g. from the knowledge of some Gopakumar-Vafa invariants.

Let us now apply this procedure to the three-section geometries that we constructed from $\Delta_n^{(3)}$ in the previous section. The toric data of the four-dimensional ambient space is

as follows:

$$\begin{array}{lcl}
 u & \left(\begin{array}{cccc|ccc} 1 & 0 & 0 & 0 & 1 & * & * \end{array} \right) & \leftarrow \text{3-section } E_0 \\
 v & \left(\begin{array}{cccc|ccc} 0 & 1 & 0 & 0 & 1 & * & * \end{array} \right) & \leftarrow \text{3-section } E_0 - n \cdot D_1 + (\nu_2 - \nu_1) \cdot D_2 \\
 w & \left(\begin{array}{cccc|ccc} -1 & -1 & 0 & 0 & 1 & * & * \end{array} \right) & \leftarrow \text{3-section } E_0 - n \cdot D_1 - \nu_1 \cdot D_2 \\
 a_1 & \left(\begin{array}{cccc|ccc} n & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \right) & \leftarrow \text{vertical divisor } D'_2 = \pi^{-1}(B - \nu_3 F) \\
 a_2 & \left(\begin{array}{cccc|ccc} 0 & 0 & -1 & 0 & 0 & 1 & \nu_3 \end{array} \right) & \leftarrow \text{vertical divisor } D_1 = \pi^{-1}(B) \\
 b_1 & \left(\begin{array}{cccc|ccc} \nu_1 & \nu_2 & \nu_3 & 1 & 0 & 0 & 1 \end{array} \right) & \leftarrow \text{vertical divisor } D_2 = D'_1 = \pi^{-1}(F) \\
 b_2 & \left(\begin{array}{cccc|ccc} 0 & 0 & 0 & -1 & 0 & 0 & 1 \end{array} \right) & \leftarrow \text{---"} \\
 & \left(\begin{array}{cccc|ccc} 0 & 0 & 0 & 0 & -3 & -1 & * \end{array} \right) &
 \end{array}$$

In the first column we list the homogeneous coordinates that are associated to the rays spanned by the vertices of the polytope. Recall that we used lattice automorphisms to fix $\nu_3 \leq 0$ such that B and F are the base and fiber of the Hirzebruch surface $\mathbb{F}_{-\nu_3}$ that is the base of the genus one fibration. While one can give closed expressions for the empty values in the Mori-cone, they are somewhat bulky and we do not require them in the following discussion. For every value of n and ν we pick the unique triangulation that is compatible with the genus one fibration over $\mathbb{F}_{-\nu_3}$.

We choose $\{u = 0\}$ as the “zero” three-section and denote the corresponding divisor class by $E_0 = [u]$. Using standard toric intersection calculus we then find

$$E_0^2 \cdot D_1 = -4 \cdot \nu_1 + 2 \cdot \nu_2 + \nu_3 + 2, \quad E_0^2 \cdot D_2 = 2 - 4 \cdot n. \quad (4.28)$$

as intersections on the generic Calabi-Yau hypersurface. This implies that

$$\tilde{E}_0 = E_0 + \frac{2}{3}(2 \cdot n - 1)D_1 + \frac{1}{3}(4 \cdot n + 4 \cdot \nu_1 - 2 \cdot \nu_2 - \nu_3 - 4)D_2, \quad (4.29)$$

is orthogonal to the curves $E_0 \cdot D_{1/2}$ and we calculate

$$\tilde{a}_1 = \tilde{E}_0^2 \cdot D_2 = 4 \cdot n - 2, \quad \tilde{a}_2 = \tilde{E}^2 \cdot D_1 = 4 \cdot \nu_1 - 2 \cdot \nu_2 - \nu_3 - 2. \quad (4.30)$$

The correct parametrization of the Kähler form to obtain modular amplitudes is then

$$\omega = \tau \cdot \tilde{E}_0 + \left(t_1 - \frac{1}{6}\tilde{a}_1\right) D'_1 + \left(t_2 - \frac{1}{6}\tilde{a}_2\right) D'_2. \quad (4.31)$$

We want to apply the modular bootstrap to obtain all genus amplitudes with respect to the $K3$ fiber, i.e. we are interested in Z_β for $\beta = k \cdot F$. The exponent r_β in (4.26) is then given by

$$1 - \frac{r_{k \cdot F}}{3} \equiv \frac{k}{3}(2 \cdot n - 1) \pmod{1}. \quad (4.32)$$

For $n \in \{-1, 2\}$ and $k = 1$ we use genus zero Gromov-Witten invariants to fix the ansatz

$$Z_1(\tau, \lambda) = \frac{1}{48} \frac{\Delta_6(\tau)}{\eta(3\tau)^{24} \phi_{-2,1}(3\tau, \lambda)} \left[(120 + \chi) E_4(\tau) - 9(152 + \chi) \mathcal{E}_3(\tau)^2 \right], \quad (4.33)$$

where we refer to the appendix A for the definition of the modular and Jacobi forms. From this we can extract Gopakumar-Vafa invariants for arbitrary fiber degrees and genera and the results for low orders are listed in table 10. This matches with the predictions from the heterotic calculation and provides another strong check of the duality.

$d_F \backslash g$	0	1	2	3	4	5
0	$\frac{\chi}{6} + 26$	0	0	0	0	0
1	$\frac{\chi}{2} + 240$	0	0	0	0	0
2	$1962 - 3\chi$	0	0	0	0	0
3	$\frac{10\chi}{3} + 18016$	$-\frac{\chi}{3} - 52$	0	0	0	0
4	$\frac{15\chi}{2} + 95454$	$-\chi - 480$	0	0	0	0
5	$413280 - 30\chi$	$6\chi - 3924$	0	0	0	0
6	$\frac{88\chi}{3} + 1627330$	$-\frac{23\chi}{3} - 36188$	$\frac{\chi}{2} + 78$	0	0	0
7	$54\chi + 5694624$	$-18\chi - 192348$	$\frac{3\chi}{2} + 720$	0	0	0
8	$18353988 - 195\chi$	$78\chi - 838332$	$5886 - 9\chi$	0	0	0
9	$170\chi + 55646304$	$-80\chi - 3362964$	$\frac{38\chi}{3} + 54464$	$-\frac{2\chi}{3} - 104$	0	0
10	$291\chi + 159217686$	$-157\chi - 11963892$	$\frac{61\chi}{2} + 290202$	$-2\chi - 960$	0	0

Table 10. Gopakumar-Vafa invariants of degree zero with respect to the base of the $K3$ fibration and degree one with respect to the base of the $K3$ fiber for the families of Calabi-Yau manifolds $M_{-1,\nu}^{(3)}$ and $M_{2,\nu}^{(3)}$.

5 Conclusions

In this paper we have explicitly calculated the new supersymmetric index for several heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_3$. To this end we considered a $T^4/\mathbb{Z}_3$ orbifold limit of the $K3$ with all of the five possible embeddings of the $\mathbb{Z}_3$ action into the gauge group. The index encodes the gauge and gravitational coupling of the $\mathcal{N} = 2$ effective theory that arise from heterotic one-loop amplitudes and leads to predictions for certain Gopakumar-Vafa invariants of dual Calabi-Yau compactification spaces. This generalizes earlier calculations for compactifications on $K3 \times T^2$ [1–3] as well as the subsequent extensions to $(K3 \times T^2)/\mathbb{Z}_N$, where for $N = 3$ only the standard embedding has been considered [5, 6]. In particular, we obtain two vacua where the hypermultiplet spectrum enables us to maximally higgs the gauge group to $U(1)^3$.

Let us note that for all choices of embeddings except for the standard embedding, the new supersymmetric index depends only on the difference $N_h - N_v$ of the number of vector- and hypermultiplets. This is the same behaviour that was observed for *all* embeddings when the heterotic string is compactified on $K3 \times T^2$ [3] or $(K3 \times T^2)/\mathbb{Z}_2$ [6]. It would be interesting to understand the reason why the standard embedding for $\mathbb{Z}_3$ orbifolds is special. We hope to come back to this question in future work.

We then systematically constructed candidate Calabi-Yau duals that exhibit both a $K3$ and a genus one fibration structure, as well as three sections with respect to the latter. In this way we could show that the enumerative invariants exhibit the correct structure if and only if the intersection form on the polarization lattice is three times the intersection form on the Narain lattice $\Gamma^{1,1}$. Physically, this can be interpreted as the requirement that the dual heterotic compactification exhibits T-duality. For each of the two heterotic models that can be higgsed to $U(1)^3$ we find Calabi-Yau manifolds that reproduce the predicted enumerative invariants. We apply the modular bootstrap [9] to obtain all genus results for

a subset of the topological string amplitudes and in that way provide a strong check of the duality. Our discussion generalizes and provides a more systematic understanding of earlier constructions of Calabi-Yau duals for heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_2$ [7–9].

Multiple geometries that match the same new supersymmetric index are distinguished by their classical intersection numbers and by those enumerative invariants that lead to predictions for what from the heterotic perspective are non-perturbative corrections. This situation is analogous to that encountered already for heterotic compactifications on $K3 \times T^2$. It is well known that the embeddings of the gauge connection with instanton numbers $(12 - n, 12 + n)$ for $n = 0, 1, 2$ produce the same new supersymmetric index and are respectively dual to type IIA compactifications on generic elliptic fibrations over the Hirzebruch surfaces $\mathbb{F}_n$ [58]. However, it is quite difficult to derive the heterotic instanton numbers from the geometry of the Calabi-Yau or the classical intersection numbers of the Calabi-Yau from a heterotic calculation, see e.g. [37] for a discussion of this issue. It will be very interesting to study this problem in the context of heterotic compactifications on $(K3 \times T^2)/\mathbb{Z}_N$ and their dual Calabi-Yau manifolds. We leave this as another question that we hope to address in future work.

A Conventions for modular forms

In this appendix we summarize the definitions of the various modular and Jacobi forms that appear throughout the paper. The Dedekind eta function is defined as

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n), \quad (\text{A.1})$$

where $q = \exp(2\pi i\tau)$. It is a modular form of weight $1/2$ with a multiplier system and transforms as

$$\eta(\tau + 1) = e^{\frac{\pi i}{12}} \eta(\tau), \quad \eta(-1/\tau) = \sqrt{-i\tau} \eta(\tau). \quad (\text{A.2})$$

The classical Eisenstein series can be written as

$$E_4(\tau) = 1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n}, \quad E_6(\tau) = 1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n}, \quad (\text{A.3})$$

and are modular forms of respective weight four and six.

We will also need to consider modular forms for the congruence subgroup $\Gamma_1(3)$ of the modular group. The corresponding ring can be generated by three functions E_4, E_6 and $\mathcal{E}_3$, where $\mathcal{E}_3$ is a holomorphic modular form of weight two that can be expressed in terms of the Dedekind eta function via

$$\mathcal{E}_3(\tau) = -\frac{24}{2\pi i} \partial_{\tau} \log \left(\frac{\eta(\tau)}{\eta(3\tau)} \right). \quad (\text{A.4})$$

Moreover, the unique weight six modular form for $\Gamma_1(3)$ with a second order zero at $\tau \rightarrow i\infty$ is given by

$$\Delta_6(\tau) = \frac{\eta(3\tau)^{18}}{\eta(\tau)^6}. \quad (\text{A.5})$$

The generalized Jacobi theta function can be written as

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} (\tau, z) = \sum_{k \in \mathbb{Z}} q^{\frac{1}{2}(k + \frac{a}{2})^2} e^{\pi i (k + \frac{a}{2})b} e^{(2\pi i z)(k + \frac{a}{2})}, \quad (\text{A.6})$$

and one introduces

$$\begin{aligned} \theta_1(\tau, z) &= \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (\tau, z), & \theta_2(\tau, z) &= \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (\tau, z), \\ \theta_3(\tau, z) &= \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (\tau, z), & \theta_4(\tau, z) &= \theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} (\tau, z). \end{aligned} \quad (\text{A.7})$$

This determines also the weak Jacobi form $\phi_{-2,1}(\tau, z)$ of weight -2 and index 1 via

$$\phi_{-2,1}(\tau, z) = -\frac{\theta_1(\tau, z)^2}{\eta(\tau)^6}, \quad (\text{A.8})$$

that we need for the modular bootstrap.

B Higgsing of the gauge group

In this appendix we will explain in some detail how the gauge group of the models of table 3 can be higgsed by giving vacuum expectation values (vev's) to scalars in the hypermultiplets. We take the necessary branching rules from [60]. For more on higgsing see for example [19, 61–63].

For concreteness sake we will study the third model of table 3. Before higgsing we have the following gauge group and matter content, coming from the twisted and untwisted sector

$$\begin{aligned} &\text{SO}(14) \times \text{SO}(14) \times \text{U}(1)^2 \\ &4(\mathbf{14}, \mathbf{1}) + 4(\mathbf{1}, \mathbf{14}) + (\mathbf{64}, \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 8(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.1})$$

The matter content is labelled by the representations under the two $\text{SO}(14)$ groups, i.e. we have left out the $\text{U}(1)$ charges. Counting degrees of freedom we find the numbers of vector- and hypermultiplets $N_v = 184$ and $N_h = 248$. We will start by explaining how to higgs the first $\text{SO}(14)$ factor.

For this we notice the following branching rules

$$\text{SO}(14) \supset \text{SO}(13) : \mathbf{14} \rightarrow \mathbf{13} + \mathbf{1}, \mathbf{64} \rightarrow \mathbf{64}. \quad (\text{B.2})$$

Giving a vev to one $\mathbf{14}$ will break $\text{SO}(14)$ to $\text{SO}(13)$. One $\mathbf{13}$ will get ‘eaten’ by the broken generators of the gauge fields (turning them massive) and the result is the following gauge group and matter spectrum

$$\begin{aligned} &\text{SO}(13) \times \text{SO}(14) \times \text{U}(1)^2, \\ &3(\mathbf{13}, \mathbf{1}) + 4(\mathbf{1}, \mathbf{14}) + (\mathbf{64}, \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 12(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.3})$$

We may check that $N_h - N_v = 235 - 171 = 64$ is unchanged. In the next step we give a vev to a **13** and use the branching rules

$$\mathrm{SO}(13) \supset \mathrm{SO}(12) : \mathbf{13} \rightarrow \mathbf{12} + \mathbf{1}, \mathbf{64} \rightarrow \mathbf{32} + \mathbf{32}'. \quad (\text{B.4})$$

Similar to before a **12** gets ‘eaten’ by the broken generators of $\mathrm{SO}(13)$ and we find the following gauge group and matter spectrum after higgsing

$$\begin{aligned} & \mathrm{SO}(12) \times \mathrm{SO}(14) \times \mathrm{U}(1)^2, \\ & 2(\mathbf{12}, \mathbf{1}) + 4(\mathbf{1}, \mathbf{14}) + (\mathbf{32}, \mathbf{1}) + (\mathbf{32}', \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 15(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.5})$$

Two more similar steps lead to the following gauge group and matter spectrum

$$\begin{aligned} & \mathrm{SO}(10) \times \mathrm{SO}(14) \times \mathrm{U}(1)^2, \\ & 4(\mathbf{1}, \mathbf{14}) + 2(\mathbf{16}, \mathbf{1}) + 2(\overline{\mathbf{16}}, \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 18(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.6})$$

The branching rules

$$\begin{aligned} \mathrm{SO}(10) \supset \mathrm{SU}(5) \times \mathrm{U}(1) : \mathbf{16} & \rightarrow \mathbf{10}(\mathbf{1}) + \bar{\mathbf{5}}(-\mathbf{3}) + \mathbf{1}(\mathbf{5}), \\ \overline{\mathbf{16}} & \rightarrow \bar{\mathbf{10}}(-\mathbf{1}) + \mathbf{5}(\mathbf{3}) + \mathbf{1}(-\mathbf{5}) \end{aligned} \quad (\text{B.7})$$

(the number in brackets give the $\mathrm{U}(1)$ charge) indicate that we can give a vev to **16** and $\overline{\mathbf{16}}$ thereby breaking $\mathrm{SO}(10)$ to $\mathrm{SU}(5)$ where $\mathbf{10}, \bar{\mathbf{10}}$ and one scalar will get ‘eaten’ by the broken generators. The gauge group and spectrum after this step of higgsing are thus

$$\begin{aligned} & \mathrm{SU}(5) \times \mathrm{SO}(14) \times \mathrm{U}(1)^2, \\ & 4(\mathbf{1}, \mathbf{14}) + 2(\mathbf{5}, \mathbf{1}) + 2(\bar{\mathbf{5}}, \mathbf{1}) + (\mathbf{10}, \mathbf{1}) + (\bar{\mathbf{10}}, \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 21(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.8})$$

In the next step we give a vev to **5**, $\bar{\mathbf{5}}$ and use the branching rules

$$\begin{aligned} \mathrm{SU}(5) \supset \mathrm{SU}(4) \times \mathrm{U}(1) : \mathbf{5} & \rightarrow \mathbf{4}(\mathbf{1}) + \mathbf{1}(-\mathbf{4}), \\ \mathbf{10} & \rightarrow \mathbf{4}(-\mathbf{3}) + \mathbf{6}(\mathbf{2}) \end{aligned} \quad (\text{B.9})$$

to obtain the gauge group and spectrum

$$\begin{aligned} & \mathrm{SU}(4) \times \mathrm{SO}(14) \times \mathrm{U}(1)^2, \\ & 4(\mathbf{1}, \mathbf{14}) + 2(\mathbf{4}, \mathbf{1}) + 2(\bar{\mathbf{4}}, \mathbf{1}) + (\mathbf{6}, \mathbf{1}) + (\bar{\mathbf{6}}, \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 24(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.10})$$

We proceed by breaking $\mathrm{SU}(4)$ to $\mathrm{SU}(3)$ by giving a vev to **4**, $\bar{\mathbf{4}}$ and using the branching rules

$$\begin{aligned} \mathrm{SU}(4) \supset \mathrm{SU}(3) \times \mathrm{U}(1) : \mathbf{4} & \rightarrow \mathbf{3}(\mathbf{1}) + \mathbf{1}(-\mathbf{3}), \\ \mathbf{6} & \rightarrow \mathbf{3}(-\mathbf{2}) + \bar{\mathbf{3}}(\mathbf{2}). \end{aligned} \quad (\text{B.11})$$

We obtain the following gauge group and spectrum

$$\begin{aligned} & \mathrm{SU}(3) \times \mathrm{SO}(14) \times \mathrm{U}(1)^2, \\ & 4(\mathbf{1}, \mathbf{14}) + 3(\mathbf{3}, \mathbf{1}) + 3(\bar{\mathbf{3}}, \mathbf{1}) + (\mathbf{1}, \mathbf{64}) + 27(\mathbf{1}, \mathbf{1}). \end{aligned} \quad (\text{B.12})$$

We can continue in similar manner using $3(\mathbf{3}, \mathbf{1})$ and $3(\bar{\mathbf{3}}, \mathbf{1})$ to higgs $SU(3)$ completely and end up with

$$\begin{aligned} &SO(14) \times U(1)^2, \\ &4(\mathbf{1}, \mathbf{14}) + (\mathbf{1}, \mathbf{64}) + 37(\mathbf{1}, \mathbf{1}). \end{aligned} \tag{B.13}$$

In the same way we may higgs the second $SO(14)$. The two $U(1)$ factors in the gauge group may be higgsed by any of the charged scalars. Thus we end up with a completely higgsed gauge group and 64 neutral scalars.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.A. Harvey and G.W. Moore, *Algebras, BPS states and strings*, *Nucl. Phys. B* **463** (1996) 315 [[hep-th/9510182](#)] [[INSPIRE](#)].
- [2] M. Mariño and G.W. Moore, *Counting higher genus curves in a Calabi-Yau manifold*, *Nucl. Phys. B* **543** (1999) 592 [[hep-th/9808131](#)] [[INSPIRE](#)].
- [3] S. Stieberger, *(0, 2) heterotic gauge couplings and their M-theory origin*, *Nucl. Phys. B* **541** (1999) 109 [[hep-th/9807124](#)] [[INSPIRE](#)].
- [4] S. Cecotti, P. Fendley, K.A. Intriligator and C. Vafa, *A new supersymmetric index*, *Nucl. Phys. B* **386** (1992) 405 [[hep-th/9204102](#)] [[INSPIRE](#)].
- [5] A. Chattopadhyaya and J.R. David, *Gravitational couplings in $\mathcal{N} = 2$ string compactifications and Mathieu Moonshine*, *JHEP* **05** (2018) 211 [[arXiv:1712.08791](#)] [[INSPIRE](#)].
- [6] A. Chattopadhyaya and J.R. David, *$\mathcal{N} = 2$ heterotic string compactifications on orbifolds of $K3 \times T^2$* , *JHEP* **01** (2017) 037 [[arXiv:1611.01893](#)] [[INSPIRE](#)].
- [7] S. Kachru, A. Klemm and Y. Oz, *Calabi-Yau duals for CHL strings*, *Nucl. Phys. B* **521** (1998) 58 [[hep-th/9712035](#)] [[INSPIRE](#)].
- [8] A. Banlaki, A. Chowdhury, A. Kidambi and M. Schimpf, *On Mathieu moonshine and Gromov-Witten invariants*, *JHEP* **02** (2020) 082 [[arXiv:1811.11619](#)] [[INSPIRE](#)].
- [9] C.F. Cota, A. Klemm and T. Schimannek, *Topological strings on genus one fibered Calabi-Yau 3-folds and string dualities*, *JHEP* **11** (2019) 170 [[arXiv:1910.01988](#)] [[INSPIRE](#)].
- [10] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Holomorphic anomalies in topological field theories*, *Nucl. Phys. B* **405** (1993) 279 [[hep-th/9302103](#)] [[INSPIRE](#)].
- [11] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes*, *Commun. Math. Phys.* **165** (1994) 311 [[hep-th/9309140](#)] [[INSPIRE](#)].
- [12] I. Antoniadis, E. Gava, K.S. Narain and T.R. Taylor, *Topological amplitudes in string theory*, *Nucl. Phys. B* **413** (1994) 162 [[hep-th/9307158](#)] [[INSPIRE](#)].
- [13] A. Klemm, W. Lerche and P. Mayr, *$K3$ fibrations and heterotic type-II string duality*, *Phys. Lett. B* **357** (1995) 313 [[hep-th/9506112](#)] [[INSPIRE](#)].

- [14] S. Mukai, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*, *Invent. Math.* **94** (1988) 183.
- [15] M.R. Gaberdiel, S. Hohenegger and R. Volpato, *Symmetries of K3 σ -models*, *Commun. Num. Theor. Phys.* **6** (2012) 1 [[arXiv:1106.4315](#)] [[INSPIRE](#)].
- [16] J.R. David, D.P. Jatkar and A. Sen, *Product representation of Dyon partition function in CHL models*, *JHEP* **06** (2006) 064 [[hep-th/0602254](#)] [[INSPIRE](#)].
- [17] A. Chattopadhyaya and J.R. David, *Dyon degeneracies from Mathieu moonshine symmetry*, *Phys. Rev. D* **96** (2017) 086020 [[arXiv:1704.00434](#)] [[INSPIRE](#)].
- [18] L. Dixon, J. Harvey, C. Vafa and E. Witten, *Strings on orbifolds II*, *Nucl. Phys. B* **274** (1986) 285.
- [19] G. Honecker and M. Trapletti, *Merging heterotic orbifolds and K3 compactifications with line bundles*, *JHEP* **01** (2007) 051 [[hep-th/0612030](#)] [[INSPIRE](#)].
- [20] J.R. David, D.P. Jatkar and A. Sen, *Dyon spectrum in generic $N = 4$ supersymmetric $Z(N)$ orbifolds*, *JHEP* **01** (2007) 016 [[hep-th/0609109](#)] [[INSPIRE](#)].
- [21] J.R. David and A. Sen, *CHL dyons and statistical entropy function from D1-D5 system*, *JHEP* **11** (2006) 072 [[hep-th/0605210](#)] [[INSPIRE](#)].
- [22] M.R. Gaberdiel, S. Hohenegger and R. Volpato, *Mathieu twining characters for K3*, *JHEP* **09** (2010) 058 [[arXiv:1006.0221](#)] [[INSPIRE](#)].
- [23] T. Eguchi, H. Ooguri and Y. Tachikawa, *Notes on the K3 surface and the Mathieu group M_{24}* , *Exper. Math.* **20** (2011) 91 [[arXiv:1004.0956](#)].
- [24] T. Eguchi and K. Hikami, *Note on twisted elliptic genus of K3 surface*, *Phys. Lett. B* **694** (2011) 446 [[arXiv:1008.4924](#)] [[INSPIRE](#)].
- [25] M.C.N. Cheng, *K3 surfaces, $N = 4$ Dyons and the Mathieu group M_{24}* , *Commun. Num. Theor. Phys.* **4** (2010) 623 [[arXiv:1005.5415](#)] [[INSPIRE](#)].
- [26] M.A. Walton, *The heterotic string on the simplest Calabi-Yau manifold and its orbifold limits*, *Phys. Rev. D* **37** (1988) 377 [[INSPIRE](#)].
- [27] G. Aldazabal, A. Font, L.E. Ibáñez and F. Quevedo, *Chains of $N = 2$, $D = 4$ heterotic type-II duals*, *Nucl. Phys. B* **461** (1996) 85 [[hep-th/9510093](#)] [[INSPIRE](#)].
- [28] M.B. Green, J.H. Schwarz and P.C. West, *Anomaly free chiral theories in six-dimensions*, *Nucl. Phys. B* **254** (1985) 327 [[INSPIRE](#)].
- [29] M. Henningson and G.W. Moore, *Threshold corrections in $K3 \times T^2$ heterotic string compactifications*, *Nucl. Phys. B* **482** (1996) 187 [[hep-th/9608145](#)] [[INSPIRE](#)].
- [30] R. Gopakumar and C. Vafa, *M theory and topological strings. 1*, [hep-th/9809187](#) [[INSPIRE](#)].
- [31] R. Gopakumar and C. Vafa, *M theory and topological strings. 2*, [hep-th/9812127](#) [[INSPIRE](#)].
- [32] R.E. Borcherds, *Automorphic forms with singularities on Grassmannians*, *Invent. Math.* **132** (1998) 491 [[alg-geom/9609022](#)] [[INSPIRE](#)].
- [33] I. Antoniadis, E. Gava and K.S. Narain, *Moduli corrections to gravitational couplings from string loops*, *Phys. Lett. B* **283** (1992) 209 [[hep-th/9203071](#)] [[INSPIRE](#)].
- [34] I. Antoniadis, E. Gava, K.S. Narain and T.R. Taylor, *$N = 2$ type-II heterotic duality and higher derivative F terms*, *Nucl. Phys. B* **455** (1995) 109 [[hep-th/9507115](#)] [[INSPIRE](#)].

- [35] I. Antoniadis, E. Gava and K.S. Narain, *Moduli corrections to gauge and gravitational couplings in four-dimensional superstrings*, *Nucl. Phys. B* **383** (1992) 93 [[hep-th/9204030](#)] [[INSPIRE](#)].
- [36] P.S. Aspinwall, *K3 surfaces and string duality*, in *Differential geometry inspired by string theory*, S.T. Yau ed., International Press, U.S.A. (1996), [[hep-th/9611137](#)] [[INSPIRE](#)].
- [37] A.P. Braun and T. Watari, *Heterotic-type IIA duality and degenerations of K3 surfaces*, *JHEP* **08** (2016) 034 [[arXiv:1604.06437](#)] [[INSPIRE](#)].
- [38] C.M. Hull and P.K. Townsend, *Unity of superstring dualities*, *Nucl. Phys. B* **438** (1995) 109 [[hep-th/9410167](#)] [[INSPIRE](#)].
- [39] E. Witten, *String theory dynamics in various dimensions*, *Nucl. Phys. B* **443** (1995) 85 [[hep-th/9503124](#)] [[INSPIRE](#)].
- [40] S. Kachru and C. Vafa, *Exact results for $N = 2$ compactifications of heterotic strings*, *Nucl. Phys. B* **450** (1995) 69 [[hep-th/9505105](#)] [[INSPIRE](#)].
- [41] C. Vafa and E. Witten, *Dual string pairs with $N = 1$ and $N = 2$ supersymmetry in four-dimensions*, *Nucl. Phys. Proc. Suppl.* **46** (1996) 225 [[hep-th/9507050](#)] [[INSPIRE](#)].
- [42] S. Kachru et al., *Nonperturbative results on the point particle limit of $N = 2$ heterotic string compactifications*, *Nucl. Phys. B* **459** (1996) 537 [[hep-th/9508155](#)] [[INSPIRE](#)].
- [43] A. Klemm, M. Kreuzer, E. Riegler and E. Scheidegger, *Topological string amplitudes, complete intersection Calabi-Yau spaces and threshold corrections*, *JHEP* **05** (2005) 023 [[hep-th/0410018](#)] [[INSPIRE](#)].
- [44] T. Schimannek, *Modularity from monodromy*, *JHEP* **05** (2019) 024 [[arXiv:1902.08215](#)] [[INSPIRE](#)].
- [45] V. Braun and D.R. Morrison, *F-theory on genus-one fibrations*, *JHEP* **08** (2014) 132 [[arXiv:1401.7844](#)] [[INSPIRE](#)].
- [46] D. Huybrechts, *Lectures on K3 surfaces*, *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge U.K. (2016).
- [47] A. Grassi, *On minimal models of elliptic threefolds*, *Math. Ann.* **290** (1991) 287.
- [48] V.V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, (1993).
- [49] M. Kreuzer and H. Skarke, *Classification of reflexive polyhedra in three-dimensions*, *Adv. Theor. Math. Phys.* **2** (1998) 853 [[hep-th/9805190](#)] [[INSPIRE](#)].
- [50] M. Kreuzer and H. Skarke, *Complete classification of reflexive polyhedra in four-dimensions*, *Adv. Theor. Math. Phys.* **4** (2002) 1209 [[hep-th/0002240](#)] [[INSPIRE](#)].
- [51] V. Braun, *Toric elliptic fibrations and F-theory compactifications*, *JHEP* **01** (2013) 016 [[arXiv:1110.4883](#)] [[INSPIRE](#)].
- [52] The Sage developers, *SageMath, the Sage Mathematics Software System*, version 8.2 (2019).
- [53] M. Kreuzer and H. Skarke, *PALP: a package for analyzing lattice polytopes with applications to toric geometry*, *Comput. Phys. Commun.* **157** (2004) 87 [[math/0204356](#)] [[INSPIRE](#)].
- [54] A.P. Braun et al., *PALP — A user manual*, in *Strings, gauge fields and the geometry behind: the legacy of Maximilian Kreuzer*, A. Rebhan et al. eds., World Scientific, Singapore (2012), [[arXiv:1205.4147](#)] [[INSPIRE](#)].

- [55] C. Wall, *Classification problems in differential topology. v. on certain 6-manifolds.*, *Invent. Math.* **1** (1966) 355.
- [56] A.-m. He and P. Candelas, *On the number of complete intersection Calabi-Yau manifolds*, *Commun. Math. Phys.* **135** (1990) 193 [[INSPIRE](#)].
- [57] S. Hosono, A. Klemm, S. Theisen and S.-T. Yau, *Mirror symmetry, mirror map and applications to Calabi-Yau hypersurfaces*, *Commun. Math. Phys.* **167** (1995) 301 [[hep-th/9308122](#)] [[INSPIRE](#)].
- [58] D.R. Morrison and C. Vafa, *Compactifications of F-theory on Calabi-Yau threefolds. 2.*, *Nucl. Phys. B* **476** (1996) 437 [[hep-th/9603161](#)] [[INSPIRE](#)].
- [59] M.-x. Huang, S. Katz and A. Klemm, *Topological string on elliptic CY 3-folds and the ring of Jacobi forms*, *JHEP* **10** (2015) 125 [[arXiv:1501.04891](#)] [[INSPIRE](#)].
- [60] N. Yamatsu, *Finite-dimensional Lie algebras and their representations for unified model building*, [arXiv:1511.08771](#) [[INSPIRE](#)].
- [61] M. Srednicki, *Quantum field theory*, Cambridge University Press, Cambridge U.K. (2007).
- [62] R. Slansky, *Group theory for unified model building*, *Phys. Rept.* **79** (1981) 1 [[INSPIRE](#)].
- [63] C. Nazaroglu, *Jacobi forms of higher index and paramodular groups in $\mathcal{N} = 2, D = 4$ compactifications of string theory*, *JHEP* **12** (2013) 074 [[arXiv:1309.6404](#)] [[INSPIRE](#)].