Recoupling mechanism for exotic mesons and baryons

Yu.A. Simonov

*NRC “Kurchatov Institute” — ITEP,
B. Cheremushkinskaya 25, Moscow, 117259, Russia*

E-mail: simonov@itep.ru

ABSTRACT: The infinite chain of transitions of one pair of mesons (channel I) into another pair of mesons (channel II) can produce bound states and resonances in both channels even if no interactions inside channels exist. These resonances which can occur also in meson-baryon channels are called channel-coupling (CC) resonances. A new mechanism of CC resonances is proposed where transitions occur due to a rearrangement of confining strings inside each channel — the recoupling mechanism. The amplitude of this recoupling mechanism is expressed via overlap integrals of the wave functions of participating mesons (baryons). The explicit calculation with the known wave functions yields the peak at $E = 4.12 \text{ GeV}$ for the transitions $J/\psi + \phi \leftrightarrow D_s^* + \bar{D}_s^*$, which can be associated with $\chi_{c1}(4140)$, and a narrow peak at 3.98 GeV with the width 10 MeV for the transitions $D_s^- + D_0^* \leftrightarrow J/\psi + K^{*-}$, which can be associated with the recently discovered $Z_{cs}(3985)$.

KEYWORDS: Phenomenological Models

ARXIV EPRINT: [2011.12326](https://arxiv.org/abs/2011.12326)

Contents

1	Introduction	1
2	The simplest case: only separable CC interaction	4
3	Equations for two channel amplitudes in the recoupling formalism	5
4	Recoupling mechanism for the meson-meson amplitudes	8
5	Recoupling mechanism for meson-baryon systems	9
6	Physical amplitudes and their singularities in the recoupling process	11
7	Numerical results and discussion	13
8	Conclusions and an outlook	15
A	The center-of-mass coordinates and average quark and antiquark energies in a hadron	16
B	The channel-coupling coefficient $\bar{y}_{1234}$	16
C	Oscillator parameters of hadron wave function	18
D	Calculation of the cross sections $f_{ik} ^2$ and pole positions	18

1 Introduction

The modern situation with the spectra of quarkonia and baryons requires the dynamical explanation of numerous extra states, which are not present in the one-channel spectra of a given meson [1].¹ A similar situation occurs in the excited baryon spectra.² To be more precise, in the case of heavy quarkonia, i.e. states, which contain $c\bar{c}$ and $b\bar{b}$ pairs, the experimental data contain a number of charged Z_c, Z_b and neutral Y_c, Y_b states, which cannot be explained by the dynamics of $c\bar{c}$, or $b\bar{b}$ pairs alone, see [2] for review.

There are theoretical suggestions of different mechanisms [3–24], which should be taken into account. E.g. poles (resonances) in the meson-meson channels can occur due to strong interaction in these systems, and appear as additional poles in the S matrix [4, 7, 8, 18, 19] — the molecular-type approach.

¹See also the update on 6th December 2019.

²M. Karliner and T. Skwarnicki, *Pentaquarks*, in [1].

A similar in the choice of the driving channels ($Q\bar{q}\bar{Q}q$), but different in the dynamics, is the approach of the tetraquark model [5, 10, 12–14, 17, 20, 24], see [25–27] for reviews. A more general approach contains features of both molecular and tetraquark models — the hybridized tetraquark model [28]. One of the basic features of these models (as well as of the hadron-hadron interaction in general) is the determination of the dynamics in the system of two white objects. In the case of a deep tetraquark state one can disregard the two-body white asymptotics with some accuracy, however in the molecular case and in general one needs a classification of possible dynamical exchanges V_{hh} between two white hadrons, taking into account the full gauge invariance and the confinement via general Wilson loop representation, which is now standard. It is clear that one-gluon exchange and its spin versions (spin-spin, spin-orbit, tensor) are not present in V_{hh} , while two-gluon exchange is the glueball exchange. The “exchange of confining interaction” is not a local potential but the transition of two white objects into a common white one with $q\bar{q}$ annihilations before and after, e.g. $\pi + \pi \rightarrow \rho \rightarrow \pi + \pi$. In a similar way one obtains (actually nonlocal) t -channel and u -channel hadron exchanges. A special role is played by the Coulomb (not color Coulomb) interaction, which always exists locally in addition to listed above interactions. It is important that all types of interaction between white objects (except for Coulomb) are short ranged for massive exchanged hadrons and do not produce narrow singularities by themselves. Our purpose in this paper is to find the dynamical origin of relatively narrow peaks nearby threshold which occur due to transformation of one white pair into another arbitrary number of times and to find what is the transformation itself. As we shall see, we shall meet a completely different type of transformation — we shall call it the recoupling and explain this mechanism in some detail. As a result we shall have here a new theoretical treatment of hadron-hadron interaction, suggesting a simple and quite general mechanism for exotic peaks in mesons and baryons — the recoupling mechanism.

In contrast to the approaches, where white-white interaction in the one-channel system, (e.g. in meson-meson) is generating resonances, we propose the dynamical picture, where the summed up transitions from one channel to another (without interaction inside channels) can be strong enough to produce resonances nearby thresholds. The specific feature of this interaction is that it depends strongly on the wave functions of both channels, entering in the overlap integral of the transition matrix element, which measures the amplitude of the transition between the initial $q\bar{Q} + \bar{q}Q$ and final $q\bar{q} + Q\bar{Q}$ states.

In what follows we are exploiting the channel-coupling (CC) interaction in the form of the energy-dependent recoupling Green’s functions as a possible origin of extra states — the recoupling mechanism.

Indeed, more than 30 years ago, the present author participated in the systematic study of CC effects in the spectra of hadrons, nuclei and atoms [29]. It was found there, that the CC interaction defined by the Transition Matrix Element (TME) is able to produce resonances (poles) of its own, if TME is strong enough, i.e. if the corresponding TME satisfies certain conditions, similar to that for one-channel potential.

We show below, that at the basis of this recoupling process lies a simple picture of the string recoupling between the same systems of quarks and antiquarks, which does not need neither energy nor additional interaction, and is simply a kind of topological transformation of two confining strings with fixed ends into another pair of strings — the string recoupling.

Figure 1. The transition of the mesons $(q_1\bar{q}_2) + (q_3\bar{q}_4) \leftrightarrow (q_1\bar{q}_4) + (\bar{q}_2q_3)$ via recoupling of the confining strings.

Figure 2. The mechanism of string recoupling via double string-breaking process at the points shown by thick dots.

One can see in figure 1 the confining regions (the crossed areas) for the bound states of quark-antiquark (mesons) $q_1\bar{q}_2$ and $q_3\bar{q}_4$ in the l.h.s. of the figure 1, which is transformed in the middle part of figure 1 into the confining region between $q_1\bar{q}_4$ on the plane of the figure, and “the confining bridge” — the double-crossed area between $\bar{q}_2q_3$. The r.h.s. of the figure 1 is the same as the l.h.s. As the result the transition is $(q_1\bar{q}_2) + (q_3\bar{q}_4) \leftrightarrow (q_1\bar{q}_4) + (\bar{q}_2q_3)$. It is interesting to understand what kind of vertices are responsible for this transition, and to this end we demonstrate in figure 2 below the possible construction of the “confining bridge” in the figure 1 by cutting the confining film and turning up the middle piece.

We show in this way that topologically this process is equivalent to the double string breaking, and numerically is defined by the overlap integral of participating hadron wave functions. This mechanism is quite general and can work for meson-meson, meson-baryon, baryon-baryon states. In particular it can work for some of X, Y and Z states of heavy quarkonia, like $Z_c(3900)$ and $Z_c(4020)$.

It is a purpose of the present paper to exploit this formalism for the case of extra states in meson-meson or meson-baryon spectra and define possible resonances and thresholds, and further on to apply this formalism to the case of pentaquark states like $P_c(4312), P_c(4440), P_c(4457)$.

Our main procedure will be the calculation of TME using realistic wave functions of $c\bar{c}, b\bar{b}, c\bar{u}$ systems, as well as approximate for baryon systems. Using those we calculate the

resulting Green's functions and resonance positions and compare them with experiment. The plan of the paper is as follows.

In the next section we introduce the reader to the method by solving a simplified two-channel problem with a separable potential. Section 3 is devoted to the explicit formulation of the recoupling mechanism, section 4 contains application to the meson-meson channel, and section 5 considers the meson-baryon case. Section 6, 7 are devoted to the analysis of the physical structure and numerical results and discussion, while section 7 contains conclusions and an outlook.

2 The simplest case: only separable CC interaction

Suppose we have two channels 1 and 2 with thresholds E_1 and E_2 and the CC interaction is separable

$$V_{12}(p_1, p_2) = -\lambda v_1(p_1)v_2(p_2) = V_{21}. \quad (2.1)$$

The Schroedinger-like (possibly relativistic) equations are

$$(T_1 - E)\varphi_1 + V_{12}\varphi_2 = 0, \quad (T_2 - E)\varphi_2 + V_{21}\varphi_1 = 0 \quad (2.2)$$

and can be reduced to the equation

$$(T_1 - E)\varphi_1 + V_{121}(E)\varphi_1 = 0, \quad (2.3)$$

where V_{121} is

$$V_{121}(p_1, p'_1 E) = -\lambda^2 v_1(p_1)v_1(p'_1) \int \frac{v_2^2(k)d^3k/(2\pi)^3}{T_2(k) + E_2 - E}. \quad (2.4)$$

Solving (2.3) one obtains the equation for the eigenvalue E

$$1 = \lambda^2 \int \frac{v_1^2(p)d^3p/(2\pi)^3}{T_1(p) + E_1 - E} \int \frac{v_2^2(k)d^3k/(2\pi)^3}{T_2(k) + E_2 - E}, \quad (2.5)$$

or

$$1 = \lambda^2 I_1(E)I_2(E). \quad (2.6)$$

For $E_1 < E_2$ one can put $E = E_1$ and get a condition for the existence of a bound state in our two-channel system [29].

$$\lambda^2 I_1(E)I_2(E) \geq 1. \quad (2.7)$$

One of the intriguing points now is how the bound state poles, or more generally, any poles appear when the interaction strength λ is large enough. To this end we make a simplifying assumption about the form of $v_i(k)$ and write

$$a) \quad v_i^2(k) = \frac{1}{k^2 + \nu_i^2} \quad b) \quad v_i^2(k) = \exp\left(-\frac{k^2}{4\beta_i^2}\right) \quad (2.8)$$

where ν_1, ν_2 and β_1, β_2 are some constants. Assuming also the nonrelativistic kinematics $T_i = \frac{k_i^2}{2\mu_i}$ one obtains in the case a)

$$I_i(E) = \frac{\mu_i}{2\pi(\nu_i - i\sqrt{2\mu_i\Delta_i})}, \quad \Delta_i = E - E_i, \quad (2.9)$$

and we have taken square root (2.9) on the physical Riemann sheet, $E = E + i\delta$. Hence the equation (2.5) for the poles (energy eigenvalues) is

$$\left(\nu_1 - i\sqrt{2\mu_1(E - E_1)}\right) \left(\nu_2 - i\sqrt{2\mu_2(E - E_2)}\right) = C = \frac{\mu_1\mu_2\lambda^2}{4\pi^2}. \quad (2.10)$$

We shall be mostly interested in the poles around the threshold E_2 and therefore in the first approximation we replace the first factor on the l.h.s. of (2.10) by a constant, assuming, that $E_2 - E_1$ has a large positive value, hence one can write for $k_2 = \sqrt{\mu_2(E - E_2)}$ using (2.10)

$$k_2 \cong -i\nu_2 + i\lambda'^2, \quad \lambda'^2 = \frac{\mu_1\mu_2\lambda^2}{4\pi^2(\nu_1 - i\sqrt{2\mu_1(E_2 - E_1)})}. \quad (2.11)$$

From (2.1) one can see that the pole is originally (at $\lambda' = 0$) on the second E_2 sheet, $k_2 = -i\nu_2$ and remains on the second E_2 sheet with increasing λ' . Note, however, that since originally we have been on the E_1 first sheet, then $\text{Im } \lambda'^2 > 0$, and therefore $\text{Re } k_2 < 0$, implying that the pole can be of the Breit-Winger type for $\text{Re } \lambda'^2 > \nu_2$.

As will be shown below in section 4, resonance production cross sections are proportional to the function

$$\frac{d\sigma^*(E)}{dE} = \left| \frac{1}{1 - \lambda^2 I_1(E - E_1) I_2(E - E_2)} \right|^2 k_2(E). \quad (2.12)$$

We can generalize this separable form to the relativistic case, when two hadrons with masses m_3, m_4 , so that the denominators in (2.5) look as follows:

$$\begin{aligned} T_1(p) + E_1 - E &\rightarrow \sqrt{p^2 + m_1^2} + \sqrt{p^2 + m_2^2} - E, \\ T_2(p) + E_2 - E &\rightarrow \sqrt{k^2 + m_3^2} + \sqrt{k^2 + m_4^2} - E. \end{aligned} \quad (2.13)$$

Here we have two thresholds $m_1 + m_2$ and $m_3 + m_4$, and we shall assume that $m_1 + m_2 < m_3 + m_4$.

Making the replacement (2.13) in $I_1(E), I_2(E)$ one can calculate these functions and find the behaviour of the approximate cross section in (2.12).

3 Equations for two channel amplitudes in the recoupling formalism

In this section we discuss the Green's function of the system of two white (noninteracting) hadrons h_1, h_2 , which can transform into another system of white hadrons H_3, H_4 and this transformation can occur infinite number of times $h_1 h_2 \rightarrow H_3 H_4 \rightarrow h_1 h_2 \rightarrow H_3 H_4 \rightarrow \dots$

Denoting the transition amplitude $V(h_1 h_2 \rightarrow H_3 H_4) = V^+(H_3 H_4 \rightarrow h_1 h_2)$, and the corresponding Green's functions as G_h, G_H , we obtain the total Green's function $G_{\alpha\beta}$, e.g. G_{hh}

$$\begin{aligned} G_{hh} &= G_h + G_h V_{hH} G_H V_{Hh} G_h + G_h V_{hH} G_H V_{Hh} G_h V_{hH} G_H V_{Hh} G_h + \dots = \\ &= G_h \frac{1}{1 - V_{hH} G_H V_{Hh} G_h}; \end{aligned} \quad (3.1)$$

as a result one obtains the equation, which defines all possible singularities of the physical amplitudes, including the resonance poles.

$$1 = V_{hH} G_H V_{Hh} G_h. \quad (3.2)$$

Note, that the described above method of the channel coupling was proposed before in the nonrelativistic form by the Cornell group [30–33], and exploited for the analytic calculation of the charmonium spectra, where the $h_1 h_2$ are strongly interacting quarks $c\bar{c}$. The subsequent development of this method in [34–36] has allowed to understand the nature of the $X(3872)$ [35] and Z_b states [36]. For the light quarks this method requires the explicit knowledge of $q\bar{q}$ spectrum and wave functions, which are available in the QCD string approach [37–46].

Recently the same approach, called the relativistic Cornell-type formalism successfully explained the spectrum of light scalars [47, 48]. In our present case we disregard the interaction of hadrons h_1 with h_2 and H_3 with H_4 .

Both Green’s function G_H, G_h describe propagation of two noninteracting subsystems, but each of these hadrons can have its own nontrivial spectrum.

In the simplest case, e.g. $h = \pi\bar{\pi}, H = K\bar{K}$, the Green’s functions of noninteracting particles are well known, see e.g. [47, 48] for the scalar $\pi\pi, K\bar{K}$ Green’s functions with the fixed spatial distance between $\pi\pi$ or $K\bar{K}$, needed to define the transition matrix element.

Since each of h_i or H_j is a composite system consisting of $q\bar{q}$ or qqq one must write the corresponding relativistic composite Green’s function, using the path integral formalism, see [44–46] for a recent review.

As it is seen from (2.13), one needs the explicit form of the relativistic Green’s function, consisting of two quark-antiquark mesons $h_1(q_1, \bar{Q}_1)$ and $h_2(\bar{q}_2, Q_2)$ with the zero total momentum $\mathbf{P} = 0$, so that the c.m. momentum of $q_1 \bar{Q}_1$ is $\mathbf{p}_1$, while for $\bar{q}_2 Q_2$ it is $-\mathbf{p}_1$. As a result the wave function of the $h_1 h_2$ system with $\mathbf{P} = 0$ and c.m. coordinates $\mathbf{R}$ can be written as

$$\Psi_{h_1 h_2}(\mathbf{u} - \mathbf{x}; \mathbf{y} - \mathbf{v}) = e^{i\mathbf{p}_1 \mathbf{R}(\mathbf{u}, \mathbf{x}) - i\mathbf{p}_1 \mathbf{R}(\mathbf{y}, \mathbf{v})} \psi_1(\mathbf{u} - \mathbf{x}) \psi_2(\mathbf{y} - \mathbf{v}). \quad (3.3)$$

At the same time the relativistic wave function of the hadrons $H_3, H_4, h_H(q_1 \bar{q}_2), H_4(\bar{Q}_1 Q_2)$ has the form

$$\Psi_{h_3 h_4}(\mathbf{u} - \mathbf{v}; \mathbf{v} - \mathbf{y}) = e^{i\mathbf{p}_2 \mathbf{R}(\mathbf{u}, \mathbf{v}) - i\mathbf{p}_2 \mathbf{R}(\mathbf{x}, \mathbf{y})} \psi_3(\mathbf{u} - \mathbf{v}) \psi_4(\mathbf{x} - \mathbf{y}). \quad (3.4)$$

Here we have introduced the c.m. coordinates $\mathbf{R}$ of the hadrons, expressed via the average energies ω_i, Ω_i of the quarks and antiquarks in the hadron [49]

$$\mathbf{R}(\mathbf{u}, \mathbf{x}) = \frac{\omega_1 \mathbf{u} + \bar{\Omega}_1 \mathbf{x}}{\omega_1 + \bar{\Omega}_1}, \quad \mathbf{R}(\mathbf{y}, \mathbf{v}) = \frac{\bar{\omega}_2 \mathbf{v} + \Omega_2 \mathbf{y}}{\bar{\omega}_2 + \Omega_2}, \quad (3.5)$$

$$\mathbf{R}(\mathbf{u}, \mathbf{v}) = \frac{\omega_3 \mathbf{u} + \bar{\omega}_4 \mathbf{v}}{\omega_3 + \bar{\omega}_4}, \quad \mathbf{R}(\mathbf{x}, \mathbf{y}) = \frac{\bar{\Omega}_3 \mathbf{x} + \Omega_4 \mathbf{y}}{\bar{\Omega}_3 + \Omega_4}. \quad (3.6)$$

Here ω_i, Ω_i are given in the appendix A of this paper.

Next we must calculate the overlap matrix element of $\Psi_{h_1 h_2}$ and $\Psi_{h_3 h_4}$

$$V_{12|34}(\mathbf{p}_1, \mathbf{p}_2) = \int \bar{y}_{1234} d^3(u-x) d^3(y-v) d^3(u-v) \Psi_{h_1 h_2} \Psi_{h_3 h_4}^+. \quad (3.7)$$

Introducing the Fourier component of the wave functions e.g. $\psi_1(\mathbf{u} - \mathbf{x}) = \int \tilde{\psi}_1(\mathbf{q}_1) e^{i\mathbf{q}_1(\mathbf{u}-\mathbf{x})} \frac{d^3 q_1}{(2\pi)^3}$, one obtains in the simple case when $q_2 = q_1, Q_2 = Q_1$

$$V_{12|34}(\mathbf{p}_1, \mathbf{p}_2) = \int \frac{d^3 \mathbf{q}_1}{(2\pi)^3} \bar{y}_{1234} \tilde{\psi}_1(\mathbf{q}_1) \tilde{\psi}_2(\mathbf{q}_1 + \mathbf{p}_2) \tilde{\psi}_3\left(-\mathbf{q}_1 - \frac{\mathbf{p}_2}{2} - \mathbf{p}_1 \frac{\omega_1}{\omega_1 + \Omega_1}\right) \times \tilde{\psi}_4\left(\mathbf{q}_1 - \frac{\mathbf{p}_2}{2} - \mathbf{p}_1 \frac{\Omega_1}{\omega_1 + \Omega_1}\right). \quad (3.8)$$

In (3.7), (3.8) we introduced the numerical recoupling coefficient $\bar{y}_{1234}$, which is discussed in appendix B.

The transition element (3.7) with the factor $\bar{y}_{1234}$, responsible for the recoupling of hadrons, shown in figure 1, has a simple structure. Indeed, as one can see in figure 1, the creation of two string configurations in the intermediate confining strings position and back into the original configuration. One may wonder what is the explicit mechanism of this recoupling, and what are the vertices denoted by thick points in the figure 2. To this end we note, that we have two strings on the r.h.s. of figure 2: string from Q_1 to $\bar{Q}_2$ and another from $\bar{q}_1$ to q_2 ; this position results from the double string decay (the l.h.s. of figure 2) with the subsequent rotation of the string between Q_1 and $\bar{Q}_2$ to the right, where this string is at some distance above the string between $\bar{q}_1, q_2$. One can associate the quantity $M(x, y)$ with this process and we must add this factor to $V_{12|34}$. Writing $M(x, y) = \sigma|\mathbf{x} - \mathbf{y}|$ in analogy with the one-point string decay described by the effective Lagrangian [50] for the string decay,

$$\mathcal{L}_{sd} = \int d^4 x \bar{\psi}(x) M(x) \psi(x) \quad (3.9)$$

and replacing it with the numerical value M_ω , similarly to [34–36], (see appendix B for details) one can write $y_{1234} = M_\omega \chi_{1234}$ in (3.8), with χ_{1234} describing the spin-isospin recoupling. Finally one obtains the expression for the whole combination

$$N(E) = G_{h_1 h_2} V_{12|34} G_{h_3 h_4} V_{34|12} \quad (3.10)$$

$$N(E) = \int \frac{d^3 \mathbf{p}_1}{(2\pi)^3} \frac{d^3 \mathbf{p}_2}{(2\pi)^3} \frac{V_{12|34}(\mathbf{p}_1, \mathbf{p}_2) V_{34|12}(\mathbf{p}_1, \mathbf{p}_2)}{(E_1(\mathbf{p}_1) + E_2(\mathbf{p}_1) - E)(E_3(\mathbf{p}_2) + E_4(\mathbf{p}_2) - E)}. \quad (3.11)$$

The resulting singularities (square root threshold singularities and possible poles from the equation $N(E) = 1$) can be found in the integral (3.11).

One can see, that the structure of the expression (3.11) is the same as in eq. (2.5), provided $V_{12|34}$ factorizes in factors $v_1(p_1)v_2(p_2)$, and consequently one expects the same behaviour of the cross sections as in (2.12).

At this point it is useful to introduce the approximate form of the wave functions in (3.8), which is discussed in [36]. Here we only give the simplest form of the Gaussian wave functions for the ground states of light, heavy-light and heavy quarkonia. One can write

$$\tilde{\psi}_i(q) = c_i \exp\left(-\frac{q^2}{2\beta_i^2}\right), \quad c_i^2 = \frac{8\pi^{3/2}}{\beta_i^3}; \quad \int \tilde{\psi}_i^2(q) \frac{d^3 q}{(2\pi)^3} = 1 \quad (3.12)$$

where β_i was found in [34–36], see appendix C, e.g. for ground states of bottomonium $\beta = 1.27$ GeV, for charmonium $\beta = 0.7$ GeV and for D, B mesons $\beta = 0.48, 0.49$ GeV.

Inserting $\tilde{\psi}_i(q)$ in (3.12) into (3.8) and integrating over $d^3\mathbf{q}_1$ one obtains

$$V_{12|34}(\mathbf{p}_1\mathbf{p}_2) = \bar{y}_{1234} \left(\prod_{i=1}^4 c_i \right) \frac{\exp(-AP_2^2 - Bp_1^2 - C\mathbf{p}_1\mathbf{p}_2)}{(2\pi)^3 8\pi^{3/2} a^{3/2}}, \quad (3.13)$$

where a, A, B, C are

$$a = \frac{1}{2} \sum_{i=1}^4 \frac{1}{\beta_i^2}; \quad A = \frac{1}{2} \left(\frac{1}{\beta_2^2} + \frac{1}{4\beta_3^2} + \frac{1}{4\beta_4^2} \right) - \frac{1}{4a} \left(\frac{1}{\beta_2^2} + \frac{1}{2\beta_3^2} + \frac{1}{2\beta_4^2} \right)^2 \quad (3.14)$$

$$B = \frac{1}{2\beta_3^2} \left(\frac{\omega_1}{\omega_1 + \Omega_1} \right)^2 + \frac{1}{2\beta_4^2} \left(\frac{\Omega_1}{\omega_1 + \Omega_1} \right)^2 - \frac{1}{4a} \left(\frac{1}{\beta_3^2} \frac{\omega_1}{\omega_1 + \Omega_1} - \frac{1}{\beta_4^2} \frac{\Omega_1}{\omega_1 + \Omega_1} \right)^2 \quad (3.15)$$

$$C = \frac{1}{2} \left(\frac{1}{\beta_3^2} \frac{\omega_1}{\omega_1 + \Omega_1} - \frac{1}{\beta_4^2} \frac{\Omega_1}{\omega_1 + \Omega_1} \right) - \frac{1}{2a} \left(\frac{1}{\beta_2^2} + \frac{1}{2\beta_3^2} + \frac{1}{2\beta_4^2} \right) \times \left(\frac{\omega_1}{\beta_3^2(\omega_1 + \Omega_1)} - \frac{\Omega_1}{\beta_4^2(\omega_1 + \Omega_1)} \right). \quad (3.16)$$

The resulting $N(E)$ has the form

$$N(E) = \frac{M_\omega^2 \bar{\chi}_{1234}^2}{a^3 (\prod_i \beta_i)^3 (\pi)^3} \int \frac{d^3p_1 d^3p_2 \exp(-2Ap_2^2 - 2Bp_1^2 - 2C\mathbf{p}_1\mathbf{p}_2)}{(E_1(p_1) + E_2(p_1) - E)(E_3(p_2) + E_4(p_2) - E)} \quad (3.17)$$

and the differential cross section with the final second channel is proportional to

$$\frac{d\sigma}{dE} \sim \frac{p(E)}{|1 - N(E)|^2} \quad (3.18)$$

where $p(E) \sim \sqrt{E^2 - (m_3 + m_4)^2}$. It is interesting, that for the fully symmetric case, when all β_i are equal, and $\omega_1 = \Omega_1$, one obtains for the exponent in (3.13) $\exp\left(-\frac{p_1^2 + p_2^2}{2\beta^2}\right)$, and $V_{12|34} = V_{34|12}$ and $N(E)$ are

$$V_{12|34}^{\text{symm}}(\mathbf{p}_1, \mathbf{p}_2) = \frac{2^{5/2} \sqrt{\pi}}{\beta^3} \bar{y}_{1234} \exp\left(-\frac{p_1^2 + p_2^2}{4\beta^2}\right) \quad (3.19)$$

$$N(E) = \frac{2M_\omega^2 \bar{\chi}_{1234}^2}{\pi \beta^6} \int \frac{p_1^2 dp_1 p_2^2 dp_2 \exp\left(-\frac{p_1^2 + p_2^2}{4\beta^2}\right)}{(E_1(p_1) + E_2(p_1) - E)(E_3(p_2) + E_4(p_2) - E)}. \quad (3.20)$$

4 Recoupling mechanism for the meson-meson amplitudes

The formalism introduced on the previous section can be directly applied to the amplitudes, containing two meson-meson thresholds, $m_1 + m_2 \leftrightarrow m_3 + m_4$ with the singularities given by the equation

$$1 - N(m_1, m_2, m_3, m_4; E) = 0. \quad (4.1)$$

As we discussed in section II, the conditions for the appearance of visible singularities require that the threshold difference $\Delta M = m_3 + m_4 - m_1 - m_2$ should be comparable or smaller than average size $\langle\beta\rangle$ of the hadron wave functions in momentum space, while

the recoupling coefficient $\bar{y}_{1234}^2$ is of the order of unity, i.e. there should be no angular momentum excitation or spin flip process.

An additional requirement is the relatively small widths of participating hadrons, otherwise all singularities would be smoothed out.

One can choose several examples in this respect.

- 1) The set of transformations

$$J/\psi + \phi \leftrightarrow D_s^* + \bar{D}_s^* \rightarrow J/\psi + \phi \quad (4.2)$$

with masses $m_1 = 3097 \text{ MeV}$, $m_2 = 1020 \text{ MeV}$, $m_3 = m_4 = 2112 \text{ MeV}$, and the corresponding thresholds $m_1 + m_2 = 4117 \text{ MeV}$ and $m_3 + m_4 = 4224 \text{ MeV}$. One can see no spin flip in the sequence $c_+ \bar{c}_+ + s_+ \bar{s}_- \rightarrow c_+ \bar{s}_- + \bar{c}_+ s_+$ for (4.1), where lower indices denote spin projections, and therefore no damping of transition probability. One can expect, that the yield of the reaction (4.2) would have the form similar to that of $\chi_{c1}(4140)$ with the mass (4147 MeV with the width $\Gamma = (22 \pm 8) \text{ MeV}$ [51–54].

- 2) One of the best studied exotic resonances $Z_c(3900)$ [55–58] was found in the reaction $e^+e^- \rightarrow \pi^+\pi^- J/\psi \rightarrow \pi^\pm Z_c(3900)$. It can be associated with the recoupling process $D\bar{D}^* \leftrightarrow \pi J/\psi$, where the higher threshold is $M_2 = 3874 \text{ MeV}$, and the spin, charge and isospin recombination agrees with this recoupling. One expects the peak above M_2 in agreement with experiment.

A similar situation can be in the case of the $Z_c(4020)$ observed in the reaction $e^+e^- \rightarrow \pi\pi h_c$ [59], which can be associated with the recoupling $\pi h_c \rightarrow D^* \bar{D}^*$ with threshold $M_1 = 3665 \text{ MeV}$ and $M_2 = 4020 \text{ MeV}$. One can envisage the yield of the reaction to be described by the equation (3.18), with $p(E) \rightarrow p^3(E)$, since one needs the P -wave in $D^* \bar{D}^*$ near threshold, as in h_c .

Note, that in general the recoupling can easily produce both Z_b, Z_c resonance peaks, when a charged particle (like ρ) is participating in the sequence of transformations.

5 Recoupling mechanism for meson-baryon systems

One can consider the transformation sequence for baryons of the form, e.g.

$$(qqq) + Q\bar{Q} \leftrightarrow (qqQ) + (q\bar{Q}) \quad (5.1)$$

and apply the same formalism as the used above for the meson-meson recoupling transformations.

In principle it implies the new degrees of freedom, associated with the additional quark in (qqq) as compared to the meson $(q\bar{q})$. To simplify the matter, we start below with the assumption, that the diquark combination can be factorized out in the baryon $(qqq) \rightarrow q(qq)$ and does not change during the recoupling process, which can now be written as

$$q(qq) + (Q\bar{Q}) \leftrightarrow Q(qq) + (q\bar{Q}). \quad (5.2)$$

In doing so we neglect also the internal structure of the diquark (qq) system, which stays unchanged during the recoupling process, so that only its total spin, spin projection and its relative motion with the quark q or $\bar{Q}$ in the baryon bound state is present in the matrix element (3.11), while the norm of (qq) is factored out. As a result one can use eqs. (3.11), (3.13), where we need the wave functions of the relative motion of quark and diquark in the baryons $q(qq)$ and $Q(qq)$. Using our notations $h_1(q_1\bar{Q}_1) + h_2(\bar{q}_2Q_2) \leftrightarrow h_3(q_1\bar{q}_2) + h_4(\bar{Q}_1Q_2)$ we are replacing $\bar{q}_2$ by the diquark (qq'). The accuracy of this replacement was discussed in literature [61–65] and the interactions are discussed and compared in [66]. In what follows we need the approximate baryon wave functions as $\psi_2(\mathbf{y} - \mathbf{v})e^{-i\mathbf{p}_1\mathbf{R}(\mathbf{y},\mathbf{v})}$ and $\psi_3(\mathbf{u} - \mathbf{v})e^{-i\mathbf{p}_2\mathbf{R}(\mathbf{u},\mathbf{v})}$ in (3.2), (3.4), where $\mathbf{v}$ denotes the center-of-mass of the diquark, and $\mathbf{R}(\mathbf{y},\mathbf{v})$ is the c.m. of the quark-diquark combination, i.e. actually is the c.m. of the baryon $Q_2(qq)$. The same for the $\psi_3(\mathbf{u} - \mathbf{v})$ and its c.m. $\mathbf{R}(\mathbf{u},\mathbf{v})$. Using the oscillator forms for ψ_2, ψ_3 one is actually exploiting the description of the only one part (factor) of the baryon wave function, which can be associated with only one leaf of the three-leaf baryon configuration. As a result, one can approximate this part of the wave function with the wave function of the heavy-light meson for the $Q_2(qq)$ baryon ($Q_2(qq) \rightarrow Q_2\bar{q}$) or with the light baryon for the $h_3(q_1\bar{q}_2 \rightarrow q_1(qq))$.

As a first example one can take the transitions $p + \phi \rightarrow \bar{K}^* + \Lambda$ with thresholds $M_1 = 1960$ MeV and $M_2 = 2005$ MeV, where the role of quarks $\bar{Q}_1, Q_2$ is played by $\bar{s}, s$ and one has a transition $u(ud) + \bar{s}s \rightarrow \bar{s}u + s(ud)$, where all β parameters have a similar magnitude, and one can expect a peak nearby M_2 .

As a concrete example one can take the case of the triad $P_c(4312), P_c(4440)$ and $P_c(4457)$, found experimentally in [67, 68], with a vast literature devoted to this phenomenon, called pentaquark, see a review in PDG,² and for the latest pentaquark papers see [69–86].

The most part of the literature considers pentaquarks as a result of molecular interaction between a white baryon and a white meson, which creates a bound state nearby the threshold of this system. In what follows we shall exploit the recoupling mechanism and we shall show, that it can provide the observed peaks without an assumption of the white-white strong interaction.

We shall have in mind the recoupling transformations of the type

$$J/\psi + P \leftrightarrow (\Sigma, \Sigma^*) + (D, D^*) \quad (5.3)$$

and we impose the requirement of s -wave recoupling without spin flip processes and parity conservation, which excludes $\Lambda_c^*(2595)$ with $(IJ^P) = (0, \frac{1}{2}^-)$ and includes $\Lambda_c(2286)(0, \frac{1}{2}^+)$, $\Sigma_c(2455)(1, \frac{1}{2}^+)$, $\Gamma \approx 2$ MeV, $\Sigma_c^*(2529)(1, \frac{3}{2}^+)$, $\Gamma \approx 15$ MeV, in addition to $D(1864), (\frac{1}{2}, 0^+)$ and $D^*(2010), (\frac{1}{2}, 1^-)$ with $\Gamma_D, \Gamma_{D^*} < 1$ MeV.

thresholds (MeV)	4150	4319	4465	4384
pairs	$\Lambda_c \bar{D}$	$\Sigma_c \bar{D}$	$\Sigma_c \bar{D}^*$	$\Sigma_c^* \bar{D}$
pentaquarks		$P_c(4312)$	$P_c(4457)$	$P_c(4440)$
width, MeV		$\Gamma = 9.8$	$\Gamma = 6.4$	$\Gamma = 20.6$

Table 1. Meson-baryon thresholds and the associated pentaquarks.

As a result one obtains the thresholds $M_2 = m_3 + m_4$ in the table 1 together with P_c . Following the table, we can consider two types of reactions,

$$I. \quad (c(ud))(\Sigma_c, \Sigma_c^*) + (\bar{c}u)(\bar{D}, \bar{D}^*) \leftrightarrow (c\bar{c})(J/\psi) + (u(ud))(p) \quad (5.4)$$

$$II. \quad (c(uu))(\Sigma_c^*) + \bar{c}d(\bar{D}, \bar{D}^*) \leftrightarrow (c\bar{c}) + (u(ud))(p). \quad (5.5)$$

To proceed one needs the values of $\beta_i, i = 1, 2, 3, 4$ and A, B, C and a in (3.17). Using appendix A one finds the values of ω, Ω in (3.14), (3.15) $\Omega_1 = 1509$ MeV, $\omega_1 = 507$ MeV. From appendix C one finds the values of β_i :

$$\beta_i(D, D^*) \cong \beta_2(\Sigma) = 0.48 \text{ GeV}, \quad \beta_3(p) \approx 0.26 \text{ GeV}, \quad (5.6)$$

and $\beta_4(J/\psi) = 0.7$ GeV. Note, that for Σ and p we have used the principle of replacement of light diquark by a light antiquark, $(ud) \rightarrow \bar{q}$. Therefore $\beta(\sigma) = \beta(c(ud)) = \beta(c\bar{u}) = \beta(D)$.

As a consequence one obtains the values given in (5.6). Using those we get the numerical values for a, A, B, C .

$$a = 12.76 \text{ GeV}^{-2}, \quad A = 4.02 \text{ GeV}^{-2}, \quad B = 0.94 \text{ GeV}^{-2}, \quad |C| < 0.03 \text{ GeV}^{-2}. \quad (5.7)$$

As a result one can neglect the $C \mathbf{p}_1 \mathbf{p}_2$ in (3.17) and the integrals $d^3 \mathbf{p}_1, d^3 \mathbf{p}_2$ factorize. We turn now to the recoupling coefficients $M_\omega, \bar{y}_{1234}$.

As it was shown in [50], the effective parameter M_ω can be expressed via the wave functions of objects, produced by the string breaking, in our case it is heavy-light mesons with the coefficient $\beta(D) \cong \beta(B) = 0.48$ GeV, and from eq. (35) of [50] one has

$$M_\omega \cong \frac{2\sigma}{\beta(D)} \cong 0.8 \text{ GeV}. \quad (5.8)$$

Finally, the coefficient $\bar{\chi}_{1234}$ for the transition into $(\Sigma_c \bar{D})$ and $(\Sigma_c \bar{D}^*)$ can be estimated as in the appendix B to be equal to 1.

6 Physical amplitudes and their singularities in the recoupling process

As was discussed in section 3, (3.18), the differential cross section for the production of hadrons in channel 1 can be written as

$$\frac{d\sigma}{dE} = |F_1(E) f_{12}(E)|^2 p_1(E) \quad (6.1)$$

where $F_1(E)$ is the production amplitude of channel 1 particles without final state FS interaction and f_{12} is the FS interaction, which we take as an infinite sum of transitions from channel 1 to channel 2-the Cornell-type mechanism [30–35].

$f_{12}(E)$ can be written as

$$f(E) = \frac{1}{1 - N(E)} \quad (6.2)$$

where $N(E)$ has the form

$$N(E) = \lambda I_1(E) I_2(E), \quad (6.3)$$

where $I_i(E)$ has the form

$$I_i(E) = \int \frac{d^3 p_i}{(2\pi)^3} \frac{v_i^2(p_i)}{E'(p_i) + E''(p_i) - E}. \quad (6.4)$$

Here v_i is proportional to the product of wave functions in momentum space (see (ref 42)) and can be written in two forms: a/ as a Gaussian of p and b/ as an inverse of $(p^2 + \nu^2)$. To simplify matter we shall consider situation close to nonrelativistic for the energies in the denominator of (6.4) and write

$$E'(p_1) + E''(p_1) = m_1 + m_2 + \frac{p_1^2}{2\mu_1}, \quad E'(p_2) + E''(p_2) = m_3 + m_4 + \frac{p_2^2}{2\mu_2}. \quad (6.5)$$

We have considered above in the paper $v_i^2(p)$ as a Gaussian $\exp(-b_i p_i^2)$ with $b_1 = 2B, b_2 = 2A$, see (3.17). We simplify below this expression, writing $\exp(-b_i p^2) = \frac{1}{\exp(b_i p^2)} \approx b_i^{-1} \frac{1}{p^2 + \nu_i^2}$, where $\nu_i = \frac{1}{\sqrt{b_i}}$. As a result one can write for $I_i(E)$ in the region $E > E_i(th) = m_1 + m_2 (i = 1)$ or $m_3 + m_4 (i = 2)$

$$I_i(E) = \frac{\mu_i}{2\pi b_i} \nu_i - i \sqrt{2\mu_i \Delta_i}, \quad \Delta_i = E - E_i(th). \quad (6.6)$$

As a result one obtains a simple expression for the amplitude $f_{12}(E)$

$$f_{12}(E) = \frac{1}{1 - \frac{\lambda' \mu_1 \mu_2}{(\nu_1 - i \sqrt{2\mu_1 \Delta_1})(\nu_2 - i \sqrt{2\mu_2 \Delta_2})}} = \frac{1}{1 - z t_1(E) t_2(E)}. \quad (6.7)$$

Here $z = \frac{\lambda}{\mu_1 \mu_2 b_1 b_2 (2\pi)^2}$ and the (6.7) refers to the region $E > E_1(th), E_2(th)$ otherwise one should replace $-i$ before the $\sqrt{}$ terms by $(+1)$ for the roots, where $\Delta(E)$ is negative.

We now consider the channel coupling constant λ' , which enters (6.7). From (3.17) one obtains

$$\lambda' = \frac{M_\omega^2 \chi_{1234}^2 b_1 b_2}{4\pi^5 a^3 (\prod_i \beta_i)^3}. \quad (6.8)$$

The resulting z is larger than unity for the recoupling coefficient χ_{1234} of the order of 1, and one can vary z in the interval from one to larger values. At this point we shall discuss the physical structure of the obtained expression for the amplitude $f_{12}(E)$ (6.7). One can see that f_{12} contains the product of two unitary single channel amplitudes t_1, t_2 which describe separately scattering amplitudes in channel 1 and 2 characterizing by numbers $\nu_i, i = 1, 2$. The latter are obtained from the hadron wave functions and reflect their

structure given by the Gaussian coefficients β_i . Now let us study the structure of the possible singularities of f_{12} in the E plane, which is given by zeros of the denominator of f_{12} in (6.7) and we concentrate on the k_1 plane while the second channel is characterized by the value of $B(E) = \frac{1}{\nu_2 - ik_2}$ and $-ik_2 > 0$ below higher threshold E_2 . As a result one obtains an equation for the pole of f_{12}

$$k_1 = -i\nu_1 + izB(E), B(E) > 0, \nu_1 > 0. \quad (6.9)$$

It is evident that originally (at $zB = 0$) virtual pole of k_1 is proceeding to the upper half plane of k_1 with increasing zB , and at some zB it appears at $k_1 = 0$ i.e. at the threshold E_1 , and at larger zB one obtains a real pole, i.e. the effect of recoupling with the channel 2 increases the attraction and creates a real pole from the original distant virtual pole. However for the physical amplitude $t_1(E)$ of the hadron-hadron interaction the virtual pole can be not a good approximation and more generally one should write for the amplitude $t_1(E) = \frac{1}{\nu_1(E) - ik_1}$ the $\nu_1(E)$ in the form $\nu_1(E) = a + b \ln(\frac{m^2 + 4k_1^2}{m^2})$, where a, b are positive numbers. This construction occurs from the t-channel exchange of the hadron with mass m . One can see from (6.9) that the logarithmic singularity is moving towards the threshold with the increasing z and can appear on the first sheet ($E < E_1$). We can now study the situation with the singularities associated with the threshold E_2 and find that here the situation is completely different because when $E \rightarrow E_2$ the momentum $k_1(E)$ as a rule is large and positive imaginary. As a result there are no pole solutions of the equation (6.7) for all values of E nearby the threshold E_2 and all resulting widths are of the order of parameters μ_i, ν_i , i.e. $(0.5 - 1)$ GeV, and therefore cannot be seen in experiment. Nevertheless we give in the Appendix 4 full quartic equation and its analysis for the finding of resonance poles in the whole combination of k_1, k_2 planes. As a special case one can consider the situation when the distance $\Delta = E_2 - E_1$ is small as compared with $\nu_i, i = 1, 2$. In this case the amplitude $f_{12}(E)$ written as

$$f_{12} = \frac{k_2 - i\nu_2}{k_2 + \frac{zk_1}{\nu_1^2 + k_1^2} + i(\nu_2 - \frac{z\nu_1}{\nu_1^2 + k_1^2})}. \quad (6.10)$$

One can see that for $k_1 \ll \nu_1$ and $\nu_2 \frac{z}{\nu_1}$ one obtains a resonance pole slightly shifted above the threshold E_2 , however this situation is coincidental. Summarizing we expect that the resulting singularities of the recoupling amplitude are produced by the singularities of the lower threshold process (process 1 in our definition) shifted to the lower threshold due to recoupling interaction with the process 2). In the next section we shall study this effect numerically in real physical examples.

7 Numerical results and discussion

We can now consider 3 transitions, partly discussed above:

1)

$$J/\psi + \phi \rightarrow D_s^* + \bar{D}_s^*$$

$$E_1(\text{th}) = 4.12 \text{ GeV}, E_2(\text{th}) = 4.224 \text{ GeV}, \mu_1 = 0.767, \mu_2 = 1.056, \nu_1 = 0.96, \nu_2 = 0.87, z = 1 \text{ (all GeV)}.$$

$E(\text{GeV})$	4.0	4.05	4.12	4.17	4.224	4.3
$ f_{12}(E) ^2$	3.43	4.49	17.36	10.76	5.6	1.24

Table 2. The values of the $|f_{12}(E)|^2$ near the channel thresholds for the transition **1**).

2)

$$J/\psi + \phi \rightarrow D_s + \bar{D}_s$$

$$E_1(\text{th}) = 3.936, E_2(\text{th}) = 4.12, \mu_1 = 0.767, \mu_2 = 0.984, \nu_1 = 0.96, \nu_2 = 0.87, z = 1.$$

As a special interesting case we consider below the recent experiment of BES III [87],

$$e^+ + e^- \rightarrow K^+(D_s^- D^* + D_s^{*-} D).$$

Applying here our recoupling mechanism, shown in the figure 1, one easily finds that the second channel obtained from the first channel $D_s^- D^*$ by recoupling is the channel $J/\psi + K^{*-}$ which creates the chain of reactions possibly generating a peak in the system $D_s^- D^*$ or $D_s^{*-} D$, namely we consider as the third example

3)

$$D_s^- + D_s^* \rightarrow J/\psi + K^{*-}$$

$$E_1(\text{th}) = 3.975, E_2(\text{th}) = 3.992, \mu_1 = 0.9936, \mu_2 = 0.692, \nu_1 = 0.87, \nu_2 = 0.96 \text{ (all GeV)}.$$

One can easily find that in all cases the values of ν_i and the values of μ_i are in the range 0.35 – 1.06 GeV. Applying the (6.7) one can find the f_{12} in all cases and hence $\frac{d\sigma}{dE}$ in the cases **1**), **2**), **3**), assuming z as a positive number typically larger or equal 1. However for the **3**) one could use also the sum: $f_{12} \rightarrow f_{12} + \alpha f'_{12}$, implying possible superposition of intermediate states in the rescattering series.

We proceed now with the cases **1**)-**3**) and insert the values of $E_i(\text{th})$, μ_i , ν_i in (6.3), (6.7) and fixing the value of z one obtains the form of the recoupling amplitude shown in the tables 2–4 below.

For the case **1**) the resulting values of $|f_{12}(E)|^2$ can be seen in the table 2.

One can see in the table 2 a strong enhancement around $E = 4.12$ GeV with the width around 10 MeV which can be associated with the resonance $\chi_{c1}(4140)$ having the mass 4147 MeV and the width $\Gamma = 22$ MeV.

It is now interesting how these numerical data are explained by the exact solution of the equation (6.7), (6.9). From these equations one obtains the exact position of the pole $k_1 = i0.068$ GeV, $E = E_1 - 0.003$ GeV which is unphysical result due to too small values of ν_1 , and in a more realistic case of the virtual pole its position moves to $k_1 = 0$, $E = E_1$ for $\nu_1 = 0.84$ GeV. Three more complex poles are O(1 GeV) far from the thresholds, as can be found from the solution of the quartic equation in appendix D. Looking at table 2 one can see a good agreement with this result. In a similar way we obtain the results for the $J/\psi + p$ transitions of **2**).

$E(\text{GeV})$	3.8	3.85	3.936	4.0	4.05	4.12	4.2
$ f_{12}(E) ^2$	3.12	3.84	18.9	6.15	5.33	298	1.17

Table 3. The values of the $|f_{12}(E)|^2$ near the ΣD threshold.

$E(\text{GeV})$	3.96	3.975	3.98	3.985	3.992	4.0
$ f_{12} ^2(z=1)$	5.43	322	67.56	31.84	12.95	5.54
$ f_{12} ^2(z=1.5)$	19.9	2.93	3.19	2.58	1.61	2.71

Table 4. The values of the transition probability as a function of energy in the transition **3**).

One can see in table 3 a strong peak near the lower threshold $E = 3.94 \text{ GeV}$ and it is easy to check that this peak is stable when one varies z in the region around $z = 1$. One can associate this peak with the virtual pole appearing at 50 MeV below the threshold $E_1 = 3.94 \text{ GeV}$. The exact form of the solution for $f_{12}(E)$ near the peak has the form $f_{12}(E) = \frac{k_1 + i0.96}{k_1 + i0.28}$. In this way our method can support the origin of the $X(3915)$ state as due to the $J/\psi + \phi \leftrightarrow D_s + \bar{D}_s$ transitions.

We come now to the recent interesting discovery of the new state $Z_{cs}(3985)$ [87], where we take for simplicity only the first chain denoted as the **3**) above. Similarly to the previous cases one obtains

One can see in table 4 a narrow peak with the summit at $E = 3.975 \text{ GeV}$ for $z = 1$ with the width around 10 MeV, which closely corresponds to the experimental data from [87] $E = 3.982.5, \Gamma = 12.8 \text{ MeV}$. In our case the resonance parameters weakly depend on z . As seen in the table 4, for $z = 1.5$ the peak shifts down to $E = 3.960 \text{ GeV}$ with much larger width. Its form is exactly the same as in the previous two examples and corresponds to virtual pole almost exactly at the lower threshold. In this way we can explain the newly discovered resonance $Z_{cs}(3985)$ by the recoupling mechanism in the rescattering series of transitions

$$D_s^- + D_s^* \rightarrow J/\psi + K^{*-}.$$

8 Conclusions and an outlook

As it was shown above, the new mechanism having the only parameter z is able to predict and explain the resonances in different systems, as it was shown above, and possibly in other systems which can transfer one into another via the recoupling of the confining strings. The necessary conditions for the realization of these transitions and the appearance of a resonance are connected to the value of the transition coefficient z , which should be of the order of unity or larger. Therefore the transition should be strong, i.e. without serious restructuring of the hadrons involved, since otherwise the transition will be strongly suppressed e.g. in the case when not only strings are recoupled, but also spins, orbital momenta, isospins should be exchanged. In any case the suggested mechanism provides an alternative to the popular tetra — and pentaquark mechanisms, which dominate in the literature. One should stress at this point, that the independent and objective checks, e.g.

the lattice calculations do not give strong support for the molecular or tetraquark models and the existence of an independent mechanism is welcome. It is necessary that this method should be studied more carefully. As to the recoupling mechanism, it is strongly associated with the thresholds participating in the transitions, and the best situation for its application is when both thresholds are close by. In all 3 cases considered above the distance between threshold was less than 200 MeV, and in all cases one could see a strong enhancement in the transition coefficient and hence in the resulting cross section. The necessary improvements of the present study are 1) a more accurate calculation of the coefficient z (originally λ'), and 2) the use of a more realistic Gaussian approximation for the wave functions instead of approximate ν_i parametrizations to define f_{12} with good accuracy in the future.

Acknowledgments

The author is grateful to Lu Meng for an important remark, to R.A.Abramchuk for useful discussions and to A.M.Badalian for discussions and suggestions. This work was done in the frame of the scientific project supported by the Russian Science Foundation Grant No. 16-12-10414.

A The center-of-mass coordinates and average quark and antiquark energies in a hadron

Following [49] one can define the c.m. coordinate of a hadron consisting of a quark Q at the point $\mathbf{x}$ and an antiquark $\bar{q}$ at the point $\mathbf{u}$ via average energies Ω and ω of Q and $\bar{q}$ correspondingly as

$$\mathbf{R}_{Q\bar{q}} = \frac{\Omega\mathbf{x} + \omega\mathbf{u}}{\Omega + \omega}, \quad \Omega = \langle \sqrt{\mathbf{p}_Q^2 + m_Q^2} \rangle, \quad \omega = \langle \sqrt{\mathbf{p}_q^2 + m_q^2} \rangle \quad (\text{A.1})$$

where $\sqrt{\mathbf{p}_Q^2 + m_Q^2} + \sqrt{\mathbf{p}_q^2 + m_q^2}$ is the kinetic part of the $Q\bar{q}$ Hamiltonian in the so-called spinless Salpeter formalism or an equivalent form in the so-called einbein formalism.³

As a result one obtains the following value of $\omega = \Omega$ for $q\bar{q}$ mesons, shown in table 2.

B The channel-coupling coefficient $\bar{y}_{1234}$

We discuss here two topics: 1) the problem of the double string decay vertex contribution to the recoupling coefficient M_ω in (3.11), 2) the construction of the recoupling vertex $\bar{y}_{1234}$.

We start with the topic 1), and following [50] define the relativistic expression for the string decay vertex as in (3.9), but without free parameters namely $M(x)$ in (3.9) is

$$M(x) = \sigma(|\mathbf{x} - \mathbf{x}_Q| + |\mathbf{x} - \mathbf{x}_{\bar{Q}}|). \quad (\text{B.1})$$

As one can see in figure 2, in our case the structure of the recoupling process can be explained by the double string breaking, which we can write as a product

$$S = \int d^4x \bar{\psi}(x) \bar{M}(x) d^4y \psi(x) \bar{\psi}(y) \bar{M}(y) \psi(y) \quad (\text{B.2})$$

³See a short recent review in the last refs. in [44–46].

State	J/ψ	$\psi(2S)$	$\psi(3770)$	$\psi(3S)$	$\psi(4S)$
Ω , GeV	1.58	1.647	1.640	1.711	1.17
State	$\Upsilon(1S)$	$\Upsilon(2S)$	$\Upsilon(3S)$	$\Upsilon(4S)$	$\Upsilon(5S)$
Ω , GeV	5.021	5.026	5.056	5.088	5.120
Sate	D	D_s	B	B_s	ρ
Ω , GeV	1.509	1.515	4.827	4.830	0.4
ω , GeV	0.507	0.559	0.587	0.639	0.4

*) Note, that the difference in Ω, ω obtained in these approaches is less or around 1%.

Table 5. Average values of quark and antiquark kinetic energies in different mesons.

and one must take into account, that the energy minimum of the resulting broken string occurs when both time moments x_0, y_0 of string breaking are equal. Indeed, taking the integral in (B.2) with account of the string action in the exponent of the path integral,

$$\Delta S_{\text{string}} = \int \sigma \sqrt{r_{xy}^2 + (x_4 - y_4)^2} d \frac{x_4 + y_4}{2},$$

which produces a factor on (B.2) $\langle r_{xy} \rangle \sqrt{2\pi}$, which denoted as M_ω in (3.11). The resulting double string breaking action can be written as,

$$S \cong \int d^3x \bar{\psi} i \not{\partial} \psi(x) d^3y \bar{\psi}(y) \psi(y) M_\omega(\mathbf{x}, \mathbf{y}) \frac{d(x_4 + y_4)}{2} \quad (\text{B.3})$$

using the notation

$$\int d(x_4 - y_4) \langle \bar{M}(x) \bar{M}(y) \rangle = M_\omega(\mathbf{x}, \mathbf{y}).$$

In what follows one can estimate $\langle M_\omega(\mathbf{x}, \mathbf{y}) \rangle \equiv M_\omega$ in the same way, as it was done in [50], with the result $M_\omega \approx \frac{2\sigma}{\beta}$, where β is the oscillator parameter for the $(Q\bar{Q})$ meson. We now turn to the point 2) above, the recoupling vertex $\bar{y}_{1234}$.

To define $\bar{y}_{1234}$ we notice that all 4 quarks $q, \bar{q}, Q, \bar{Q}$ keep their identity and spin polarization during the whole process of transformations, provided we neglect the spin dependent corrections. This can be also seen in the structure of the recoupling process: in (B.2) one does not see spin dependence, and this means, that the spin projection of each quark or antiquark is kept unchanged during recoupling. As a result one can write the nonrelativistic spin part of the matrix element $V_{12/34}$ as

$$V_{\text{meson}}^{\text{spin}} = C_{\mu_1 \mu_2}^{Lm} \chi_{\mu_1}^{(1)} \bar{\chi}_{\mu_2}^{(2)} C_{\mu'_1 \mu'_2}^{L'm'} \chi_{\mu'_1}^{(3)} \bar{\chi}_{\mu'_2}^{(4)} (C_{\nu_1 \nu_2}^{JM} \chi_{\nu_1}^{(5)} \bar{\chi}_{\nu_2}^{(6)} C_{\nu'_1 \nu'_2}^{J'M'} \chi_{\nu'_1}^{(7)} \bar{\chi}_{\nu'_2}^{(8)}) \quad (\text{B.4})$$

where the Klebsch-Gordon coefficient $C_{\nu_1 \nu_2}^{JM} \equiv C_{\frac{1}{2} \nu_1 \frac{1}{2} \nu_2}$ and $\chi_\mu^{(i)}, \bar{\chi}_\lambda^{(k)}$ are quark and anti-quark spinors.

As was told above, due to the spin conservation in recoupling, the matrix element (B.4) should be proportional to $\delta_{26} \delta_{48} \delta_{17} \delta_{35}$, implying the recoupling of quarks.

State	$\Upsilon(1S)$	$\Upsilon(2S)$	$\Upsilon(3S)$	$\Upsilon(4S)$	$\Upsilon(5S)$
β , GeV	1.27	0.88	0.76	0.64	0.6
State	J/ψ	$\psi(2S)$	$\psi(3S)$	$\psi(4S)$	$\psi(5S)$
β , GeV	0.7	0.53	0.48	0.43	0.41
Sate	D	B	ρ		
β , GeV	0.48	0.49	0.26		

Table 6. The Gaussian parameters β of different mesons.

As a result one obtains

$$V^{\text{spin}} = \sum_{\mu_i, \mu'_k} C_{\mu_1 \mu_2}^{Lm} C_{\mu'_1 \mu'_2}^{L'm'} C_{\mu'_1 \mu_2}^{JM} C_{\mu_1 \mu'_2}^{J'M'}. \quad (\text{B.5})$$

Here L, L', J, J' correspond to the spin values of hadrons $L + L' \rightarrow J + J'$ and we have assumed zero orbital momenta for all hadrons. Finally for the final expression in (3.11), $(V^{\text{spin}})^2$ should be summed up over all m, m', M, M' , so that for $\bar{y}_{1234}^2$ one has

$$\langle (\bar{y}_{1234})^2 \rangle = \sum_{m, m', M, M'} (V^{\text{spin}})^2. \quad (\text{B.6})$$

In a similar way one can find the recoupling coefficient $\bar{y}_{1234}$ for the ensemble transformation

$$p + J/\psi \rightarrow \Sigma(\Sigma^*) + D(D^*) \rightarrow p + J/\psi. \quad (\text{B.7})$$

In this case we write the baryon wave function as $\psi_B = u(ud)_0$, where the lower indices imply the total spin of the diquark (ud). In the simplest approximation one can approximate the proton as the quark-diquark combination $p = u(ud) \cong u\tilde{d}$, with the diquark $\tilde{d}$ kept unchanged d during recoupling.

C Oscillator parameters of hadron wave function

The oscillator parameters for the bottomonium, charmonium and B, D mesons have been obtained in [37–46], using the expansion of relativistic wave functions, obtained from the solutions of the relativistic string Hamiltonian [49], in the full set of the oscillator wave functions. As a result one obtains

The accuracy of the oscillator one-term approximation can be judged by the relative value of the sum of squared coefficients of four higher term of expansion as compared to the square of the main term. This amounts to the accuracy of the order or less than 10% for lowest states of charmonia and bottomonia and few percent for D, B, ρ .

D Calculation of the cross sections $|f_{ik}|^2$ and pole positions

As it is written in (6.7) the cross section is defined as $\frac{d\sigma}{dE} = |f_{ik}|^2$ where $f_{ik} = \frac{1}{1-z_{AB}}$, and z is numerical parameter proportional to λ' , where for three numerical examples (1), (2), (3)

in that section $z = 1, 5, 1$ and $A = \frac{1}{\nu_1 - ik_1}$ and $B = \frac{1}{\nu_2 - ik_2}$. Moreover $k_i = \sqrt{2\mu_i(E - E_i)}$ and parameters μ_i, ν_i, E_i are given in the section. As it is seen in tables 2, 3, 4 the cross sections have peaks and our purpose here is to calculate the positions of the poles both in E plane and in k_i plane. To do this one starts with two equations

$$(\nu_1 - ik_1)(\nu_2 - ik_2) = z, \frac{k_1^2}{2\mu_1} - \frac{k_2^2}{2\mu_2} = E_2 - E_1. \quad (\text{D.1})$$

One of possible strategy is to insert k_1 from the first equation into second equation and one obtains an equation of 4-th power for k_2 . Note that both k_i are positive on the real axis for $E > E_i$ and positive imaginary below E_i . Among many roots of eq. we need the closest to the real axis. The pole positions are below in the table. Now defining from the first equation in (D.1) k_1 via k_2 and submitting it into second equation one obtains the quartic equation for $x = k_2 + i\nu_2$

$$x^4 + ax^3 + bx^2 + cx + d = 0, \quad (\text{D.2})$$

where the coefficients are

$$a = -2i\nu_2, b = 2\mu_2\Delta - \nu_2^2 + \frac{\mu_2\nu_1^2}{\mu_1}, c = -2i\frac{\nu_1\mu_2z}{\mu_1}, d = -\frac{m u_2 z^2}{\mu_1}. \quad (\text{D.3})$$

Now taking values of these coefficients from item i), for three transitions $i = 1, 2, 3$ discussed in section Numerical examples and discussion with $z = 1, 0.2, 1$ respectively for these transitions, and $\Delta = E_2 - E_1$, one obtains the roots of this equation together with the corresponding values of the resonance energy. The results of this computation for the examples 1)-3) are given in the text of the section 7.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *Phys. Rev. D* **98** (2018) 030001 [[INSPIRE](#)].
- [2] N. Brambilla et al., *The XYZ states: experimental and theoretical status and perspectives*, *Phys. Rept.* **873** (2020) 1 [[arXiv:1907.07583](#)] [[INSPIRE](#)].
- [3] J.L. Rosner, *Threshold effect and $\pi^\pm\psi(2S)$ peak*, *Phys. Rev. D* **76** (2007) 114002 [[arXiv:0708.3496](#)] [[INSPIRE](#)].
- [4] E. Braaten and M. Lu, *Line shapes of the $Z(4430)$* , *Phys. Rev. D* **79** (2009) 051503 [[arXiv:0712.3885](#)] [[INSPIRE](#)].
- [5] K.-M. Cheung, W.-Y. Keung and T.-C. Yuan, *Bottomed analog of $Z^+(4433)$* , *Phys. Rev. D* **76** (2007) 117501 [[arXiv:0709.1312](#)] [[INSPIRE](#)].
- [6] C. Meng and K.-T. Chao, *$Z^+(4430)$ as a resonance in the $D_1(D'_1)D^*$ channel*, [arXiv:0708.4222](#) [[INSPIRE](#)].

- [7] G.-J. Ding, *Understanding the charged meson $Z(4430)$* , [arXiv:0711.1485](#) [INSPIRE].
- [8] G.-J. Ding, W. Huang, J.-F. Liu and M.-L. Yan, $Z^+(4430)$ and analogous heavy flavor molecules, *Phys. Rev. D* **79** (2009) 034026 [[arXiv:0805.3822](#)] [INSPIRE].
- [9] Y. Li, C.-D. Lu and W. Wang, *Partners of $Z(4430)$ and productions in B decays*, *Phys. Rev. D* **77** (2008) 054001 [[arXiv:0711.0497](#)] [INSPIRE].
- [10] L. Maiani, A.D. Polosa and V. Riquer, *The charged $Z(4433)$: towards a new spectroscopy*, [arXiv:0708.3997](#) [INSPIRE].
- [11] C.-F. Qiao, *A uniform description of the states recently observed at B -factories*, *J. Phys. G* **35** (2008) 075008 [[arXiv:0709.4066](#)] [INSPIRE].
- [12] X.-H. Liu, Q. Zhao and F.E. Close, *Search for tetraquark candidate $Z(4430)$ in meson photoproduction*, *Phys. Rev. D* **77** (2008) 094005 [[arXiv:0802.2648](#)] [INSPIRE].
- [13] X. Liu, B. Zhang and S.-L. Zhu, *The two-body open charm decays of $Z^+(4430)$* , *Phys. Rev. D* **77** (2008) 114021 [[arXiv:0803.4270](#)] [INSPIRE].
- [14] L. Maiani, A.D. Polosa and V. Riquer, *The charged $Z(4430)$ in the diquark-antidiquark picture*, *New J. Phys.* **10** (2008) 073004 [INSPIRE].
- [15] D.V. Bugg, *How resonances can synchronise with thresholds*, *J. Phys. G* **35** (2008) 075005 [[arXiv:0802.0934](#)] [INSPIRE].
- [16] T. Matsuki, T. Morii and K. Sudoh, *Is the $Z^+(4430)$ a radially excited state of D_s ?*, *Phys. Lett. B* **669** (2008) 156 [[arXiv:0805.2442](#)] [INSPIRE].
- [17] M. Cardoso and P. Bicudo, *Microscopic calculation of the decay of Jaffe-Wilczek tetraquarks, and the $Z(4433)$* , *AIP Conf. Proc.* **1030** (2008) 352 [[arXiv:0805.2260](#)] [INSPIRE].
- [18] Y.-R. Liu and Z.-Y. Zhang, *A chiral quark model study of $Z^+(4430)$ in the molecular picture*, [arXiv:0908.1734](#) [INSPIRE].
- [19] T. Branz, T. Gutsche and V.E. Lyubovitskij, *Hidden-charm and radiative decays of the $Z(4430)$ as a hadronic $D_1\bar{D}^*$ bound state*, *Phys. Rev. D* **82** (2010) 054025 [[arXiv:1005.3168](#)] [INSPIRE].
- [20] G. Galata, *Photoproduction of $Z(4430)$ through mesonic Regge trajectories exchange*, *Phys. Rev. C* **83** (2011) 065203 [[arXiv:1102.2070](#)] [INSPIRE].
- [21] M. Nielsen and F.S. Navarra, *Charged exotic charmonium states*, *Mod. Phys. Lett. A* **29** (2014) 1430005 [[arXiv:1401.2913](#)] [INSPIRE].
- [22] X.-W. Kang, Z.-H. Guo and J.A. Oller, *General considerations on the nature of $Z_b(10610)$ and $Z_b(10650)$ from their pole positions*, *Phys. Rev. D* **94** (2016) 014012 [[arXiv:1603.05546](#)] [INSPIRE].
- [23] F.K. Guo et al., *Interplay of quark and meson degrees of freedom in near-threshold states: a practical parametrization for line shapes*, *Phys. Rev. D* **93** (2016) 074031 [[arXiv:1602.00940](#)] [INSPIRE].
- [24] C. Deng, J. Ping, H. Huang and F. Wang, *Systematic study of Z_c^+ family from a multiquark color flux-tube model*, *Phys. Rev. D* **92** (2015) 034027 [[arXiv:1507.06408](#)] [INSPIRE].
- [25] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao and B.-S. Zou, *Hadronic molecules*, *Rev. Mod. Phys.* **90** (2018) 015004 [[arXiv:1705.00141](#)] [INSPIRE].

- [26] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu and S.-L. Zhu, *Pentaquark and tetraquark states*, *Prog. Part. Nucl. Phys.* **107** (2019) 237 [[arXiv:1903.11976](#)] [[INSPIRE](#)].
- [27] Y.S. Kalashnikova and A.V. Nefediev, *$X(3872)$ in the molecular model*, *Phys. Usp.* **62** (2019) 568 [*Usp. Fiz. Nauk* **189** (2019) 603] [[arXiv:1811.01324](#)] [[INSPIRE](#)].
- [28] A. Esposito, A. Pilloni and A.D. Polosa, *Hybridized tetraquarks*, *Phys. Lett. B* **758** (2016) 292 [[arXiv:1603.07667](#)] [[INSPIRE](#)].
- [29] A.M. Badalian, L.P. Kok, M.I. Polikarpov and Y.A. Simonov, *Resonances in coupled channels in nuclear and particle physics*, *Phys. Rept.* **82** (1982) 31 [[INSPIRE](#)].
- [30] E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.-M. Yan, *Charmonium: the model*, *Phys. Rev. D* **17** (1978) 3090 [*Erratum ibid.* **21** (1980) 313] [[INSPIRE](#)].
- [31] E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.-M. Yan, *Charmonium: comparison with experiment*, *Phys. Rev. D* **21** (1980) 203 [[INSPIRE](#)].
- [32] E.J. Eichten, K. Lane and C. Quigg, *Charmonium levels near threshold and the narrow state $X(3872) \rightarrow \pi^+ \pi^- J/\psi$* , *Phys. Rev. D* **69** (2004) 094019 [[hep-ph/0401210](#)] [[INSPIRE](#)].
- [33] Y.S. Kalashnikova, *Coupled-channel model for charmonium levels and an option for $X(3872)$* , *Phys. Rev. D* **72** (2005) 034010 [[hep-ph/0506270](#)] [[INSPIRE](#)].
- [34] I.V. Danilkin and Y.A. Simonov, *Channel coupling in heavy quarkonia: energy levels, mixing, widths and new states*, *Phys. Rev. D* **81** (2010) 074027 [[arXiv:0907.1088](#)] [[INSPIRE](#)].
- [35] I.V. Danilkin and Y.A. Simonov, *Dynamical origin and the pole structure of $X(3872)$* , *Phys. Rev. Lett.* **105** (2010) 102002 [[arXiv:1006.0211](#)] [[INSPIRE](#)].
- [36] I.V. Danilkin, V.D. Orlovsky and Y.A. Simonov, *Hadron interaction with heavy quarkonia*, *Phys. Rev. D* **85** (2012) 034012 [[arXiv:1106.1552](#)] [[INSPIRE](#)].
- [37] A.M. Badalian and B.L.G. Bakker, *Dynamics of the quark-antiquark interaction and the universality of Regge trajectories*, *Phys. Rev. D* **100** (2019) 034010 [[arXiv:1901.10280](#)] [[INSPIRE](#)].
- [38] A.M. Badalian and B.L.G. Bakker, *Radial and orbital Regge trajectories in heavy quarkonia*, *Phys. Rev. D* **100** (2019) 054036 [[arXiv:1902.09174](#)] [[INSPIRE](#)].
- [39] A.Y. Dubin, A.B. Kaidalov and Y.A. Simonov, *The QCD string with quarks. 1. Spinless quarks*, *Phys. Atom. Nucl.* **56** (1993) 1745 [*Yad. Fiz.* **56** (1993) 213] [[hep-ph/9311344](#)] [[INSPIRE](#)].
- [40] A.Y. Dubin, A.B. Kaidalov and Y.A. Simonov, *Dynamical regimes of the QCD string with quarks*, *Phys. Lett. B* **323** (1994) 41 [[INSPIRE](#)].
- [41] Y.A. Simonov, *Exact relativistic Green's functions for the time-independent potentials*, *Phys. Rev. D* **99** (2019) 096025 [[arXiv:1902.05364](#)] [[INSPIRE](#)].
- [42] S.S. Afonin, *Towards understanding spectral degeneracies in nonstrange hadrons. Part I. Mesons as hadron strings versus phenomenology*, *Mod. Phys. Lett. A* **22** (2007) 1359 [[hep-ph/0701089](#)] [[INSPIRE](#)].
- [43] S.S. Afonin and T.D. Solomko, *The large- N_c limit of borelized spectral sum rules and the slope of radial Regge trajectories*, *Int. J. Mod. Phys. A* **33** (2018) 1850115 [[arXiv:1805.03089](#)] [[INSPIRE](#)].
- [44] Y.A. Simonov, *Relativistic path integral and relativistic Hamiltonians in QCD and QED*, *Phys. Rev. D* **88** (2013) 025028 [[arXiv:1303.4952](#)] [[INSPIRE](#)].

- [45] Y.A. Simonov, *Spin interactions in mesons in strong magnetic field*, *Phys. Rev. D* **88** (2013) 053004 [[arXiv:1304.0365](#)] [[INSPIRE](#)].
- [46] Y.A. Simonov, *Exact relativistic Green's functions for the time-independent potentials*, *Phys. Rev. D* **99** (2019) 096025 [[arXiv:1902.05364](#)] [[INSPIRE](#)].
- [47] M.S. Lukashov and Y.A. Simonov, *Scalar mesons in a chiral theory with quark degrees of freedom*, *Phys. Rev. D* **101** (2020) 094028 [[arXiv:1909.10384](#)] [[INSPIRE](#)].
- [48] A.M. Badalian, M.S. Lukashov and Y.A. Simonov, *Relativistic Cornell-type mechanism of exotic scalar resonances*, *Phys. Rev. D* **102** (2020) 094012 [[arXiv:2001.07113](#)] [[INSPIRE](#)].
- [49] Y.A. Simonov, *Di-pion decays of heavy quarkonium in the field correlator method*, *Phys. Atom. Nucl.* **71** (2008) 1048 [[arXiv:0711.3626](#)] [[INSPIRE](#)].
- [50] Y.A. Simonov, *Relativistic theory of string breaking in QCD*, *Phys. Rev. D* **84** (2011) 065013 [[arXiv:1103.4028](#)] [[INSPIRE](#)].
- [51] LHCb collaboration, *Observation of $J/\psi\phi$ structures consistent with exotic states from amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ decays*, *Phys. Rev. Lett.* **118** (2017) 022003 [[arXiv:1606.07895](#)] [[INSPIRE](#)].
- [52] CDF collaboration, *Observation of the $Y(4140)$ structure in the $J/\psi\phi$ mass spectrum in $B^\pm \rightarrow J/\psi\phi K^\pm$ decays*, *Mod. Phys. Lett. A* **32** (2017) 1750139 [[arXiv:1101.6058](#)] [[INSPIRE](#)].
- [53] D0 collaboration, *Inclusive production of the $X(4140)$ state in $p\bar{p}$ collisions at D0*, *Phys. Rev. Lett.* **115** (2015) 232001 [[arXiv:1508.07846](#)] [[INSPIRE](#)].
- [54] CMS collaboration, *Observation of a peaking structure in the $J/\psi\phi$ mass spectrum from $B^\pm \rightarrow J/\psi\phi K^\pm$ decays*, *Phys. Lett. B* **734** (2014) 261 [[arXiv:1309.6920](#)] [[INSPIRE](#)].
- [55] BESIII collaboration, *Observation of a charged charmonium-like structure in $e^+e^-\pi^+\pi^-J/\psi$ at $\sqrt{s} = 4.26$ GeV*, *Phys. Rev. Lett.* **110** (2013) 252001 [[arXiv:1303.5949](#)] [[INSPIRE](#)].
- [56] BELLE collaboration, *Study of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and observation of a charged charmonium-like state at Belle*, *Phys. Rev. Lett.* **110** (2013) 252002 [Erratum *ibid.* **111** (2013) 019901] [[arXiv:1304.0121](#)] [[INSPIRE](#)].
- [57] T. Xiao, S. Dobbs, A. Tomaradze and K.K. Seth, *Observation of the charged hadron $Z_c^\pm(3900)$ and evidence for the neutral $Z_c^0(3900)$ in $e^+e^- \rightarrow \pi\pi J/\psi$ at $\sqrt{s} = 4170$ MeV*, *Phys. Lett. B* **727** (2013) 366 [[arXiv:1304.3036](#)] [[INSPIRE](#)].
- [58] BESIII collaboration, *Observation of $e^+e^- \rightarrow \pi^0\pi^0 h_c$ and a neutral charmonium-like structure $Z_c(4020)^0$* , *Phys. Rev. Lett.* **113** (2014) 212002 [[arXiv:1409.6577](#)] [[INSPIRE](#)].
- [59] BESIII collaboration, *Observation of a charged charmonium-like structure $Z_c(4020)$ and search for the $Z_c(3900)$ in $e^+e^- \rightarrow \pi^+\pi^- h_c$* , *Phys. Rev. Lett.* **111** (2013) 242001 [[arXiv:1309.1896](#)] [[INSPIRE](#)].
- [60] G.C. Rossi and G. Veneziano, *Tetra-quarks, penta-quarks and the like: old and new views, in The hadronic vacuum polarization from lattice QCD at high precision: a topical workshop of the muon $g-2$ theory initiative*, (2020) [[arXiv:2011.09774](#)] [[INSPIRE](#)].
- [61] M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson and D.B. Lichtenberg, *Diquarks*, *Rev. Mod. Phys.* **65** (1993) 1199 [[INSPIRE](#)].
- [62] S. Fleck, B. Silvestre-Brac and J.M. Richard, *Search for diquark clustering in baryons*, *Phys. Rev. D* **38** (1988) 1519 [[INSPIRE](#)].

- [63] E. Santopinto and J. Ferretti, *Strange and nonstrange baryon spectra in the relativistic interacting quark-diquark model with a Gürsey and Radicati-inspired exchange interaction*, *Phys. Rev. C* **92** (2015) 025202 [[arXiv:1412.7571](#)] [[INSPIRE](#)].
- [64] C. Gutierrez and M. De Sanctis, *A study of a relativistic quark-diquark model for the nucleon*, *Eur. Phys. J. A* **50** (2014) 169 [[INSPIRE](#)].
- [65] C. Alexandrou, P. de Forcrand and B. Lucini, *Evidence for diquarks in lattice QCD*, *Phys. Rev. Lett.* **97** (2006) 222002 [[hep-lat/0609004](#)] [[INSPIRE](#)].
- [66] E. Megías, E. Ruiz Arriola and L.L. Salcedo, *Quark-diquark string tension, excited baryonic resonances and thermal fluctuations*, *Phys. Scripta* **95** (2020) 074005 [[arXiv:2002.10871](#)] [[INSPIRE](#)].
- [67] LHCb collaboration, *Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays*, *Phys. Rev. Lett.* **115** (2015) 072001 [[arXiv:1507.03414](#)] [[INSPIRE](#)].
- [68] LHCb collaboration, *Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$* , *Phys. Rev. Lett.* **122** (2019) 222001 [[arXiv:1904.03947](#)] [[INSPIRE](#)].
- [69] U. Skerbis and S. Prelovsek, *Nucleon- J/ψ and nucleon- η_c scattering in P_c pentaquark channels from LQCD*, *Phys. Rev. D* **99** (2019) 094505 [[arXiv:1811.02285](#)] [[INSPIRE](#)].
- [70] H.-X. Chen, W. Chen and S.-L. Zhu, *Possible interpretations of the $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$* , *Phys. Rev. D* **100** (2019) 051501 [[arXiv:1903.11001](#)] [[INSPIRE](#)].
- [71] M.-Z. Liu et al., *Emergence of a complete heavy-quark spin symmetry multiplet: seven molecular pentaquarks in light of the latest LHCb analysis*, *Phys. Rev. Lett.* **122** (2019) 242001 [[arXiv:1903.11560](#)] [[INSPIRE](#)].
- [72] F.-K. Guo, H.-J. Jing, U.-G. Meißner and S. Sakai, *Isospin breaking decays as a diagnosis of the hadronic molecular structure of the $P_c(4457)$* , *Phys. Rev. D* **99** (2019) 091501 [[arXiv:1903.11503](#)] [[INSPIRE](#)].
- [73] J. He, *Study of $P_c(4457)$, $P_c(4440)$, and $P_c(4312)$ in a quasipotential Bethe-Salpeter equation approach*, *Eur. Phys. J. C* **79** (2019) 393 [[arXiv:1903.11872](#)] [[INSPIRE](#)].
- [74] Z.-H. Guo and J.A. Oller, *Anatomy of the newly observed hidden-charm pentaquark states: $P_c(4312)$, $P_c(4440)$ and $P_c(4457)$* , *Phys. Lett. B* **793** (2019) 144 [[arXiv:1904.00851](#)] [[INSPIRE](#)].
- [75] JPAC collaboration, *Interpretation of the LHCb $P_c(4312)^+$ signal*, *Phys. Rev. Lett.* **123** (2019) 092001 [[arXiv:1904.10021](#)] [[INSPIRE](#)].
- [76] Z.-G. Wang, *Analysis of the $P_c(4312)$, $P_c(4440)$, $P_c(4457)$ and related hidden-charm pentaquark states with QCD sum rules*, *Int. J. Mod. Phys. A* **35** (2020) 2050003 [[arXiv:1905.02892](#)] [[INSPIRE](#)].
- [77] F.-L. Wang, R. Chen, Z.-W. Liu and X. Liu, *Probing new types of P_c states inspired by the interaction between S -wave charmed baryon and anti-charmed meson in a $\bar{T}$ doublet*, *Phys. Rev. C* **101** (2020) 025201 [[arXiv:1905.03636](#)] [[INSPIRE](#)].
- [78] J.-B. Cheng and Y.-R. Liu, *$P_c(4457)^+$, $P_c(4440)^+$, and $P_c(4312)^+$: molecules or compact pentaquarks?*, *Phys. Rev. D* **100** (2019) 054002 [[arXiv:1905.08605](#)] [[INSPIRE](#)].
- [79] J.-J. Wu, T.S.H. Lee and B.-S. Zou, *Nucleon resonances with hidden charm in γp reactions*, *Phys. Rev. C* **100** (2019) 035206 [[arXiv:1906.05375](#)] [[INSPIRE](#)].

- [80] C.W. Xiao, J. Nieves and E. Oset, *Heavy quark spin symmetric molecular states from $\bar{D}^{(*)}\Sigma_c^{(*)}$ and other coupled channels in the light of the recent LHCb pentaquarks*, *Phys. Rev. D* **100** (2019) 014021 [[arXiv:1904.01296](#)] [[INSPIRE](#)].
- [81] M.B. Voloshin, *Some decay properties of hidden-charm pentaquarks as baryon-meson molecules*, *Phys. Rev. D* **100** (2019) 034020 [[arXiv:1907.01476](#)] [[INSPIRE](#)].
- [82] Y.-J. Xu, C.-Y. Cui, Y.-L. Liu and M.-Q. Huang, *Partial decay widths of $P_c(4312)$ as a $\bar{D}\Sigma_c$ molecular state*, *Phys. Rev. D* **102** (2020) 034028 [[arXiv:1907.05097](#)] [[INSPIRE](#)].
- [83] M.-Z. Liu, T.-W. Wu, M. Sánchez Sánchez, M.P. Valderrama, L.-S. Geng and J.-J. Xie, *Spin-parities of the $P_c(4440)$ and $P_c(4457)$ in the one-boson-exchange model*, *Phys. Rev. D* **103** (2021) 054004 [[arXiv:1907.06093](#)] [[INSPIRE](#)].
- [84] M.-L. Du et al., *Interpretation of the LHCb P_c states as hadronic molecules and hints of a narrow $P_c(4380)$* , *Phys. Rev. Lett.* **124** (2020) 072001 [[arXiv:1910.11846](#)] [[INSPIRE](#)].
- [85] B.-T. Zhang, J.-S. Wang and Y.-L. Ma, *Pentaquark states from bound state approach with chiral partner structure*, [arXiv:2002.10954](#) [[INSPIRE](#)].
- [86] S.-Q. Kuang, L.-Y. Dai, X.-W. Kang and D.-L. Yao, *Pole analysis on the hadron spectroscopy of $\Lambda_b \rightarrow J/\Psi p K^-$* , *Eur. Phys. J. C* **80** (2020) 433 [[arXiv:2002.11959](#)] [[INSPIRE](#)].
- [87] BESIII collaboration, *Observation of a near-threshold structure in the K^+ recoil-mass spectra in $e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$* , *Phys. Rev. Lett.* **126** (2021) 102001 [[arXiv:2011.07855](#)] [[INSPIRE](#)].