

The 750 GeV diphoton resonance as an sgoldstino: a reappraisal

Debjoyti Bardhan,^a Pritibhajan Byakti,^b Diptimoy Ghosh^c and Tarun Sharma^c

^a*Department of Theoretical Physics, Tata Institute of Fundamental Research,
1 Homi Bhabha Road, Mumbai 400005, India*

^b*Center for High Energy Physics, Indian Institute of Science,
Bangalore 560012, India*

^c*Department of Particle Physics and Astrophysics, Weizmann Institute of Science,
Rehovot 76100, Israel*

E-mail: debjyoti@theory.tifr.res.in, prtibhajan@cts.iisc.ernet.in,
diptimoy.ghosh@weizmann.ac.il, tarun.sharma@weizmann.ac.il

ABSTRACT: Among the various explanations of the possible 750 GeV diphoton resonance, the possibility of it being an sgoldstino is an attractive one, as it is related to the spontaneous breaking of global supersymmetry. We discuss this possibility in this paper and point out the various theoretical issues associated with it. In particular, we indicate the difficulties of this explanation in realistic models of gauge mediated supersymmetry breaking.

KEYWORDS: Supersymmetry Phenomenology

ARXIV EPRINT: [1603.05251](https://arxiv.org/abs/1603.05251)

Contents

1	Introduction	2
2	Generalities	3
2.1	Theoretical framework	3
2.2	Explaining the excess	5
3	Ordinary gauge mediation	5
3.1	Possibility of larger λ	9
3.2	Estimate of the mass of \mathcal{S}	10
4	Extra ordinary gauge mediation	11
5	Way out?	13
5.1	D -term contribution to the gaugino mass	13
5.2	Metastable SUSY breaking	14
5.3	Quark anti-quark initiated production of the sgoldstino	15
6	Conclusion	18
A	Calculation of the partial decay widths	18
A.1	$\phi \rightarrow \gamma\gamma$	19
A.2	$a \rightarrow \gamma\gamma$	20
A.3	$\phi \rightarrow ZZ$	20
A.4	$a \rightarrow ZZ$	21
B	Calculation of $\mathcal{A}_{\text{LHC energy}}^{ii}$	22
B.1	Production by gluon fusion	22
B.2	Production by quarks	24
C	Calculation of the sgoldstino mass	24
C.1	Diagrammatic calculation	25
C.2	Coleman-Weinberg potential	27
C.3	Tree level sgoldstino mass	28
D	Calculation of the gaugino mass	29

1 Introduction

The ATLAS and the CMS collaborations have recently reported some excess of events in the diphoton invariant mass ($m_{\gamma\gamma}$) distribution based on 3.2 and 2.6 fb⁻¹ of proton-proton collision data respectively collected at a center-of-mass energy of 13 TeV. ATLAS observed the most significant deviation from the background hypothesis at $m_{\gamma\gamma} \approx 750$ GeV, corresponding to a local (global) significance of 3.6 (2.0)¹ [1]. The largest excess in the CMS data was seen around the 760 GeV mass bin with a local (global) significance of 2.6 ($\lesssim 1.2$) standard deviations [2]. This excess is also found consistent with the constraints from the run 1 data [3]. It was also reported by ATLAS that the properties of the events in the signal region were found to be compatible with those in the invariant mass regions above and below the excess. As suggested by many authors, the most simple-minded explanation of this excess is to propose the existence of a resonance (\mathcal{S}) of mass ~ 750 GeV. In order to generate the correct amount of signal, the resonance must have couplings that produce $\sigma^{\text{signal}} \equiv \sigma(pp \rightarrow \mathcal{S})\text{Br}(\mathcal{S} \rightarrow \gamma\gamma)$ about 5 fb [3–5].

In this article, we consider the possibility of this resonance being an sgoldstino,² the “superpartner” of the goldstino, the goldstone fermion of spontaneous global supersymmetry (SUSY) breaking. This possibility has been discussed by [7–11] using an effective description of how the SUSY breaking is mediated to the MSSM sector. In this article, we scrutinise the viability of this proposal when realistic models for the mediation of SUSY breaking are considered. But before we start discussing that, we would like to make a few general comments about SUSY breaking in order to put things in perspective.

Unlike other symmetries, there are some interesting limitations on the possibility of spontaneous global SUSY breaking. For example, neither a pure super Yang-Mills (SYM) nor a SYM theory with massive matter in real representations of the gauge group breaks SUSY spontaneously.³ In particular, global $N = 2$ SYM theories (that have matter in real representations) cannot have SUSY spontaneously broken. This is one of the reasons why one needs global $N = 1$ SUSY with complex representation for phenomenology (i.e., MSSM) as there is a hope that SUSY can be spontaneously broken as required by experiments.

However, even in MSSM, it turns out to be impossible to break SUSY spontaneously. In fact, with the minimal field content of MSSM both the SUSY and the EW symmetry remain unbroken.⁴ Hence, adding more fields to the MSSM is unavoidable. However, even after adding many heavy fields, the gaugino masses cannot arise in a renormalisable SUSY theory at tree-level. This is because SUSY does not contain any (gaugino)-(gaugino)-(scalar) coupling that could give rise to a gaugino mass term when the scalar gets a vacuum expectation value (VEV). Moreover, the tree level supertrace rules do not allow a phenomenologically acceptable spectrum.

Hence, one possibility for breaking SUSY spontaneously in the MSSM is to have tree level SUSY breaking in a so-called “hidden sector” and radiatively mediate the information

¹This was obtained using a narrow width of the signal component. The statistical significance increases slightly once the possibility of larger width is taken into account. See [1] for more details.

²To our knowledge, the name “sgoldstino” was first used in [6].

³This follows from the fact that Witten Index of these theories is non-zero [12]. See also [13, 14].

⁴A Fayet-Iliopoulos D -term breaking also turns out to be phenomenologically unacceptable [15].

of SUSY breaking to the MSSM sector.⁵ This also helps in finding a solution of the SUSY flavour problem. As the pattern of SUSY breaking interactions in the visible MSSM sector is determined by the interactions of the messenger particles with the MSSM, a natural way to avoid additional flavour violation in the MSSM is to have flavour symmetries in the messenger interactions. The models of gauge mediation, where the information of SUSY breaking is communicated to the MSSM sector by gauge interactions, achieve this goal in a natural way.⁶ In the gauge mediation scenarios, one assumes the existence of “messenger fields” that are charged both under the SM gauge group as well as the hidden sector quantum numbers. The mass scale of these messengers is arbitrary and, in principle, can be as low as ~ 10 TeV. These models are often called “low scale SUSY breaking” scenarios and, as we will see later, are the only ones (among the different SUSY breaking scenarios) relevant for the diphoton excess.

In the following section, we review the general framework that leads to the sgoldstino explanation of the diphoton excess and present the necessary formulae to study the phenomenology. In section 3, we will discuss the ordinary gauge mediation (OGM) scenario and point out the various theoretical issues it confronts in connection to the diphoton excess. The generalisation of the OGM framework, called the extraordinary gauge mediation (EOGM), will be discussed in section 4. In section 5, we will investigate whether there is some way out of the difficulties raised in the previous sections. We will conclude in section 6.

2 Generalities

2.1 Theoretical framework

In order to parameterise the effect of SUSY breaking in the visible sector, it is usually assumed that SUSY is broken in the hidden sector by the VEV of the F component of a chiral superfield X . In particular, the gaugino masses are generated by the following terms,

$$\begin{aligned} \mathcal{L} \subset & -\frac{1}{2} \frac{c_1}{M_1} \int d^2\theta X W^{1\alpha} W_\alpha^1 - \frac{1}{2} \frac{c_2}{M_2} \int d^2\theta X W^{2\alpha A} W_\alpha^{2A} \\ & - \frac{1}{2} \frac{c_3}{M_3} \int d^2\theta X W^{3\alpha A} W_\alpha^{3A} + \text{h.c.} \end{aligned} \quad (2.1)$$

where the superscripts $\{1,2,3\}$ refer to the U(1), SU(2) and SU(3) gauge groups respectively (the adjoint indices for both the gauge groups SU(2) and SU(3) are denoted by A), and α is the spinor index. The scale M_i denotes the mass scale of the messenger fields which have been integrated out to get the above Lagrangian terms.⁷ The chiral superfield X and W_α

⁵Note that, in four space-time dimensions, if supersymmetry is not broken spontaneously at the tree level, then it can not be broken by radiative Coleman-Weinberg mechanism [16].

⁶This is however not true in general, as the messenger fields can have renormalisable superpotential couplings to the MSSM [17–22].

⁷In models of gravity mediation, the scale M_i is of the order of the planck scale. It is then clear that gravity mediation models are not relevant for the diphoton excess.

have the following expansion in terms of the ordinary fields,

$$X = \mathcal{S} + \sqrt{2}\theta\xi(y) + \theta\theta F_x(y) \quad (2.2)$$

$$= \frac{1}{\sqrt{2}}(\phi(y) + ia(y)) + \sqrt{2}\theta\xi(y) + \theta\theta F_x(y) \quad (2.3)$$

$$W_\alpha^A = -i\lambda_\alpha^A(y) + D^A(y)\theta_\alpha - (\sigma^{\mu\nu}\theta)_\alpha F_{\mu\nu}^A(y) - \theta\theta\sigma_{\alpha\dot{\beta}}^\mu D_\mu^{(y)}\lambda^{\dagger A\dot{\beta}}(y), \quad (2.4)$$

where, $y^\mu = x^\mu - i\theta\sigma^\mu\theta^\dagger$.

Once the F term of X gets a VEV, say $\langle F_x \rangle$, the above Lagrangian terms generate the following Majorana masses for the gauginos,

$$m_i = c_i \frac{\langle F_x \rangle}{M_i}. \quad (2.5)$$

The Lagrangian of eq. (2.1) also generates couplings of the scalar components of X to the gauge bosons,

$$\mathcal{L}_{gg} = \frac{1}{2\sqrt{2}} \frac{c_3}{M_3} \left(\phi G_{\mu\nu}^a G^{\mu\nu a} - a G_{\mu\nu} \tilde{G}^{\mu\nu} \right) \quad (2.6)$$

$$\mathcal{L}_{WW} = \frac{1}{2\sqrt{2}} \frac{2c_2}{M_2} \left(\phi W_{\mu\nu}^+ W^{-\mu\nu} - a W_{\mu\nu}^+ \tilde{W}^{-\mu\nu} \right) \quad (2.7)$$

$$\mathcal{L}_{\gamma\gamma} = \frac{1}{2\sqrt{2}} \left(\frac{c_1}{M_1} c_W^2 + \frac{c_2}{M_2} s_W^2 \right) \left(\phi F_{\mu\nu} F^{\mu\nu} - a F_{\mu\nu} \tilde{F}^{\mu\nu} \right) \quad (2.8)$$

$$\mathcal{L}_{ZZ} = \frac{1}{2\sqrt{2}} \left(\frac{c_1}{M_1} s_W^2 + \frac{c_2}{M_2} c_W^2 \right) \left(\phi Z_{\mu\nu} Z^{\mu\nu} - a Z_{\mu\nu} \tilde{Z}^{\mu\nu} \right) \quad (2.9)$$

$$\mathcal{L}_{Z\gamma} = \frac{1}{2\sqrt{2}} 2s_W c_W \left(\frac{c_2}{M_2} - \frac{c_1}{M_1} \right) \left(\phi Z_{\mu\nu} F^{\mu\nu} - a Z_{\mu\nu} \tilde{F}^{\mu\nu} \right). \quad (2.10)$$

The scalars ϕ and a can decay to the gauge bosons through these couplings. The corresponding partial decay rates are given by (see appendix A for details)⁸

$$\Gamma_{\gamma\gamma} \equiv \Gamma(\phi \rightarrow \gamma\gamma) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \right] \left[\frac{1}{8} \left(\frac{c_1}{M_1} c_W^2 + \frac{c_2}{M_2} s_W^2 \right)^2 \right] [8m_\phi^4] \left[\frac{1}{2} \right] \quad (2.11)$$

$$\Gamma_{gg} \equiv \Gamma(\phi \rightarrow gg) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \right] \left[\frac{1}{8} \left(\frac{c_3}{M_3} \right)^2 \right] [64m_\phi^4] \left[\frac{1}{2} \right] \quad (2.12)$$

$$\Gamma_{z\gamma} \equiv \Gamma(\phi \rightarrow Z\gamma) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \left(1 - \frac{m_Z^2}{m_\phi^2} \right) \right] \left[\frac{1}{8} \left(\frac{c_2}{M_2} - \frac{c_1}{M_1} \right)^2 4s_W^2 c_W^2 \right] \times \left[2m_\phi^4 \left(1 - \frac{m_Z^2}{m_\phi^2} \right)^2 \right] \quad (2.13)$$

$$\Gamma_{zz} \equiv \Gamma(\phi \rightarrow ZZ) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \left(1 - 4 \frac{m_Z^2}{m_\phi^2} \right)^{1/2} \right] \left[\frac{1}{8} \left(\frac{c_1}{M_1} s_W^2 + \frac{c_2}{M_2} c_W^2 \right)^2 \right] \times \left[8m_\phi^4 \left(1 - 4 \frac{m_Z^2}{m_\phi^2} + 6 \frac{m_Z^4}{m_\phi^4} \right) \right] \left[\frac{1}{2} \right] \quad (2.14)$$

⁸Signatures of sgoldstino at the e^+e^- and hadron colliders were first studied in [23, 24] where the formulae for the decay rates can also be found.

$$\Gamma_{ww} \equiv \Gamma(\phi \rightarrow WW) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \left(1 - 4 \frac{m_W^2}{m_\phi^2} \right)^{1/2} \right] \left[\frac{1}{8} \left(\frac{2c_2}{M_2} \right)^2 \right] \times \left[8m_\phi^4 \left(1 - 4 \frac{m_W^2}{m_\phi^2} + 6 \frac{m_W^4}{m_\phi^4} \right) \right] \quad (2.15)$$

Here s_W and c_W denote the sine and cosine of the Weinberg angle respectively. The partial decay rates for the scalar a can be obtained from the above expressions by replacing m_ϕ by m_a . There is slight difference between the decay rates of $\phi \rightarrow ZZ(W^+W^-)$ and $a \rightarrow ZZ(W^+W^-)$; however, that is numerically insignificant (see appendix A).

2.2 Explaining the excess

The total cross section for the diphoton production via the resonance \mathcal{S} is given by,⁹

$$\begin{aligned} \sigma_{\text{LHC energy}} &= \sigma(pp \rightarrow \mathcal{S})_{\text{LHC energy}} \text{Br}(\mathcal{S} \rightarrow \gamma\gamma) \\ &= \sum_i \mathcal{A}_{\text{LHC energy}}^{ii} \Gamma(\mathcal{S} \rightarrow p_i p_i) \frac{\Gamma(\mathcal{S} \rightarrow \gamma\gamma)}{\Gamma_{\mathcal{S}}}, \end{aligned} \quad (2.16)$$

where $\{p_i p_i\}$ refers to the initial state partons i.e., $\{gg\}$, $\{\bar{u}u\}$, $\{\bar{d}d\}$ and so on. The total width of \mathcal{S} is denoted by $\Gamma_{\mathcal{S}}$. The numerical values of the quantities $\mathcal{A}_{\text{LHC energy}}^{ii}$ are calculated in appendix B and are given by,

$$\begin{aligned} \mathcal{A}_{13}^{gg} &\equiv \mathcal{A}_{13 \text{ TeV LHC}}^{gg} = \frac{5.44 \text{ pb}}{\text{GeV}} & \mathcal{A}_8^{gg} &\equiv \mathcal{A}_{8 \text{ TeV LHC}}^{gg} = \frac{1.15 \text{ pb}}{\text{GeV}} \\ \mathcal{A}_{13}^{\bar{u}u} &\equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{u}u} = \frac{2.94 \text{ pb}}{\text{GeV}} & \mathcal{A}_8^{\bar{u}u} &\equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{u}u} = \frac{1.2 \text{ pb}}{\text{GeV}} \\ \mathcal{A}_{13}^{\bar{d}d} &\equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{d}d} = \frac{1.73 \text{ pb}}{\text{GeV}} & \mathcal{A}_8^{\bar{d}d} &\equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{d}d} = \frac{0.66 \text{ pb}}{\text{GeV}} \end{aligned} \quad (2.17)$$

In order to explain the signal, $\sigma_{13 \text{ TeV}}$ must be approximately in the range 3 – 8 fb, assuming that the resonance has a small width \lesssim few GeV [3]. A larger width of ~ 40 GeV requires $\sigma_{13 \text{ TeV}}$ to be slightly higher: $\sigma_{13 \text{ TeV}} \approx 5 - 14$ fb [3]. As the sgoldstino typically has a narrow width, in our estimates we will use the range 3 – 8 fb for the required cross section.

We will first consider the production by gluon fusion only, as the production by $u\bar{u}$ and $d\bar{d}$ initial states is slightly disfavoured [3–5]. In section 5.3, we will comment on the possibility of quark initiated production.

3 Ordinary gauge mediation

In the OGM framework, the hidden sector is parameterised by a single chiral superfield X . Both the scalar and auxiliary components of X are assumed to get VEVs that are denoted by $\langle \mathcal{S} \rangle$ and $\langle F_x \rangle$ respectively. In addition to this, OGM also includes N_5 vector like pairs

⁹Here we use the approximation that $\Gamma_{\mathcal{S}}/m_{\mathcal{S}}$ is small. This is a very good approximation even for the case when $\Gamma = 45$ GeV, which gives $\Gamma_{\mathcal{S}}/m_{\mathcal{S}} = 0.06$.

of messenger fields, $(\Phi_i, \tilde{\Phi}_i)$, transforming under $\mathbf{5} + \bar{\mathbf{5}}$ of $SU(5)$.¹⁰ The corresponding superpotential reads,

$$W_{\text{OGM}} = \lambda_{ij} X \tilde{\Phi}_i \Phi_j, \quad (3.1)$$

where the indices $\{i, j\}$ run from 1 to N_5 . Note that the matrix λ_{ij} can always be brought to a diagonal form with real entries by independent unitary rotations on Φ and $\tilde{\Phi}$ (the Kähler potential remain unchanged). Hence, in the rest of this section, we will assume that λ_{ij} is diagonal with $\lambda_{ii} \equiv \lambda_i$.

The fermions of each $\{\Phi_i, \tilde{\Phi}_i\}$ pair has a Dirac mass $m_F^i = \lambda_i \langle \mathcal{S} \rangle$. The mass eigenstates of the complex scalars, on the other hand, have squared masses $m_{\pm}^i = \lambda_i^2 \langle \mathcal{S} \rangle^2 \pm \lambda_i \langle F_x \rangle$. The gaugino masses are generated at the one loop level and are given by [26],

$$m_a = \frac{\alpha_a}{4\pi} \sum_{i=1}^{N_5} d_i^a \frac{\lambda_i \langle F_x \rangle}{m_F^i} g(x_i) \quad (a = 1, 2, 3) \quad (3.2)$$

where, $x_i = \frac{\lambda_i \langle F_x \rangle}{(m_F^i)^2}$ and the function $g(x)$ is given by [26],

$$g(x) = \frac{1}{x^2} [(1+x)\text{Log}(1+x) + (1-x)\text{Log}(1-x)]. \quad (3.3)$$

The symbol d_i denotes twice the Dynkin index for a particular representation. For example, in the case of $\mathbf{5} + \bar{\mathbf{5}}$ of $SU(5)$, $d = 1$. In eq. (3.2), we have used the GUT normalisation of the hypercharge gauge coupling.

Note that the SUSY breaking F -term VEV $\langle F_x \rangle$ must satisfy $\langle F_x \rangle \leq \lambda_i \langle \mathcal{S} \rangle^2, \forall i$ in order to avoid the messenger scalar masses from becoming tachyonic. For simplicity, we assume all the λ_i couplings to be equal and set them to a common value λ . We define the ratio $\lambda \langle F_x \rangle / m_F^2$ to be κ . With these definitions, the formula for the gaugino mass takes the form (for messengers in $\mathbf{5} + \bar{\mathbf{5}}$ of $SU(5)$),

$$m_a = \frac{\alpha_a}{4\pi} \kappa m_F N_5 g(\kappa) \quad (a = 1, 2, 3). \quad (3.4)$$

The c_a couplings (see eq. (2.5)) which control the signal strength are given by,

$$\frac{c_a}{M_a} = \frac{m_a}{\langle F_x \rangle} = \frac{\alpha_a}{4\pi} \frac{\lambda}{m_F} N_5 g(\kappa) \quad (a = 1, 2, 3). \quad (3.5)$$

Similarly, the scalar masses can be written as [27, 28],

$$\tilde{m}_a^2 = 2N_5 \kappa^2 m_F^2 \left[C_3^a \left(\frac{\alpha_3}{4\pi} \right)^2 + C_2^a \left(\frac{\alpha_2}{4\pi} \right)^2 + C_1^a \left(\frac{\alpha_1}{4\pi} \right)^2 \right] f(\kappa) \quad (3.6)$$

where C^a are the quadratic Casimirs and the function $f(x)$ is given by [26],

$$f(x) = \frac{1+x}{x^2} \left[\text{Log}(1+x) - 2\text{Li}_2 \left(\frac{x}{1+x} \right) + \frac{1}{2} \text{Li}_2 \left(\frac{2x}{1+x} \right) \right] + (x \rightarrow -x). \quad (3.7)$$

¹⁰Complete representations of a GUT group are normally used in order to keep the unification of the gauge couplings intact. However, in general, complete representations are not necessary. The use of incomplete representations often also have interesting phenomenology, see for example, [25] and the references therein.

In order to calculate the gaugino masses at the \sim TeV scale, we use the values of α_a at 1 TeV, which we compute using the one loop SM running equations,

$$\frac{1}{\alpha_a(\mu)} \simeq \frac{1}{\alpha_a(m_Z)} + \frac{b_a}{\pi} \text{Ln} \left(\frac{\mu}{m_Z} \right)$$

$$\left\{ \frac{1}{\alpha_1(m_Z)}, \frac{1}{\alpha_2(m_Z)}, \frac{1}{\alpha_3(m_Z)} \right\} = \{59, 30, 8.5\} \quad (3.8)$$

$$\{b_1^{\text{SM}}, b_2^{\text{SM}}, b_3^{\text{SM}}\} = \left\{ -\frac{41}{20}, \frac{19}{12}, \frac{7}{2} \right\} \quad (3.9)$$

We now examine the requirements on m_F , $\langle F_x \rangle$ and N_5 in order to generate the correct cross section for the excess. In order to have a feeling for the messenger mass scale required for the excess, we first consider a single pair of SU(5) messengers $\{\mathbf{5} + \bar{\mathbf{5}}\}$ i.e., $N_5 = 1$ and also set $\lambda = 1$. Following the discussion of the previous section, the explanation of the diphoton excess requires,¹¹

$$\mathcal{A}_{13}^{gg} \frac{\Gamma_{gg} \Gamma_{\gamma\gamma}}{\Gamma_{gg} + \Gamma_{\gamma\gamma} + \Gamma_{ww} + \Gamma_{zz} + \Gamma_{z\gamma}} \gtrsim 3 \text{ fb}. \quad (3.10)$$

This gives,

$$m_F \lesssim 175 \text{ GeV}. \quad (3.11)$$

The messenger scale can be raised if the number of messenger fields is increased. In figure 1 we show the allowed region in the m_F - N_5 plane for $\lambda = 1$ and $\kappa = 0.8$. In the left panel, only the contribution of ϕ to the signal is considered, while in the upper right panel contributions from both ϕ and a are taken into account. As discussed before, κ should satisfy $\kappa \leq 1$ to avoid tachyonic states in the messenger sector. For κ very close to unity, one of the complex scalars in every pair of messenger fields becomes too light (its squared mass is $m_F^2(1 - \kappa)$). Also, the function $f(\kappa)$ decreases rapidly for $\kappa \gtrsim 0.8$ [26] reducing the MSSM squark masses. Hence, we have chosen a value $\kappa = 0.8$ in figure 1.

The light green shaded region reproduces the correct amount of signal to explain the excess. In the light red shaded region, the gaugino masses are what is required by the exclusion limits of the LHC. In particular, the gluino mass is set to more than 1.5 TeV and a conservative lower bound of 200 GeV is considered for the bino and wino masses (we also show the region satisfying a stricter lower bound of 650 GeV on the bino and wino masses [29]). Similarly, in the light blue region the squarks are heavier than a TeV. It can be seen that a very large number of messengers $\gtrsim 60$ is required in order to both successfully explain the signal as well as produce sufficiently large gaugino and squark masses.

However, for such a large number of messenger fields, the gauge couplings lose asymptotic freedom. The one-loop running of the gauge couplings above the messenger fermion mass m_F is shown in figure 2 for two sets of values of $\{m_F, N_5\}$, shown as black dots in

¹¹Here we have neglected any decay mode other than the gauge boson final states. However, existence of other decay modes will increase the total width of the resonance, hence adding an extra contribution to the denominator of eq. (3.10). This means that the required signal cross section will be even higher, as pointed out also in the end of section 2.2. Thus our estimate is on the conservative side.




Figure 1. Allowed region in the OGM parameter space that successfully explains the signal and satisfy LHC bounds on squark and gaugino masses. While in the left panel the contribution from only ϕ is considered, the right panel takes into account both ϕ and a contributions.

figure 1. The point $\{m_F, N_5\} = \{14 \text{ TeV}, 65\}$ is chosen such that all the requirements namely, correct amount of the signal cross section and heavy enough gaugino and squark masses are satisfied. It can be seen from the left panel of figure 2 that the SU(3) gauge coupling in this case hits a one-loop Landau pole below $\sim 50 \text{ TeV}$. The right panel of figure 2 shows the renormalisation group (RG) running for $\{m_F, N_5\} = \{7 \text{ TeV}, 35\}$ i.e., when the constraint from the squark masses is relaxed. This is relevant for example, in models where the squark masses are generated at the tree level [30, 31]. However, even in this case, the required number of messenger pairs is $\gtrsim 35$ and the one-loop Landau pole is encountered below $\sim 80 \text{ TeV}$.

Before concluding this section, we would like to make two final comments:

- i) Although we have presented our results for messengers transforming under $\{\mathbf{5} + \bar{\mathbf{5}}\}$ of SU(5), our general conclusions hold for other representations also and even in the case when the possibility of doublet-triplet splitting is considered (this will be more clear in section 5.3).
- ii) The formula in eq. (3.5) is strictly valid only if the SUSY breaking VEV is small namely, $\kappa \ll 1$. For $\kappa \sim 1$, one has to compute the separate loop contributions from the messenger scalar with masses $m_{\pm}^2 = m_F^2(1 \pm \kappa)$. This gives a correction factor $\sim \left(\frac{1-2/3\kappa^2}{1-\kappa^2}\right)^2$ in the decay rates for the scalar ϕ (here we have assumed $\lambda = 1$ for simplicity). This factor is only ≈ 2.5 for $\kappa = 0.8$ which we use for our analysis¹² and is absent for a . Hence, this does not affect our numerical analysis.

¹²The paper [32] which appeared *after* the first version of our paper considered the very fine tuned possibility of κ being extremely close to unity which may somewhat mitigate the problem, however, at the cost of very large trilinear coupling between the sgoldstino and some of the light messenger scalars. We do not consider this extremely fine-tuned possibility in this paper.




Figure 2. RG running of the SM gauge couplings above m_F for the two representative sets of values of $\{m_F, N_5\}$ shown as black dots in figure 1, see text for more details. The values of the couplings at the scale m_F is obtained using the SM evolution from m_Z to 2 TeV and the MSSM evolution from 2 TeV to m_F .




Figure 3. RG evolution of λ for $N_5 = 5$ and for two initial values of λ : $\lambda(10 \text{ TeV}) = 1$ (left) and $\lambda(10 \text{ TeV}) = 2$ (right).

3.1 Possibility of larger λ

It can be seen from eq. (3.5) that, for a given gaugino mass, the c_i coefficients (hence, diphoton signal cross section) can be increased by increasing λ . However, one should first check the RG running of λ in order to see the maximum value of λ that is safe.

As the fundamental representation of SU(5) can be decomposed into representations of SU(3) \otimes SU(2) \otimes U(1) in the following way,

$$\mathbf{5} \rightarrow (3, 1)_{-1/3} \oplus (1, 2)_{1/2} \tag{3.12}$$

the superpotential can be rewritten as,

$$W = \lambda_i^{D^c} X \Phi_i^{D^c} \tilde{\Phi}_i^{D^c} + \lambda_i^L X \Phi_i^L \tilde{\Phi}_i^L \quad (3.13)$$

Note that, the notation D^c and L have been used just for notational convenience and they do not represent the MSSM fields. The beta functions of these couplings are given by

$$\beta_{\lambda_i^{D^c}} = \lambda_i^{D^c} \left(\gamma(\Phi_i^{D^c}) + \gamma(\tilde{\Phi}_i^{D^c}) + \gamma(X) \right), \quad (3.14)$$

$$\beta_{\lambda_i^L} = \lambda_i^L \left(\gamma(\Phi_i^L) + \gamma(\tilde{\Phi}_i^L) + \gamma(X) \right), \quad (3.15)$$

where

$$\gamma(\Phi_i^{D^c}) = \gamma(\tilde{\Phi}_i^{D^c}) = \frac{1}{4\pi} \left(-\frac{2\alpha_1}{15} + \alpha_{di} - \frac{8\alpha_3}{3} \right) \quad (3.16)$$

$$\gamma(\Phi_i^L) = \gamma(\tilde{\Phi}_i^L) = \frac{1}{4\pi} \left(-\frac{3\alpha_1}{10} + \alpha_{li} - \frac{3\alpha_2}{2} \right) \quad (3.17)$$

$$\gamma(X) = \sum_i \frac{1}{4\pi} (3\alpha_{di} + 2\alpha_{li}) \quad (3.18)$$

We have used the notation, $\alpha_{di} \equiv \frac{(\lambda_i^{D^c})^2}{4\pi}$ and $\alpha_{li} \equiv \frac{(\lambda_i^L)^2}{4\pi}$.

Hence, the RG equations for the λ couplings are,

$$\frac{d\lambda_i^{D^c}}{dt} = \frac{1}{16\pi^2} \lambda_i^{D^c} \left[(3N+2) (\lambda_i^{D^c})^2 - \frac{16}{3} g_3^2 - \frac{4}{15} g_1^2 + 2N (\lambda_i^L)^2 \right], \quad (3.19)$$

$$\frac{d\lambda_i^L}{dt} = \frac{1}{16\pi^2} \lambda_i^L \left[(2N+2) (\lambda_i^L)^2 - 3g_2^2 - \frac{3}{5} g_1^2 + 3N (\lambda_i^{D^c})^2 \right], \quad (3.20)$$

In figure 3 we show the running of these λ couplings for five pairs of $\{\mathbf{5} + \bar{\mathbf{5}}\}$ messengers and for two initial values of λ at the scale 10 TeV, $\lambda(10 \text{ TeV}) = 1$ and 2. It can be seen from the right panel of figure 3 that even for $\lambda(10 \text{ TeV}) = 2$, it grows very fast and hits a one-loop Landau-pole below ~ 25 TeV. Needless to say, the situation gets worse if a larger number of messenger pairs is considered. Hence, we conclude from this analysis that values of λ much larger than unity at the messenger scale is not a possibility.

3.2 Estimate of the mass of \mathcal{S}

It was shown in [33] that in renormalizable Wess-Zumino models with canonical Kähler potential, the existence of a massless fermion implies that the complex scalar in the same chiral multiplet remains massless at the tree level even if SUSY is spontaneously broken. As the fermion component of X is the goldstino in our case (which is exactly massless even at loop level), the scalar component of X , the sgoldstino will be massless at the tree level. However, in general, the sgoldstino is expected to acquire non-zero mass when loop corrections are included.

In our scenario, the sgoldstino mass gets contribution from the loops of messenger fields (apart from possible contributions from the hidden sector). The messenger contribution is computed in appendix C. The final result is given by (for N_5 pairs of $\mathbf{5} + \bar{\mathbf{5}}$ of SU(5)),

$$\Pi(p^2 = 0) = - \left(\frac{\lambda}{g_3^2} \right)^2 \left(4\pi \sqrt{\frac{5}{N_5}} F(x) \right)^2 m_g^2 \quad (3.21)$$

Hence, the potential for the sgoldstino gets a one-loop negative quadratic contribution from the messenger fields and this contribution is considerably larger in magnitude than the squared gluino mass.¹³ This means that a large contribution from the hidden sector is required to stabilise the sgoldstino potential and somehow generate a small mass ~ 750 GeV for the sgoldstino.

At this point, we would like to remind the readers that, in our discussions till now, we have completely ignored specifying the details of the hidden sector and how SUSY is broken there. We just assumed that the chiral superfield X gets a SUSY breaking F -term VEV from the dynamics of the hidden sector without specifying the hidden sector at all. However, in order to understand whether a light sgoldstino can be obtained without too much tuning, we are now forced to consider the hidden sector as part of our model and think about the problem in its entirety. We postpone any further investigation of this issue to section 5.

4 Extra ordinary gauge mediation

We have seen in the previous section that the OGM framework needs a very large number of messengers in order to explain the diphoton signal and avoid the strong constraints on the gluino and squark masses from LHC. We have also seen that such a large number of messengers renders the theory non-perturbative at scales as low as ~ 50 TeV, much below the GUT scale.

In this section we will consider a generalisation of the OGM framework namely, the Extra Ordinary Gauge Mediation (EOGM) where the OGM Lagrangian (eq. (3.1)) is supplemented with vector-like mass terms for the chiral superfields $\tilde{\Phi}_i$ and Φ_j [35]. Hence, we now have the EOGM superpotential

$$W_{\text{EOGM}} = (\lambda_{ij}X + m_{ij}) \tilde{\Phi}_i \Phi_j, \tag{4.1}$$

where, λ_{ij} and m_{ij} are arbitrary complex matrices. As in the OGM scenario, the auxiliary field of X is assumed to get a VEV to break SUSY spontaneously. The fermion components of the messenger fields have the Dirac mass matrix,

$$m_F = \lambda_{ij} \langle S \rangle + m_{ij}. \tag{4.2}$$

Without loss of generality, one can always go to the basis of $\tilde{\Phi}$ and Φ (by independent unitary rotations on them that do not affect their Kähler potential) where m_F is diagonal with real eigenvalues $(m_F)_i$. Hence, from now on we will assume that the matrix m_F is diagonal and the matrices λ_{ij} and m_{ij} are defined in the basis where m_F is diagonal. The scalar mass-squared matrix in this basis can now be written as,

$$\tilde{m}^2 = \begin{pmatrix} m_F^2 & -\lambda \langle F_x \rangle \\ -\lambda \langle F_x \rangle & m_F^2 \end{pmatrix}. \tag{4.3}$$

¹³Note that, models with non-polynomial superpotential can give rise to tree level sgoldstino mass. We compute the goldstino mass in one such model [34] in appendix C.3, however, again it turns out to be in general much larger than the gluino mass.

We will assume the matrix λ to be real and symmetric in order to impose invariance under CP and messenger parity (i.e., $\Phi_i \rightarrow \tilde{\Phi}_i$ in the basis where m_F is diagonal) in the messenger sector [36, 37].

The matrix \tilde{m}^2 can be block diagonalised by a suitable change of basis of the scalar fields, the block diagonalised matrix being,

$$\mathcal{M}^2 = \begin{pmatrix} m_+^2 & 0 \\ 0 & m_-^2 \end{pmatrix}, \tag{4.4}$$

where $m_{\pm}^2 = m_F^2 \pm \lambda \langle F_x \rangle$. Now assuming that the matrices m_{\pm}^2 are diagonalised by the unitary matrices U_{\pm} , the gaugino masses can be written as [38],

$$m_a = \frac{\alpha_a}{4\pi} \sum_{\pm} \sum_{i,j=1}^N (\pm) (U_{\pm}^{\dagger})_{ij} (U_{\pm})_{ji} m_j \frac{m_{\pm i}^2 \text{Log}(m_{\pm i}^2/m_j^2)}{m_{\pm i}^2 - m_j^2}. \tag{4.5}$$

Let us now consider only one pair of messengers to simplify the discussion. In this case the expressions of the gaugino masses and couplings c_a take the same form as the OGM case,

$$m_a = \frac{\alpha_a}{4\pi} \kappa m_F g(\kappa) \tag{4.6}$$

$$\frac{c_a}{M_a} = \frac{\alpha_a}{4\pi} \frac{\lambda}{m_F} g(\kappa) \tag{4.7}$$

the only difference being in the definition of m_F which now has the form,

$$m_F = \lambda \langle \mathcal{S} \rangle + m. \tag{4.8}$$

Hence, for fixed values of the messenger fermion masses, the situation is exactly the same as OGM. In the presence of many pair of messengers, if $[m_F, \lambda] = 0$ then the matrix λ can be diagonalised simultaneously with m_F and hence, the situation is again exactly the same as OGM with many messenger fields. In the case when $[m_F, \lambda] \neq 0$, in general, one has to analyse the situation numerically. Analytic results are known even in this case for $\lambda \langle F_x \rangle \ll m_F^2$ [35, 39]:

- The R charge for the field X , $R(X) \neq 0$: in this case the expression of the gaugino mass can be written as,

$$m_a = \frac{\alpha_a}{4\pi} n_{\text{eff}} \frac{\langle F_x \rangle}{\langle \mathcal{S} \rangle} \tag{4.9}$$

where,

$$n_{\text{eff}} = \frac{1}{R(X)} \sum_i \left(2 - R(\Phi_i) - R(\tilde{\Phi}_i) \right). \tag{4.10}$$

As n_{eff} is less than the total number of messengers, the gaugino mass in this case is always less than that in the OGM case.

- $R(X) = 0$, even in this case the expression of the gaugino mass simplifies to,

$$m_a = \frac{\alpha_a}{4\pi} \langle F_x \rangle \sum_i \frac{\lambda_{ii}}{m_F^i}, \tag{4.11}$$

If $\min(m_F^i) = m$, then

$$m_a \leq \frac{\alpha_a}{4\pi} \frac{\langle F_x \rangle}{m} \text{Tr} \lambda. \tag{4.12}$$

Hence, the situation is again the same as the OGM case.

We have checked numerically that the situation does not improve for the case when $\lambda \langle F_x \rangle \sim m_F^2$.

5 Way out?

We have seen in the previous sections that an sgoldstino explanation of the diphoton excess faces two major issues: i) the gaugino masses, and in particular the gluino mass, turn out to be rather low unless a very large number of messenger fields is considered; ii) the messenger particles yield a large negative one loop contribution to the sgoldstino potential. In this section, our goal is to look for potential solutions of the above problems.

5.1 D -term contribution to the gaugino mass

We have only considered F -term contribution to the gaugino mass in the previous sections. We will now assume that the messenger fields are also charged under some new $U(1)$ gauge group. The Φ fields have charge $+1$ and the $\tilde{\Phi}$ fields carry a charge -1 under this new $U(1)$. The relevant part of the Lagrangian is given by,

$$\mathcal{L} \subset \int d^4\theta \left(\Phi_i^\dagger e^{gV} \Phi_i + \tilde{\Phi}_i^\dagger e^{-gV} \tilde{\Phi}_i \right) + \int d^2\theta (\lambda_{ij} X + m_{ij}) \tilde{\Phi}_i \Phi_j + \text{h.c.} \tag{5.1}$$

The F -term of the chiral superfield X and the D -term of the vector superfield V are assumed to have VEVs $\langle F_x \rangle$ and $\langle D \rangle$ respectively.¹⁴ However, since the above Lagrangian possesses an $U(1)$ R -symmetry, the charges being $R(\Phi) = 1, R(\tilde{\Phi}) = 1, R(X) = 0$ and $R(V) = 0$, it follows that the F -term and the D -term have the R -charges $R(F) = 2$ and $R(D) = 0$. Hence, $\langle F_x \rangle \neq 0$ breaks R -symmetry spontaneously, while $\langle D \rangle \neq 0$ does not. It is then clear that the gaugino masses must be associated with non-zero $\langle F_x \rangle$.

As we discussed previously, the leading F -term contribution to the gaugino mass comes from the term

$$-\frac{1}{2} \frac{c_F}{\Lambda} X W_A W^A. \tag{5.2}$$

As the gaugino mass is always associated with $\langle F_x \rangle$, the D -term contribution must always be suppressed by higher powers of Λ and hence, subdominant compared to the leading F -term contribution. That there is no D -term contribution at the leading order in the F -term VEV can also be understood diagrammatically. It can be seen from figure 4 that, in order to join the scalar lines, one needs a term $\phi_1 \phi_2$ in the Lagrangian (refer to appendix C for the notation) which does not arise from the D -term.

¹⁴Note that the existence of non-zero $\langle D \rangle$ breaks the messenger parity spontaneously.




Figure 4. Diagram showing that a D -term does not contribute to the Majorana gaugino mass at the leading order in the F -term VEV.

In models with explicitly broken R -symmetry, the lowest dimensional operators that can give rise to the gaugino mass should be,

$$-\frac{1}{2} \frac{c_D}{\Lambda_D^3} \widetilde{W} \widetilde{W} W_A W^A \tag{5.3}$$

which generates a contribution,

$$m_\lambda = c_D \frac{\langle D_{\widetilde{W}} \rangle^2}{\Lambda_D^3}, \tag{5.4}$$

which is subleading compared to (5.2). The chiral superfield \widetilde{W} belongs to the hidden sector and corresponds to either an abelian or a non-abelian gauge group. Note that, as mentioned before, the term in (5.3) breaks R -symmetry explicitly. We thus conclude that D -term contribution can not enhance the gaugino mass considerably.

We would like to comment in passing on the problem of vanishing leading order (in SUSY breaking F term VEV) gaugino masses in models of direct gauge mediation [40, 41] and semi-direct gauge mediation [42], regardless of how the R -symmetry is broken. The authors of [33] proved this in generalised renormalizable O’Raifeartaigh models assuming a locally stable pseudomoduli space. This problem can be avoided with non-polynomial superpotential which naturally appears in many models of dynamical/non-perturbative SUSY breaking (DSB) [43–45]. Hence, the gaugino mass to leading order in $\langle F_x \rangle$ that were considered in the previous sections should indeed be thought in the framework of DSB models.

5.2 Metastable SUSY breaking

Before going to the discussion of metastable SUSY breaking, it is worth reviewing briefly the relation between R -symmetry and spontaneous SUSY breaking.

Consider a generic model of gauge mediated supersymmetry breaking in which a *Hidden* sector (HS) consisting of the superfields (Y_a, X) breaks supersymmetry and then *messenger* fields $(\Phi_i, \tilde{\Phi}_i)$ communicate the supersymmetry breaking to the *visible* MSSM sector via loop effects. The hidden sector fields are neutral under the Standard Model gauge group but could have its own gauge dynamics while the messenger fields $(\Phi_i, \tilde{\Phi}_i)$ transform in a vector like representation of SM gauge group and could also be charged under the HS gauge group.

Let us write the full superpotential of the theory as follows

$$\begin{aligned}
 W &= W_{\text{HS}}(\{Y_a\}, X) + W_M(X, \Phi_i, \tilde{\Phi}_i) + W_{\text{MSSM}}, \\
 \text{with } W_M &= \lambda_{ij} X \Phi_i \tilde{\Phi}_j + m_{ij} \Phi_i \tilde{\Phi}_j.
 \end{aligned}
 \tag{5.5}$$

Here W_{MSSM} is the MSSM superpotential and W_{HS} is hidden sector superpotential which spontaneously breaks SUSY.¹⁵

What can one say about the R -symmetry in W_{HS} ? Note that, for generic superpotential without R -symmetry, Nelson and Seiberg showed that a supersymmetric vacuum always exists [46]. In other words, R -symmetry is a necessary (but not sufficient) condition for spontaneous breaking of supersymmetry. However, unbroken R -symmetry forbids (Majorana) masses for the gauginos. Thus, it must be broken spontaneously which, in turn, would lead to a massless R -axion that may be dangerous for phenomenology.¹⁶

Another possibility is to break R -symmetry explicitly in hidden sector (W_{HS}). Now it is possible to write down models with no R -symmetry which break SUSY spontaneously but these models have a non-generic superpotential in the sense that it doesn't allow all renormalisable terms allowed by symmetries. As superpotential couplings are protected from renormalisation and hence are not generated at loop levels, a non generic superpotential is technically natural. However, it is tuned and not satisfactory.

One scenario which avoids these problems is metastable supersymmetry breaking [48]. It is based on the idea that though the true vacuum is supersymmetric, our universe lies in a metastable vacuum. In this picture, there is no need to keep R -symmetry but one does need to worry about decay rates from the metastable vacuum to the true vacuum and arrange for a long lived universe.

As mentioned in the previous section, the problem of vanishing leading order (in SUSY breaking F -term) gaugino masses can be avoided in models of DSB. Hence, DSB in a metastable vacuum is an attractive phenomenological possibility. In fact, some of these models can potentially solve the problem mentioned in section 3.2 and give rise to a light sgoldstino [49–51]. However, detailed exploration of these models is necessary to see whether they can indeed serve as natural models for a light sgoldstino and avoid the problems mentioned in section 3.

5.3 Quark anti-quark initiated production of the sgoldstino

In this section, we consider the possibility that the production cross section of sgoldstino has a significant contribution from quark anti-quark initial state. The coupling of the sgoldstino to the quark anti-quark pair can arise from the same effective Lagrangian that generates the trilinear A -terms namely,

$$\mathcal{L}_{\text{trilinear}} \subset \frac{A_u}{\langle F_x \rangle} \int d^2\theta X H_u Q U^c + \frac{A_d}{\langle F_x \rangle} \int d^2\theta X H_d Q D^c + \text{h.c.}
 \tag{5.6}$$

¹⁵Note that the R -parity conserving MSSM has three parameter worth of R -symmetries. However, R -symmetry has gauge anomalies in the MSSM.

¹⁶ R -symmetry may be broken by Gravity effects, thus giving mass to the R -axion [47].




Figure 5. Allowed region in the case of quark anti-quark initiated production of the sgoldstino. See text for more details.

which generates following couplings for the sgoldstino,

$$\frac{v \sin \beta A_u}{\sqrt{2} \langle F_x \rangle} \mathcal{S} \bar{u} P_L u, \quad \frac{v \cos \beta A_d}{\sqrt{2} \langle F_x \rangle} \mathcal{S} \bar{d} P_L d. \quad (5.7)$$

The decay rates $\Gamma(\phi \rightarrow \bar{u} u)$ and $\Gamma(\phi \rightarrow \bar{d} d)$ can now be calculated from the above Lagrangian and read,

$$\Gamma(\phi \rightarrow \bar{u} u) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \right] \left[\left(\frac{v \sin \beta A_u}{\sqrt{2} \langle F_x \rangle} \right)^2 3m_\phi^2 \right], \quad (5.8)$$

$$\Gamma(\phi \rightarrow \bar{d} d) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \right] \left[\left(\frac{v \cos \beta A_d}{\sqrt{2} \langle F_x \rangle} \right)^2 3m_\phi^2 \right], \quad (5.9)$$

where we have neglected the quark masses. In this limit, the corresponding decay rates of a have the same expressions with m_ϕ replaced by m_a .

We now assume that the production of sgoldstino is mostly by the $\bar{u} u$ and $\bar{d} d$ initial states so that a large coupling to gluons is not necessary. We define the number of messengers with quantum numbers $(1, 2)_{1/2}$ to be N_{12} . Their mass will be denoted by m_F^{weak} . In the left panel of figure 5 we show the allowed region in the $N_{12} - m_F^{\text{weak}}$ plane when two sets of values for A_u and A_d are chosen.¹⁷ Similarly, in the right panel the allowed region in the $N_{12} - A_u/A_d$ plane is shown for $m_F^{\text{weak}} = 8 \text{ TeV}$. It can be seen that even for very large value of $A_u = A_d \sim 10 \text{ TeV}$,¹⁸ quite low masses for the electroweak messenger fields $m_F^{\text{weak}} \lesssim 10 \text{ TeV}$ with a very large multiplicity $\gtrsim 50$ are necessary. Consequently, the SU(2)

¹⁷In general, A -terms are generated at 1-loop level in the models of messenger matter interactions. Thus they are of same order of the gaugino masses. Larger A -terms can be obtained from model where A -terms are generated at the tree level [52]. These models have the advantage of being free from A/m^2 problem [20].

¹⁸Note that very large A -terms may give rise to electric charge and SU(3) colour breaking minima in the potential [53, 54], thus we restrict them to 10 TeV in our analysis.




Figure 6. Allowed region in the case of quark anti-quark initiated production of the sgoldstino with only light hypercharge messengers.

and U(1) couplings (i.e., $g_i^2/4\pi$) hit Landau poles typically below few hundred TeV. For example, for $m_F^{\text{weak}} = 8$ TeV and $N_{12} = 100$, the one loop Landau poles for SU(2) and U(1) appear around 50 TeV and 200 TeV respectively.

As the SUSY breaking F -term VEV $\langle F_x \rangle$ must be less than $(m_F^{\text{weak}})^2$ in order to avoid tachyons in the messenger sector, it also turns out that a gluino mass of more than 1.5 TeV again requires a very large number of SU(3) messengers, exactly as in the OGM scenario discussed earlier.

However, one could consider a scenario where the X superfields that couple to the SU(3) messengers (denoted by Φ_3 and $\tilde{\Phi}_3$ below) are different from the X superfields that couple to the SU(2) messengers (denoted by Φ_2 and $\tilde{\Phi}_2$ below) so that,

$$W = (X_2 + m_2)\tilde{\Phi}_2\Phi_2 + (X_3 + m_3)\tilde{\Phi}_3\Phi_3, \quad (5.10)$$

The X_2 and X_3 superfields get VEVs given by,

$$\langle X_2 \rangle = \langle \mathcal{S}_2 \rangle + \theta\theta\langle F_2 \rangle, \quad (5.11)$$

$$\langle X_3 \rangle = \langle \mathcal{S}_3 \rangle + \theta\theta\langle F_3 \rangle. \quad (5.12)$$

One can define two complex scalars that are linear combinations of \mathcal{S}_2 and \mathcal{S}_3 ,

$$\mathcal{S}_h = \frac{F_2\mathcal{S}_2 + F_3\mathcal{S}_3}{\sqrt{F_2^2 + F_3^2}} \quad (5.13)$$

$$\mathcal{S}_l = \frac{-F_3\mathcal{S}_2 + F_2\mathcal{S}_3}{\sqrt{F_2^2 + F_3^2}} \quad (5.14)$$

In the limit of $F_3 \gg F_2$, $\mathcal{S}_h \approx \mathcal{S}_3$ and $\mathcal{S}_l \approx \mathcal{S}_2$. If we now assume that the scalar \mathcal{S}_l is actually the 750 GeV resonance and the other scalar \mathcal{S}_h is much heavier than the diphoton

signal can be explained. Moreover, as F_3 is now assumed to be much larger than F_2 , large gluino mass can also be easily obtained.

However, it should be mentioned that the scalar \mathcal{S}_l is actually not the sgoldstino. It is actually \mathcal{S}_h which appears in the goldstino multiplet, hence, \mathcal{S}_h should be identified as the sgoldstino. In this sense, we have not solved the original problem with sgoldstino being the candidate for the 750 GeV resonance.

Before concluding this section, we would also like to point out that one can also consider the extreme case when there are three different superfields X_1 , X_2 and X_3 that couple to the U(1), SU(2) and SU(3) messengers respectively. In this case, both the SU(2) and SU(3) messenger masses can be very high. In figure 6 we show the number of U(1) messengers (N_1) and their mass (m_F^Y) required for the correct amount of signal and also mass of Bino more than 200 GeV. It can be seen that for $m_F^Y \sim 5$ TeV one needs $N_1 \gtrsim 50$. The Landau pole in the U(1) gauge coupling only appears around 2000 TeV in this case.

6 Conclusion

In this paper we have carefully studied the possibility of an sgoldstino being a candidate for the signal of a possible new resonance with mass ~ 750 GeV recently reported by the ATLAS and CMS collaborations. We have found that the explanation of the signal is in tension with the lower bound on masses from direct searches of gauginos, in particular, the gluino. In order to achieve a large enough gluino mass, a very large number of messenger fields is required, which, in turn, renders the theory non-perturbative at a rather low scale of order few tens of TeV. Moreover, we find that the one-loop messenger contribution to the sgoldstino potential is negative and large in magnitude (larger than the gluino mass squared). Hence, a large positive contribution from the hidden sector is required to tune this away and get a small mass ~ 750 GeV for the sgoldstino.

While there exist examples of models with dynamically broken SUSY where a light sgoldstino can, in principle, be achieved, perhaps without large tuning, getting both the correct amount of signal cross-section and also large enough gluino and squark masses (without spoiling the calculability of the theory at a rather low scale) seems to be a stubborn problem. It would be interesting to find explicit models where these problems can be overcome in a satisfactory way. We postpone investigation in this direction to future studies.

We have also considered the possibility of the resonance being produced by quark anti-quark initial state. While in this case the problem of Landau poles can be delayed beyond few thousand TeV, the scalar resonance can not be the sgoldstino.

Acknowledgments

We thank Zohar Komargodski for many useful discussions and comments on this work. DG thanks Yevgeny Kats, Paride Paradisi, Fabrizio Senia and Fabio Zwirner for fruitful discussions. DB thanks Sabyasachi Chakraborty for several useful discussions.

A Calculation of the partial decay widths


In this appendix we will calculate the partial decay rate of ϕ and a to two vector bosons.

A.1 $\phi \rightarrow \gamma\gamma$

We start with the decay $\phi \rightarrow \gamma\gamma$ which arises from the following term in the Lagrangian,

$$\mathcal{L} \subset \frac{1}{\Lambda} \phi F_{\mu\nu} F^{\mu\nu}. \quad (\text{A.1})$$

This yields the following Feynman rule,



$$[i] [2!] \left[-\frac{2}{\Lambda} (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) \right] \quad (\text{A.2})$$

Thus, the matrix element is given by,

$$i\mathcal{M} = -\frac{4}{\Lambda} i (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) \epsilon^{*\nu}(p_1) \epsilon^{*\mu}(p_2) \quad (\text{A.3})$$

This gives,

$$|\mathcal{M}|^2 = \frac{16}{\Lambda^2} (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) (p_1 \cdot p_2 g_{\alpha\beta} - p_{1\alpha} p_{2\beta}) \epsilon^{*\nu}(p_1) \epsilon^{*\mu}(p_2) \epsilon^\beta(p_1) \epsilon^\alpha(p_2) \quad (\text{A.4})$$

Summing over the polarizations, i.e.,

$$\sum \epsilon^\mu(p) \epsilon^{*\nu}(p) = -g^{\mu\nu}$$

we get,

$$\begin{aligned} |\mathcal{M}|^2 &= \frac{16}{\Lambda^2} (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) (p_1 \cdot p_2 g_{\alpha\beta} - p_{1\alpha} p_{2\beta}) g^{\alpha\mu} g^{\beta\nu} \\ &= \frac{16}{\Lambda^2} (p_1^2 p_2^2 + 4(p_1 \cdot p_2)^2 - 2(p_1 \cdot p_2)^2) \\ &= \frac{32}{\Lambda^2} (p_1 \cdot p_2)^2 \\ &= \frac{32}{\Lambda^2} \left(\frac{m_\phi^2}{2} \right)^2 \\ &= \frac{8m_\phi^4}{\Lambda^2} \end{aligned} \quad (\text{A.5})$$

Hence,

$$\Gamma(\phi \rightarrow \gamma\gamma) = \frac{1}{\Lambda^2} \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \right] [8m_\phi^4] \left[\frac{1}{2} \right]. \quad (\text{A.6})$$

The factor of 1/2 in the end is due to the presence of two identical particles in the final state.

A.2 $a \rightarrow \gamma\gamma$

The decay $a \rightarrow \gamma\gamma$ arises from the Lagrangian

$$\mathcal{L} \subset \frac{1}{\Lambda} a F_{\mu\nu} \tilde{F}^{\mu\nu} = \frac{1}{2\Lambda} a F_{\mu\nu} F_{\alpha\beta} \varepsilon^{\mu\nu\alpha\beta} \quad (\text{A.7})$$

The Feynman rule for this vertex is given by

$$[i] [2!] \left[-\frac{2}{\Lambda} \varepsilon^{\mu\nu\rho\sigma} p_{1\mu} p_{2\rho} \right] \quad (\text{A.8})$$

The matrix element and its square are given by,

$$\begin{aligned} i\mathcal{M} &= -\frac{4}{\Lambda} i \varepsilon^{\mu\nu\rho\sigma} p_{1\mu} p_{2\rho} \epsilon_\nu^*(p_1) \epsilon_\sigma^*(p_2) \\ |\mathcal{M}|^2 &= \frac{16}{\Lambda^2} \varepsilon^{\mu\nu\rho\sigma} \varepsilon^{\alpha\beta\gamma\delta} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} \epsilon_\nu^*(p_1) \epsilon_\sigma^*(p_2) \epsilon_\beta(p_1) \epsilon_\delta(p_2) \end{aligned} \quad (\text{A.9})$$

Summing over the polarisations we get,

$$\begin{aligned} \sum |\mathcal{M}|^2 &= \frac{16}{\Lambda^2} \varepsilon^{\mu\nu\rho\sigma} \varepsilon^{\alpha\beta\gamma\delta} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} g_{\nu\beta} g_{\sigma\delta} \\ &= \frac{16}{\Lambda^2} \varepsilon_\beta^{\mu\rho\sigma} \varepsilon_\sigma^{\beta\alpha\gamma} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} g_{\nu\beta} g_{\sigma\delta} \\ &= \frac{32}{\Lambda^2} (-g^{\mu\alpha} g^{\rho\gamma} + g^{\mu\gamma} g^{\rho\alpha}) p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} \\ &= \frac{32}{\Lambda^2} (p_1 \cdot p_2)^2 \\ &= \frac{32}{\Lambda^2} \left(\frac{m_a^2}{2} \right)^2 \\ &= \frac{8m_a^4}{\Lambda^2} \end{aligned} \quad (\text{A.10})$$

Hence, finally we get

$$\Gamma(a \rightarrow \gamma\gamma) = \frac{1}{\Lambda^2} \left[\frac{1}{2m_a} \right] \left[\frac{1}{8\pi} \right] [8m_a^4] \left[\frac{1}{2} \right]. \quad (\text{A.11})$$

A.3 $\phi \rightarrow ZZ$

The relevant part of the Lagrangian is

$$\mathcal{L} \subset \frac{1}{\Lambda} \phi Z_{\mu\nu} Z^{\mu\nu}. \quad (\text{A.12})$$

The Feynman rule is same as the decay $\phi \rightarrow \gamma\gamma$ (eq. (A.2)).

The squared matrix element is given by,

$$|\mathcal{M}|^2 = \frac{16}{\Lambda^2} (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) (p_1 \cdot p_2 g_{\alpha\beta} - p_{1\alpha} p_{2\beta}) \epsilon^{*\nu}(p_1) \epsilon^{*\mu}(p_2) \epsilon^\beta(p_1) \epsilon^\alpha(p_2) \quad (\text{A.13})$$

Summation over the polarization vectors,

$$\sum \epsilon^\mu(p) \epsilon^{*\nu}(p) = -g^{\mu\nu} + \frac{p^\mu p^\nu}{m_Z^2}$$

we get,

$$\begin{aligned} |\mathcal{M}|^2 &= 16 (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) (p_1 \cdot p_2 g_{\alpha\beta} - p_{1\alpha} p_{2\beta}) \times \\ &\quad \times \left(-g^{\mu\alpha} + \frac{p_2^\mu p_2^\alpha}{m_Z^2} \right) \left(-g^{\nu\beta} + \frac{p_1^\nu p_1^\beta}{m_Z^2} \right) \\ &= 16 (p_1 \cdot p_2 g_{\mu\nu} - p_{1\mu} p_{2\nu}) (p_1 \cdot p_2 g_{\alpha\beta} - p_{1\alpha} p_{2\beta}) (-g^{\mu\alpha}) (-g^{\nu\beta}) \\ &= 16 (2(p_1 \cdot p_2)^2 + p_1^2 p_2^2) \\ &= 8m_\phi^4 \left(1 - 4 \frac{m_Z^2}{m_\phi^2} + 6 \frac{m_Z^4}{m_\phi^4} \right) \end{aligned} \quad (\text{A.14})$$

$$\Gamma(\phi \rightarrow Z Z) = \left[\frac{1}{2m_\phi} \right] \left[\frac{\lambda^{1/2}(1, m_Z^2/m_\phi^2, m_Z^2/m_\phi^2)}{8\pi} \right] |\mathcal{M}|^2 \left[\frac{1}{2} \right] \quad (\text{A.15})$$

where,

$$\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2ac - 2bc \quad (\text{A.16})$$

$$\Gamma(\phi \rightarrow Z Z) = \frac{1}{\Lambda^2} \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \left(1 - 4 \frac{m_Z^2}{m_\phi^2} \right)^{1/2} \right] |\mathcal{M}|^2 \left[\frac{1}{2} \right] \quad (\text{A.17})$$

$$\begin{aligned} &= \frac{1}{\Lambda^2} \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \left(1 - \frac{4m_Z^2}{m_\phi^2} \right)^{1/2} \right] \\ &\quad \times \left[8m_\phi^4 \left(1 - 4 \frac{m_Z^2}{m_\phi^2} + 6 \frac{m_Z^4}{m_\phi^4} \right) \right] \left[\frac{1}{2} \right] \end{aligned} \quad (\text{A.18})$$

A.4 $a \rightarrow Z Z$

The relevant part of the Lagrangian is

$$\mathcal{L} \subset \frac{1}{\Lambda} a Z_{\mu\nu} \tilde{Z}^{\mu\nu}. \quad (\text{A.19})$$

The Feynman rule is same as the decay $a \rightarrow \gamma\gamma$ (eq. (A.8)).

The squared matrix element is given by,

$$|\mathcal{M}|^2 = \frac{16}{\Lambda^2} \varepsilon^{\mu\nu\rho\sigma} \varepsilon^{\alpha\beta\gamma\delta} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} \epsilon_\nu^*(p_1) \epsilon_\beta(p_1) \epsilon_\sigma^*(p_2) \epsilon_\delta(p_2) \quad (\text{A.20})$$

Summing over the polarisations, we have,

$$|\mathcal{M}|^2 = \frac{16}{\Lambda^2} \varepsilon^{\mu\nu\rho\sigma} \varepsilon^{\alpha\beta\gamma\delta} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} \left(-g_{\nu\beta} + \frac{p_{1\nu} p_{1\beta}}{M_Z^2} \right) \left(-g_{\sigma\delta} + \frac{p_{2\sigma} p_{2\delta}}{M_Z^2} \right) \quad (\text{A.21})$$

$$= \frac{16}{\Lambda^2} \varepsilon^{\mu\nu\rho\sigma} \varepsilon^{\alpha\beta\gamma\delta} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} g_{\nu\beta} g_{\sigma\delta} \quad (\text{A.22})$$

where we have used the fact that the second terms in each of the parenthesis vanish due to the antisymmetry of the Levi-civita symbols. We thus have,

$$|\mathcal{M}|^2 = \frac{16}{\Lambda^2} \varepsilon_\beta^{\mu\rho\sigma} \varepsilon_\sigma^{\beta\alpha\gamma} p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} \quad (\text{A.23})$$

Using the relation,

$$\varepsilon_\beta^{\mu\rho\sigma} \varepsilon_\sigma^{\beta\alpha\gamma} = 2(-g^{\mu\alpha} g^{\rho\gamma} + g^{\mu\gamma} g^{\rho\alpha}) \quad (\text{A.24})$$

we get,

$$|\mathcal{M}|^2 = \frac{32}{\Lambda^2} (-g^{\mu\alpha} g^{\rho\gamma} + g^{\mu\gamma} g^{\rho\alpha}) p_{1\mu} p_{1\alpha} p_{2\rho} p_{2\gamma} \quad (\text{A.25})$$

$$= \frac{32}{\Lambda^2} (-p_1^2 p_2^2 + (p_1 \cdot p_2)^2) \quad (\text{A.26})$$

$$= \frac{32}{\Lambda^2} \left(-m_Z^4 + \frac{(m_a^2 - 2m_Z^2)^2}{4} \right) \quad (\text{A.27})$$

$$= \frac{8m_a^4}{\Lambda^2} \left(1 - \frac{4m_Z^2}{m_a^2} \right) \quad (\text{A.28})$$

$$\Gamma(a \rightarrow Z Z) = \left[\frac{1}{2m_a} \right] \left[\frac{\lambda^{1/2}(1, m_Z^2/m_a^2, m_Z^2/m_a^2)}{8\pi} \right] |\mathcal{M}|^2 \left[\frac{1}{2} \right] \quad (\text{A.29})$$

$$= \frac{1}{\Lambda^2} \left[\frac{1}{2m_a} \right] \left[\frac{1}{8\pi} \left(1 - 4 \frac{m_Z^2}{m_a^2} \right)^{1/2} \right] |\mathcal{M}|^2 \left[\frac{1}{2} \right] \quad (\text{A.30})$$

$$= \frac{1}{\Lambda^2} \left[\frac{1}{2m_a} \right] \left[\frac{1}{8\pi} \left(1 - \frac{4m_Z^2}{m_a^2} \right)^{1/2} \right] \times \left[8m_a^4 \left(1 - 4 \frac{m_Z^2}{m_a^2} \right) \right] \left[\frac{1}{2} \right] \quad (\text{A.31})$$

B Calculation of $\mathcal{A}_{\text{LHC energy}}^{ii}$

In this appendix we will calculate the quantities $\mathcal{A}_{\text{LHC energy}}^{ii}$ defined in section 2.2 for two LHC energies 8 TeV and 13 TeV, and for the initial states $\{gg\}$, $\{\bar{u}u\}$ and $\{\bar{d}d\}$.

B.1 Production by gluon fusion

The partonic cross section for the process $g(p) g(k) \rightarrow \phi(q)$ is given by

$$\hat{\sigma}(g(p) g(k) \rightarrow \phi(q)) \quad (\text{B.1})$$

$$= \frac{1}{2^2} \frac{1}{8^2} \frac{1}{2E_p} \frac{1}{2E_k} \frac{1}{|v_p - v_k|} \int \frac{d^3q}{(2\pi)^3} \frac{1}{2E_q} |\mathcal{M}|^2 (2\pi)^4 \delta^{(4)}(p + k - q) \quad (\text{B.2})$$

$$\Rightarrow \text{using the identity } \int d^4q \delta(q^2 - m_\phi^2) \Theta(q^0) = \frac{1}{2E_q}, \text{ we get} \quad (\text{B.3})$$

$$= \frac{1}{2^2} \frac{1}{8^2} \frac{2\pi}{2E_p} \frac{1}{2E_k} \frac{1}{|v_p - v_k|} \int d^4q \delta(q^2 - m_\phi^2) \Theta(q^0) |\mathcal{M}|^2 \delta^{(4)}(p + k - q) \quad (\text{B.4})$$

$$= \frac{1}{2^2} \frac{1}{8^2} \frac{2\pi}{2E_p} \frac{1}{2E_k} \frac{1}{|v_p - v_k|} |\mathcal{M}|^2 \delta((p + k)^2 - m_\phi^2) \quad (\text{B.5})$$

$$= \frac{1}{2^2} \frac{1}{8^2} \frac{2\pi}{x_1 x_2 S} \frac{1}{2} |\mathcal{M}|^2 \delta(x_1 x_2 S - m_\phi^2) \quad (\text{B.6})$$

$$= \frac{\pi}{256} \frac{1}{x_1 x_2 S} |\mathcal{M}|^2 \delta(x_1 x_2 S - m_\phi^2) \quad (\text{B.7})$$

where the following definitions have been used,

$$p = x_1 P_1, \quad k = x_2 P_2, \quad P_1 = \frac{\sqrt{S}}{2}(1, 0, 0, 1) \quad \text{and} \quad P_2 = \frac{\sqrt{S}}{2}(1, 0, 0, -1).$$

Here, P_1 and P_2 are the 4-momenta of the two protons and \sqrt{S} is their centre-of-mass energy.

We now proceed to compute the hadronic cross section which is given by

$$\sigma_{\sqrt{S}} = \int_0^1 dx_1 \int_0^1 dx_2 f_{g/p}(x_1) f_{g/p}(x_2) \hat{\sigma}(x_1, x_2) \quad (\text{B.8})$$

\Rightarrow using the change of variables $\{x_1, x_2\} \rightarrow \{x = x_1, z = x_1 x_2\}$, we get

$$= \int_0^1 \frac{dx}{x} \int_0^x dz f_{g/p}(x) f_{g/p}(z/x) \hat{\sigma}(z) \quad (\text{B.9})$$

$$= \int_0^1 \frac{dx}{x} \int_0^x dz f_{g/p}(x) f_{g/p}(z/x) \times \frac{\pi}{256} \frac{1}{z S} |\mathcal{M}|^2 \delta(z S - m_\phi^2) \quad (\text{B.10})$$

We now use the expression for $\Gamma_{\phi \rightarrow gg}$ (following appendix A),

$$\Gamma(\phi \rightarrow gg) = \left[\frac{1}{2m_\phi} \right] \left[\frac{1}{8\pi} \right] |\mathcal{M}|^2 \left[\frac{1}{2} \right], \quad (\text{B.11})$$

to get

$$\sigma_{\sqrt{S}} = \frac{\pi}{256} \frac{32\pi m_\phi \Gamma_{\phi \rightarrow gg}}{S} \int_0^1 \frac{dx}{x} \int_0^x dz f_{g/p}(x) f_{g/p}(z/x) \frac{1}{z} \delta(z S - m_\phi^2) \quad (\text{B.12})$$

$$= \frac{\pi}{256} \frac{32\pi m_\phi \Gamma_{\phi \rightarrow gg}}{S^2} \int_0^1 \frac{dx}{x} \int_0^x dz f_{g/p}(x) f_{g/p}(z/x) \frac{1}{z} \delta\left(z - \frac{m_\phi^2}{S}\right) \quad (\text{B.13})$$

$$= \frac{\pi}{256} \frac{32\pi m_\phi \Gamma_{\phi \rightarrow gg}}{S^2} \int_{\frac{m_\phi^2}{S}}^1 \frac{dx}{x} f_{g/p}(x) f_{g/p}\left(\frac{m_\phi^2}{Sx}\right) \frac{S}{m_\phi^2} \quad (\text{B.14})$$

$$= \frac{\pi^2}{8} \frac{\Gamma_{\phi \rightarrow gg}}{m_\phi S} \int_{\frac{m_\phi^2}{S}}^1 \frac{dx}{x} f_{g/p}(x) f_{g/p}\left(\frac{m_\phi^2}{Sx}\right) \quad (\text{B.15})$$

Hence,

$$\mathcal{A}_{\text{LHC energy}}^{gg} = \frac{\pi^2}{8} \frac{1}{m_\phi S} \int_{\frac{m_\phi^2}{S}}^1 \frac{dx}{x} f_{g/p}(x) f_{g/p}\left(\frac{m_\phi^2}{Sx}\right) \quad (\text{B.16})$$

Using the MSTW 2008 LO parton distribution functions (PDF) we get,

$$\mathcal{A}_{13 \text{ TeV}}^{gg} = \frac{5.44 \text{ pb}}{\text{GeV}} \quad (\text{B.17})$$

$$\mathcal{A}_{8 \text{ TeV}}^{gg} = \frac{1.15 \text{ pb}}{\text{GeV}}. \quad (\text{B.18})$$

B.2 Production by quarks

The cross section of the process $\bar{q}q \rightarrow \phi$ can be calculated in the same way as above, except for the following changes,

- The colour factor is different, so we must have $1/3^2$ instead of $1/8^2$ as in the case for gluons
- The symmetry factor (1/2) for identical particle used in eq. (B.11) no longer applies
- The PDF are different — we now have quark PDF instead of the gluon PDF.

Applying the above changes, we finally get,

$$\sigma_{\sqrt{S}} = \frac{4\pi^2}{9} \frac{\Gamma_{\phi \rightarrow q\bar{q}}}{m_\phi S} \int_{\frac{m_\phi^2}{S}}^1 \frac{dx}{x} (f_{q/p}(x)f_{\bar{q}/p}(m_\phi^2/Sx) + f_{\bar{q}/p}(x)f_{q/p}(m_\phi^2/Sx))$$

Hence,

$$\mathcal{A}_{\text{LHC energy}}^{q\bar{q}} = \frac{4\pi^2}{9} \frac{1}{m_\phi S} \int_{\frac{m_\phi^2}{S}}^1 \frac{dx}{x} (f_{q/p}(x)f_{\bar{q}/p}(m_\phi^2/Sx) + f_{\bar{q}/p}(x)f_{q/p}(m_\phi^2/Sx))$$

Using again the MSTW 2008 LO parton distribution functions (PDF) we get,

$$\mathcal{A}_{13}^{\bar{u}u} \equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{u}u} = \frac{2.94 \text{ pb}}{\text{GeV}} \quad \mathcal{A}_8^{\bar{u}u} \equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{u}u} = \frac{1.2 \text{ pb}}{\text{GeV}} \quad (\text{B.19})$$

$$\mathcal{A}_{13}^{\bar{d}d} \equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{d}d} = \frac{1.73 \text{ pb}}{\text{GeV}} \quad \mathcal{A}_8^{\bar{d}d} \equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{d}d} = \frac{0.66 \text{ pb}}{\text{GeV}} \quad (\text{B.20})$$

$$\mathcal{A}_{13}^{\bar{c}c} \equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{c}c} = \frac{0.11 \text{ pb}}{\text{GeV}} \quad \mathcal{A}_8^{\bar{c}c} \equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{c}c} = \frac{0.03 \text{ pb}}{\text{GeV}} \quad (\text{B.21})$$

$$\mathcal{A}_{13}^{\bar{s}s} \equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{s}s} = \frac{0.21 \text{ pb}}{\text{GeV}} \quad \mathcal{A}_8^{\bar{s}s} \equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{s}s} = \frac{0.05 \text{ pb}}{\text{GeV}} \quad (\text{B.22})$$

$$\mathcal{A}_{13}^{\bar{b}b} \equiv \mathcal{A}_{13 \text{ TeV LHC}}^{\bar{b}b} = \frac{0.05 \text{ pb}}{\text{GeV}} \quad \mathcal{A}_8^{\bar{b}b} \equiv \mathcal{A}_{8 \text{ TeV LHC}}^{\bar{b}b} = \frac{0.01 \text{ pb}}{\text{GeV}} \quad (\text{B.23})$$

C Calculation of the sgoldstino mass

In this appendix, we want to compute the 1-loop contribution to the sgoldstino mass from the term,

$$\mathcal{L} \subset \int d^2\theta \lambda X \Phi_1 \Phi_2 + \text{h.c.} \quad (\text{C.1})$$

We will ignore the gauge indices of Φ_1 and Φ_2 for the time being. The following notation will be used for the chiral superfields:

$$X = \mathcal{S} + \sqrt{2}\theta \psi_x + \theta\theta F_x \quad (\text{C.2})$$

$$\Phi_1 = \phi_1 + \sqrt{2}\theta \xi_1 + \theta\theta F_1 \quad (\text{C.3})$$

$$\Phi_2 = \phi_2 + \sqrt{2}\theta \xi_2 + \theta\theta F_2 \quad (\text{C.4})$$




Figure 7. One loop contributions to the sgoldstino mass from the messengers.

A Dirac fermion Ψ is constructed out of the two Weyl fermions ξ_1 and ξ_2 ,

$$\Psi = \begin{pmatrix} \xi_{1\alpha} \\ \xi_2^{\dagger\dot{\alpha}} \end{pmatrix} \quad (\text{C.5})$$

whose Dirac mass will be denoted by $m_\Psi = \lambda \langle S \rangle$. The scalar mass eigenstates will be denoted by ϕ_+ and ϕ_- with their mass squared given by $m_\pm^2 = m_\Psi^2 \pm \lambda \langle F_x \rangle$.

C.1 Diagrammatic calculation

The relevant vertex factors are given by,

$$\mathcal{S} \bar{\Psi} \Psi : -\lambda P_L \quad (\text{C.6})$$

$$\mathcal{S}^* \mathcal{S} \phi_+^* \phi_+ : -\lambda^2 \quad (\text{C.7})$$

$$\mathcal{S}^* \mathcal{S} \phi_-^* \phi_- : -\lambda^2 \quad (\text{C.8})$$

$$\mathcal{S} \phi_+^* \phi_+ : -\lambda m_\Psi \quad (\text{C.9})$$

$$\mathcal{S} \phi_-^* \phi_- : -\lambda m_\Psi \quad (\text{C.10})$$

The Feynman rules can be obtained by multiplying the above vertex factors by i and appropriate symmetry factors.

The relevant diagrams are,

We will now compute the diagrams one-by-one.

Fermion loop:

$$\begin{aligned} -i \Pi(p^2 = 0) &= -(-i\lambda)(-i\lambda) \int \frac{d^4 q}{(2\pi)^4} \text{Tr} \left[P_L \frac{i}{\not{q} - m_\Psi} P_R \frac{i}{\not{q} - m_\Psi} \right] \\ &= -2\lambda^2 \int \frac{d^4 q}{(2\pi)^4} \frac{q^2}{(q^2 - m_\Psi^2)^2} \\ &= -2\lambda^2 \int \frac{d^4 q}{2\pi^4} \left[\frac{1}{q^2 - m_\Psi^2} + \frac{m_\Psi^2}{(q^2 - m_\Psi^2)^2} \right] \end{aligned} \quad (\text{C.11})$$

First scalar loop:

$$\begin{aligned} -i \Pi(p^2 = 0) &= (-i\lambda^2) \sum_{\phi=\phi_\pm} \int \frac{d^4 q}{(2\pi)^4} \frac{i}{q^2 - m_\phi^2} \\ &= \lambda^2 \int \frac{d^4 q}{(2\pi)^4} \left[\frac{1}{q^2 - m_+^2} + \frac{1}{q^2 - m_-^2} \right] \end{aligned} \quad (\text{C.12})$$

Second scalar loop:

$$-i \Pi(p^2 = 0) = (-i\lambda m_\Psi)(-i\lambda m_\Psi) \sum_{\phi=\phi_\pm} \int \frac{d^4 q}{(2\pi)^4} \frac{i^2}{(q^2 - m_\phi^2)^2} \quad (\text{C.13})$$

$$= \lambda^2 m_\Psi^2 \int \frac{d^4 q}{(2\pi)^4} \left[\frac{1}{(q^2 - m_+^2)^2} + \frac{1}{(q^2 - m_-^2)^2} \right] \quad (\text{C.14})$$

Note that the sum of the diagrams goes to zero in the limit of equal masses for the scalars and fermions, i.e. when SUSY is unbroken.

We need to evaluate integrals of two the forms:

$$A_0(m) = \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2 - m^2} \quad ; \quad B_0(0, m, m) = \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - m^2)^2} \quad (\text{C.15})$$

They are given by,

$$A_0(m) = \frac{i}{16\pi^2} m^2 \left[\frac{1}{\hat{\epsilon}} + 1 - \text{Ln} \frac{m^2}{\mu^2} \right] \quad (\text{C.16})$$

$$B_0(0, m, m) = \frac{A_0(m)}{m^2} - \frac{i}{16\pi^2} \quad (\text{C.17})$$

where,

$$\frac{1}{\hat{\epsilon}} = \frac{2}{4-D} - \gamma_E + \text{Ln}(4\pi), \quad \gamma_E \text{ being the Euler constant.} \quad (\text{C.18})$$

Putting all loop contributions in order, we have

$$-i \Pi(p^2 = 0) = -2\lambda^2 \int \frac{d^4 q}{(2\pi)^4} \left[\frac{1}{q^2 - m_\Psi^2} - \frac{1}{2} \frac{1}{q^2 - m_+^2} - \frac{1}{2} \frac{1}{q^2 - m_-^2} \right. \\ \left. + m_\Psi^2 \frac{1}{(q^2 - m_\Psi^2)^2} - \frac{m_+^2}{2} \frac{1}{(q^2 - m_+^2)^2} - \frac{m_-^2}{2} \frac{1}{(q^2 - m_-^2)^2} \right] \quad (\text{C.19})$$

$$= -2\lambda^2 \frac{i}{16\pi^2} \left[m_\Psi^2 \left(\frac{1}{\hat{\epsilon}} + 1 - \text{Ln} \frac{m_\Psi^2}{\mu^2} \right) - \frac{m_+^2}{2} \left(\frac{1}{\hat{\epsilon}} + 1 - \text{Ln} \frac{m_+^2}{\mu^2} \right) \right. \\ \left. - \frac{m_-^2}{2} \left(\frac{1}{\hat{\epsilon}} + 1 - \text{Ln} \frac{m_-^2}{\mu^2} \right) + m_\Psi^2 \left(\frac{1}{\hat{\epsilon}} - \text{Ln} \frac{m_\Psi^2}{\mu^2} \right) \right. \\ \left. - \frac{m_+^2}{2} \left(\frac{1}{\hat{\epsilon}} - \text{Ln} \frac{m_+^2}{\mu^2} \right) - \frac{m_-^2}{2} \left(\frac{1}{\hat{\epsilon}} - \text{Ln} \frac{m_-^2}{\mu^2} \right) \right] \quad (\text{C.20})$$

$$= -2\lambda^2 \frac{i}{16\pi^2} \left[\frac{m_+^2}{2} \text{Ln} \frac{m_+^2}{\mu^2} + \frac{m_-^2}{2} \text{Ln} \frac{m_-^2}{\mu^2} - m_\Psi^2 \text{Ln} \frac{m_\Psi^2}{\mu^2} \right. \\ \left. + \frac{m_\Psi^2}{2} \text{Ln} \frac{m_+^2}{\mu^2} + \frac{m_\Psi^2}{2} \text{Ln} \frac{m_-^2}{\mu^2} - m_\Psi^2 \text{Ln} \frac{m_\Psi^2}{\mu^2} \right] \quad (\text{C.21})$$

$$= -2\lambda^2 \frac{i}{16\pi^2} \left[m_\Psi^2 \text{Ln} \frac{m_+ m_-}{m_\Psi^2} + \lambda \langle F_x \rangle \text{Ln} \frac{m_+}{m_-} + m_\Psi^2 \text{Ln} \frac{m_+ m_-}{m_\Psi^2} \right] \quad (\text{C.22})$$




Figure 8. The behaviour of $F(x)$ as a function of x .

Hence, assuming Φ_1 (Φ_2) to be a $\mathbf{5}$ ($\bar{\mathbf{5}}$) of $SU(5)$, and for N_m pairs of $\{\Phi_1, \Phi_2\}$, we have,

$$\Pi(p^2 = 0) = 5 N_m \frac{2 \lambda^2}{16 \pi^2} \left[2 m_\Psi^2 \text{Ln} \frac{m_+ m_-}{m_\Psi^2} + \lambda \langle F_x \rangle \text{Ln} \frac{m_+}{m_-} \right] \quad (\text{C.23})$$

$$= 5 N_m \frac{\lambda^2}{16 \pi^2} m_\Psi^2 \left[2 \text{Ln} \frac{m_+^2 m_-^2}{m_\Psi^4} + \frac{\lambda \langle F_x \rangle}{m_\Psi^2} \text{Ln} \frac{m_+^2}{m_-^2} \right] \quad (\text{C.24})$$

$$= \left(4 \pi \sqrt{5} N_m \right)^2 \left(\frac{\lambda}{16 \pi^2} \right)^2 \frac{\lambda^2 \langle F_x \rangle^2}{m_\Psi^2} G \left(\frac{\lambda \langle F_x \rangle}{m_\Psi^2} \right) \quad (\text{C.25})$$

where the function $G(x)$ is given by,

$$G(x) = \frac{1}{x^2} [(2+x) \text{Log}(1+x) + (2-x) \text{Log}(1-x)]. \quad (\text{C.26})$$

In terms of gaugino mass, this can be written as,

$$\Pi(p^2 = 0) = - \left(\frac{\lambda}{g_a^2} \right)^2 \left(4 \pi \sqrt{\frac{5}{N_m}} F(x) \right)^2 m_a^2 \quad (\text{C.27})$$

The behaviour of the function $F(x) \equiv \sqrt{-G(x)/g(x)^2}$ is shown in figure 8.

C.2 Coleman-Weinberg potential

The Dirac mass for the fermions as a function of \mathcal{S} is given by,

$$m_F(\mathcal{S}) = \lambda \mathcal{S} \quad (\text{C.28})$$

and the scalar mass matrix is

$$\tilde{m}^2(\mathcal{S}) = \begin{pmatrix} \lambda^2 \mathcal{S}^* \mathcal{S} & -\lambda \langle F_x \rangle \\ -\lambda \langle F_x \rangle & \lambda^2 \mathcal{S}^* \mathcal{S} \end{pmatrix}, \quad (\text{C.29})$$

with the eigenvalues,

$$m_\pm^2(\mathcal{S}^* \mathcal{S}) = \lambda^2 \mathcal{S}^* \mathcal{S} \pm \lambda \langle F_x \rangle. \quad (\text{C.30})$$

Using the standard formula for the Coleman-Weinberg potential [55],

$$V_{\text{CW}} = \frac{1}{64\pi^2} \text{STr} \left(\mathcal{M}^4 \left[\log \frac{\mathcal{M}^2}{\Lambda_{\text{cut-off}}^2} - \frac{3}{2} \right] \right), \quad (\text{C.31})$$

we get,

$$\begin{aligned} V_{\text{CW}} = \frac{2}{64\pi^2} & \left[[m_+^2(\mathcal{S}, \mathcal{S}^*)]^2 \text{Ln} [m_+^2(\mathcal{S}, \mathcal{S}^*)] + [m_-^2(\mathcal{S}, \mathcal{S}^*)]^2 \text{Ln} [m_-^2(\mathcal{S}, \mathcal{S}^*)] \right. \\ & - 2 [m_F(\mathcal{S})^* m_F(\mathcal{S})]^2 \text{Ln} [m_F(\mathcal{S})^* m_F(\mathcal{S})] \\ & \left. - \lambda^2 \langle F_x \rangle^2 \left(\log \Lambda_{\text{cut-off}}^2 + \frac{3}{2} \right) \right] \end{aligned} \quad (\text{C.32})$$

After replacing $\mathcal{S} \rightarrow \langle \mathcal{S} \rangle + \mathcal{S}$, we get the coefficient of $\mathcal{S}^* \mathcal{S}$ to be,

$$\Pi(p^2 = 0) = \frac{2\lambda^2}{16\pi^2} \left[2m_\Psi^2 \text{Ln} \frac{m_+ m_-}{m_\Psi^2} + \lambda \langle F_x \rangle \text{Ln} \frac{m_+}{m_-} \right] \quad (\text{C.33})$$

C.3 Tree level goldstino mass

Here we give an example of a model where the goldstino gets tree level mass at the time of SUSY breaking [34]. The model is just an extension of the Affleck-Dine-Seiberg model (ADS) or 3-2 model of [43–45]. The field content of the ADS model is

	$SU(3)$	$SU(2)$
Q	3	2
U^c	$\bar{3}$	1
D^c	$\bar{3}$	1
L	1	$\bar{2}$

and the superpotential is given by,

$$W_{3-2} = W_{\text{cl}} + W_{\text{np}} \quad (\text{C.34})$$

$$\text{where, } W_{\text{cl}} = h Q_A^a D_a^c L^A, \quad (\text{C.35})$$

$$W_{\text{np}} = \frac{\Lambda_3^7}{\det(QQ^c)}, \quad (\text{C.36})$$

where, Q^c is defined as $Q^c \equiv (U^c, D^c)$. In this model $h \ll \tilde{g}_2, \tilde{g}_3$ which are the gauge couplings for the groups $SU(2)$ and $SU(3)$ respectively. Thus, F -term contribution to the scalar potential is subdominant compared to the D -term contribution. The minimum of the potential can be obtained perturbatively along the D -flat directions,

$$Q = \begin{pmatrix} a & 0 \\ 0 & b \\ 0 & 0 \end{pmatrix} M, \quad Q^c = \begin{pmatrix} a & 0 \\ 0 & b \\ 0 & 0 \end{pmatrix} M, \quad L = \begin{pmatrix} \sqrt{a^2 - b^2} \\ 0 \end{pmatrix} M \quad (\text{C.37})$$

where,

$$M \equiv \frac{\Lambda_3}{h^{1/7}} \gg \Lambda_3, \quad (\text{C.38})$$

and $a \approx 1.164$, $b \approx 1.132$.

Note that L_1 (the component of L getting a non-vanishing VEV) is the sgoldstino here. The $SU(2)$ D -term equation of motion gives,

$$D_2^a = \tilde{g}_2 \sum_f f^\dagger T_2^a f \tag{C.39}$$

where $T_2^a = \sigma^a/2$, σ^a being the Pauli matrices. The eq. (C.39) will get contributions from all the fields carrying $SU(2)$ charge i.e., Q and L ,

$$D^a = \tilde{g}_2 \left(L^\dagger \frac{\sigma^a}{2} L + \sum_r Q_i^{r\dagger} \frac{\sigma^a}{2} Q_i^r \right) \tag{C.40}$$

where the index r is the $SU(3)$ index. This gives, for the scalar potential,

$$V = \frac{1}{2} D^a D_a \tag{C.41}$$

$$= \frac{\tilde{g}_2^2}{8} \left(L^\dagger \frac{\sigma^a}{2} L + \sum_r Q_i^{r\dagger} \frac{\sigma^a}{2} Q_i^r \right) \left(L^\dagger \frac{\sigma_a}{2} L + \sum_r Q_i^{r\dagger} \frac{\sigma_a}{2} Q_i^r \right). \tag{C.42}$$

Noting that only the third Pauli matrix contributes, we have,

$$V = \frac{\tilde{g}_2^2}{8} \left[(L_1^\dagger L_1)^2 + 2(L_1^\dagger L_1)(Q_1^{r\dagger} Q_1^r - Q_2^{r\dagger} Q_2^r) + \dots \right], \tag{C.43}$$

where the ellipses denote terms unimportant for sgoldstino mass. This generates a mass term for L_1 which is given by,

$$M_{L_1}^2 = \frac{\tilde{g}_2^2}{8} (4(a^2 - b^2)M^2 + 2(a^2 - b^2)M^2) \tag{C.44}$$

$$= \frac{3\tilde{g}_2^2}{4} (a^2 - b^2)M^2. \tag{C.45}$$

This is, in general, much larger than the gaugino mass.

D Calculation of the gaugino mass

The relevant part of the Lagrangian is given by

$$\mathcal{L} \subset \int d^4\theta \Phi_1^\dagger e^{2gT^a V^a} \Phi_1 + \int d^4\theta \Phi_2^\dagger e^{2gT^a V^a} \Phi_2 + \left(\int d^2\theta y X \Phi_1 \Phi_2 + \text{h.c.} \right) \tag{D.1}$$

where,

$$V^a = \theta \bar{\sigma}^\mu \bar{\theta} A_\mu^a + i\theta^2 \bar{\theta} \lambda^{\dagger a} - i\theta \bar{\theta}^2 \lambda^a + \frac{1}{2} \theta^2 \bar{\theta}^2 D^a \tag{D.2}$$

A Majorana fermion Ψ_λ is constructed out of the (Weyl) gaugino field λ_a ,

$$\Psi_\lambda = \begin{pmatrix} \lambda_\alpha \\ \lambda^{\dagger\dot{\alpha}} \end{pmatrix} \tag{D.3}$$

The relevant vertex factors are given by,

$$\alpha \bar{\Psi}_\xi \Psi_\lambda^A : -igT^A \tag{D.4}$$

$$\beta \bar{\Psi}_\xi \Psi_\lambda^A : +ig\gamma_5 T^A \tag{D.5}$$

The gaugino mass is generated via the one loop diagrams shown in figure 9.




Figure 9. One-loop contribution to the gaugino mass.

Loop with the scalar α :

$$-\frac{i}{2}m_{\Psi_\lambda}^{(\alpha)AB} = \text{Tr}[T^A T^B] \int \frac{d^4 q}{(2\pi)^4} (-g) \frac{-i}{\not{q} + m_\Psi} (+g) \frac{i}{q^2 - m_\alpha^2} \quad (\text{D.6})$$

$$= -g^2 \text{Tr}[T^A T^B] \int \frac{d^4 q}{(2\pi)^4} \frac{\not{q} - m_\Psi}{q^2 - m_\Psi^2} \frac{1}{q^2 - m_\alpha^2} \quad (\text{D.7})$$

$$= g^2 m_\Psi \text{Tr}[T^A T^B] \int \frac{d^4 q}{(2\pi)^4} \frac{1}{q^2 - m_\Psi^2} \frac{1}{q^2 - m_\alpha^2} \quad (\text{D.8})$$

$$= g^2 m_\Psi \text{Tr}[T^A T^B] B_0(0, m_\Psi, m_\alpha) \quad (\text{D.9})$$

Loop with the scalar β :

$$-\frac{i}{2}m_{\Psi_\lambda}^{(\beta)AB} = \text{Tr}[T^A T^B] \int \frac{d^4 q}{(2\pi)^4} (-g\gamma_5) \frac{-i}{\not{q} + m_\Psi} (-g\gamma_5) \frac{i}{q^2 - m_\beta^2} \quad (\text{D.10})$$

$$= g^2 \text{Tr}[T^A T^B] \int \frac{d^4 q}{(2\pi)^4} \gamma_5 \frac{\not{q} - m_\Psi}{q^2 - m_\Psi^2} \gamma_5 \frac{1}{q^2 - m_\beta^2} \quad (\text{D.11})$$

$$= g^2 \text{Tr}[T^A T^B] \int \frac{d^4 q}{(2\pi)^4} \frac{(-\not{q} - m_\Psi)}{q^2 - m_\Psi^2} \frac{1}{q^2 - m_\beta^2} \quad (\text{D.12})$$

$$= -g^2 m_\Psi \text{Tr}[T^A T^B] B_0(0, m_\Psi, m_\beta) \quad (\text{D.13})$$

where, the B_0 function is given by,

$$B_0(0, m_1, m_2) = \frac{A_0(m_1) - A_0(m_2)}{m_1^2 - m_2^2} \quad (\text{D.14})$$

$$-\frac{i}{2}m_{\Psi_\lambda}^{AB} = g^2 m_\Psi \text{Tr}[T^A T^B] (B_0(0, m_\Psi, m_\alpha) - B_0(0, m_\Psi, m_\beta)) \quad (\text{D.15})$$

$$= -i \frac{g^2 m_\Psi}{16\pi^2} \text{Tr}[T^A T^B] \frac{(1+x) \ln(1+x) + (1-x) \ln(1-x)}{x} \quad (\text{D.16})$$

$$= -i \frac{g^2}{16\pi^2} \text{Tr}[T^A T^B] \frac{\langle F_x \rangle}{m_\Psi} \frac{(1+x) \ln(1+x) + (1-x) \ln(1-x)}{x^2} \quad (\text{D.17})$$

$$m_{\Psi_\lambda}^{AB} = \frac{g^2}{16\pi^2} 2\text{Tr}[T^A T^B] \frac{\langle F_x \rangle}{m_\Psi} g(x) \quad (\text{D.18})$$

$$m_{\Psi_\lambda}^{AB} = \frac{g^2}{16\pi^2} \frac{\langle F_x \rangle}{m_\Psi} g(x) \delta^{AB}. \quad (\text{D.19})$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, *Search for resonances decaying to photon pairs in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*, [ATLAS-CONF-2015-081](#) (2015).
- [2] CMS Collaboration, *Search for new physics in high mass diphoton events in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$* , [CMS-PAS-EXO-15-004](#) (2015).
- [3] A. Falkowski, O. Slone and T. Volansky, *Phenomenology of a 750 GeV singlet*, [JHEP 02 \(2016\) 152](#) [[arXiv:1512.05777](#)] [[INSPIRE](#)].
- [4] R.S. Gupta, S. Jäger, Y. Kats, G. Perez and E. Stamou, *Interpreting a 750 GeV diphoton resonance*, [arXiv:1512.05332](#) [[INSPIRE](#)].
- [5] D. Aloni, K. Blum, A. Dery, A. Efrati and Y. Nir, *On a possible large width 750 GeV diphoton resonance at ATLAS and CMS*, [arXiv:1512.05778](#) [[INSPIRE](#)].
- [6] A. Brignole, F. Feruglio and F. Zwirner, *Four-fermion interactions and sgoldstino masses in models with a superlight gravitino*, [Phys. Lett. B 438 \(1998\) 89](#) [[hep-ph/9805282](#)] [[INSPIRE](#)].
- [7] B. Bellazzini, R. Franceschini, F. Sala and J. Serra, *Goldstones in diphotons*, [JHEP 04 \(2016\) 072](#) [[arXiv:1512.05330](#)] [[INSPIRE](#)].
- [8] C. Petersson and R. Torre, *750 GeV diphoton excess from the goldstino superpartner*, [Phys. Rev. Lett. 116 \(2016\) 151804](#) [[arXiv:1512.05333](#)] [[INSPIRE](#)].
- [9] S.V. Demidov and D.S. Gorbunov, *On the sgoldstino interpretation of the diphoton excess*, [JETP Lett. 103 \(2016\) 219](#) [[arXiv:1512.05723](#)] [[INSPIRE](#)].
- [10] J.A. Casas, J.R. Espinosa and J.M. Moreno, *The 750 GeV diphoton excess as a first light on supersymmetry breaking*, [Phys. Lett. B 759 \(2016\) 159](#) [[arXiv:1512.07895](#)] [[INSPIRE](#)].
- [11] R. Ding et al., *Systematic study of diphoton resonance at 750 GeV from sgoldstino*, [arXiv:1602.00977](#) [[INSPIRE](#)].
- [12] E. Witten, *Constraints on supersymmetry breaking*, [Nucl. Phys. B 202 \(1982\) 253](#) [[INSPIRE](#)].
- [13] K.-I. Izawa and T. Yanagida, *Dynamical supersymmetry breaking in vector-like gauge theories*, [Prog. Theor. Phys. 95 \(1996\) 829](#) [[hep-th/9602180](#)] [[INSPIRE](#)].
- [14] K.A. Intriligator and S.D. Thomas, *Dynamical supersymmetry breaking on quantum moduli spaces*, [Nucl. Phys. B 473 \(1996\) 121](#) [[hep-th/9603158](#)] [[INSPIRE](#)].
- [15] S.P. Martin, *A supersymmetry primer*, [hep-ph/9709356](#) [[INSPIRE](#)].
- [16] S.J. Gates, M.T. Grisaru, M. Roček and W. Siegel, *Superspace or one thousand and one lessons in supersymmetry*, [Front. Phys. 58 \(1983\) 1](#) [[hep-th/0108200](#)] [[INSPIRE](#)].
- [17] T. Han and R.-J. Zhang, *Direct messenger-matter interactions in gauge-mediated supersymmetry breaking models*, [Phys. Lett. B 428 \(1998\) 120](#) [[hep-ph/9802422](#)] [[INSPIRE](#)].
- [18] Y. Shadmi and P.Z. Szabo, *Flavored gauge-mediation*, [JHEP 06 \(2012\) 124](#) [[arXiv:1103.0292](#)] [[INSPIRE](#)].

- [19] Z. Kang, T. Li, T. Liu, C. Tong and J.M. Yang, *A heavy SM-like Higgs and a light stop from Yukawa-Deflected gauge mediation*, *Phys. Rev. D* **86** (2012) 095020 [[arXiv:1203.2336](#)] [[INSPIRE](#)].
- [20] N. Craig, S. Knapen, D. Shih and Y. Zhao, *A complete model of low-scale gauge mediation*, *JHEP* **03** (2013) 154 [[arXiv:1206.4086](#)] [[INSPIRE](#)].
- [21] P. Byakti and T.S. Ray, *Burgeoning the Higgs mass to 125 GeV through messenger-matter interactions in GMSB models*, *JHEP* **05** (2013) 055 [[arXiv:1301.7605](#)] [[INSPIRE](#)].
- [22] J.A. Evans and D. Shih, *Surveying extended GMSB models with $m_h = 125$ GeV*, *JHEP* **08** (2013) 093 [[arXiv:1303.0228](#)] [[INSPIRE](#)].
- [23] E. Perazzi, G. Ridolfi and F. Zwirner, *Signatures of massive sgoldstinos at e^+e^- colliders*, *Nucl. Phys. B* **574** (2000) 3 [[hep-ph/0001025](#)] [[INSPIRE](#)].
- [24] E. Perazzi, G. Ridolfi and F. Zwirner, *Signatures of massive sgoldstinos at hadron colliders*, *Nucl. Phys. B* **590** (2000) 287 [[hep-ph/0005076](#)] [[INSPIRE](#)].
- [25] P. Byakti and D. Ghosh, *Magic messengers in gauge mediation and signal for 125 GeV boosted Higgs boson*, *Phys. Rev. D* **86** (2012) 095027 [[arXiv:1204.0415](#)] [[INSPIRE](#)].
- [26] G.F. Giudice and R. Rattazzi, *Theories with gauge mediated supersymmetry breaking*, *Phys. Rept.* **322** (1999) 419 [[hep-ph/9801271](#)] [[INSPIRE](#)].
- [27] S. Dimopoulos, G.F. Giudice and A. Pomarol, *Dark matter in theories of gauge mediated supersymmetry breaking*, *Phys. Lett. B* **389** (1996) 37 [[hep-ph/9607225](#)] [[INSPIRE](#)].
- [28] S.P. Martin, *Generalized messengers of supersymmetry breaking and the sparticle mass spectrum*, *Phys. Rev. D* **55** (1997) 3177 [[hep-ph/9608224](#)] [[INSPIRE](#)].
- [29] ATLAS collaboration, *Search for photonic signatures of gauge-mediated supersymmetry in 8 TeV pp collisions with the ATLAS detector*, *Phys. Rev. D* **92** (2015) 072001 [[arXiv:1507.05493](#)] [[INSPIRE](#)].
- [30] M. Nardecchia, A. Romanino and R. Ziegler, *Tree level gauge mediation*, *JHEP* **11** (2009) 112 [[arXiv:0909.3058](#)] [[INSPIRE](#)].
- [31] M. Nardecchia, A. Romanino and R. Ziegler, *General aspects of tree level gauge mediation*, *JHEP* **03** (2010) 024 [[arXiv:0912.5482](#)] [[INSPIRE](#)].
- [32] P. Baratella, J. Elias-Miro, J. Penedo and A. Romanino, *A closer look to the sgoldstino interpretation of the diphoton excess*, [arXiv:1603.05682](#) [[INSPIRE](#)].
- [33] Z. Komargodski and D. Shih, *Notes on SUSY and R-symmetry breaking in Wess-Zumino models*, *JHEP* **04** (2009) 093 [[arXiv:0902.0030](#)] [[INSPIRE](#)].
- [34] F. Caracciolo and A. Romanino, *Simple and direct communication of dynamical supersymmetry breaking*, *JHEP* **12** (2012) 109 [[arXiv:1207.5376](#)] [[INSPIRE](#)].
- [35] C. Cheung, A.L. Fitzpatrick and D. Shih, *(Extra)ordinary gauge mediation*, *JHEP* **07** (2008) 054 [[arXiv:0710.3585](#)] [[INSPIRE](#)].
- [36] G.R. Dvali, G.F. Giudice and A. Pomarol, *The Mu problem in theories with gauge mediated supersymmetry breaking*, *Nucl. Phys. B* **478** (1996) 31 [[hep-ph/9603238](#)] [[INSPIRE](#)].
- [37] S. Dimopoulos and G.F. Giudice, *Multimessenger theories of gauge mediated supersymmetry breaking*, *Phys. Lett. B* **393** (1997) 72 [[hep-ph/9609344](#)] [[INSPIRE](#)].

- [38] D. Marques, *Generalized messenger sector for gauge mediation of supersymmetry breaking and the soft spectrum*, *JHEP* **03** (2009) 038 [[arXiv:0901.1326](#)] [[INSPIRE](#)].
- [39] T.T. Dumitrescu, Z. Komargodski, N. Seiberg and D. Shih, *General messenger gauge mediation*, *JHEP* **05** (2010) 096 [[arXiv:1003.2661](#)] [[INSPIRE](#)].
- [40] E. Poppitz and S.P. Trivedi, *New models of gauge and gravity mediated supersymmetry breaking*, *Phys. Rev. D* **55** (1997) 5508 [[hep-ph/9609529](#)] [[INSPIRE](#)].
- [41] N. Arkani-Hamed, J. March-Russell and H. Murayama, *Building models of gauge mediated supersymmetry breaking without a messenger sector*, *Nucl. Phys. B* **509** (1998) 3 [[hep-ph/9701286](#)] [[INSPIRE](#)].
- [42] N. Seiberg, T. Volansky and B. Wecht, *Semi-direct gauge mediation*, *JHEP* **11** (2008) 004 [[arXiv:0809.4437](#)] [[INSPIRE](#)].
- [43] I. Affleck, M. Dine and N. Seiberg, *Dynamical supersymmetry breaking in four-dimensions and its phenomenological implications*, *Nucl. Phys. B* **256** (1985) 557 [[INSPIRE](#)].
- [44] M.E. Peskin, *Duality in supersymmetric Yang-Mills theory*, in the proceedings of *Fields, strings and duality, Theoretical Advanced Study Institute in Elementary Particle Physics (TASI96)*, June 2–28, Boulder U.S.A. (1996), [hep-th/9702094](#) [[INSPIRE](#)].
- [45] E. Poppitz and S.P. Trivedi, *Dynamical supersymmetry breaking*, *Ann. Rev. Nucl. Part. Sci.* **48** (1998) 307 [[hep-th/9803107](#)] [[INSPIRE](#)].
- [46] A.E. Nelson and N. Seiberg, *R symmetry breaking versus supersymmetry breaking*, *Nucl. Phys. B* **416** (1994) 46 [[hep-ph/9309299](#)] [[INSPIRE](#)].
- [47] J. Bagger, E. Poppitz and L. Randall, *The R axion from dynamical supersymmetry breaking*, *Nucl. Phys. B* **426** (1994) 3 [[hep-ph/9405345](#)] [[INSPIRE](#)].
- [48] K.A. Intriligator, N. Seiberg and D. Shih, *Supersymmetry breaking, R-symmetry breaking and metastable vacua*, *JHEP* **07** (2007) 017 [[hep-th/0703281](#)] [[INSPIRE](#)].
- [49] A. Giveon, A. Katz, Z. Komargodski and D. Shih, *Dynamical SUSY and R-symmetry breaking in SQCD with massive and massless flavors*, *JHEP* **10** (2008) 092 [[arXiv:0808.2901](#)] [[INSPIRE](#)].
- [50] A. Giveon, A. Katz and Z. Komargodski, *Uplifted metastable vacua and gauge mediation in SQCD*, *JHEP* **07** (2009) 099 [[arXiv:0905.3387](#)] [[INSPIRE](#)].
- [51] A. Amariti and A. Mariotti, *Two loop R-symmetry breaking*, *JHEP* **07** (2009) 071 [[arXiv:0812.3633](#)] [[INSPIRE](#)].
- [52] A. Basirnia, D. Egana-Ugrinovic, S. Knapen and D. Shih, *125 GeV Higgs from tree-level A-terms*, *JHEP* **06** (2015) 144 [[arXiv:1501.00997](#)] [[INSPIRE](#)].
- [53] A. Kusenko, P. Langacker and G. Segre, *Phase transitions and vacuum tunneling into charge and color breaking minima in the MSSM*, *Phys. Rev. D* **54** (1996) 5824 [[hep-ph/9602414](#)] [[INSPIRE](#)].
- [54] D. Chowdhury, R.M. Godbole, K.A. Mohan and S.K. Vempati, *Charge and color breaking constraints in MSSM after the Higgs discovery at LHC*, *JHEP* **02** (2014) 110 [[arXiv:1310.1932](#)] [[INSPIRE](#)].
- [55] S.R. Coleman and E.J. Weinberg, *Radiative corrections as the origin of spontaneous symmetry breaking*, *Phys. Rev. D* **7** (1973) 1888 [[INSPIRE](#)].