

RECEIVED: April 11, 2017

REVISED: May 29, 2017

ACCEPTED: June 7, 2017

PUBLISHED: June 15, 2017

Extraction of partonic transverse momentum distributions from semi-inclusive deep-inelastic scattering, Drell-Yan and Z-boson production

Alessandro Bacchetta,^{a,b} Filippo Delcarro,^{a,b} Cristian Pisano,^{a,b} Marco Radici^b and Andrea Signori^c

^a*Dipartimento di Fisica, Università di Pavia,
via Bassi 6, I-27100 Pavia, Italy*

^b*INFN Sezione di Pavia,
via Bassi 6, I-27100 Pavia, Italy*

^c*Theory Center, Thomas Jefferson National Accelerator Facility,
12000 Jefferson Avenue, Newport News, VA 23606, U.S.A.*

E-mail: alessandro.bacchetta@unipv.it, filippo.delcarro@pv.infn.it,
cristian.pisano@unipv.it, marco.radici@pv.infn.it, asignori@jlab.org

ABSTRACT: We present an extraction of unpolarized partonic transverse momentum distributions (TMDs) from a simultaneous fit of available data measured in semi-inclusive deep-inelastic scattering, Drell-Yan and Z boson production. To connect data at different scales, we use TMD evolution at next-to-leading logarithmic accuracy. The analysis is restricted to the low-transverse-momentum region, with no matching to fixed-order calculations at high transverse momentum. We introduce specific choices to deal with TMD evolution at low scales, of the order of 1 GeV^2 . This could be considered as a first attempt at a global fit of TMDs.

KEYWORDS: Deep Inelastic Scattering (Phenomenology), QCD Phenomenology

ARXIV EPRINT: [1703.10157](https://arxiv.org/abs/1703.10157)

Contents

1	Introduction	1
2	Formalism	2
2.1	Semi-inclusive DIS	2
2.2	Drell-Yan and Z production	5
2.3	TMDs and their evolution	7
3	Data analysis	10
3.1	Semi-inclusive DIS data	10
3.1.1	HERMES data	12
3.1.2	COMPASS data	12
3.2	Low-energy Drell-Yan data	13
3.3	Z-boson production data	14
3.4	The replica method	14
4	Results	15
4.1	Agreement between data and theory	16
4.2	Transverse momentum dependence at 1 GeV	22
4.3	Stability of our results	25
5	Conclusions	26

1 Introduction

Parton distribution functions describe the internal structure of the nucleon in terms of its elementary constituents (quarks and gluons). They cannot be easily computed from first principles, because they require the ability to carry out Quantum Chromodynamics (QCD) calculations in its nonperturbative regime. Many experimental observables in hard scattering experiments involving hadrons are related to parton distribution functions (PDFs) and fragmentation functions (FFs), in a way that is specified by factorization theorems (see, e.g., refs. [1, 2]). These theorems also elucidate the universality properties of PDFs and FFs (i.e., the fact that they are the same in different processes) and their evolution equations (i.e., how they get modified by the change in the hard scale of the process). Availability of measurements of different processes in different experiments makes it possible to test factorization theorems and extract PDFs and FFs through so-called global fits. On the other side, the knowledge of PDFs and FFs allows us to make predictions for other hard hadronic processes. These general statements apply equally well to standard collinear PDFs and FFs and to transverse-momentum-dependent parton distribution functions (TMD PDFs) and fragmentation functions (TMD FFs). Collinear PDFs describe the distribution of partons integrated over all components of partonic momentum except the one collinear to the parent hadron; hence, collinear PDFs are functions of the parton longitudinal momentum fraction x . TMD PDFs (or TMDs for short) include also the dependence on the transverse

momentum \mathbf{k}_\perp . They can be interpreted as three-dimensional generalizations of collinear PDFs. Similar arguments apply to collinear FFs and TMD FFs [3].

There are several differences between collinear and TMD distributions. From the formal point of view, factorization theorems for the two types of functions are different, implying also different universality properties and evolution equations [4]. From the experimental point of view, observables related to TMDs require the measurement of some transverse momentum component much smaller than the hard scale of the process [5, 6]. For instance, Deep-Inelastic Scattering (DIS) is characterized by a hard scale represented by the 4-momentum squared of the virtual photon ($-Q^2$). In inclusive DIS this is the only scale of the process, and only collinear PDFs and FFs can be accessed. In semi-inclusive DIS (SIDIS) also the transverse momentum of the outgoing hadron (P_{hT}) can be measured [7, 8]. If $P_{hT}^2 \ll Q^2$, TMD factorization can be applied and the process is sensitive to TMDs [2].

If polarization is taken into account, several TMDs can be introduced [7, 9–12]. Attempts to extract some of them have already been presented in the past [13–21]. In this work, we focus on the simplest ones, i.e., the unpolarized TMD PDF $f_1^q(x, k_\perp^2)$ and the unpolarized TMD FF $D_1^{q \rightarrow h}(z, P_\perp^2)$, where z is the fractional energy carried by the detected hadron h , k_\perp is the transverse momentum of the parton with respect to the parent hadron, and P_\perp is the transverse momentum of the produced hadron with respect to the parent parton. Despite their simplicity, the phenomenology of these unpolarized TMDs presents several challenges [22]: the choice of a functional form for the nonperturbative components of TMDs, the inclusion of a possible dependence on partonic flavor [23], the implementation of TMD evolution [4, 24], the matching to fixed-order calculations in collinear factorization [25].

We take into consideration three kinds of processes: SIDIS, Drell-Yan processes (DY) and the production of Z bosons. To date, they represent all possible processes where experimental information is available for unpolarized TMD extractions. The only important process currently missing is electron-positron annihilation, which is particularly important for the determination of TMD FFs [24]. This work can therefore be considered as the first attempt at a global fit of TMDs.

The paper is organized as follows. In section 2, the general formalism for TMDs in SIDIS, DY processes, and Z production is briefly outlined, including a description of the assumptions and approximations in the phenomenological implementation of TMD evolution equations. In section 3, the criteria for selecting the data analyzed in the fit are summarized and commented. In section 4, the results of our global fit are presented and discussed. In section 5, we summarize the results and present an outlook for future improvements.

2 Formalism

2.1 Semi-inclusive DIS

In one-particle SIDIS, a lepton ℓ with momentum l scatters off a hadron target N with mass M and momentum P . In the final state, the scattered lepton momentum l' is measured together with one hadron h with mass M_h and momentum P_h . The corresponding reaction

formula is

$$\ell(l) + N(P) \rightarrow \ell(l') + h(P_h) + X. \quad (2.1)$$

The space-like momentum transfer is $q = l - l'$, with $Q^2 = -q^2$. We introduce the usual invariants

$$x = \frac{Q^2}{2P \cdot q}, \quad y = \frac{P \cdot q}{P \cdot l}, \quad z = \frac{P \cdot P_h}{P \cdot q}, \quad \gamma = \frac{2Mx}{Q}. \quad (2.2)$$

The available data refer to SIDIS hadron multiplicities, namely to the differential number of hadrons produced per corresponding inclusive DIS event. In terms of cross sections, we define the multiplicities as

$$m_N^h(x, z, |\mathbf{P}_{hT}|, Q^2) = \frac{d\sigma_N^h/(dx dz d|\mathbf{P}_{hT}| dQ^2)}{d\sigma_{\text{DIS}}/(dx dQ^2)}, \quad (2.3)$$

where $d\sigma_N^h$ is the differential cross section for the SIDIS process and $d\sigma_{\text{DIS}}$ is the corresponding inclusive one, and where \mathbf{P}_{hT} is the component of \mathbf{P}_h transverse to \mathbf{q} (we follow here the notation suggested in ref. [26]). In the single-photon-exchange approximation, the multiplicities can be written as ratios of structure functions (see ref. [8] for details):

$$m_N^h(x, z, |\mathbf{P}_{hT}|, Q^2) = \frac{2\pi |\mathbf{P}_{hT}| F_{UU,T}(x, z, \mathbf{P}_{hT}^2, Q^2) + 2\pi \varepsilon |\mathbf{P}_{hT}| F_{UU,L}(x, z, \mathbf{P}_{hT}^2, Q^2)}{F_T(x, Q^2) + \varepsilon F_L(x, Q^2)}, \quad (2.4)$$

where

$$\varepsilon = \frac{1 - y - \frac{1}{4}\gamma^2 y^2}{1 - y + \frac{1}{2}y^2 + \frac{1}{4}\gamma^2 y^2}. \quad (2.5)$$

In the numerator of eq. (2.4) the structure function $F_{XY,Z}$ corresponds to a lepton with polarization X scattering on a target with polarization Y by exchanging a virtual photon in a polarization state Z . In the denominator, only the photon polarization is explicitly written (T, L), as usually done in the literature.

The semi-inclusive cross section can be expressed in a factorized form in terms of TMDs only in the kinematic limits $M^2 \ll Q^2$ and $\mathbf{P}_{hT}^2 \ll Q^2$. In these limits, the structure function $F_{UU,L}$ of eq. (2.4) can be neglected [27]. The structure function F_L in the denominator contains contributions involving powers of the strong coupling constant α_S at an order that goes beyond the level reached in this analysis; hence, it will be consistently neglected (for measurements and estimates of the F_L structure function see, e.g., refs. [28, 29] and references therein).

To express the structure functions in terms of TMD PDFs and FFs, we rely on the factorized formula for SIDIS [2, 30–37] (see figure 1 for a graphical representation of the involved transverse momenta):

$$\begin{aligned} F_{UU,T}(x, z, \mathbf{P}_{hT}^2, Q^2) &= \sum_a \mathcal{H}_{UU,T}^a(Q^2) \\ &\times x \int d^2\mathbf{k}_\perp d^2\mathbf{P}_\perp f_1^a(x, \mathbf{k}_\perp^2; Q^2) D_1^{a \rightarrow h}(z, \mathbf{P}_\perp^2; Q^2) \delta^{(2)}(z\mathbf{k}_\perp - \mathbf{P}_{hT} + \mathbf{P}_\perp) \\ &+ Y_{UU,T}(Q^2, \mathbf{P}_{hT}^2) + \mathcal{O}(M^2/Q^2). \end{aligned} \quad (2.6)$$




Figure 1. Diagram describing the relevant momenta involved in a semi-inclusive DIS event (see also ref. [38]): a virtual photon (defining the reference axis) strikes a parton inside a proton. The parton has a transverse momentum \mathbf{k}_{\perp} (not measured). The struck parton fragments into a hadron, which acquires a further transverse momentum \mathbf{P}_{\perp} (not measured). The total measured transverse-momentum of the final hadron is \mathbf{P}_{hT} . When Q^2 is very large, the longitudinal components are all much larger than the transverse components. In this regime, $\mathbf{P}_{hT} \approx z\mathbf{k}_{\perp} + \mathbf{P}_{\perp}$.

Here, $\mathcal{H}_{UU,T}$ is the hard scattering part; $f_1^a(x, \mathbf{k}_{\perp}^2; Q^2)$ is the TMD PDF of unpolarized partons with flavor a in an unpolarized proton, carrying longitudinal momentum fraction x and transverse momentum \mathbf{k}_{\perp} . The $D_1^{a \rightarrow h}(z, \mathbf{P}_{\perp}^2; Q^2)$ is the TMD FF describing the fragmentation of an unpolarized parton with flavor a into an unpolarized hadron h carrying longitudinal momentum fraction z and transverse momentum \mathbf{P}_{\perp} (see figure 1). TMDs generally depend on two energy scales [2], which enter via the renormalization of ultraviolet and rapidity divergencies. In this work we choose them to be equal and set them to Q^2 . The term $Y_{UU,T}$ is introduced to ensure a matching to the perturbative fixed-order calculations at higher transverse momenta.

In our analysis, we neglect any correction of the order of M^2/Q^2 or higher to eq. (2.6). At large Q^2 this is well justified. However, fixed-target DIS experiments typically collect a large amount of data at relatively low Q^2 values, where these assumptions should be all tested in future studies. The reliability of the theoretical description of SIDIS at low Q^2 has been recently discussed in refs. [39, 40].

Eq. (2.6) can be expanded in powers of α_S . In the present analysis, we will consider only the terms at order α_S^0 . In this case $\mathcal{H}_{UU,T}^a(Q^2) \approx e_a^2$ and $Y_{UU,T} \approx 0$. However, perturbative corrections include large logarithms $L \equiv \log(z^2 Q^2 / P_{hT}^2)$, so that $\alpha_S L \approx 1$. In the present analysis, we will take into account all leading and Next-to-Leading Logarithms (NLL).¹

In these approximations (α_S^0 and NLL), only the first term in eq. (2.6) is relevant (often in the literature this has been called W term). We expect this term to provide a good description of the structure function only in the region where $P_{hT}^2 \ll Q^2$. We leave a detailed treatment of the matching to the high $P_{hT}^2 \approx Q^2$ region to future investigations (see, e.g., ref. [25]).

¹We remark that formulas at NNLL are available in the literature [41].

To the purpose of applying TMD evolution equations, we need to calculate the Fourier transform of the part of eq. (2.6) involving TMDs. The structure function thus reduces to

$$F_{UU,T}(x, z, \mathbf{P}_{hT}^2, Q^2) \approx 2\pi \sum_a e_a^2 x \int_0^\infty d\xi_T \xi_T J_0(\xi_T |\mathbf{P}_{hT}|/z) \tilde{f}_1^a(x, \xi_T^2; Q^2) \tilde{D}_1^{a-h}(z, \xi_T^2; Q^2), \quad (2.7)$$

where we introduced the Fourier transforms of the TMD PDF and FF according to

$$\tilde{f}_1^a(x, \xi_T^2; Q^2) = \int_0^\infty d|\mathbf{k}_\perp| |\mathbf{k}_\perp| J_0(\xi_T |\mathbf{k}_\perp|) f_1^a(x, \mathbf{k}_\perp^2; Q^2), \quad (2.8)$$

$$\tilde{D}_1^{a-h}(z, \xi_T^2; Q^2) = \int_0^\infty \frac{d|\mathbf{P}_\perp|}{z^2} |\mathbf{P}_\perp| J_0(\xi_T |\mathbf{P}_\perp|/z) D_1^{a-h}(z, \mathbf{P}_\perp^2; Q^2). \quad (2.9)$$

2.2 Drell-Yan and Z production

In a Drell-Yan process, two hadrons A and B with momenta P_A and P_B collide at a center-of-mass energy squared $s = (P_A + P_B)^2$ and produce a virtual photon or a Z boson plus hadrons. The boson decays into a lepton-antilepton pair. The reaction formula is

$$A(P_A) + B(P_B) \rightarrow [\gamma^*/Z + X \rightarrow] \ell^+(l) + \ell^-(l') + X. \quad (2.10)$$

The invariant mass of the virtual photon is $Q^2 = q^2$ with $q = l + l'$. We introduce the rapidity of the virtual photon/ Z boson

$$\eta = \frac{1}{2} \log \left(\frac{q^0 + q_z}{q^0 - q_z} \right), \quad (2.11)$$

where the z direction is defined along the momentum of hadron A (see figure 2).

The cross section can be written in terms of structure functions [42, 43]. For our purposes, we need the unpolarized cross section integrated over $d\Omega$ and over the azimuthal angle of the virtual photon,

$$\frac{d\sigma}{dQ^2 dq_T^2 d\eta} = \sigma_0^{\gamma,Z} \left(F_{UU}^1 + \frac{1}{2} F_{UU}^2 \right). \quad (2.12)$$

The elementary cross sections are

$$\sigma_0^\gamma = \frac{4\pi^2 \alpha_{\text{em}}^2}{3Q^2 s}, \quad \sigma_0^Z = \frac{\pi^2 \alpha_{\text{em}}}{s \sin^2 \theta_W \cos^2 \theta_W} B_R(Z \rightarrow \ell^+ \ell^-) \delta(Q^2 - M_Z^2), \quad (2.13)$$

where θ_W is Weinberg's angle, M_Z is the mass of the Z boson, and $B_R(Z \rightarrow \ell^+ \ell^-)$ is the branching ratio for the Z boson decay in two leptons. We adopted the narrow-width approximation, i.e., we neglect contributions for $Q^2 \neq M_Z^2$. We used the values $\sin^2 \theta_W = 0.2313$, $M_Z = 91.18 \text{ GeV}$, and $B_R(Z \rightarrow \ell^+ \ell^-) = 3.366$ [44]. Similarly to the SIDIS case, in the kinematic limit $q_T^2 \ll Q^2$ the structure function F_{UU}^2 can be neglected (for measurement and estimates of this structure function see, e.g., ref. [45] and references therein).

The longitudinal momentum fractions of the annihilating quarks can be written in terms of rapidity in the following way

$$x_A = \frac{Q}{\sqrt{s}} e^\eta, \quad x_B = \frac{Q}{\sqrt{s}} e^{-\eta}. \quad (2.14)$$




Figure 2. Diagram describing the relevant momenta involved in a Drell-Yan event: two partons from two hadrons collide. They have transverse momenta $\mathbf{k}_{\perp A}$ and $\mathbf{k}_{\perp B}$ (not measured). They produce a virtual photon with (measured) transverse momentum $\mathbf{q}_T = \mathbf{k}_{\perp A} + \mathbf{k}_{\perp B}$ with respect to the hadron collision axis.

Some experiments use the variable x_F , which is connected to the other variables by the following relations

$$\eta = \sinh^{-1} \left(\frac{\sqrt{s} x_F}{Q} \right), \quad x_A = \sqrt{\frac{Q^2}{s} + \frac{x_F^2}{4}} + \frac{x_F}{2}, \quad x_B = x_A - x_F. \quad (2.15)$$

The structure function F_{UU}^1 can be written as (see figure 2 for a graphical representation of the involved transverse momenta)

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T^2, Q^2) = \sum_a \mathcal{H}_{UU}^{1a}(Q^2) \times \int d^2\mathbf{k}_{\perp A} d^2\mathbf{k}_{\perp B} f_1^a(x_A, \mathbf{k}_{\perp A}^2; Q^2) f_1^{\bar{a}}(x_B, \mathbf{k}_{\perp B}^2; Q^2) \delta^{(2)}(\mathbf{k}_{\perp A} - \mathbf{q}_T + \mathbf{k}_{\perp B}) + Y_{UU}^1(Q^2, \mathbf{q}_T^2) + \mathcal{O}(M^2/Q^2). \quad (2.16)$$

As in the SIDIS case, in our analysis we neglect the Y_{UU} term and we consider the hard coefficients only up to leading order in the couplings, i.e.,

$$\mathcal{H}_{UU,\gamma}^{1a}(Q^2) \approx \frac{e_a^2}{N_c}, \quad \mathcal{H}_{UU,Z}^{1a}(Q^2) \approx \frac{V_a^2 + A_a^2}{N_c}, \quad (2.17)$$

where²

$$V_a = I_{3a} - 2e_a \sin^2 \theta_W, \quad A_a = I_{3a}. \quad (2.18)$$

The structure function can be conveniently expressed as a Fourier transform of the right-hand side of eq. (2.16) as

$$F_{UU}^1(x_A, x_B, \mathbf{q}_T^2, Q^2) \approx 2\pi \sum_a \mathcal{H}_{UU}^{1a} \int_0^\infty d\xi_T \xi_T J_0(\xi_T |\mathbf{q}_T|) \tilde{f}_1^a(x_A, \xi_T^2; Q^2) \tilde{f}_1^{\bar{a}}(x_B, \xi_T^2; Q^2). \quad (2.19)$$

²We remind the reader that the value of weak isospin I_3 is equal to $+1/2$ for u, c, t and $-1/2$ for d, s, b .

2.3 TMDs and their evolution

Evolution equations quantitatively describe the connection between different values for the energy scales. In the following we will set their initial values to μ_b^2 and their final values as Q^2 , so that only Q^2 and μ_b^2 need to be specified in a TMD distribution. Following the formalism of refs. [2, 34], the unpolarized TMD distribution and fragmentation functions in configuration space for a parton with flavor a at a certain scale Q^2 can be written as

$$\begin{aligned} \tilde{f}_1^a(x, \xi_T^2; Q^2) &= \sum_{i=q, \bar{q}, g} (C_{a/i} \otimes f_1^i)(x, \bar{\xi}_*, \mu_b^2) \\ &\quad \times e^{S(\mu_b^2, Q^2)} \left(\frac{Q^2}{\mu_b^2}\right)^{-K(\bar{\xi}_*; \mu_b)} \left(\frac{Q^2}{Q_0^2}\right)^{g_K(\xi_T)} \tilde{f}_{\text{INP}}^a(x, \xi_T^2), \end{aligned} \quad (2.20)$$

$$\begin{aligned} \tilde{D}_1^{a \rightarrow h}(z, \xi_T^2; Q^2) &= \sum_{i=q, \bar{q}, g} (\hat{C}_{a/i} \otimes D_1^{i \rightarrow h})(z, \bar{\xi}_*, \mu_b^2) \\ &\quad \times e^{S(\mu_b^2, Q^2)} \left(\frac{Q^2}{\mu_b^2}\right)^{-K(\bar{\xi}_*; \mu_b)} \left(\frac{Q^2}{Q_0^2}\right)^{g_K(\xi_T)} \tilde{D}_{\text{INP}}^{a \rightarrow h}(z, \xi_T^2). \end{aligned} \quad (2.21)$$

We choose the scale μ_b to be

$$\mu_b = \frac{2e^{-\gamma_E}}{\bar{\xi}_*}, \quad (2.22)$$

where γ_E is the Euler constant and

$$\bar{\xi}_* \equiv \bar{\xi}_*(\xi_T; \xi_{\min}, \xi_{\max}) = \xi_{\max} \left(\frac{1 - e^{-\xi_T^4/\xi_{\max}^4}}{1 - e^{-\xi_T^4/\xi_{\min}^4}} \right)^{1/4}. \quad (2.23)$$

This variable replaces the simple dependence upon ξ_T in the perturbative parts of the TMD definitions of eqs. (2.20), (2.21). In fact, at large ξ_T these parts are no longer reliable. Therefore, the $\bar{\xi}_*$ is chosen to saturate on the maximum value ξ_{\max} , as suggested by the CSS formalism [2, 34]. On the other hand, at small ξ_T the TMD formalism is not valid and should be matched to the fixed-order collinear calculations. The way the matching is implemented is not unique. In any case, the TMD contribution can be arbitrarily modified at small ξ_T . In our approach, we choose to saturate $\bar{\xi}_*$ at the minimum value $\xi_{\min} \propto 1/Q$. With the appropriate choices, for $\xi_T = 0$ the Sudakov exponent vanishes, as it should [46, 47]. Our choice partially corresponds to modifying the resummed logarithms as in ref. [48] and to other similar modifications proposed in the literature [25, 49]. One advantage of these kind of prescriptions is that by integrating over the impact parameter ξ_T , the collinear expression for the cross section in terms of collinear PDFs is recovered, at least at leading order [25]. The choice of the functional form in eq. (2.23) is arbitrary. In the original CSS article [31], a functional form based on a square-root saturation was adopted. With our choice, $\bar{\xi}_*$ rapidly saturates to its limiting values ξ_{\min} and ξ_{\max} , leaving a reasonably wide region where it is equal to ξ_T . This is particularly critical for low values of Q , where ξ_{\min} and ξ_{\max} are very close. We remind the reader that there are different schemes available to deal with the high- ξ_T region, such as the so-called ‘‘complex- ξ prescription’’ [50] or an extrapolation of the perturbative small- ξ_T calculation to the large ξ_T region based on dynamical power corrections [51].

The values of ξ_{\max} and ξ_{\min} could be regarded as boundaries between the regions of “small” and “large” distances. They are not completely arbitrary, but there is also no way to determine them precisely. We choose to fix them to the values

$$\xi_{\max} = 2e^{-\gamma_E} \text{ GeV}^{-1} \approx 1.123 \text{ GeV}^{-1}, \quad \xi_{\min} = 2e^{-\gamma_E}/Q. \quad (2.24)$$

The motivations are the following:

- with the above choices, the scale μ_b is constrained between 1 GeV and Q , so that the collinear PDFs are never computed at a scale lower than 1 GeV and the lower limit of the integrals contained in the definition of the perturbative Sudakov factor (see eq. (2.30)) can never become larger than the upper limit;
- at $Q = Q_0 = 1 \text{ GeV}$, $\xi_{\max} = \xi_{\min}$ and there are no evolution effects; the TMD is simply given by the corresponding collinear function multiplied by a nonperturbative contribution depending on k_{\perp} (plus possible corrections of order α_S from the Wilson coefficients).

At NLL accuracy, for our choice of scales $K(\bar{\xi}_*, \mu_b) = 0$. Similarly, the C and \hat{C} are perturbatively calculable Wilson coefficients for the TMD distribution and fragmentation functions, respectively. They are convoluted with the corresponding collinear functions according to

$$(C_{a/i} \otimes f_1^i)(x, \bar{\xi}_*, \mu_b^2) = \int_x^1 \frac{du}{u} C_{a/i}\left(\frac{x}{u}, \bar{\xi}_*, \alpha_S(\mu_b^2)\right) f_1^i(u; \mu_b^2), \quad (2.25)$$

$$(\hat{C}_{a/i} \otimes D_1^{i \rightarrow h})(z, \bar{\xi}_*, \mu_b^2) = \int_z^1 \frac{du}{u} \hat{C}_{a/i}\left(\frac{z}{u}, \bar{\xi}_*, \alpha_S(\mu_b^2)\right) D_1^{i \rightarrow h}(u; \mu_b^2). \quad (2.26)$$

In the present analysis, we consider only the leading-order term in the α_S expansion for C and \hat{C} , i.e.,

$$C_{a/i}\left(\frac{x}{u}, \bar{\xi}_*, \alpha_S(\mu_b^2)\right) \approx \delta_{ai} \delta(1 - x/u), \quad \hat{C}_{a/i}\left(\frac{z}{u}, \bar{\xi}_*, \alpha_S(\mu_b^2)\right) \approx \delta_{ai} \delta(1 - z/u). \quad (2.27)$$

As a consequence of the choices we made, the expression for the evolved TMD functions reduces to

$$\tilde{f}_1^a(x, \xi_T^2; Q^2) = f_1^a(x; \mu_b^2) e^{S(\mu_b^2, Q^2)} e^{g_K(\xi_T) \ln(Q^2/Q_0^2)} \tilde{f}_{1\text{NP}}^a(x, \xi_T^2), \quad (2.28)$$

$$\tilde{D}_1^{a \rightarrow h}(z, \xi_T^2; Q^2) = D_1^{a \rightarrow h}(z; \mu_b^2) e^{S(\mu_b^2, Q^2)} e^{g_K(\xi_T) \ln(Q^2/Q_0^2)} \tilde{D}_{1\text{NP}}^{a \rightarrow h}(z, \xi_T^2). \quad (2.29)$$

The Sudakov exponent S can be written as

$$S(\mu_b^2, Q^2) = - \int_{\mu_b^2}^{Q^2} \frac{d\mu^2}{\mu^2} \left[A(\alpha_S(\mu^2)) \ln\left(\frac{Q^2}{\mu^2}\right) + B(\alpha_S(\mu^2)) \right], \quad (2.30)$$

where the functions A and B have a perturbative expansions of the form

$$A(\alpha_S(\mu^2)) = \sum_{k=1}^{\infty} A_k \left(\frac{\alpha_S}{\pi}\right)^k, \quad B(\alpha_S(\mu^2)) = \sum_{k=1}^{\infty} B_k \left(\frac{\alpha_S}{\pi}\right)^k. \quad (2.31)$$

To NLL accuracy, we need the following terms [31, 52]

$$A_1 = C_F, \quad A_2 = \frac{1}{2}C_F \left[C_A \left(\frac{67}{18} - \frac{\pi^2}{6} \right) - \frac{5}{9}N_f \right], \quad B_1 = -\frac{3}{2}C_F. \quad (2.32)$$

We use the approximate analytic expression for α_S at NLO with the $\Lambda_{\text{QCD}} = 340 \text{ MeV}$, 296 MeV , 214 MeV for three, four, five flavors, respectively, corresponding to a value of $\alpha_S(M_Z) = 0.117$. We fix the flavor thresholds at $m_c = 1.5 \text{ GeV}$ and $m_b = 4.7 \text{ GeV}$. The integration of the Sudakov exponent in eq. (2.30) can be done analytically (for the complete expressions see, e.g., refs. [36, 53, 54]).

Following refs. [55–57], for the nonperturbative Sudakov factor we make the traditional choice

$$g_K(\xi_T) = -g_2 \xi_T^2 / 2 \quad (2.33)$$

with g_2 a free parameter. Recently, several alternative forms have been proposed [58, 59]. Also, recent theoretical studies aimed at calculating this term using nonperturbative methods [60]. All these choices should be tested in future studies. In ref. [61], a good agreement with data was achieved even without this term, but this is not possible when including data at low Q^2 .

In this analysis, for the collinear PDFs f_1^a we adopt the GJR08FFnloE set [62] through the LHAPDF library [63], and for the collinear fragmentation functions the DSS14 NLO set for pions [64] and the DSS07 NLO set for kaons [65].³ We will comment on the use of other PDF sets in section 4.3.

We parametrize the intrinsic nonperturbative parts of the TMDs in the following ways

$$\tilde{f}_{\text{1NP}}^a(x, \xi_T^2) = \frac{1}{2\pi} e^{-g_{1a} \frac{\xi_T^2}{4}} \left(1 - \frac{\lambda g_{1a}^2}{1 + \lambda g_{1a}} \frac{\xi_T^2}{4} \right), \quad (2.34)$$

$$\tilde{D}_{\text{1NP}}^{a \rightarrow h}(z, \xi_T^2) = \frac{g_{3a \rightarrow h} e^{-g_{3a \rightarrow h} \frac{\xi_T^2}{4z^2}} + (\lambda_F/z^2) g_{4a \rightarrow h}^2 \left(1 - g_{4a \rightarrow h} \frac{\xi_T^2}{4z^2} \right) e^{-g_{4a \rightarrow h} \frac{\xi_T^2}{4z^2}}}{2\pi z^2 \left(g_{3a \rightarrow h} + (\lambda_F/z^2) g_{4a \rightarrow h}^2 \right)}. \quad (2.35)$$

After performing the anti-Fourier transform, the f_{1NP} and D_{1NP} in momentum space correspond to

$$f_{\text{1NP}}^a(x, \mathbf{k}_\perp^2) = \frac{1}{\pi} \frac{(1 + \lambda \mathbf{k}_\perp^2)}{g_{1a} + \lambda g_{1a}^2} e^{-\frac{\mathbf{k}_\perp^2}{g_{1a}}}, \quad (2.36)$$

$$D_{\text{1NP}}^{a \rightarrow h}(z, \mathbf{P}_\perp^2) = \frac{1}{\pi} \frac{1}{g_{3a \rightarrow h} + (\lambda_F/z^2) g_{4a \rightarrow h}^2} \left(e^{-\frac{\mathbf{P}_\perp^2}{g_{3a \rightarrow h}}} + \lambda_F \frac{\mathbf{P}_\perp^2}{z^2} e^{-\frac{\mathbf{P}_\perp^2}{g_{4a \rightarrow h}}} \right). \quad (2.37)$$

The TMD PDF at the starting scale is therefore a normalized sum of a Gaussian with variance g_1 and the same Gaussian weighted by a factor $\lambda \mathbf{k}_\perp^2$. The TMD FF at the starting scale is a normalized sum of a Gaussian with variance g_3 and a second Gaussian with variance g_4 weighted by a factor $\lambda_F \mathbf{P}_\perp^2/z^2$. The choice of this particular functional forms is motivated by model calculations: the weighted Gaussian in the TMD PDF could arise

³After the completion of our analysis, a new set of kaon fragmentation function was presented in ref. [66].

from the presence of components of the quark wave function with angular momentum $L = 1$ [67–71]. Similar features occur in models of fragmentation functions [38, 67, 72].

The Gaussian width of the TMD distributions may depend on the parton flavor a [23, 38, 73]. In the present analysis, however, we assume they are flavor independent. The justification for this choice is that most of the data we are considering are not sufficiently sensitive to flavor differences, leading to unclear results. We will devote attention to this issue in further studies.

Finally, we assume that the Gaussian width of the TMD depends on the fractional longitudinal momentum x according to

$$g_1(x) = N_1 \frac{(1-x)^\alpha x^\sigma}{(1-\hat{x})^\alpha \hat{x}^\sigma}, \quad (2.38)$$

where α , σ , and $N_1 \equiv g_1(\hat{x})$ with $\hat{x} = 0.1$, are free parameters. Similarly, for fragmentation functions we have

$$g_{3,4}(z) = N_{3,4} \frac{(z^\beta + \delta)(1-z)^\gamma}{(\hat{z}^\beta + \delta)(1-\hat{z})^\gamma}, \quad (2.39)$$

where β , γ , δ , and $N_{3,4} \equiv g_{3,4}(\hat{z})$ with $\hat{z} = 0.5$ are free parameters.

The average transverse momentum squared for the distributions in eq. (2.36) and (2.37) can be computed analytically:

$$\langle \mathbf{k}_\perp^2 \rangle(x) = \frac{g_1(x) + 2\lambda g_1^2(x)}{1 + \lambda g_1(x)}, \quad \langle \mathbf{P}_\perp^2 \rangle(z) = \frac{g_3^2(z) + 2\lambda_F g_4^3(z)}{g_3(z) + \lambda_F g_4^2(z)}. \quad (2.40)$$

3 Data analysis

The main goals of our work are to extract information about intrinsic transverse momenta, to study the evolution of TMD parton distributions and fragmentation functions over a large enough range of energy, and to test their universality among different processes. To achieve this we included measurements taken from SIDIS, Drell-Yan and Z boson production from different experimental collaborations at different energy scales. In this section we describe the data sets considered for each process and the applied kinematic cuts.

Table 1 refers to the data sets for SIDIS off proton target (HERMES experiment) and presents their kinematic ranges. The same holds for table 2, table 3, table 4 for SIDIS off deuteron (HERMES and COMPASS experiments), Drell-Yan events at low energy and Z boson production respectively. If not specified otherwise, the theoretical formulas are computed at the average values of the kinematic variables in each bin.

3.1 Semi-inclusive DIS data

The SIDIS data are taken from HERMES [74] and COMPASS [75] experiments. Both data sets have already been analyzed in previous works, e.g., refs. [23, 76], however they have never been fitted together, including also the contributions deriving from TMD evolution.

The application of the TMD formalism to SIDIS depends on the capability of identifying the current fragmentation region. This task has been recently discussed in ref. [39], where the authors point out a possible overlap among different fragmentation regions when the

	HERMES $p \rightarrow \pi^+$	HERMES $p \rightarrow \pi^-$	HERMES $p \rightarrow K^+$	HERMES $p \rightarrow K^-$
Reference	[74]			
Cuts	$Q^2 > 1.4 \text{ GeV}^2$ $0.20 < z < 0.74$ $P_{hT} < \text{Min}[0.2 Q, 0.7 Qz] + 0.5 \text{ GeV}$			
Points	190	190	189	187
Max. Q^2	9.2 GeV ²			
x range	$0.04 < x < 0.4$			

Table 1. SIDIS proton-target data (HERMES experiment).

	HERMES $D \rightarrow \pi^+$	HERMES $D \rightarrow \pi^-$	HERMES $D \rightarrow K^+$	HERMES $D \rightarrow K^-$	COMPASS $D \rightarrow h^+$	COMPASS $D \rightarrow h^-$
Reference	[74]				[75]	
Cuts	$Q^2 > 1.4 \text{ GeV}^2$ $0.20 < z < 0.74$ $P_{hT} < \text{Min}[0.2 Q, 0.7 Qz] + 0.5 \text{ GeV}$					
Points	190	190	189	189	3125	3127
Max. Q^2	9.2 GeV ²				10 GeV ²	
x range	$0.04 < x < 0.4$				$0.005 < x < 0.12$	
Notes					Observable: $m_{\text{norm}}(x, z, \mathbf{P}_{hT}^2, Q^2)$, eq. (3.1)	

Table 2. SIDIS deuteron-target data (HERMES and COMPASS experiments).

	E288 200	E288 300	E288 400	E605
Reference	[79]	[79]	[79]	[80]
Cuts	$q_T < 0.2 Q + 0.5 \text{ GeV}$			
Points	45	45	78	35
\sqrt{s}	19.4 GeV	23.8 GeV	27.4 GeV	38.8 GeV
Q range	4-9 GeV	4-9 GeV	5-9, 11-14 GeV	7-9, 10.5-11.5 GeV
Kin. var.	$\eta=0.40$	$\eta=0.21$	$\eta=0.03$	$x_F = 0.1$

Table 3. Low energy Drell-Yan data collected by the E288 and E605 experiments at Tevatron, with different center-of-mass energies.

hard scale Q is sufficiently low. In this paper we do not tackle this problem and we leave it to future studies. As described in tables 1 and 2, we identify the current fragmentation region operating a cut on z only, namely $0.2 < z < 0.74$.

	CDF Run I	D0 Run I	CDF Run II	D0 Run II
Reference	[81]	[82]	[83]	[84]
Cuts	$q_T < 0.2 Q + 0.5 \text{ GeV} = 18.7 \text{ GeV}$			
Points	31	14	37	8
\sqrt{s}	1.8 TeV	1.8 TeV	1.96 TeV	1.96 TeV
Normalization	1.114	0.992	1.049	1.048

Table 4. Z boson production data collected by the CDF and D0 experiments at Tevatron, with different center-of-mass energies.

Another requirement for the applicability of TMD factorization is the presence of two separate scales in the process. In SIDIS, those are the Q^2 and P_{hT}^2 , which should satisfy the condition $P_{hT}^2 \ll Q^2$, or more precisely $P_{hT}^2/z^2 \ll Q^2$. We implement this condition by imposing $P_{hT} < \min[0.2 Q, 0.7 Qz] + 0.5 \text{ GeV}$. With this choice, P_{hT}^2 is always smaller than $Q^2/3$, but in a few bins (at low Q^2 and z) P_{hT}^2/z^2 may become larger than Q^2 . The applicability of TMD factorization in this case could be questioned. However, as we will explain further in section 4.3, we can obtain a fit that can describe a wide region of P_{hT} and can also perform very well in a restricted region, where TMD factorization certainly holds.

All these choices are summarized in tables 1 and 2.

3.1.1 HERMES data

HERMES hadron multiplicities are measured in a fixed target experiment, colliding a 27.6 GeV lepton beam on a hydrogen (p) or deuterium (D) gas target, for a total of 2688 points. These are grouped in bins of (x, z, Q^2, P_{hT}) with the average values of (x, Q^2) ranging from about $(0.04, 1.25 \text{ GeV}^2)$ to $(0.4, 9.2 \text{ GeV}^2)$. The collinear energy fraction z in eq. (2.2) ranges in $0.1 \leq z \leq 0.9$. The transverse momentum of the detected hadron satisfies $0.1 \text{ GeV} \leq |P_{hT}| \leq 1.3 \text{ GeV}$. The peculiarity of HERMES SIDIS experiment lies in the ability of its detector to distinguish between pions and kaons in the final state, in addition to determining their momenta and charges. We consider eight different combinations of target (p, D) and detected charged hadron (π^\pm, K^\pm). The HERMES collaboration published two distinct sets, characterized by the inclusion or subtraction of the vector meson contribution. In our work we considered only the data set where this contribution has been subtracted.

3.1.2 COMPASS data

The COMPASS collaboration extracted multiplicities for charge-separated but unidentified hadrons produced in SIDIS off a deuteron (${}^6\text{LiD}$) target [75]. The number of data points is an order of magnitude higher compared to the HERMES experiment. The data are organized in multidimensional bins of (x, z, Q^2, P_{hT}) , they cover a range in (x, Q^2) from about $(0.005, 1.11 \text{ GeV}^2)$ to $(0.09, 7.57 \text{ GeV}^2)$ and the interval $0.2 \leq z \leq 0.8$. The multiplicities published by COMPASS are affected by normalization errors (see the *erratum* to ref. [75]). In order to avoid this issue, we divide the data in each bin in (x, z, Q^2) by the data point

with the lowest P_{hT}^2 in the bin. As a result, we define the *normalized* multiplicity as

$$m_{\text{norm}}(x, z, \mathbf{P}_{hT}^2, Q^2) = \frac{m_N^h(x, z, \mathbf{P}_{hT}^2, Q^2)}{m_N^h(x, z, \min[\mathbf{P}_{hT}^2], Q^2)}, \quad (3.1)$$

where the multiplicity m_N^h is defined in eq. (2.3). When fitting normalized multiplicities, the first data point of each bin is considered as a fixed constraint and excluded from the degrees of freedom.

3.2 Low-energy Drell-Yan data

We analyze Drell-Yan events collected by fixed-target experiments at low-energy. These data sets have been considered also in previous works, e.g., in refs. [56, 57, 77, 78]. We used data sets from the E288 experiment [79], which measured the invariant dimuon cross section $Ed^3\sigma/dq^3$ for the production of $\mu^+\mu^-$ pairs from the collision of a proton beam with a fixed target, either composed of Cu or Pt. The measurements were performed using proton incident energies of 200, 300 and 400 GeV, producing three different data sets. Their respective center of mass energies are $\sqrt{s} = 19.4, 23.8, 27.4$ GeV. We also included the set of measurements $Ed^3\sigma/dq^3$ from E605 [80], extracted from the collision of a proton beam with an energy of 800 GeV ($\sqrt{s} = 38.8$ GeV) on a copper fixed target.

The explored Q values are higher compared to the SIDIS case, as can be seen in table 3. E288 provides data at fixed rapidity, whereas E605 provides data at fixed $x_F = 0.1$. We can apply TMD factorization if $q_T^2 \ll Q^2$, where q_T is the transverse momentum of the intermediate electroweak boson, reconstructed from the kinematics of the final state leptons. We choose $q_T < 0.2 Q + 0.5$ GeV. As suggested in ref. [79], we consider the target nuclei as an incoherent ensemble composed 40% by protons and 60% by neutrons.

As we already observed, results from E288 and E605 experiments are reported as $\frac{Ed^3\sigma}{d^3q}$; this variable is related to the differential cross section of eq. (2.12) in the following way:

$$\frac{Ed^3\sigma}{d^3q} = \frac{d^3\sigma}{d\phi d\eta q_T dq_T} \Rightarrow \frac{d^2\sigma}{\pi d\eta d(q_T^2)}, \quad (3.2)$$

where ϕ is the polar angle of q_T and the third term is the average over ϕ . Therefore, the invariant dimuon cross section can be obtained from eq. (2.12) integrating over Q^2 and adding a factor $1/\pi$ to the result

$$\frac{Ed^3\sigma}{d^3q} = \frac{1}{\pi} \int dQ^2 \frac{d\sigma}{dQ^2 dq_T^2 d\eta}. \quad (3.3)$$

Numerically we checked that integrating in Q^2 only the prefactor σ_q^γ (see eq. (2.13)) introduces only a negligible error in the theoretical estimates. We also assume that α_{em} does not change within the experimental bin. Therefore, for Drell-Yan we obtain

$$\frac{1}{\pi} \int dQ^2 \frac{d\sigma}{dQ^2 dq_T^2 d\eta} \approx \frac{4\pi\alpha_{\text{em}}^2}{3s} \ln\left(\frac{Q_f^2}{Q_i^2}\right) F_{UU}^1, \quad (3.4)$$

where $Q_{i,f}$ are the lower and upper values in the experimental bin.

3.3 Z-boson production data

In order to reach higher Q and q_T values, we also consider Z boson production in collider experiments at Tevatron. We analyze data from CDF and D0, collected during Tevatron Run I [81, 82] at $\sqrt{s} = 1.8$ TeV and Run II [83, 84] at $\sqrt{s} = 1.96$ TeV. CDF and D0 collaborations studied the differential cross section for the production of an e^+e^- pair from $p\bar{p}$ collision through an intermediate Z vector boson, namely $p\bar{p} \rightarrow Z \rightarrow e^+e^- + X$.

The invariant mass distribution peaks at the Z -pole, $Q \approx M_Z$, while the transverse momentum of the exchanged Z ranges in $0 < q_T < 20$ GeV. We use the same kinematic cut applied to Drell-Yan events: $q_T < 0.2 Q + 0.5$ GeV = 18.7 GeV, since Q is fixed to M_Z .

The observable measured in CDF and D0 is

$$\frac{d\sigma}{dq_T} = \int dQ^2 d\eta 2q_T \frac{d\sigma}{dQ^2 dq_T^2 d\eta} \approx \frac{\pi^2 \alpha_{\text{em}}}{s \sin^2 \theta_W \cos^2 \theta_W} B_R(Z \rightarrow \ell^+ \ell^-) 2q_T \int d\eta F_{UU}^1, \quad (3.5)$$

apart from the case of D0 Run II, for which the published data refer to $1/\sigma \times d\sigma/dq_T$. In order to work with the same observable, we multiply the D0 Run II data by the total cross section of the process $\sigma_{\text{exp}} = 255.8 \pm 16$ pb [85]. In this case, we add in quadrature the uncertainties of the total cross section and of the published data.

We normalize our functional form with the factors listed in table 4. These are the same normalization factors used in ref. [78], computed by comparing the experimental total cross section with the theoretical results based on the code of ref. [86]. These factors are not precisely consistent with our formulas. In fact, as we will discuss in section 4.3 a 5% increase in these factors would improve the agreement with data, without affecting the TMD parameters.

3.4 The replica method

Our fit is based on the replica method. In this section we describe it and we give a definition of the χ^2 function minimized by the fit procedure. The fit and the error analysis are carried out using a similar Monte Carlo approach as in refs. [23, 87, 88] and taking inspiration from the work of the Neural-Network PDF (NNPDF) collaboration (see, e.g., refs. [89–91]). The approach consists in creating \mathcal{M} replicas of the data points. In each replica (denoted by the index r), each data point i is shifted by a Gaussian noise with the same variance as the measurement. Each replica, therefore, represents a possible outcome of an independent experimental measurement, which we denote by $m_{N,r}^h(x, z, \mathbf{P}_{hT}^2, Q^2)$. The number of replicas is chosen so that the mean and standard deviation of the set of replicas accurately reproduces the original data points. In this case 200 replicas are sufficient for the purpose. The error for each replica is taken to be equal to the error on the original data points. This is consistent with the fact that the variance of the \mathcal{M} replicas should reproduce the variance of the original data points.

A minimization procedure is applied to each replica separately, by minimizing the following error function:

$$E_r^2(\{p\}) = \sum_i \frac{\left(m_{N,r}^h(x_i, z_i, \mathbf{P}_{hT}^2, Q_i^2) - m_{N,\text{theo}}^h(x_i, z_i, \mathbf{P}_{hT}^2; \{p\}) \right)^2}{\left(\Delta m_{N,\text{stat}}^{h,2} + \Delta m_{N,\text{sys}}^{h,2} \right)(x_i, z_i, \mathbf{P}_{hT}^2, Q_i^2) + \left(\Delta m_{N,\text{theo}}^h(x_i, z_i, \mathbf{P}_{hT}^2) \right)^2}. \quad (3.6)$$

The sum runs over the i experimental points, including all species of targets N and final-state hadrons h . In each z bin for each replica the values of the collinear fragmentation functions D_1^{a-h} are independently modified with a Gaussian noise with standard deviation equal to the theoretical error ΔD_1^{a-h} . In this work we rely on different parametrizations for D_1^{a-h} : for pions we use the DSEHS analysis [64] at NLO in α_S ; for kaons we use the DSS parametrization [65] at LO in α_S . The uncertainties ΔD_1^{a-h} are estimated from the plots in ref. [92]; they represents the only source of uncertainty in $\Delta m_{N,\text{theo}}^h$. Statistical and systematic experimental uncertainties $\Delta m_{N,\text{stat}}^h$ and $\Delta m_{N,\text{sys}}^h$ are taken from the experimental collaborations. We do not take into account the covariance among different kinematic bins.

We minimize the error function in eq. (3.6) with MINUIT [93]. In each replica we randomize the starting point of the minimization, to better sample the space of fit parameters. The final outcome is a set of \mathcal{M} different vectors of best-fit parameters, $\{p_{0r}\}$, $r = 1, \dots, \mathcal{M}$, with which we can calculate any observable, its mean, and its standard deviation. The distribution of these values needs not to be necessarily Gaussian. In fact, in this case the 1σ confidence interval is different from the 68% interval. The latter can simply be computed for each experimental point by rejecting the largest and the lowest 16% of the \mathcal{M} values.

Although the minimization is performed on the function defined in eq. (3.6), the agreement of the \mathcal{M} replicas with the original data is expressed in terms of a χ^2 function defined as in eq. (3.6) but with the replacement $m_{N,r}^h \rightarrow m_N^h$, i.e., with respect to the original data set. If the model is able to give a good description of the data, the distribution of the \mathcal{M} values of $\chi^2/\text{d.o.f.}$ should be peaked around one.

4 Results

Our work aims at simultaneously fitting for the first time data sets related to different experiments. In the past, only fits related either to SIDIS or hadronic collisions have been presented. Here we mention a selection of recent existing analyses.

In ref. [23], the authors fitted HERMES multiplicities only (taking into account a total of 1538 points) without taking into account QCD evolution. In that work, a flavor decomposition in transverse momentum of the unpolarized TMDs and an analysis of the kinematic dependence of the intrinsic average square transverse momenta were presented. In ref. [76] the authors fitted HERMES and COMPASS multiplicities separately (576 and 6284 points respectively), without TMD evolution and introducing an ad-hoc normalization for COMPASS data. A fit of SIDIS data including TMD evolution was performed on measurements by the H1 collaboration of the so-called transverse energy flow [55, 94].

Looking at data from hadronic collisions, Konychev and Nadolsky [57] fitted data of low-energy Drell-Yan events and Z -boson production at Tevatron, taking into account TMD evolution at NLL accuracy (this is the most recent of a series of important papers on the subject [56, 77, 95]). They fitted in total 98 points. Contrary to our approach, Konychev and Nadolsky studied the quality of the fit as a function of ξ_{max} . They found that the best value for ξ_{max} is 1.5 GeV^{-1} (to be compared to our choice $\xi_{\text{max}} \approx 1.123 \text{ GeV}^{-1}$, see section 2.3). Comparisons of best-fit values in the nonperturbative Sudakov form factors are delicate, since the functional form is different from ours. In 2014 D'Alesio, Echevarria, Melis, Scimemi performed a fit [78] of Drell-Yan data and Z -boson production data at Tevatron,

Points	Parameters	χ^2	$\chi^2/\text{d.o.f.}$
8059	11	12629 ± 363	1.55 ± 0.05

Table 5. Total number of points analyzed, number of free parameters and χ^2 values.

focusing in particular on the role of the nonperturbative contribution to the kernel of TMD evolution. This is the fit with the highest accuracy in TMD evolution performed up to date (NNLL in the Sudakov exponent and $\mathcal{O}(\alpha_S)$ in the Wilson coefficients). In the same year Echevarria, Idilbi, Kang and Vitev [15] presented a parametrization of the unpolarized TMD that described qualitatively well some bins of HERMES and COMPASS data, together with Drell-Yan and Z-production data. A similar result was presented by Sun, Isaacson, Yuan and Yuan [96].

In the following, we detail the results of a fit to the data sets described in section 3 with a flavor-independent configuration for the transverse momentum dependence of unpolarized TMDs. In table 5 we present the total χ^2 . The number of degrees of freedom (d.o.f.) is given by the number of data points analyzed reduced by the number of free parameters in the error function. The overall quality of the fit is good, with a global $\chi^2/\text{d.o.f.} = 1.55 \pm 0.05$. Uncertainties are computed as the 68% confidence level (C.L.) from the replica methodology.

4.1 Agreement between data and theory

The partition of the global χ^2 among SIDIS off a proton, SIDIS off a deuteron, Drell-Yan and Z production events is given in table 6, 7, 8, 9 respectively.

Semi-inclusive DIS. For SIDIS at HERMES off a proton, most of the contribution to the χ^2 comes from events with a π^+ in the final state. In ref. [23] the high χ^2 was attributed to the poor agreement between experiment and theory at the level of the collinear multiplicities. In this work we use a newer parametrization of the collinear FFs (DSEHS [64]), based on a fit which includes HERMES collinear pion multiplicities. In spite of this improvement, the contribution to χ^2 from HERMES data is higher than in ref. [23], because the present fit includes data from other experiments (HERMES represents less than 20% of the whole data set). The bins with the worst agreement are at low Q^2 . As we will discuss in section 4.3, we think that the main reason for the large χ^2 at HERMES is a normalization difference. This may also be due to the fact that we are computing our theoretical estimates at the average values of the kinematic variables, instead of integrating the multiplicities in each bin. Kaon multiplicities have in general a lower χ^2 , due to the bigger statistical errors and the large uncertainties for the kaon FFs.

For pion production off a deuteron at HERMES the χ^2 is lower with respect to the production off a proton, but still compatible within uncertainties. For kaon production off a deuteron the χ^2 is higher with respect to the scattering off a proton. The difference is especially large for K^- .

SIDIS at COMPASS involves scattering off deuteron only, $D \rightarrow h^\pm$, and we identify $h \equiv \pi$. The quality of the agreement between theory and COMPASS data is better than in the case of pion production at HERMES. This depends on at least two factors: first, our fit is essentially driven by the COMPASS data, which represent about 75% of the whole data

	HERMES $p \rightarrow \pi^+$	HERMES $p \rightarrow \pi^-$	HERMES $p \rightarrow K^+$	HERMES $p \rightarrow K^-$
Points	190	190	189	187
χ^2/points	4.83 ± 0.42	2.47 ± 0.28	0.91 ± 0.14	0.82 ± 0.17

Table 6. Number of points analyzed and χ^2 values for SIDIS off a proton target.

	HERMES $D \rightarrow \pi^+$	HERMES $D \rightarrow \pi^-$	HERMES $D \rightarrow K^+$	HERMES $D \rightarrow K^-$	COMPASS $D \rightarrow h^+$	COMPASS $D \rightarrow h^-$
Points	190	190	189	189	3125	3127
χ^2/points	3.46 ± 0.32	2.00 ± 0.17	1.31 ± 0.26	2.54 ± 0.57	1.11 ± 0.03	1.61 ± 0.04

Table 7. Number of points analyzed and χ^2 values for SIDIS off a deuteron target.

set; second, the observable that we fit in this case is the normalized multiplicity, defined in eq. (3.1). This automatically eliminates most of the discrepancy between theory and data due to normalization.

Figure 3 presents the agreement between the theoretical formula in (2.3) and the HERMES multiplicities for production of pions off a proton and a deuteron. Different $\langle x \rangle$, $\langle z \rangle$ and $\langle Q^2 \rangle$ bins are displayed as a function of the transverse momentum of the detected hadron P_{hT} . The grey bands are an envelope of the 200 replica of best-fit curves. For every point in P_{hT} we apply a 68% C.L. selection criterion. Points marked with different symbols and colors correspond to different $\langle z \rangle$ values. There is a strong correlation between $\langle x \rangle$ and $\langle Q^2 \rangle$ that does not allow us to explore the x and Q^2 dependence of the TMDs separately. Studying the contributions to the χ^2/points as a function of the kinematics, we notice that the $\chi^2(Q^2)$ tends to improve as we move to higher Q^2 values, where the kinematic approximations of factorization are more reliable. Moreover, usually the $\chi^2(z)$ increases at lower z values.

Figure 4 has same contents and notation as in figure 3 but for kaons in the final state. In this case, the trend of the agreement as a function of Q^2 is not as clear as for the case of pions: good agreement is found also at low Q^2 .

In figure 5 we present COMPASS normalized multiplicities (see eq. (3.1)) for production of π^- off a deuteron for different $\langle x \rangle$, $\langle z \rangle$, and $\langle Q^2 \rangle$ bins as a function of the transverse momentum of the detected hadron P_{hT} . The open marker around the first P_{hT} point in each panel indicates that the first value is fixed and not fitted. The correlation between x and Q^2 is less strong than at HERMES and this allows us to study different $\langle x \rangle$ bins at fixed $\langle Q^2 \rangle$. For the highest Q^2 bins, the agreement is good for all $\langle x \rangle$, $\langle z \rangle$ and P_{hT}^2 . In bins at lower Q^2 , the descriptions gets worse, especially at low and high z . For fixed $\langle Q^2 \rangle$ and high $\langle z \rangle$, a good agreement is recovered moving to higher $\langle x \rangle$ bins (see, e.g., the third line from the top in figure 5).

Figure 6 has same content and notation as in figure 5, but for $h^+ \equiv \pi^+$. The same comments on the agreement between theory and the data apply.




Figure 3. HERMES multiplicities for production of pions off a proton and a deuteron for different $\langle x \rangle$, $\langle z \rangle$, and $\langle Q^2 \rangle$ bins as a function of the transverse momentum of the detected hadron P_{hT} . For clarity, each $\langle z \rangle$ bin has been shifted by an offset indicated in the legend.

Drell-Yan and Z production. The low energy Drell-Yan data collected by the E288 and E605 experiments at Fermilab have large error bands (see figure 7). This is why the χ^2 values in table 8 are rather low compared to the other data sets.

The agreement is also good for Z boson production, see table 9. The statistics from Run-II is higher, which generates smaller experimental uncertainties and higher χ^2 , especially for the CDF experiment.

Figure 7 displays the cross section for DY events differential with respect to the transverse momentum q_T of the virtual photon, its invariant mass Q^2 and rapidity y . As for the




Figure 4. HERMES multiplicities for production of kaons off a proton and a deuteron for different $\langle x \rangle$, $\langle z \rangle$, and $\langle Q^2 \rangle$ bins as a function of the transverse momentum of the detected hadron P_{hT} . For clarity, each $\langle z \rangle$ bin has been shifted by an offset indicated in the legend.

case of SIDIS, the grey bands are the 68% C.L. envelope of the 200 replicas of the fit function. The four panels represents different values for the rapidity y or x_F (see eq. (2.15)). In each panel, we have plots for different Q^2 values. The lower is Q , the less points in q_T we fit (see also section 3.2). The hard scale lies in the region $4.5 < \langle Q \rangle < 13.5$ GeV. This region is of particular importance, since these “moderate” Q values should be high enough to safely apply factorization and, at the same time, low enough in order for the nonperturbative effects to not be shaded by transverse momentum resummation.




Figure 5. COMPASS multiplicities for production of negative hadrons (π^-) off a deuteron for different $\langle x \rangle$, $\langle z \rangle$, and $\langle Q^2 \rangle$ bins as a function of the transverse momentum of the detected hadron P_{hT} . Multiplicities are normalized to the first bin in P_{hT} for each $\langle z \rangle$ value (see (3.1)). For clarity, each $\langle z \rangle$ bin has been shifted by an offset indicated in the legend.

In figure 8 we compare the cross section differential with respect to the transverse momentum q_T of the virtual Z (namely eq. (2.12) integrated over η) with data from CDF and D0 at Tevatron Run I and II. Due to the higher $Q = M_Z$, the range explored in q_T is much larger compared to all the other observables considered. The tails of the distributions deviate from a Gaussian behavior, as it is also evident in the bins at higher Q^2 in figure 7. The band from the replica methodology in this case is much narrower, due to the reduced




Figure 6. COMPASS multiplicities for production of positive hadrons (π^+) off a deuteron for different $\langle x \rangle$, $\langle z \rangle$, and $\langle Q^2 \rangle$ bins as a function of the transverse momentum of the detected hadron P_{hT} . Multiplicities are normalized to the first bin in P_{hT} for each $\langle z \rangle$ value (see (3.1)). For clarity, each $\langle z \rangle$ bin has been shifted by an offset indicated in the legend.

sensitivity to the intrinsic transverse momenta at $Q = M_Z$ and to the limited range of best-fit values for the parameter g_2 , which controls soft-gluon emission. As an effect of TMD evolution, the peak shifts from ~ 1 GeV for Drell-Yan events in figure 7 to ~ 5 GeV in figure 8. The position of the peak is affected both by the perturbative and the nonperturbative part of the Sudakov exponent (see section 2.3 and [22]). Most of the contributions to the χ^2 comes from normalization effects and not from the shape in q_T (see section 4.3).




Figure 7. Drell-Yan differential cross section for different experiments and different values of \sqrt{s} and for different $\langle Q \rangle$ bins. For clarity, each $\langle Q \rangle$ bin has been normalized (the first data point has been set always equal to 1) and then shifted by an offset indicated in the legend.

	E288 [200]	E288 [300]	E288 [400]	E605
Points	45	45	78	35
χ^2/points	0.99 ± 0.09	0.84 ± 0.10	0.32 ± 0.01	1.12 ± 0.08

Table 8. Number of points analyzed and χ^2 values for fixed-target Drell-Yan experiments at low energy. The labels in square brackets were introduced in section 3.2.

	CDF Run I	D0 Run I	CDF Run II	D0 Run II
Points	31	14	37	8
χ^2/points	1.36 ± 0.00	1.11 ± 0.02	2.00 ± 0.02	1.73 ± 0.01

Table 9. Number of points analyzed and χ^2 values for Z boson production at Tevatron.

4.2 Transverse momentum dependence at 1 GeV

The variables ξ_{\min} and ξ_{\max} delimit the range in ξ_T where transverse momentum resummation is computed perturbatively. The g_2 parameter enters the nonperturbative Sudakov exponent and quantifies the amount of transverse momentum due to soft gluon radiation that is not included in the perturbative part of the Sudakov form factor. As already explained in section 2.3, in this work we fix the value for ξ_{\min} and ξ_{\max} in such a way that at $Q = 1$ GeV the unpolarized TMDs coincide with their nonperturbative input. We leave g_2 as a fit parameter.

Table 10 summarizes the chosen values of ξ_{\min} , ξ_{\max} and the best-fit value for g_2 . The latter is given as an average with 68% C.L. uncertainty computed over the set of 200 replicas. We also quote the results obtained from replica 105, since its parameters are very




Figure 8. Cross section differential with respect to the transverse momentum q_T of a Z boson produced from $p\bar{p}$ collisions at Tevatron. The four panels refer to different experiments (CDF and D0) with two different values for the center-of-mass energy ($\sqrt{s} = 1.8$ TeV and $\sqrt{s} = 1.96$ TeV). In this case the band is narrow due to the narrow range for the best-fit values of g_2 .

	ξ_{\max} [GeV $^{-1}$] (fixed)	ξ_{\min} [GeV $^{-1}$] (fixed)	g_2 [GeV 2]
All replicas	$2e^{-\gamma E}$	$2e^{-\gamma E}/Q$	0.13 ± 0.01
Replica 105	$2e^{-\gamma E}$	$2e^{-\gamma E}/Q$	0.128

Table 10. Values of parameters common to TMD PDFs and TMD FFs.

close to the mean values of all replicas. We obtain a value $g_2 = 0.13 \pm 0.01$, smaller than the value ($g_2 = 0.184 \pm 0.018$) obtained in ref. [57], where however no SIDIS data was taken into consideration, and smaller than the value ($g_2 = 0.16$) chosen in ref. [15]. We stress however that our prescriptions involving both ξ_{\min} and ξ_{\max} are different from previous works.

Table 11 collects the best-fit values of parameters in the nonperturbative part of the TMDs at $Q = 1$ GeV (see eqs. (2.34) and (2.35)); as for g_2 , we give the average value over the full set of replicas and the standard deviation based on a 68% C.L. (see section 3.4), and we also quote the value of replica 105.

In figure 9 we compare different extractions of partonic transverse momenta. The horizontal axis shows the value of the average transverse momentum squared for the incoming parton, $\langle k_{\perp}^2 \rangle(x = 0.1)$ (see eq. (2.40)). The vertical axis shows the value of $\langle P_{\perp}^2 \rangle(z = 0.5)$, the average transverse momentum squared acquired during the fragmentation process (see eq. (2.40)). The white square (label 1) indicates the average values of the two quantities obtained in the present analysis at $Q^2 = 1$ GeV 2 . Each black dot around the white square is an outcome of one replica. The red region around the white square contains the 68% of the replicas that are closest to the average value. The same applies to the white circle and the

TMD PDFs	N_1 [GeV ²]	α	σ		λ [GeV ⁻²]	
All replicas	0.28 ± 0.06	2.95 ± 0.05	0.17 ± 0.02		0.86 ± 0.78	
Replica 105	0.285	2.98	0.173		0.39	
TMD FFs	N_3 [GeV ²]	β	δ	γ	λ_F [GeV ⁻²]	N_4 [GeV ²]
All replicas	0.21 ± 0.02	1.65 ± 0.49	2.28 ± 0.46	0.14 ± 0.07	5.50 ± 1.23	0.13 ± 0.01
Replica 105	0.212	2.10	2.52	0.094	5.29	0.135

Table 11. 68% confidence intervals of best-fit values for parametrizations of TMDs at $Q = 1$ GeV.




Figure 9. Correlation between transverse momenta in TMD FFs, $\langle P_{\perp}^2 \rangle(z = 0.5)$, and in TMD PDFs, $\langle k_{\perp}^2 \rangle(x = 0.1)$, in different phenomenological extractions. (1): average values (white square) obtained in the present analysis, values obtained from each replica (black dots) and 68% C.L. area (red); (2) results from ref. [23], (3) results from ref. [97], (4) results from ref. [76] for HERMES data, (5) results from ref. [76] for HERMES data at high z , (6) results from ref. [76] for normalized COMPASS data, (7) results from ref. [76] for normalized COMPASS data at high z , (8) results from ref. [15].

orange region around it (label 2), related to the flavor-independent version of the analysis in ref. [23], obtained by fitting only HERMES SIDIS data at an average $\langle Q^2 \rangle = 2.4$ GeV² and neglecting QCD evolution. A strong anticorrelation between the transverse momenta is evident in this older analysis. In our new analysis, the inclusion of Drell-Yan and Z production data adds physical information about TMD PDFs, free from the influence of TMD FFs. This reduces significantly the correlation between $\langle k_{\perp}^2 \rangle(x = 0.1)$ and $\langle P_{\perp}^2 \rangle(z = 0.5)$. The 68% confidence region is smaller than in the older analysis. The average values of $\langle k_{\perp}^2 \rangle(x = 0.1)$ are similar and compatible within error bands. The values of $\langle P_{\perp}^2 \rangle(z = 0.5)$ in the present analysis turn out to be larger than in the older analysis, an effect that is due mainly to COMPASS data. It must be kept in mind that the two analyses lead also to differences in the x and z dependence of the transverse momentum squared. This dependence is




Figure 10. Kinematic dependence of $\langle k_{\perp}^2 \rangle(x)$ (a) and of $\langle P_{\perp}^2 \rangle(z)$ (b). The bands are the 68% C.L. envelope of the full sets of best-fit curves. The data used in the fit approximately cover the range $5 \times 10^{-3} \lesssim x \lesssim 0.5$ and $0.2 \lesssim z \lesssim 0.7$.

	$p \rightarrow \pi^+$	$p \rightarrow \pi^-$	$p \rightarrow K^+$	$p \rightarrow K^-$	$D \rightarrow \pi^+$	$D \rightarrow \pi^-$	$D \rightarrow K^+$	$D \rightarrow K^-$
Original	5.18	2.67	0.75	0.78	3.63	2.31	1.12	2.27
Normalized	1.94	1.13	0.57	0.29	1.59	0.80	0.47	0.97

Table 12. $\chi^2/\text{d.o.f.}$ for HERMES data with and without normalization to the value of the first bin in P_{hT} .

shown in figure 10 (a) for $\langle k_{\perp}^2 \rangle(x)$ and figure 10 (b) for $\langle P_{\perp}^2 \rangle(z)$. The bands are computed as the 68% C.L. envelope of the full sets of curves from the 200 replicas. Comparison with other extractions are presented and the legend is detailed in the caption of figure 9.

4.3 Stability of our results

In this subsection we discuss the effect of modifying some of the choices we made in our default fit. Instead of repeating the fitting procedure with different choices, we limit ourselves to checking how the χ^2 of a single replica is affected by the modifications.

As starting point we choose replica 105, which, as discussed above, is one of the most representative among the whole replica set. The global $\chi^2/\text{d.o.f.}$ of replica 105 is 1.51. We keep all parameters fixed, without performing any new minimization, and we compute the $\chi^2/\text{d.o.f.}$ after the modifications described in the following.

First of all, we analyze HERMES data with the same strategy as COMPASS, i.e., we normalize HERMES data to the value of the first bin in P_{hT} . In this case, the global $\chi^2/\text{d.o.f.}$ reduces sharply to 1.27. The partial χ^2 for the different SIDIS processes measured at HERMES are shown in table 12. This confirms that normalization effects are the main contribution to the χ^2 of SIDIS data and have minor effects on TMD-related parameters. In fact, even if we perform a new fit with this modification, the χ^2 does not improve significantly and parameters do not change much.

We consider the effect of changing the normalization of the Z -boson data: if we increase the normalization factors quoted in the last row of table 4 by 5%, the χ^2 quoted in the last row of table 9 drops to 0.66, 0.52, 0.65, 0.68. This effect is also already visible by eye in figure 8: the theoretical curves are systematically below the experimental data points, but the shape is reproduced very well.

We consider the sensitivity of our results to the parameterizations adopted for the collinear quark PDFs. The $\chi^2/\text{d.o.f.}$ varies from its original value 1.51, obtained with the NLO GJR 2008 parametrization [62], to 1.84 using NLO MSTW 2008 [98], and 1.85 using NLO CJ12 [99]. In both cases, the agreement with HERMES and Z boson data is not affected significantly, the agreement with COMPASS data becomes slightly worse, and the agreement with DY data becomes clearly worse.

An extremely important point is the choice of kinematic cuts. Our default choices are listed in tables 1–4. We consider also more stringent kinematic cuts on SIDIS data: $Q^2 > 1.5 \text{ GeV}^2$ and $0.25 < z < 0.6$ instead of $Q^2 > 1.4 \text{ GeV}^2$ and $0.2 < z < 0.7$, leaving the other ones unchanged. The number of bins with these cuts reduces from 8059 to 5679 and the $\chi^2/\text{d.o.f.}$ decreases to the value 1.23. In addition, if we replace the constraint $P_{hT} < \text{Min}[0.2Q, 0.7Qz] + 0.5 \text{ GeV}$ with $P_{hT} < \text{Min}[0.2Q, 0.5Qz] + 0.3 \text{ GeV}$, the number of bins reduces to 3380 and the $\chi^2/\text{d.o.f.}$ decreases further to 0.96. By adopting the even stricter cut $P_{hT} < 0.2Qz$, the number of bins drops to only 477, with a $\chi^2/\text{d.o.f.} = 1.02$. We can conclude that our fit, obtained by fitting data in an extended kinematic region, where TMD factorization may be questioned, works extremely well also in a narrower region, where TMD factorization is expected to be under control.

5 Conclusions

In this work we demonstrated for the first time that it is possible to perform a simultaneous fit of unpolarized TMD PDFs and FFs to data of SIDIS, Drell-Yan and Z boson production at small transverse momentum collected by different experiments. This constitutes the first attempt towards a global fit of $f_1^a(x, k_\perp^2)$ and $D_1^{a \rightarrow h}(z, P_\perp^2)$ in the context of TMD factorization and with the implementation of TMD evolution at NLL accuracy.

We extracted unpolarized TMDs using 8059 data points with 11 free parameters using a replica methodology. We selected data with $Q^2 > 1.4 \text{ GeV}^2$ and $0.2 < z < 0.7$. We restricted our fit to the small transverse momentum region, selecting the maximum value of transverse momentum on the basis of phenomenological considerations (see section 3). With these choices, we included regions where TMD factorization could be questioned, but we checked that our results describe very well the regions where TMD factorization is supposed to hold. The average $\chi^2/\text{d.o.f.}$ is 1.55 ± 0.05 and can be improved up to 1.02 restricting the kinematic cuts, without changing the parameters (see section 4.3). Most of the discrepancies between experimental data and theory comes from the normalization and not from the transverse momentum shape.

Our fit is performed assuming that the intrinsic transverse momentum dependence of TMD PDFs and FFs can be parametrized by a normalized linear combination of a Gaussian and a weighted Gaussian. We considered that the widths of the Gaussians depend on the longitudinal momenta. We neglected a possible flavor dependence. For the nonperturbative component of TMD evolution, we adopted the choice most often used in the literature (see section 2.3).

We plan to release grids of the parametrizations studied in this work via TMDlib [100] to facilitate phenomenological studies for present and future experiments.

In future studies, different functional forms for all the nonperturbative ingredients should be explored, including also a possible flavor dependence of the intrinsic transverse momenta. A more precise analysis from the perturbative point of view is also needed, which should in principle make it possible to relax the tension in the normalization and to describe data at higher transverse momenta. Moreover, the description at low transverse momentum should be properly matched to the collinear fixed-order calculations at high transverse momentum.

Together with an improved theoretical framework, in order to better understand the formalism more experimental data is needed. It would be particularly useful to extend the coverage in x , z , rapidity, and Q^2 . The 12 GeV physics program at Jefferson Lab [101] will be very important to constrain TMD distributions at large x . Additional data from SIDIS (at COMPASS, at a future Electron-Ion Collider), Drell-Yan (at COMPASS, at Fermilab), Z/W production (at LHC, RHIC, and at A Fixed-Target Experiment at the LHC [102]) will be very important. Measurements related to unpolarized TMD FFs at e^+e^- colliders (at Belle-II, BES-III, at a future International Linear Collider) will be invaluable, since they are presently missing.

Our work focused on quark TMDs. We remark that at present almost nothing is known experimentally about gluon TMDs [11, 103], because they typically require higher-energy scattering processes and they are harder to isolate as compared to quark distributions. Several promising measurements have been proposed in order to extract both the unpolarized and linearly polarized gluon TMDs inside an unpolarized proton. The cleanest possibility would be to look at dijet and heavy quark pair production in electron-proton collisions at a future EIC [104, 105]. Other proposals include isolated photon-pair production at RHIC [106] and quarkonium production at the LHC [107–110].

Testing the formalism of TMD factorization and understanding the structure of unpolarized TMDs is only the first crucial step in the exploration of the 3D proton structure in momentum space and this work opens the way to global determinations of TMDs. Building on this, we can proceed to deepen our understanding of hadron structure via polarized structure function and asymmetries (see, e.g., refs. [111, 112] and references therein) and, at the same time, to test the impact of hadron structure in precision measurements at high-energies, such as at the LHC. A detailed mapping of hadron structure is essential to interpret data from hadronic collisions, which are among the most powerful tools to look for footprints of new physics.

Acknowledgments

Discussions with Giuseppe Bozzi and Alexey Vladimirov are gratefully acknowledged. This work is supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No. 647981, 3DSPIN). AS acknowledges support from U.S. Department of Energy contract DE-AC05-06OR23177, under which Jefferson Science Associates, LLC, manages and operates Jefferson Lab. The work of AS has been funded partly also by the program of the Stichting voor Fundamenteel Onderzoek der Materie (FOM), which is financially supported by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.C. Collins, D.E. Soper and G.F. Sterman, *Factorization of Hard Processes in QCD*, *Adv. Ser. Direct. High Energy Phys.* **5** (1989) 1 [[hep-ph/0409313](#)] [[INSPIRE](#)].
- [2] J. Collins, *Foundations of perturbative QCD*, Cambridge University Press (2011) [<http://www.cambridge.org/de/knowledge/isbn/item5756723>] [[INSPIRE](#)].
- [3] R. Angeles-Martinez et al., *Transverse Momentum Dependent (TMD) parton distribution functions: status and prospects*, *Acta Phys. Polon.* **B 46** (2015) 2501 [[arXiv:1507.05267](#)] [[INSPIRE](#)].
- [4] T.C. Rogers, *An Overview of Transverse Momentum Dependent Factorization and Evolution*, *Eur. Phys. J. A* **52** (2016) 153 [[arXiv:1509.04766](#)] [[INSPIRE](#)].
- [5] A. Bacchetta, *Where do we stand with a 3-D picture of the proton?*, *Eur. Phys. J. A* **52** (2016) 163 [[INSPIRE](#)].
- [6] M. Radici, *Imaging the proton*, *AIP Conf. Proc.* **1735** (2016) 020003 [[INSPIRE](#)].
- [7] P.J. Mulders and R.D. Tangerman, *The Complete tree level result up to order $1/Q$ for polarized deep inelastic leptonproduction*, *Nucl. Phys. B* **461** (1996) 197 [Erratum *ibid.* **B 484** (1997) 538] [[hep-ph/9510301](#)] [[INSPIRE](#)].
- [8] A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P.J. Mulders and M. Schlegel, *Semi-inclusive deep inelastic scattering at small transverse momentum*, *JHEP* **02** (2007) 093 [[hep-ph/0611265](#)] [[INSPIRE](#)].
- [9] D. Boer and P.J. Mulders, *Time reversal odd distribution functions in leptonproduction*, *Phys. Rev. D* **57** (1998) 5780 [[hep-ph/9711485](#)] [[INSPIRE](#)].
- [10] A. Bacchetta and P.J. Mulders, *Deep inelastic leptonproduction of spin-one hadrons*, *Phys. Rev. D* **62** (2000) 114004 [[hep-ph/0007120](#)] [[INSPIRE](#)].
- [11] P.J. Mulders and J. Rodrigues, *Transverse momentum dependence in gluon distribution and fragmentation functions*, *Phys. Rev. D* **63** (2001) 094021 [[hep-ph/0009343](#)] [[INSPIRE](#)].
- [12] D. Boer, S. Cotogno, T. van Daal, P.J. Mulders, A. Signori and Y.-J. Zhou, *Gluon and Wilson loop TMDs for hadrons of spin ≤ 1* , *JHEP* **10** (2016) 013 [[arXiv:1607.01654](#)] [[INSPIRE](#)].
- [13] A. Bacchetta and M. Radici, *Constraining quark angular momentum through semi-inclusive measurements*, *Phys. Rev. Lett.* **107** (2011) 212001 [[arXiv:1107.5755](#)] [[INSPIRE](#)].
- [14] M. Anselmino, M. Boglione and S. Melis, *A Strategy towards the extraction of the Sivers function with TMD evolution*, *Phys. Rev. D* **86** (2012) 014028 [[arXiv:1204.1239](#)] [[INSPIRE](#)].
- [15] M.G. Echevarria, A. Idilbi, Z.-B. Kang and I. Vitev, *QCD Evolution of the Sivers Asymmetry*, *Phys. Rev. D* **89** (2014) 074013 [[arXiv:1401.5078](#)] [[INSPIRE](#)].
- [16] M. Anselmino, M. Boglione, U. D'Alesio, F. Murgia and A. Prokudin, *Study of the sign change of the Sivers function from STAR Collaboration W/Z production data*, *JHEP* **04** (2017) 046 [[arXiv:1612.06413](#)] [[INSPIRE](#)].

- [17] Z. Lu and I. Schmidt, *Updating Boer-Mulders functions from unpolarized pd and pp Drell-Yan data*, *Phys. Rev. D* **81** (2010) 034023 [[arXiv:0912.2031](#)] [[INSPIRE](#)].
- [18] V. Barone, M. Boglione, J.O. Gonzalez Hernandez and S. Melis, *Phenomenological analysis of azimuthal asymmetries in unpolarized semi-inclusive deep inelastic scattering*, *Phys. Rev. D* **91** (2015) 074019 [[arXiv:1502.04214](#)] [[INSPIRE](#)].
- [19] C. Lefky and A. Prokudin, *Extraction of the distribution function h_{1T}^\perp from experimental data*, *Phys. Rev. D* **91** (2015) 034010 [[arXiv:1411.0580](#)] [[INSPIRE](#)].
- [20] M. Anselmino, M. Boglione, U. D'Alesio, S. Melis, F. Murgia and A. Prokudin, *Simultaneous extraction of transversity and Collins functions from new SIDIS and e^+e^- data*, *Phys. Rev. D* **87** (2013) 094019 [[arXiv:1303.3822](#)] [[INSPIRE](#)].
- [21] Z.-B. Kang, A. Prokudin, P. Sun and F. Yuan, *Extraction of Quark Transversity Distribution and Collins Fragmentation Functions with QCD Evolution*, *Phys. Rev. D* **93** (2016) 014009 [[arXiv:1505.05589](#)] [[INSPIRE](#)].
- [22] A. Signori, *Flavor and Evolution Effects in TMD Phenomenology: Manifestation of Hadron Structure in High-Energy Scattering Processes*, Ph.D. Thesis, Vrije U., Amsterdam (2016), [[INSPIRE](#)].
- [23] A. Signori, A. Bacchetta, M. Radici and G. Schnell, *Investigations into the flavor dependence of partonic transverse momentum*, *JHEP* **11** (2013) 194 [[arXiv:1309.3507](#)] [[INSPIRE](#)].
- [24] A. Bacchetta, M.G. Echevarria, P.J.G. Mulders, M. Radici and A. Signori, *Effects of TMD evolution and partonic flavor on e^+e^- annihilation into hadrons*, *JHEP* **11** (2015) 076 [[arXiv:1508.00402](#)] [[INSPIRE](#)].
- [25] J. Collins, L. Gamberg, A. Prokudin, T.C. Rogers, N. Sato and B. Wang, *Relating Transverse Momentum Dependent and Collinear Factorization Theorems in a Generalized Formalism*, *Phys. Rev. D* **94** (2016) 034014 [[arXiv:1605.00671](#)] [[INSPIRE](#)].
- [26] D. Boer et al., *Gluons and the quark sea at high energies: Distributions, polarization, tomography*, [arXiv:1108.1713](#) [[INSPIRE](#)].
- [27] A. Bacchetta, D. Boer, M. Diehl and P.J. Mulders, *Matches and mismatches in the descriptions of semi-inclusive processes at low and high transverse momentum*, *JHEP* **08** (2008) 023 [[arXiv:0803.0227](#)] [[INSPIRE](#)].
- [28] ZEUS collaboration, S. Chekanov et al., *Measurement of the Longitudinal Proton Structure Function at HERA*, *Phys. Lett. B* **682** (2009) 8 [[arXiv:0904.1092](#)] [[INSPIRE](#)].
- [29] H1 collaboration, V. Andreev et al., *Measurement of inclusive ep cross sections at high Q^2 at $\sqrt{s} = 225$ and 252 GeV and of the longitudinal proton structure function F_L at HERA*, *Eur. Phys. J. C* **74** (2014) 2814 [[arXiv:1312.4821](#)] [[INSPIRE](#)].
- [30] J.C. Collins and D.E. Soper, *Back-To-Back Jets in QCD*, *Nucl. Phys. B* **193** (1981) 381 [*Erratum* *ibid.* **B 213** (1983) 545] [[INSPIRE](#)].
- [31] J.C. Collins, D.E. Soper and G.F. Sterman, *Transverse Momentum Distribution in Drell-Yan Pair and W and Z Boson Production*, *Nucl. Phys. B* **250** (1985) 199 [[INSPIRE](#)].
- [32] X.-d. Ji and F. Yuan, *Parton distributions in light cone gauge: Where are the final state interactions?*, *Phys. Lett. B* **543** (2002) 66 [[hep-ph/0206057](#)] [[INSPIRE](#)].
- [33] X.-d. Ji, J.-p. Ma and F. Yuan, *QCD factorization for semi-inclusive deep-inelastic scattering at low transverse momentum*, *Phys. Rev. D* **71** (2005) 034005 [[hep-ph/0404183](#)] [[INSPIRE](#)].

- [34] S.M. Aybat and T.C. Rogers, *TMD Parton Distribution and Fragmentation Functions with QCD Evolution*, *Phys. Rev. D* **83** (2011) 114042 [[arXiv:1101.5057](#)] [[INSPIRE](#)].
- [35] M.G. Echevarria, A. Idilbi and I. Scimemi, *Factorization Theorem For Drell-Yan At Low q_T And Transverse Momentum Distributions On-The-Light-Cone*, *JHEP* **07** (2012) 002 [[arXiv:1111.4996](#)] [[INSPIRE](#)].
- [36] M.G. Echevarria, A. Idilbi, A. Schäfer and I. Scimemi, *Model-Independent Evolution of Transverse Momentum Dependent Distribution Functions (TMDs) at NNLL*, *Eur. Phys. J. C* **73** (2013) 2636 [[arXiv:1208.1281](#)] [[INSPIRE](#)].
- [37] J.C. Collins and T.C. Rogers, *Equality of Two Definitions for Transverse Momentum Dependent Parton Distribution Functions*, *Phys. Rev. D* **87** (2013) 034018 [[arXiv:1210.2100](#)] [[INSPIRE](#)].
- [38] H.H. Matevosyan, W. Bentz, I.C. Cloet and A.W. Thomas, *Transverse Momentum Dependent Fragmentation and Quark Distribution Functions from the NJL-jet Model*, *Phys. Rev. D* **85** (2012) 014021 [[arXiv:1111.1740](#)] [[INSPIRE](#)].
- [39] M. Boglione, J. Collins, L. Gamberg, J.O. Gonzalez-Hernandez, T.C. Rogers and N. Sato, *Kinematics of Current Region Fragmentation in Semi-Inclusive Deeply Inelastic Scattering*, *Phys. Lett. B* **766** (2017) 245 [[arXiv:1611.10329](#)] [[INSPIRE](#)].
- [40] E. Moffat, W. Melnitchouk, T.C. Rogers and N. Sato, *What are the low- Q and large- x boundaries of collinear QCD factorization theorems?*, *Phys. Rev. D* **95** (2017) 096008 [[arXiv:1702.03955](#)] [[INSPIRE](#)].
- [41] M.G. Echevarria, I. Scimemi and A. Vladimirov, *Unpolarized Transverse Momentum Dependent Parton Distribution and Fragmentation Functions at next-to-next-to-leading order*, *JHEP* **09** (2016) 004 [[arXiv:1604.07869](#)] [[INSPIRE](#)].
- [42] D. Boer and W. Vogelsang, *Drell-Yan lepton angular distribution at small transverse momentum*, *Phys. Rev. D* **74** (2006) 014004 [[hep-ph/0604177](#)] [[INSPIRE](#)].
- [43] S. Arnold, A. Metz and M. Schlegel, *Dilepton production from polarized hadron hadron collisions*, *Phys. Rev. D* **79** (2009) 034005 [[arXiv:0809.2262](#)] [[INSPIRE](#)].
- [44] PARTICLE DATA GROUP collaboration, C. Patrignani et al., *Review of Particle Physics*, *Chin. Phys. C* **40** (2016) 100001 [[INSPIRE](#)].
- [45] M. Lambertsen and W. Vogelsang, *Drell-Yan lepton angular distributions in perturbative QCD*, *Phys. Rev. D* **93** (2016) 114013 [[arXiv:1605.02625](#)] [[INSPIRE](#)].
- [46] G. Parisi and R. Petronzio, *Small Transverse Momentum Distributions in Hard Processes*, *Nucl. Phys. B* **154** (1979) 427 [[INSPIRE](#)].
- [47] G. Altarelli, R.K. Ellis, M. Greco and G. Martinelli, *Vector Boson Production at Colliders: A Theoretical Reappraisal*, *Nucl. Phys. B* **246** (1984) 12 [[INSPIRE](#)].
- [48] G. Bozzi, S. Catani, G. Ferrera, D. de Florian and M. Grazzini, *Production of Drell-Yan lepton pairs in hadron collisions: Transverse-momentum resummation at next-to-next-to-leading logarithmic accuracy*, *Phys. Lett. B* **696** (2011) 207 [[arXiv:1007.2351](#)] [[INSPIRE](#)].
- [49] D. Boer and W.J. den Dunnen, *TMD evolution and the Higgs transverse momentum distribution*, *Nucl. Phys. B* **886** (2014) 421 [[arXiv:1404.6753](#)] [[INSPIRE](#)].
- [50] E. Laenen, G.F. Sterman and W. Vogelsang, *Higher order QCD corrections in prompt photon production*, *Phys. Rev. Lett.* **84** (2000) 4296 [[hep-ph/0002078](#)] [[INSPIRE](#)].

- [51] J.-w. Qiu and X.-f. Zhang, *Role of the nonperturbative input in QCD resummed Drell-Yan Q_T distributions*, *Phys. Rev. D* **63** (2001) 114011 [[hep-ph/0012348](#)] [[INSPIRE](#)].
- [52] C.T.H. Davies and W.J. Stirling, *Nonleading Corrections to the Drell-Yan Cross-Section at Small Transverse Momentum*, *Nucl. Phys. B* **244** (1984) 337 [[INSPIRE](#)].
- [53] S. Frixione, P. Nason and G. Ridolfi, *Problems in the resummation of soft gluon effects in the transverse momentum distributions of massive vector bosons in hadronic collisions*, *Nucl. Phys. B* **542** (1999) 311 [[hep-ph/9809367](#)] [[INSPIRE](#)].
- [54] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC*, *Nucl. Phys. B* **737** (2006) 73 [[hep-ph/0508068](#)] [[INSPIRE](#)].
- [55] P.M. Nadolsky, D.R. Stump and C.P. Yuan, *Semiinclusive hadron production at HERA: The effect of QCD gluon resummation*, *Phys. Rev. D* **61** (2000) 014003 [*Erratum ibid.* **D 64** (2001) 059903] [[hep-ph/9906280](#)] [[INSPIRE](#)].
- [56] F. Landry, R. Brock, P.M. Nadolsky and C.P. Yuan, *Tevatron Run-1 Z boson data and Collins-Soper-Sterman resummation formalism*, *Phys. Rev. D* **67** (2003) 073016 [[hep-ph/0212159](#)] [[INSPIRE](#)].
- [57] A.V. Konychev and P.M. Nadolsky, *Universality of the Collins-Soper-Sterman nonperturbative function in gauge boson production*, *Phys. Lett. B* **633** (2006) 710 [[hep-ph/0506225](#)] [[INSPIRE](#)].
- [58] C.A. Aidala, B. Field, L.P. Gamberg and T.C. Rogers, *Limits on transverse momentum dependent evolution from semi-inclusive deep inelastic scattering at moderate Q* , *Phys. Rev. D* **89** (2014) 094002 [[arXiv:1401.2654](#)] [[INSPIRE](#)].
- [59] J. Collins and T. Rogers, *Understanding the large-distance behavior of transverse-momentum-dependent parton densities and the Collins-Soper evolution kernel*, *Phys. Rev. D* **91** (2015) 074020 [[arXiv:1412.3820](#)] [[INSPIRE](#)].
- [60] I. Scimemi and A. Vladimirov, *Power corrections and renormalons in Transverse Momentum Distributions*, *JHEP* **03** (2017) 002 [[arXiv:1609.06047](#)] [[INSPIRE](#)].
- [61] U. D'Alesio, M.G. Echevarria, S. Melis and I. Scimemi, *Non-perturbative QCD effects in q_T spectra of Drell-Yan and Z-boson production*, *JHEP* **11** (2014) 098 [[arXiv:1407.3311](#)] [[INSPIRE](#)].
- [62] M. Gluck, P. Jimenez-Delgado and E. Reya, *Dynamical parton distributions of the nucleon and very small- x physics*, *Eur. Phys. J. C* **53** (2008) 355 [[arXiv:0709.0614](#)] [[INSPIRE](#)].
- [63] A. Buckley et al., *LHAPDF6: parton density access in the LHC precision era*, *Eur. Phys. J. C* **75** (2015) 132 [[arXiv:1412.7420](#)] [[INSPIRE](#)].
- [64] D. de Florian, R. Sassot, M. Epele, R.J. Hernández-Pinto and M. Stratmann, *Parton-to-Pion Fragmentation Reloaded*, *Phys. Rev. D* **91** (2015) 014035 [[arXiv:1410.6027](#)] [[INSPIRE](#)].
- [65] D. de Florian, R. Sassot and M. Stratmann, *Global analysis of fragmentation functions for pions and kaons and their uncertainties*, *Phys. Rev. D* **75** (2007) 114010 [[hep-ph/0703242](#)] [[INSPIRE](#)].
- [66] D. de Florian, M. Epele, R.J. Hernandez-Pinto, R. Sassot and M. Stratmann, *Parton-to-Kaon Fragmentation Revisited*, *Phys. Rev. D* **95** (2017) 094019 [[arXiv:1702.06353](#)] [[INSPIRE](#)].

- [67] A. Bacchetta, L.P. Gamberg, G.R. Goldstein and A. Mukherjee, *Collins fragmentation function for pions and kaons in a spectator model*, *Phys. Lett. B* **659** (2008) 234 [[arXiv:0707.3372](#)] [[INSPIRE](#)].
- [68] B. Pasquini, S. Cazzaniga and S. Boffi, *Transverse momentum dependent parton distributions in a light-cone quark model*, *Phys. Rev. D* **78** (2008) 034025 [[arXiv:0806.2298](#)] [[INSPIRE](#)].
- [69] H. Avakian, A.V. Efremov, P. Schweitzer and F. Yuan, *The transverse momentum dependent distribution functions in the bag model*, *Phys. Rev. D* **81** (2010) 074035 [[arXiv:1001.5467](#)] [[INSPIRE](#)].
- [70] A. Bacchetta, M. Radici, F. Conti and M. Guagnelli, *Weighted azimuthal asymmetries in a diquark spectator model*, *Eur. Phys. J. A* **45** (2010) 373 [[arXiv:1003.1328](#)] [[INSPIRE](#)].
- [71] M. Burkardt and B. Pasquini, *Modelling the nucleon structure*, *Eur. Phys. J. A* **52** (2016) 161 [[arXiv:1510.02567](#)] [[INSPIRE](#)].
- [72] A. Bacchetta, R. Kundu, A. Metz and P.J. Mulders, *Estimate of the Collins fragmentation function in a chiral invariant approach*, *Phys. Rev. D* **65** (2002) 094021 [[hep-ph/0201091](#)] [[INSPIRE](#)].
- [73] P. Schweitzer, M. Strikman and C. Weiss, *Intrinsic transverse momentum and parton correlations from dynamical chiral symmetry breaking*, *JHEP* **01** (2013) 163 [[arXiv:1210.1267](#)] [[INSPIRE](#)].
- [74] HERMES collaboration, A. Airapetian et al., *Multiplicities of charged pions and kaons from semi-inclusive deep-inelastic scattering by the proton and the deuteron*, *Phys. Rev. D* **87** (2013) 074029 [[arXiv:1212.5407](#)] [[INSPIRE](#)].
- [75] COMPASS collaboration, C. Adolph et al., *Hadron Transverse Momentum Distributions in Muon Deep Inelastic Scattering at 160 GeV/c*, *Eur. Phys. J. C* **73** (2013) 2531 [*Erratum ibid.* **C 75** (2015) 94] [[arXiv:1305.7317](#)] [[INSPIRE](#)].
- [76] M. Anselmino, M. Boglione, J.O. Gonzalez Hernandez, S. Melis and A. Prokudin, *Unpolarised Transverse Momentum Dependent Distribution and Fragmentation Functions from SIDIS Multiplicities*, *JHEP* **04** (2014) 005 [[arXiv:1312.6261](#)] [[INSPIRE](#)].
- [77] F. Landry, R. Brock, G. Ladinsky and C.P. Yuan, *New fits for the nonperturbative parameters in the CSS resummation formalism*, *Phys. Rev. D* **63** (2001) 013004 [[hep-ph/9905391](#)] [[INSPIRE](#)].
- [78] U. D'Alesio, M.G. Echevarria, S. Melis and I. Scimemi, *Non-perturbative QCD effects in qT spectra of Drell-Yan and Z-boson production*, *JHEP* **11** (2014) 098 [[arXiv:1407.3311](#)] [[INSPIRE](#)].
- [79] A.S. Ito et al., *Measurement of the Continuum of Dimuons Produced in High-Energy Proton-Nucleus Collisions*, *Phys. Rev. D* **23** (1981) 604 [[INSPIRE](#)].
- [80] G. Moreno et al., *Dimuon production in proton-copper collisions at $\sqrt{s} = 38.8\text{-GeV}$* , *Phys. Rev. D* **43** (1991) 2815 [[INSPIRE](#)].
- [81] CDF collaboration, T. Affolder et al., *The transverse momentum and total cross section of e^+e^- pairs in the Z boson region from $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$* , *Phys. Rev. Lett.* **84** (2000) 845 [[hep-ex/0001021](#)] [[INSPIRE](#)].
- [82] D0 collaboration, B. Abbott et al., *Measurement of the inclusive differential cross section for Z bosons as a function of transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8\text{ TeV}$* , *Phys. Rev. D* **61** (2000) 032004 [[hep-ex/9907009](#)] [[INSPIRE](#)].

- [83] CDF collaboration, T. Aaltonen et al., *Transverse momentum cross section of e^+e^- pairs in the Z -boson region from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. D* **86** (2012) 052010 [[arXiv:1207.7138](#)] [[INSPIRE](#)].
- [84] D0 collaboration, V.M. Abazov et al., *Measurement of the shape of the boson transverse momentum distribution in $p\bar{p} \rightarrow Z/\gamma^* \rightarrow e^+e^- + X$ events produced at $\sqrt{s} = 1.96$ -TeV*, *Phys. Rev. Lett.* **100** (2008) 102002 [[arXiv:0712.0803](#)] [[INSPIRE](#)].
- [85] CDF collaboration, A. Abulencia et al., *Measurements of inclusive W and Z cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV*, *J. Phys. G* **34** (2007) 2457 [[hep-ex/0508029](#)] [[INSPIRE](#)].
- [86] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector boson production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **103** (2009) 082001 [[arXiv:0903.2120](#)] [[INSPIRE](#)].
- [87] A. Bacchetta, A. Courtoy and M. Radici, *First extraction of valence transversities in a collinear framework*, *JHEP* **03** (2013) 119 [[arXiv:1212.3568](#)] [[INSPIRE](#)].
- [88] M. Radici, A. Courtoy, A. Bacchetta and M. Guagnelli, *Improved extraction of valence transversity distributions from inclusive dihadron production*, *JHEP* **05** (2015) 123 [[arXiv:1503.03495](#)] [[INSPIRE](#)].
- [89] S. Forte, L. Garrido, J.I. Latorre and A. Piccione, *Neural network parametrization of deep inelastic structure functions*, *JHEP* **05** (2002) 062 [[hep-ph/0204232](#)] [[INSPIRE](#)].
- [90] NNPDF collaboration, R.D. Ball et al., *A determination of parton distributions with faithful uncertainty estimation*, *Nucl. Phys. B* **809** (2009) 1 [Erratum *ibid.* **B 816** (2009) 293] [[arXiv:0808.1231](#)] [[INSPIRE](#)].
- [91] R.D. Ball et al., *A first unbiased global NLO determination of parton distributions and their uncertainties*, *Nucl. Phys. B* **838** (2010) 136 [[arXiv:1002.4407](#)] [[INSPIRE](#)].
- [92] M. Epele, R. Llubaroff, R. Sassot and M. Stratmann, *Uncertainties in pion and kaon fragmentation functions*, *Phys. Rev. D* **86** (2012) 074028 [[arXiv:1209.3240](#)] [[INSPIRE](#)].
- [93] F. James and M. Roos, *Minuit: A System for Function Minimization and Analysis of the Parameter Errors and Correlations*, *Comput. Phys. Commun.* **10** (1975) 343 [[INSPIRE](#)].
- [94] H1 collaboration, S. Aid et al., *Transverse energy and forward jet production in the low x regime at HERA*, *Phys. Lett. B* **356** (1995) 118 [[hep-ex/9506012](#)] [[INSPIRE](#)].
- [95] G.A. Ladinsky and C.P. Yuan, *The Nonperturbative regime in QCD resummation for gauge boson production at hadron colliders*, *Phys. Rev. D* **50** (1994) R4239 [[hep-ph/9311341](#)] [[INSPIRE](#)].
- [96] P. Sun, J. Isaacson, C.P. Yuan and F. Yuan, *Universal Non-perturbative Functions for SIDIS and Drell-Yan Processes*, [arXiv:1406.3073](#) [[INSPIRE](#)].
- [97] P. Schweitzer, T. Teckentrup and A. Metz, *Intrinsic transverse parton momenta in deeply inelastic reactions*, *Phys. Rev. D* **81** (2010) 094019 [[arXiv:1003.2190](#)] [[INSPIRE](#)].
- [98] A.D. Martin, W.J. Stirling, R.S. Thorne and G. Watt, *Parton distributions for the LHC*, *Eur. Phys. J. C* **63** (2009) 189 [[arXiv:0901.0002](#)] [[INSPIRE](#)].
- [99] J.F. Owens, A. Accardi and W. Melnitchouk, *Global parton distributions with nuclear and finite- Q^2 corrections*, *Phys. Rev. D* **87** (2013) 094012 [[arXiv:1212.1702](#)] [[INSPIRE](#)].
- [100] F. Hautmann et al., *TMDlib and TMDplotter: library and plotting tools for transverse-momentum-dependent parton distributions*, *Eur. Phys. J. C* **74** (2014) 3220 [[arXiv:1408.3015](#)] [[INSPIRE](#)].

- [101] J. Dudek et al., *Physics Opportunities with the 12 GeV Upgrade at Jefferson Lab*, *Eur. Phys. J. A* **48** (2012) 187 [[arXiv:1208.1244](#)] [[INSPIRE](#)].
- [102] S.J. Brodsky, F. Fleuret, C. Hadjidakis and J.P. Lansberg, *Physics Opportunities of a Fixed-Target Experiment using the LHC Beams*, *Phys. Rept.* **522** (2013) 239 [[arXiv:1202.6585](#)] [[INSPIRE](#)].
- [103] M.G. Echevarria, T. Kasemets, P.J. Mulders and C. Pisano, *QCD evolution of (un)polarized gluon TMDPDFs and the Higgs q_T -distribution*, *JHEP* **07** (2015) 158 [[arXiv:1502.05354](#)] [[INSPIRE](#)].
- [104] D. Boer, S.J. Brodsky, P.J. Mulders and C. Pisano, *Direct Probes of Linearly Polarized Gluons inside Unpolarized Hadrons*, *Phys. Rev. Lett.* **106** (2011) 132001 [[arXiv:1011.4225](#)] [[INSPIRE](#)].
- [105] C. Pisano, D. Boer, S.J. Brodsky, M.G.A. Buffing and P.J. Mulders, *Linear polarization of gluons and photons in unpolarized collider experiments*, *JHEP* **10** (2013) 024 [[arXiv:1307.3417](#)] [[INSPIRE](#)].
- [106] J.-W. Qiu, M. Schlegel and W. Vogelsang, *Probing Gluonic Spin-Orbit Correlations in Photon Pair Production*, *Phys. Rev. Lett.* **107** (2011) 062001 [[arXiv:1103.3861](#)] [[INSPIRE](#)].
- [107] D. Boer and C. Pisano, *Polarized gluon studies with charmonium and bottomonium at LHCb and AFTER*, *Phys. Rev. D* **86** (2012) 094007 [[arXiv:1208.3642](#)] [[INSPIRE](#)].
- [108] W.J. den Dunnen, J.P. Lansberg, C. Pisano and M. Schlegel, *Accessing the Transverse Dynamics and Polarization of Gluons inside the Proton at the LHC*, *Phys. Rev. Lett.* **112** (2014) 212001 [[arXiv:1401.7611](#)] [[INSPIRE](#)].
- [109] A. Signori, *Gluon TMDs in quarkonium production*, *Few Body Syst.* **57** (2016) 651 [[arXiv:1602.03405](#)] [[INSPIRE](#)].
- [110] J.-P. Lansberg, C. Pisano and M. Schlegel, *Associated production of a dilepton and a $\Upsilon(J/\psi)$ at the LHC as a probe of gluon transverse momentum dependent distributions*, *Nucl. Phys. B* **920** (2017) 192 [[arXiv:1702.00305](#)] [[INSPIRE](#)].
- [111] E.C. Aschenauer, U. D'Alesio and F. Murgia, *TMDs and SSAs in hadronic interactions*, *Eur. Phys. J. A* **52** (2016) 156 [[arXiv:1512.05379](#)] [[INSPIRE](#)].
- [112] M. Boglione and A. Prokudin, *Phenomenology of transverse spin: past, present and future*, *Eur. Phys. J. A* **52** (2016) 154 [[arXiv:1511.06924](#)] [[INSPIRE](#)].