

Integrability of conformal blocks. Part I. Calogero-Sutherland scattering theory

Mikhail Isachenkov^a and Volker Schomerus^b

^a*Department of Particle Physics and Astrophysics, Weizmann Institute of Science,
Herzl street 234, Rehovot 7610001, Israel*

^b*DESY Theory Group, DESY Hamburg,
Notkestraße 85, D-22607 Hamburg, Germany*

E-mail: mikhail.isachenkov@weizmann.ac.il, volker.schomerus@desy.de

ABSTRACT: Conformal blocks are the central ingredient of the conformal bootstrap programme. We elaborate on our recent observation that uncovered a relation with wave functions of an integrable Calogero-Sutherland Hamiltonian in order to develop a systematic theory of conformal blocks. Our main goal here is to review central ingredients of the Heckman-Opdam theory for scattering states of Calogero-Sutherland models with special emphasis to the relation with scalar 4-point blocks. We will also discuss a number of direct consequences for conformal blocks, including a new series expansion for blocks of arbitrary complex spin and a complete analysis of their poles and residues. Applications to the Froissart-Gribov formula for conformal field theory, as well as extensions to spinning blocks and defects are briefly discussed before we conclude with an outlook on forthcoming work concerning algebraic consequences of integrability.

KEYWORDS: Conformal and W Symmetry, Conformal Field Theory, Field Theories in Higher Dimensions

ARXIV EPRINT: [1711.06609](https://arxiv.org/abs/1711.06609)

Contents

1	Introduction	1
2	The BC_N Calogero-Sutherland problem	4
2.1	The Pöschl-Teller Hamiltonian	4
2.2	The Calogero-Sutherland potential	6
2.3	Affine Weyl group and the domain D_N	8
2.4	Coordinates in the CS problem	12
3	Wave functions of the Calogero-Sutherland model	12
3.1	Wave functions of the Pöschl-Teller problem	13
3.2	Harish-Chandra series expansions	17
3.3	Poles and residues of Harish-Chandra functions	20
3.4	Monodromy group and representation	24
4	Wave functions and conformal blocks	31
4.1	Conformal blocks and the Casimir equation	31
4.2	Blocks and the Pöschl-Teller problem	32
4.3	Blocks from Harish-Chandra wave functions	33
5	Some applications of scattering theory to conformal blocks	35
5.1	Poles and residues of blocks	36
5.2	On the derivation of a Gribov-Froissart formula for CFT	41
6	Conclusions	45
A	z- and x-expansions for BC_2 Harish-Chandra function	49
B	Alternative expansions for integer spin	55

1 Introduction

The conformal bootstrap programme [1–6] was initially designed as an analytical approach to the non-perturbative dynamics of critical systems. It relies on the careful separation of kinematical (or group theoretic) input from dynamical data. In spite of significant early efforts to develop the necessary background in representation theory of the conformal group, see [7] and references therein, concrete implementations of the bootstrap programme suffered from the fact that much of the relevant mathematics could not be developed at the time. It took more than a decade before the real impact of the conformal bootstrap was first demonstrated in the context of 2-dimensional conformal field theories where the

conformal symmetry is enhanced to the infinite dimensional Virasoro algebra [8]. Up until a few years ago, it was commonly assumed that such a success of the bootstrap programme was only possible in 2-dimensional systems. But since the recent numerical incarnation of the conformal bootstrap programme has delivered data e.g. on scaling weights and operator products in the $d = 3$ dimensional Ising model with unprecedented precision, see [9–15] and [16–23] for some similar results in other theories, the conformal bootstrap has attracted new attention.

The key kinematical data in the conformal bootstrap are the conformal blocks along with the so-called crossing kernel. Explicit analytical results on conformal blocks in $d > 2$ dimensional were scarce until the work of Dolan and Osborn [24–26] on scalar conformal blocks. In a few cases, such as for four scalar external fields in even dimensions, Dolan and Osborn were able to construct blocks explicitly in terms of ordinary single variable hypergeometric functions. Extensions to generic dimensions and external field with spin or general defect blocks proved more difficult, even though some remarkable progress has been achieved during the last few years, e.g. through the use of differential operators, the concept and construction of seed blocks etc., see e.g. [27–40] and references therein. In most cases, however, a construction of blocks in terms of ordinary hypergeometric functions could not be found. In order to evaluate such more general blocks, Zamolodchikov-like recurrence relations have become the most efficient tool. While these may suffice to provide the required input for the numerical bootstrap, it is fair to say that a systematic and universal theory of conformal blocks has not been developed to date. It is our main goal to fill this important gap.

In some sense, the mathematical foundations for a modern and systematic theory of conformal blocks, including those for external fields with spin, were actually laid at about the same time at which the bootstrap programme was formulated, though very much disguised at first. It gradually emerged from the systematic study of solvable Schrödinger problems starting with the work by Calogero, Moser and Sutherland [41–43]. The quantum mechanical models that were proposed in these papers describe a 1-dimensional multi-particle system whose members are subject to an external potential and exhibit pairwise interaction. It turned out that for appropriate choices of the potentials and interactions, such models can be integrable. One distinguishes two important series of such theories known as A and BC models. While the former describe particles that move on the entire real line, particles are restricted to the half-line in the case of BC -type models.

The study of eigenfunctions for Calogero-Sutherland (CS) models advanced rapidly after a very influential series of papers by Heckman and Opdam that was initiated in [44–47] (inspired by earlier papers of Koornwinder [48–51]) and provided the basis for much of the modern theory of multivariable hypergeometric functions. Subsequently, many different approaches were developed that emphasize algebraic aspects (most notably due to Cherednik [52]), combinatorial identities [53] or relations to matrix models. The general techniques that were developed in this context are fairly universal. On the other hand, explicit formulas were often only worked out for A models, with BC trailing a bit behind.

These two different strands of seemingly unrelated developments were brought together by our recent observation [54] that conformal blocks of scalar 4-point functions in a

d -dimensional conformal field theory can be mapped to eigenfunctions of a 2-particle hyperbolic Calogero-Sutherland Hamiltonian. Thereby, the modern theory of multivariable hypergeometric functions and its integrable foundation enters the court of the conformal bootstrap. This is the first of a series of papers in which we describe, and at various places advance, the mathematical theory of Calogero-Sutherland models and develop the applications to conformal blocks. Here we shall focus mainly on the classical Heckman-Opdam theory for scattering states of Calogero-Sutherland models. Algebraic consequences of integrability as well as advanced analytical features are subject of a subsequent paper [55], see also concluding section 6 for a detailed outline.

The plan of this paper is as follows. In the [next section](#) we introduce the relevant hyperbolic Calogero-Sutherland models for BC_N root systems. After setting up some notations, we spell out the [Hamiltonian](#) and describe its [symmetries](#), the associated [fundamental domain](#) and different [coordinate choices](#). Then we turn to the scattering theory. In section 3 we introduce the notion of [Harish-Chandra wave functions](#) and discuss their analytic properties both in [coordinate space](#) and in the [space of eigenvalues](#). The former are controlled by a special class of [representations of an affine braid group](#) which is discussed in detail. This will allow us to construct the true wave functions of Calogero-Sutherland models as special linear combinations of Harish-Chandra functions. We also describe the position of [poles of Harish-Chandra functions](#) in the space of eigenvalues and provide explicit formulas for their residues, at least for $N = 2$. The latter have not appeared in the mathematical literature before and they are obtained from a new series expansion for Harish-Chandra functions that we derive in appendix A. The aim of section 4 is to embed [scalar conformal blocks](#) into the general theory of Calogero-Sutherland wave functions. Our discussion includes details on the choice of boundary conditions. This will enable us to discuss a number of direct applications to scalar blocks in section 5. These include a full classification of [poles of conformal blocks](#) for arbitrary (complex) values of the spin variable l . We also calculate the corresponding residues. When the spin variable is specialized to an integer (and the dimension d is integer), our results for poles and residues agree with those that were obtained using representation theory of the conformal Lie algebra [32]. The extension to complex spins may be seen as the main new advance of this work in the context of conformal field theory. Blocks with non-integer spin play an important role in the recent conformal Froissart-Gribov formula of Caron-Huot [56] which, in particular, was used to establish existence in any CFT of operator families analytic in spin, see also [57–71]. We will sketch how such [inversion formulas](#) arise within the theory of Calogero-Sutherland models and employ the algebraic structure of the monodromy representation to explain a crucial numerical ‘coincidence’ in Caron-Huot’s derivation of the Froissart-Gribov formula. The paper concludes with an extensive outlook, in particular to the second part where we will discuss and exploit the rich algebraic structure of Calogero-Sutherland models. While some of the more advanced results we describe are geared to the root system BC_2 which appears in the context of scalar 4-point functions, most of the more general discussion is presented for general N . Larger values $N > 2$ turn out to be relevant [72] in the context of defect blocks [73, 74].

2 The BC_N Calogero-Sutherland problem

In this section we shall review the setup of the Calogero-Sutherland problem for the BC_N root system. After a short discussion the simplest example, the famous Pöschl-Teller potential, we discuss the general setup. In our description we will put some emphasis on the symmetries of the Calogero-Sutherland potential, possible domains for the associated Schrödinger problem and the singularities at the boundary of these domains.

2.1 The Pöschl-Teller Hamiltonian

The simplest example of what is now known as Calogero-Sutherland model goes back to the work of Pöschl and Teller in [75]. The so-called modified Pöschl-Teller Hamiltonian takes the form

$$H_{(a,b)}^{\text{PT}} = -\partial_u^2 + V_{(a,b)}^{\text{PT}}(u) = -\partial_u^2 - \frac{ab}{\sinh^2 \frac{u}{2}} + \frac{(a+b)^2 - \frac{1}{4}}{\sinh^2 u}. \quad (2.1)$$

and defines a 1-dimensional Schrödinger problem with a potential that depends on two continuous parameters a and b . Pöschl and Teller noticed that the corresponding eigenvalue problem can be mapped to the hypergeometric differential equation and constructed the eigenfunctions in terms of hypergeometric functions. All this is fairly standard, but there are a few things we would like to emphasize in this example that will become important for extensions to the multi-particle generalizations.

For the moment we will consider u as a complex variable. In the complex u -plane, the Pöschl-Teller potential possesses some symmetries. On the one hand it is symmetric with respect to shifts $\tau : u \rightarrow u + 2\pi i$ of u in the imaginary direction. These give rise to an action of \mathbb{Z} on the complex plane. In addition, the potential is also reflection symmetric, i.e. it is invariant under the \mathbb{Z}_2 reflection $w : u \rightarrow -u$. Together, these two transformations generate the symmetry group $\mathcal{W} = \mathbb{Z}_2 \times \mathbb{Z}$ of the Pöschl-Teller potential. The fundamental domain

$$D = \mathbb{C}/\mathcal{W}$$

for the action of \mathcal{W} in the complex u -plane is shown in figure 1. After the appropriate identifications of boundary points it has the form of a semi-infinite pillow, i.e. a semi-infinite cylinder whose end is squashed to an interval. The two corners of this semi-infinite pillow correspond to the two points $u_1 = 0$ and $u_0 = i\pi$ at which the Pöschl-Teller potential diverges. At the same time, these points are fixed under the action of a non-trivial subgroup of the symmetry \mathcal{W} on the complex u -plane.¹

The group \mathbb{W} exhausts all the symmetries of the Pöschl-Teller potential that are associated with a transformation of the variable u alone. But there exists one additional

¹When $b = 0$, the fundamental domain is further reduced to $D' = \mathbb{C}/(\mathcal{W} \rtimes \mathbb{Z}_2)$. In this case, the complex torus has non-trivial (so called) center [44, 45], which is related to existence of a *minuscule weight* for the reduced root system C_1 . The denominator $\mathcal{W} \rtimes \mathbb{Z}_2$ is called an extended affine Weyl group, and the additional \mathbb{Z}_2 accounts for a permutation of an affine and a non-affine simple root. The reduced fundamental domain D' has $\Delta \Im u = \pi$, instead of $\Delta \Im u = 2\pi$. When $b \neq 0$, there is a remnant of this symmetry which we call ϱ , see below.




Figure 1. The fundamental domain D of the Pöschl-Teller problem is a semi-infinite cylinder whose end is squashed to an interval. It is obtained from the semi-infinite strip by gluing the horizontal lines on top and bottom. Within this domain, the potential diverges at two points $u_0 = i\pi$ and $u_1 = 0$ (black dots). Different Schrödinger problems can be obtained, depending on the choice of a real subset A . The figure shows three possible choices, A^+ , $A^L = \tilde{A}^+$ and A^C . The eigenvalue equations on A^+ and A^L are related by a shift ϱ that involves the reflection of the coupling constant b and they both lead to a continuous spectrum. The 1-dimensional Schrödinger problem on the compact interval A^C possesses a discrete spectrum.

symmetry that involves a shift in u combined with a reflection of the coupling b .² More precisely, it acts as

$$\varrho : u \rightarrow u + i\pi, \quad b \rightarrow -b \tag{2.2}$$

while leaving the couplings a invariant, i.e. $\tilde{a} = a$. In fact, the Pöschl-Teller potential is invariant under these replacements

$$\begin{aligned} \varrho : V_{(a,b)}^{\text{PT}}(u) \rightarrow V_{(a,-b)}^{\text{PT}}(u + i\pi) &= -\frac{ab}{\cosh^2 \frac{u}{2}} + \frac{(a-b)^2 - \frac{1}{4}}{\sinh^2 u} \\ &= -\frac{ab}{\sinh^2 \frac{u}{2}} + \frac{(a+b)^2 - \frac{1}{4}}{\sinh^2 u} = V_{(a,b)}^{\text{PT}}(u). \end{aligned}$$

Note that the symmetry maps the two singular points $u_0 = 0$ and $u_1 = i\pi$ in the fundamental domain D onto each other.

After these comments on the symmetries of the Pöschl-Teller potential let us now discuss the possible setups for the Schrödinger problem. These correspond to different

²This symmetry is a twist of an ordinary translation symmetry of the C_1 Pöschl-Teller problem, where it acts on coordinates only since the parameter b vanishes. It may be regarded as a translation of the coordinate u by a $2\pi i$ times a minuscule coweight of the root system C_1 , see below.

1-dimensional subsets $A \subset D$ on which the Pöschl-Teller potential is real. There are essentially three such choices which we shall denote by A^+ , \tilde{A}^+ and A^C . We define the set A^+ as

$$A^+ = \{ u \in \mathbb{R} \mid u > 0 \} . \tag{2.3}$$

For this set, the Schrödinger equation reads

$$\left[-\frac{d^2}{du^2} - \frac{ab}{\sinh^2 \frac{u}{2}} + \frac{(a+b)^2 - \frac{1}{4}}{\sinh^2 u} \right] \psi(u) = \varepsilon \psi(u) . \tag{2.4}$$

Note that the potential creates a wall at $u = 0$ which shields the positive half-line $u > 0$ from the negative one. In this case, there is a continuum of states with energy $\varepsilon > 0$. We will discuss these in the [next section](#). A second possible choice is

$$\tilde{A}^+ = \{ \tilde{u} + i\pi \mid \tilde{u} \in \mathbb{R}_{>0} \} . \tag{2.5}$$

On this set, the corresponding Schrödinger equation is the same as on A^+ , except that the parameter b is sent to $-b$. Our previous comments on the structure of the spectrum apply to this case as well.

There exists a qualitatively quite different choice for a domain on which the Calogero-Sutherland Hamiltonian is real. It is given by

$$A^C = \{ u = i\varphi \mid \varphi \in [0, \pi] \} \tag{2.6}$$

where the superscript C stands for compact. On A^C the Schrödinger equation takes the form

$$\left[-\frac{d^2}{d\varphi^2} - \frac{ab}{\sin^2 \frac{\varphi}{2}} + \frac{(a+b)^2 - \frac{1}{4}}{\sin^2 \varphi} \right] \psi(\varphi) = \varepsilon \psi(\varphi) . \tag{2.7}$$

Once again, there are infinite walls at $\varphi = 0, \pi$. This setup describes a particle in a 1-dimensional box with infinitely high walls on both sides. In this case the spectrum is discrete. We will recall the precise form of the wave functions in the [next section](#).

2.2 The Calogero-Sutherland potential

Before we can spell out the Calogero-Sutherland Hamiltonian, we need a bit of notation. In general integrable Calogero-Sutherland Hamiltonians are associated with a root system Σ . Here we shall focus on BC_N root systems, see figure 2, whose positive roots are given

$$\Sigma^+ = \{ e_i, 2e_i, e_i \pm e_j \mid 1 \leq i, j \leq N; i < j \} . \tag{2.8}$$

We have used $e_i, 1 \leq i \leq N$ to denote a basis of \mathbb{R}^N that is orthonormal with respect to the canonical scalar product $\langle \cdot, \cdot \rangle$. Note that this root system is not reduced, i.e. it contains roots that are related by a factor of two.³ From time to time we will have to remove the shortest positive roots, i.e. the roots e_i . The remaining set of positive roots is denoted by

³The integrality condition in the definition of a root system demands that a projection of a root onto any other root is a half-integer multiple of the latter. One can easily see that this can indeed happen if some roots possess a collinear partner differing by a factor of two, but not more.




Figure 2. The BC_2 root system. The set Σ^+ of positive roots is shown in bold black arrows. For $N > 1$ there are three different types of roots that belong to three different orbits of the Weyl group W_N . These carry multiplicity k_α . Our basis elements α_1 and α_2 are also depicted along with the highest root $\alpha_0 = 2\alpha_1 + \alpha_2 = 2e_1$. The linear spaces that are fixed under the Weyl reflections w_1 and w_2 that are associated with α_1 and α_2 form the boundary of the Weyl chamber (shown in grey).

Σ_0^+ . Of course, Σ_0^+ are simply the positive roots of the Lie algebra C_N . Let us also select a basis of Σ_0^+ consisting of

$$\alpha_N = 2e_N, \quad \alpha_i = e_i - e_{i+1}$$

for $i = 1, \dots, N - 1$. Indeed, all elements of Σ_0^+ may be obtained as linear combinations of $\alpha_i, i = 1, \dots, N$ with non-negative integer coefficients. The unique highest root is given by $\alpha_0 = 2e_1$. For any root α , we define

$$\alpha^\vee = \frac{2\alpha}{\langle \alpha, \alpha \rangle}.$$

This concludes our short description of the root system and the special roots that will play an important role below. The potential of the associated Calogero-Sutherland model takes the form

$$V^{\text{CS}}(u_i) = \sum_{\alpha \in \Sigma^+} \frac{k_\alpha(k_\alpha + 2k_{2\alpha} - 1)\langle \alpha, \alpha \rangle}{4 \sinh^2 \frac{\langle \alpha, u \rangle}{2}}. \tag{2.9}$$

It involves the parameters k_α , often referred to as multiplicities, that are assumed to be invariant under the action of the Weyl group $W = W_N$ of Σ , i.e. $k_{w\alpha} = k_\alpha$ for $w \in W$. Since the $BC_N, N > 1$, root lattice decomposes into three orbits under the action of the Weyl group, see figure 2, the potential V contains three independent parameters, which we parametrize as

$$k_3 = k_{e_i \pm e_j} = \frac{\epsilon}{2}, \quad k_2 = k_{2e_i} = a + b + \frac{1}{2}, \quad k_1 = k_{e_i} = -2b \tag{2.10}$$

in terms of the three parameters a, b and ϵ . The reason for this choice of parameters will become clear in the [fourth section](#). In addition, we agree that $k_\beta = 0$ if $\beta \notin \Sigma$. Finally, we have also introduced $u = \sum u_i e_i$. It is easy to see that the formula (2.9) reduces to the Pöschl-Teller potential upon setting $N = 1$. Note that the root system for BC_1 consists of two orbits under the action of the Weyl group $W = W_1 = \mathbb{Z}_2$. Hence, the potential only contains two parameters, a and b . Let us also note that the case $b = 0$ is somewhat special since there are no contributions from the short roots e_i in the potential. This means that the underlying root system is C_N rather than BC_N .

Example. The case we are most interested in appears for $N = 2$. The corresponding Calogero-Sutherland potential now contains all three parameters,

$$V_{(a,b,\epsilon)}^{\text{CS}}(u_1, u_2) = V_{(a,b)}^{\text{PT}}(u_1) + V_{(a,b)}^{\text{PT}}(u_2) + \frac{\epsilon(\epsilon - 2)}{8 \sinh^2 \frac{u_1 - u_2}{2}} + \frac{\epsilon(\epsilon - 2)}{8 \sinh^2 \frac{u_1 + u_2}{2}}. \quad (2.11)$$

One may think of this potential as describing two Pöschl-Teller particles in the half-line that are interacting with an interaction strength depending on ϵ . Alternatively, one can think of a single particle that moves in an external potential on a 2-dimensional domain D_2 , see below.

2.3 Affine Weyl group and the domain D_N

The analysis of symmetries of the Calogero-Sutherland potential proceeds pretty much in the same way as for the Pöschl-Teller problem. Once again, we will think of u_i as complex variables so that the potential is a function on \mathbb{C}^N . Obviously, it is invariant under the N independent discrete shifts

$$\tau_i : u_j \longrightarrow u_j + 2\pi i \delta_{i,j}. \quad (2.12)$$

These generate the abelian group \mathbb{Z}^N . In addition, the potential is also left invariant by the action of the Weyl group W_N of the BC_N root system. This Weyl group can be generated by the N Weyl reflections that are associated with our basis $\alpha_i, i = 1, \dots, N$. We shall denote these Weyl reflections by $w_i = w(\alpha_i)$. It is not difficult to work out all relations among these generators. They are given by

$$w_i^2 = 1, \quad w_{N-1} w_N w_{N-1} w_N = w_N w_{N-1} w_N w_{N-1}, \quad w_i w_{i+1} w_i = w_{i+1} w_i w_{i+1} \quad (2.13)$$

for $i = 1, \dots, N - 2$. All other pairs of generators simply commute with each other, i.e.

$$w_i w_j = w_j w_i \quad \text{for } |i - j| \geq 2.$$

The Weyl group W_N acts on the translations τ_i by permutation and inversion. More precisely one has the following set of non-trivial relations

$$w_N \tau_N^{-1} w_N = \tau_N, \quad w_i \tau_{i+1} w_i = \tau_i \quad (2.14)$$

for $i = 1, \dots, N - 1$. Here we used a multiplicative notation for the generators τ_i of the abelian group \mathbb{Z}^N , i.e. we denote the shift of u_i by $-2\pi i$ as τ_i^{-1} rather than $-\tau_i$. Together, the elements τ_i and w_i with $i = 1, \dots, N$ generate the so-called *affine Weyl group*

$$\mathcal{W}_N = W_N \ltimes \mathbb{Z}^N. \quad (2.15)$$

The affine Weyl group \mathcal{W}_N describes all symmetries of the Calogero-Sutherland potential (2.9) that act on the coordinates u_i alone. It generalizes the group $\mathcal{W}_1 = \mathcal{W}$ we had introduced in our discussion of the Pöschl-Teller potential to the case with $N \geq 2$.

Here we have described the affine Weyl group in terms of $2N$ generators τ_i and w_i with $i = 1, \dots, N$. There exists a second description in terms of $N + 1$ generators w_i with $i = 0, \dots, N$. While w_i with $i \neq 0$ are the same Weyl reflections $w_i = w(\alpha_i)$ we used before, the new generator w_0 is given by

$$w_0^{-1} = w(\alpha_0)\tau_1 = w(2e_1)\tau_1 = (w_1 \cdots w_{N-1}w_N)(w_{N-1} \cdots w_1)\tau_1. \quad (2.16)$$

One may check by explicit computation that this new element w_0 satisfies the following relations with $w_i, i = 1, \dots, N$

$$w_0^2 = 1, \quad w_0w_1w_0w_1 = w_1w_0w_1w_0, \quad w_0w_i = w_iw_0 \quad (2.17)$$

for $i = 2, \dots, N$. Note that the relation between w_0 and w_1 is identical to the one between w_{N-1} and w_N . Obviously, one can reconstruct the generator τ_1 from the element w_0 and the Weyl reflections $w_i, i = 1, \dots, N$. The other elements $\tau_i, i > 1$ are then obtained by conjugation with $w_1 \cdots w_{i-1}$.

As in the case of the Pöschl-Teller potential there exists one additional shift symmetry that requires a combined action on the coordinates and the coupling constants. It is given by

$$\varrho : u_j \rightarrow u_j + i\pi, \quad b \rightarrow -b \quad (2.18)$$

while leaving the other two couplings a and ϵ invariant. Note that ϱ involves a simultaneous action on all N coordinates. Following the same steps as in the section 2.1 one finds that

$$\varrho : V_{(a,b,\epsilon)}^{\text{CS}}(u) \rightarrow V_{(a,-b,\epsilon)}^{\text{CS}}(u + i\pi) = V_{(a,b,\epsilon)}^{\text{CS}}(u). \quad (2.19)$$

Let us stress that this symmetry is not part of the affine Weyl group which acts only on coordinates.⁴

We are now well prepared to discuss the domain(s) on which we will consider the Calogero-Sutherland system. We start with a set of N complex coordinates $(u_i) \in \mathbb{C}^N$ and implement the identification furnished by the N discrete shifts τ_i . This leaves us with an N -dimensional complex manifold

$$\mathbb{T}_N = \mathbb{C}^N / \mathbb{Z}^N.$$

The manifold \mathbb{T}_N contains an N -dimensional real submanifold $A_N \subset \mathbb{T}_N$ that is parametrized by $u_i \in \mathbb{R}$, modulo identification with τ_i . We can shift A_N with ϱ to obtain $\tilde{A}_N = \varrho(A_N)$. The latter is parametrized by $u_i = \tilde{u}_i + i\pi$ with real $\tilde{u}_i \in \mathbb{R}$.

The Calogero-Sutherland potential diverges along the following walls of real codimension two

$$w_\alpha = \{u \in \mathbb{T}_N \mid \langle \alpha, u \rangle = 0 \text{ mod } 2\pi i\} \subset A_N \quad \text{for } \alpha \in \Sigma^+$$

⁴As in the Pöschl-Teller case, this symmetry is a twist of an ordinary translation symmetry of the C_N Pöschl-Teller problem for which it acts on the coordinates u only. The latter may be considered as a translation of u by $2\pi i$ times a minuscule coweight of the reduced root system C_N .

that are in one-to-one correspondence with the positive roots of our BC_N roots system. Note that for the long roots $2e_j$, the set ω_{2e_j} possesses two disconnected components, one of which coincides with the wall ω_{e_j} for the corresponding short root. The walls associated with the roots $e_i \pm e_j$ contain a single connected component. By construction, the walls ω_α are invariant under the action of the reflection $w(\alpha) \in W_N$.

The walls we have just described are subspaces along which the Calogero-Sutherland potential diverges so that the corresponding Schrödinger problem can be restricted to various subsets within the quotient space

$$D_N = \mathbb{T}_N/W_N = \mathbb{C}^N/W_N. \quad (2.20)$$

which describes a fundamental domain⁵ for the action of the Weyl group W_N on \mathbb{T}_N . Representatives of the quotient space in \mathbb{T}_N intersect with only $N+1$ walls $\omega_i, i = 0, \dots, N$ where $\omega_i = \omega_{\alpha_i}$ for $i = 1, \dots, N-1$, and ω_0, ω_N are the two disconnected components of ω_{α_N} with $\omega_N = \omega_{e_N}$.

Once again, the fundamental domain D_N for the Calogero-Sutherland problem possesses several subsets along which the potential is real. The most important is the Weyl chamber

$$A_N^+ = WC_N = \{u \in \mathbb{R}^N \mid \langle \alpha, u \rangle > 0 \text{ for all } \alpha \in \Sigma_0^+\} \subset D_N. \quad (2.21)$$

In this case, the eigenfunctions possess N continuous variables. As in the case of the Pöschl-Teller problem, we can also consider the shifted Weyl chamber $\tilde{A}_N^+ = \varrho(A_N^+)$. It gives rise to a similar set of wave functions except that the coupling b must be replaced by $\tilde{b} = -b$. Another extreme possibility is the case

$$A_N^C = \{u_i = i\varphi_i \mid \varphi_i \in [0, \pi]\} \quad (2.22)$$

for which the spectrum of the Calogero-Sutherland model is discrete. But in the multivariable case there are many other possibilities. We will discuss a few of them for $N = 2$.

Example. Let us give some additional details for $N = 2$. In this case, the Weyl group W_2 consists of eight elements. It can be generated from $w_2 = w(2e_2)$ and $w_1 = w(e_1 - e_2)$ subject to the relations

$$w_1^2 = 1 = w_2^2, \quad w_1 w_2 w_1 w_2 = w_2 w_1 w_2 w_1. \quad (2.23)$$

The fundamental domain D_2 for the action of the Weyl group W_2 on \mathbb{T}_2 , or rather a 3-dimensional subspace thereof that satisfies $\Im(u_1 + u_2) = 0$, is shown in figure 3.

Once again, we can consider the Schrödinger equation for the Calogero-Sutherland potential on various real subsets \mathcal{A}_2 . The most standard choice in the mathematical literature is

$$A_2^+ = \{(u_1, u_2) \mid \langle \alpha, u \rangle > 0 \text{ for } \alpha \in \Sigma^+\}. \quad (2.24)$$

⁵As in the Pöschl-Teller case, when $b = 0$ the fundamental domain becomes $D' = \mathbb{C}/(W_N \rtimes \mathbb{Z}_2)$. Again, the root system becomes of reduced, C_N type. The denominator $W_N \rtimes \mathbb{Z}_2$ is an extended affine Weyl group. The nontrivial element in the additional \mathbb{Z}_2 accounts for a permutation of an affine and a non-affine simple root that preserves the Weyl alcove. When $b \neq 0$, there is a remnant of this symmetry which we again call ϱ .




Figure 3. The three-dimensional slice of the fundamental domain D_2 for the BC_2 Calogero-Sutherland model in u -space. Front and the back side of the wedge should be identified. The fixed points (walls) under the action of w_2 and w_1 are shown as bold dashed lines. Fixed points of w_2 fall into two disconnected components which carry the labels 0 and 2. The fixed points of w_1 on the other hand form a connected set that intersects our 3-dimensional slice in two lines labeled by 1. The shaded area in front is the Weyl chamber A_2^+ . It is bounded by the walls ω_1 and ω_2 . The shifted Weyl chamber $A_2^L = \tilde{A}_2^+$ is bounded by the walls ω_0 and ω_1 . The subset A_2^E is the 2-dimensional semi-infinite strip of width π on top of the wedge. It is bounded by pieces of wall ω_1 only, except in the corners.

This is simply a Weyl chamber for the BC_2 root system. As for $N = 1$ the Calogero-Sutherland potential diverges along the walls of the chamber, see our discussion above.

There are two additional choices we want to discuss here because of their relevance for conformal field theory, see section 4 below. The first one is given by

$$A_2^L = \tilde{A}_2^+ = \{(u_1, u_2) = (\tilde{u}_1 + i\pi, \tilde{u}_2 - i\pi) \mid \tilde{u}_i > 0; , \tilde{u}_1 > \tilde{u}_2\}. \quad (2.25)$$

It may be obtained from A_2^+ by application of ρ combined with a translation by $-2\pi ie_2$. The associated Schrödinger problem has the same form as on A_2^+ , except that the parameters are changed, see our discussion above

$$V_{(a,b,\epsilon)}^{\text{CS}}(u_1, u_2) = V_{(a,-b,\epsilon)}^{\text{CS}}(\tilde{u}_1, \tilde{u}_2) \quad (2.26)$$

Another relevant possible subset that leads to real potential is the one in which the two coordinates u_1 and $u_2 = u_1^*$ are complex conjugates of each other,

$$A_2^E = \{(u_1, u_2) \mid u_1 = u_2^*, \Im u_1 \in [0, \pi]\}.$$

We shall decompose $u_1 = u + i\varphi$ into its real and imaginary part. In this case we are dealing with a particle that moves on a 2-dimensional semi-infinite strip given by $u \geq 0$

and $\varphi \in [0, \pi]$. For large u the potential becomes

$$V_{(a,b,\epsilon)}^{\text{CS}}(u, \varphi) \sim -\frac{\epsilon(\epsilon - 2)}{8 \sin^2 \varphi} \text{ for } u \rightarrow \infty.$$

So, we see that in this asymptotic regime, wave functions are given by a product of a Pöschl-Teller bound state and a plane wave in the u -direction.

2.4 Coordinates in the CS problem

Let us conclude this section with a short comment on coordinates. So far, we have described the Calogero-Sutherland problem in terms of coordinates u_i in which the kinetic term is simply the standard flat space Laplacian. Since the u_i are coordinates on \mathbb{T}_N it is tempting to apply the exponential map that sends \mathbb{T}_N to \mathbb{C}^N .⁶ Indeed, we shall often use the coordinates

$$x_i = e^{u_i} \tag{2.27}$$

instead of u_i . This has the advantage that the identification $u_i \equiv u_i + 2\pi i$ is manifest. Upon application of the exponential map (and inversion), the 3-dimensional slice of the fundamental domain D_2 that is shown in figure 3 gets mapped to a cone, see figure 4. The map sends the Weyl chamber A_2^+ and the space $A_2^L = \tilde{A}_2^+$ to one half of a section through the cone each, while the set A_2^E becomes half of the mantle of the cone. Figure 4 also keeps track of the location of the singularities. While the coordinates x_i make the identification with $l_i : u_i \rightarrow u_i + 2\pi i$ manifest so that they are proper coordinates on \mathbb{T}_N , the identification from the action of the Weyl group W_N is not built into these coordinates. It is often good to do a little better and to use coordinates that are invariant at least under the action of the Weyl reflections $w(e_i), i = 1, \dots, N$. Any function of $x_i + x_i^{-1}$ would do the job, but we shall adopt a very specific one, namely⁷

$$z_i = -\frac{1}{\sinh^2 \frac{u_i}{2}} = \frac{4}{2 - x_i - x_i^{-1}}. \tag{2.28}$$

These coordinates send the domain D_N to configurations $(z_i) \in \mathbb{C}^N$ that are symmetric under the action of the permutation group $S_N \subset W_N$. The latter is generated by the Weyl reflections $w_i = w(e_i \pm e_{i+1})$ for $i = 1, \dots, N - 1$.

3 Wave functions of the Calogero-Sutherland model

Having set up the eigenvalue equation we want to study we now turn to a discussion of the solutions. As a [warmup](#), we briefly look at the example of the Pöschl-Teller problem for which the study of wave functions involves some fairly basic facts from the theory of

⁶To be more precise, this map is injective, with the image $(\mathbb{C}^\times)^N$, and thus defines a partial compactification of the complex torus. By the action of the Weyl group, it extends to a toroidal compactification of the torus corresponding to its decomposition into Weyl chambers. This gives the toric variety of x -coordinates, see [76] (page 55), [77, 78] for details.

⁷A bit more precisely, for BC_2 we will use the branches $z_1 = \frac{e^{+i\pi}}{\sinh^2 \frac{u_1}{2}}$ $z_2 = \frac{e^{-i\pi}}{\sinh^2 \frac{u_2}{2}}$ in order to be consistent with definition (2.25).




Figure 4. The fundamental domain for the BC_2 Calogero-Sutherland model in x^{-1} -space. The shaded areas in the radial direction are the Weyl chamber A_2^+ and the shifted domain A_2^L . On the boundary of the cone, A_2^+ is bounded by the walls ω_1 and ω_2 while A_2^L is bounded by ω_1 and ω_0 . The subset A_2^E is mapped to half of the mantle.

Gauss' hypergeometric functions. Then we turn to [general \$N\$](#) and discuss a basis of scattering states that are known as [Harish-Chandra functions](#). We will discuss their definition and a new *series expansion* formula for $N = 2$ along with a few direct consequences. As a main application, in the [third subsection](#) we provide a complete analysis of *poles and residues* of Harish Chandra functions for $N = 2$. The [fourth subsection](#) finally, is devoted to the construction of physical wave functions. Through a general discussion of the *monodromy representation* for the Calogero-Sutherland eigenvalue problem we are led to consider special linear combinations of Harish-Chandra functions that possess good analytic properties along the walls of the scattering problem. The optimal choice corresponds to the so-called Heckman-Opdam multivariable hypergeometric functions and, for $N = 2$, a Euclidean analogue thereof.

3.1 Wave functions of the Pöschl-Teller problem

Here we will mostly study the Pöschl-Teller Hamiltonian on the domain A^+ that was introduced in the [previous section](#). The corresponding eigenvalue equation was stated in eq. (2.4). We shall make the following Ansatz for the wave function

$$\Psi(k; u) = \Theta(k; u)\Phi(\lambda, k; u) = 4^{a+\frac{1}{2}} \sinh^{a-b+\frac{1}{2}} \frac{u}{2} \cosh^{a+b+\frac{1}{2}} \frac{u}{2} \Phi(\lambda, k; u). \quad (3.1)$$

As before, $k = (k_1, k_2) = (-2b, a + b + 1/2)$ denotes the parameters of the Pöschl-Teller potential and λ is the momentum. In our conventions it is related to the energy eigenvalue ϵ in eq. (3.1) through $\lambda^2 = -\epsilon$ so that λ is purely imaginary for positive energy solutions.

Since the Pöschl-Teller potential tends to zero at $u = \infty$, the eigenfunctions of the Pöschl-Teller Hamiltonian are superpositions of an outgoing and an incoming plane wave

in this asymptotic regime. We can choose a basis for which one of the two wave functions of energy ϵ is purely outgoing while the other is purely incoming, i.e.

$$\Psi(\pm\lambda, k; u) \sim e^{\pm\lambda u} + \dots \quad \text{for } u \rightarrow \infty. \quad (3.2)$$

Wave functions with these asymptotic properties are also referred to as Harish-Chandra functions. These two wave functions with eigenvalue $\epsilon = -\lambda^2$ can easily be constructed in terms of Gauss' hypergeometric functions as⁸

$$\Phi(\pm\lambda, k; z) = \left(\frac{1}{4} \frac{z}{z-1}\right)^{a+1/2 \mp \lambda} {}_2F_1\left(\begin{matrix} 1/2 + a \mp \lambda, 1/2 - b \mp \lambda \\ 1 \mp 2\lambda \end{matrix}; \frac{z}{z-1}\right), \quad (3.3)$$

where z and u are related through equation (2.28). For large values of u , the argument of the hypergeometric functions approaches zero and the asymptotics of the prefactor combine with that of the gauge transformation Θ to give the desired asymptotics (3.2). Since the standard series expansion for ${}_2F_1(\alpha, \beta; \gamma|y)$ converges on the real y -line for $-1 < y < 1$ and our parameter z takes values $z \in (-\infty, 0)$ on A^+ , the usual series expansion of the function (3.3) converges in the entire Weyl chamber.

In the context of Calogero-Sutherland models, one usually requires the series expansions in the variable $x^{-1} = \exp(-u)$ to be convergent throughout the Weyl chamber, see below. We shall refer to such expansions as u -expansions, even though they are really expansions in $\exp(-u)$. To obtain such a u -expansion for the case at hand, one simply expresses z through u in the series expansion of (3.3)⁹ and then expands $(1 - e^{-u})^{-2(a+1/2 \mp \lambda + k)}$ in powers of $\exp(-u)$. This second expansion is absolutely convergent for $u > 0$. We arrive at¹⁰

$$\Phi(\pm\lambda; k; u) = \sum_{p=0}^{\infty} \frac{e^{(\pm\lambda - 1/2 - a - p)u}}{p!} (2a + 1 \mp 2\lambda)_p {}_3F_2\left(\begin{matrix} -p, 1/2 + b \mp \lambda, 1 + 2a \mp 2\lambda + p \\ 1 \mp 2\lambda, 1 + a \mp \lambda \end{matrix}; 1\right). \quad (3.4)$$

In accord with the general analysis (see next subsection), this expansion is convergent on the entire domain A^+ . It can be analytically continued to the strip $\{u | \Re u > 0, |\Im u| < \pi\}$, which is a tube-like neighborhood of A^+ . If $b = 0$ in (3.4), one can use Watson's summation for ${}_3F_2$ [79] to sum the u -expansion formula (3.4) into

$$\Phi(\pm\lambda; a, b = 0; u) = e^{(\pm\lambda - \frac{1}{2} - a)u} {}_2F_1\left(\begin{matrix} 1/2 + a, 1/2 + a \mp \lambda \\ 1 \mp \lambda \end{matrix}; e^{-2u}\right).$$

⁸Here and in the following, we choose the principal branch for $(-z)^A$ and insist on $|\Im u| < \pi$ in the u -plane for (non-twisted) Harish-Chandra functions.

⁹To avoid additional subtleties, we keep the momenta generic here, namely we assume that $\pm a + 1/2 \mp \lambda, \pm b + 1/2 \mp \lambda \neq 0, -1, -2, \dots$. Special cases can be obtained by carefully taking limits, see comments in section 3 and appendix B.

¹⁰Notice that writing this and the next formula in this manner, we explicitly break the $u \leftrightarrow -u$ symmetry which seemingly contradicts the definition of Harish-Chandra functions as W -symmetric solutions, see the next subsection. This is related to restricted range of validity of the corresponding quadratic transformation from z - to u -variables. So, to be precise, in such expansions (see also BC_2 u -expansions of appendix A) we shall always assume $e^{\pm u}$ as a shorthand for $\cosh u \pm (\cosh u + 1)\sqrt{\tanh^2 u/2}$ where the principal branch of a square root is taken, which reduces to $e^{\pm u} \mapsto (e^u + e^{-u} \pm |e^u - e^{-u}|)/2$ on real slices A^+ and A^L .

The resulting expression is well known from the theory of Calogero-Sutherland wave functions for the reduced root system $C_1 \simeq B_1 \simeq A_1$.

After this short detour on series expansions for the in- and outgoing wave functions we come back to the problem of constructing physical wave functions for the Pöschl-Teller problem. Clearly, the two wave functions we considered so far are badly behaved when we approach the wall at $u = 0$. In fact, $u = 0$ is a branch point. But there exists a unique linear combination of these two wave functions that is analytic at $u = 0$. It is given by

$$\begin{aligned} \Phi^W(\lambda, k; z) &= c(\lambda, k)\Phi(\lambda, k; z) + c(-\lambda, k)\Phi(-\lambda, k; z) \\ &= {}_2F_1\left(\begin{matrix} a + 1/2 + \lambda, a + 1/2 - \lambda \\ 1 + a - b \end{matrix}; \frac{1}{z}\right). \end{aligned} \quad (3.5)$$

with coefficients given by

$$c(\lambda, k) = 4^{-\lambda+a+1/2} \frac{\Gamma(a-b+1)\Gamma(2\lambda)}{\Gamma(1/2+\lambda+a)\Gamma(1/2+\lambda-b)}.$$

These values then allow to apply Kummer's identity in order to pass from the first to the second line in eq. (3.5). Obviously, the branch point at $u = 0$, or equivalently at $z = -\infty$, has been removed now. After multiplication with the factor Θ , Φ^W gives what we would usually consider the physical solution of the hyperbolic Pöschl-Teller system. The eigenfunctions $\Psi^W(\lambda, k; z)$ with $\lambda = ip, p \geq 0$, form a complete orthonormal basis of eigenfunctions for the hyperbolic Pöschl-Teller problem on A^+ [80], with (appropriately normalized) measure $d\mu \sim |\Theta(k; u)|^2 du$.

The branch points of $\Phi(\pm\lambda, k; z)$ at $u = 0$ prevent us from continuing the purely in- and outgoing wave functions beyond A^+ into the compact domain A^C that was defined in eq. (2.6). On the other hand, the function Φ^W can be continued into A^C . For generic choices of λ the resulting wave function of the trigonometric Pöschl-Teller problem (2.7) possesses a branch point at $u = i\pi$. This branch point can only be avoided for a discrete set of λ . In this way one obtains the usual eigenfunctions ψ_n of the trigonometric Pöschl-Teller problem

$$\psi_n(a, b; \varphi) \sim \sin^{a-b+\frac{1}{2}} \frac{\varphi}{2} \cos^{a+b+\frac{1}{2}} \frac{\varphi}{2} {}_2F_1\left(\begin{matrix} -n, 2a+1+n \\ 1+a-b \end{matrix}; \sin^2 \frac{\varphi}{2}\right) \quad (3.6)$$

for $\pm\lambda = n + a + 1/2$ where the variable φ is restricted to the interval $\varphi \in [0, \pi]$ so that it parametrizes A^C . Note that for $n = 0$, i.e. for the ground state, the hypergeometric function contributes a trivial factor. Hence, the wave function of the ground state in the compact domain coincides with the function Θ we introduced in eq. (3.1).

Let us finally also discuss the wave functions of the Pöschl-Teller problem on the shifted domain $\tilde{A}^+ = A^L$ that we defined in eq. (2.5). In this case, we select the Harish-Chandra functions such that the eigenfunctions Ψ possess the standard asymptotics in the real coordinate \tilde{u} on $\tilde{A}^+ = A^L$, i.e.

$$\tilde{\Psi}(\pm\lambda, k; u) \sim e^{\pm\lambda\tilde{u}} + \dots = e^{\mp i\pi\lambda} e^{\pm\lambda u} + \dots \quad \text{for } \tilde{u} = u - i\pi \rightarrow \infty. \quad (3.7)$$

Of course this implies that the Harish-Chandra functions $\tilde{\Phi}$ are related to Φ by a λ -dependent gauge transformation

$$\tilde{\Phi}(\pm\lambda, k; z) = e^{i\pi(\mp\lambda+a+1/2)} \Phi(\lambda, k; z). \quad (3.8)$$

Solutions of our Calogero-Sutherland problem with these asymptotics take the form

$$\tilde{\Phi}(\pm\lambda, k; z) = \left(\frac{z}{4}\right)^{a+1/2\mp\lambda} {}_2F_1\left(\begin{matrix} a \mp \lambda + 1/2, b \mp \lambda + 1/2 \\ 1 \mp 2\lambda \end{matrix}; z\right). \quad (3.9)$$

The relation (3.8) with the standard Harish-Chandra functions Φ is then a consequence of the Pfaff transformation for Gauss hypergeometric function [79]. In writing our formula for $\tilde{\Phi}$ we agree to use the principal branch for z^A so that in z -plane the twisted Harish-Chandra function has cuts along $(-\infty, 0) \cup (1, +\infty)$ for generic values of parameters. Note that on $\tilde{A}^+ = A^L$ the variable z takes values in $z = [0, 1[$, so that this usual series expansion of the functions $\tilde{\Phi}$ is convergent on the entire shifted Weyl chamber. As above, this function can be analytically continued to a semistrip $\{u | \Re u > 0, 0 < \Im u < 2\pi\}$. Once again we need to form a special linear combination of these twisted Harish-Chandra functions $\tilde{\Phi}$ to obtain the physical wave function which is regular at $u = i\pi$ or equivalently $z = 1$. It is given by

$$\begin{aligned} \tilde{\Phi}^W(\lambda, k; z) &= \tilde{c}(\lambda, k)\tilde{\Phi}(\lambda, k; z) + \tilde{c}(-\lambda, k)\tilde{\Phi}(-\lambda, k; z) \\ &= z^{a+\frac{1}{2}-\lambda} {}_2F_1\left(\begin{matrix} a + 1/2 - \lambda, b + 1/2 - \lambda \\ 1 + a + b \end{matrix}; 1 - z\right). \end{aligned} \quad (3.10)$$

with coefficients

$$\tilde{c}(\lambda, k) = 4^{-\lambda+a+1/2} \frac{\Gamma(a+b+1)\Gamma(2\lambda)}{\Gamma(1/2+\lambda+a)\Gamma(1/2+\lambda+b)} =: c_0(\lambda, k).$$

We observe that \tilde{c} can be obtained from c through the inversion $b \rightarrow -b$ of the parameter b . This replacement agrees with the action of the shift ϱ on the coupling constants in the Pöschl-Teller problem, see eq. (2.2). The functions $\tilde{\Psi}^W(\lambda, k; z)$ for $\lambda = ip, p \geq 0$, provide us with a complete orthonormal basis of eigenfunctions for the Pöschl-Teller problem on \tilde{A}^+ .

There is a different, more algebraic way to express these results on the construction of physical wave functions for the Calogero-Sutherland problems through representations of the fundamental group $\pi_1(D)$. Recall from the [previous section](#), that the domain D is a semi-infinite pillow. Hence, its fundamental group is freely generated by two elements g_0 and g_1 . These are described by loops around the singular points $u_0 = i\pi$ and $u_1 = 0$. The Harish-Chandra functions $\Phi(\pm\lambda, k; z)$ carry a 2-dimensional monodromy representation M of this fundamental group. We can easily infer the representation matrices from standard properties of hypergeometric functions, along the lines of our discussion above. For g_1 one finds that

$$M(g_1) = M_1 = C^{-1} \begin{pmatrix} 1 & 0 \\ 0 & e^{2\pi i(a-b)} \end{pmatrix} C, \quad (3.11)$$

where the matrix C is given by

$$C(\lambda, k) = \begin{pmatrix} c(\lambda, k) & c(-\lambda, k) \\ c(\lambda, k') & c(-\lambda, k') \end{pmatrix}$$

and $k' = (k'_1, k'_2) = (2b, -a - b + 1/2)$ is obtained by reversing the sign of both a and b . Our discussion of the regular solution at u_0 shows that $M(g_0) = M_0$ can be computed in

the same way after we perform a gauge transformation from Φ to $\tilde{\Phi}$, i.e.

$$M(g_0) = M_0 = \Omega^{-1} \tilde{M}_1 \Omega \tag{3.12}$$

where \tilde{M}_1 is obtained from M_1 by the substitution $b \rightarrow -b$ and the matrix Ω encodes the gauge transformation (3.8). It reads

$$\Omega(\lambda, k) = \begin{pmatrix} e^{i\pi(\lambda-a-1/2)} & 0 \\ 0 & e^{i\pi(-\lambda-a-1/2)} \end{pmatrix}.$$

We observe that both monodromy matrices possess one eigenvector with unit eigenvalue. This signals the existence of a special linear combination of Harish-Chandra functions which has trivial monodromy around u_0 and u_1 , respectively. Explicitly, these eigenvectors were given by the functions Φ^W and $\tilde{\Phi}^W$. Finally, we also want to stress that the product $M_0 M_1$ of the two monodromies describes the monodromy of the Harish-Chandra functions at $u = \infty$. The latter is fully determined by asymptotic behavior of Harish-Chandra functions, i.e.

$$M_\infty = M_0 M_1 = e^{2\pi i(a+\frac{1}{2})} \begin{pmatrix} e^{-2\pi i\lambda} & 0 \\ 0 & e^{2\pi i\lambda} \end{pmatrix}. \tag{3.13}$$

We will now explain that all this carries over to $N \geq 2$. In particular, the monodromy group and its representation on Harish-Chandra functions is explicitly known, see the section 3.4 below.

3.2 Harish-Chandra series expansions

Let us now discuss eigenstates of the hyperbolic Calogero-Sutherland Hamiltonian for $N > 1$. Our discussion will start with the Weyl chamber A_N^+ in which all the u_i are real and positive. To construct the solutions we are interested in, we note that in the region of large u where we are far away from all walls of the Weyl chamber, the Calogero-Sutherland potential goes to zero and hence, in this regime, any wave function is a superposition of plane waves. Before we give precise definitions let us split off a simple factor from the eigenfunctions Ψ and introduce a new function Φ through

$$\Psi(k; u) = \Theta(k; u) \Phi(\lambda, k; u) = \prod_{\alpha \in \Sigma^+} \left(2 \sinh \frac{\langle \alpha, u \rangle}{2} \right)^{k_\alpha} \Phi(\lambda, k; u). \tag{3.14}$$

The factor $\Theta(k; u)$ is split off for convenience, see below. In the case of $N = 1$ it reduces to the one we worked with in the [previous subsection](#) in order to map the eigenvalue equation for the Pöschl-Teller Hamiltonian to the hypergeometric differential equation in the variable $1/z$. We will often refer to the factor Θ as a *gauge transformation* and to Φ as a Calogero-Sutherland wave functions in hypergeometric gauge. Let us note in passing that the function $\Theta(k, u)$ possesses the following asymptotics for large u ,

$$\Theta(k; u) \sim e^{\langle \rho(k), u \rangle} + \dots \quad \text{with} \quad \rho(k) = \frac{1}{2} \sum_{\alpha \in \Sigma^+} \alpha k_\alpha. \tag{3.15}$$

So-called Harish-Chandra wave functions $\Phi(\lambda, k; z)$ are W_N symmetric solutions of the Calogero-Sutherland Hamiltonian for which Φ possesses the following simple asymptotic behavior

$$\Phi(\lambda, k; u) \sim e^{\langle \lambda - \rho(k), u \rangle} + \dots \quad \text{for } u \rightarrow \infty \text{ in } A_N^+ = WC_N \quad (3.16)$$

where $\lambda = \sum \lambda_i e_i$ is the vector of momenta and $u \rightarrow \infty$ in A_N^+ means that all components become large while preserving the order $u_N < u_{N-1} < \dots < u_1$. Recall that W_N symmetry means that Φ depends on the u_i through z_i and that it is symmetric under all permutations of the z_i . The condition (3.16) selects a unique solution of the scattering problem describing a single plane wave. It is analytic in the Weyl chamber A_N^+ . The corresponding eigenvalue of the Calogero-Sutherland Hamiltonian is given by

$$\varepsilon = \varepsilon(\lambda) = - \sum \lambda_i^2.$$

When we required the Harish-Chandra functions to be symmetric, we used the action of the Weyl group W_N on the coordinate space \mathbb{T}_N . On the other hand, the Weyl group also acts in a natural way on the asymptotic data λ of the Harish-Chandra functions by sending any choice of λ through a sequence of Weyl reflections to $w\lambda, w \in W_N$. Since the eigenvalue ε is invariant under all the reflections, our Harish-Chandra functions come in families. For generic choices of λ , one obtains $|W_N| = N!2^N$ solutions $\Phi(w\lambda, k; z)$ which all possess the same eigenvalue of the Hamiltonian.

At least for sufficiently generic values of the momenta,¹¹ Harish-Chandra functions can be constructed as a series expansion in the variables $x_i = \exp u_i$

$$\Phi(\lambda, k; u) = \sum_{\mu \in Q_+} \Gamma_\mu(\lambda, k) e^{\langle \lambda - \rho(k) - \mu, u \rangle}, \quad \Gamma_0(\lambda, k) = 1, \quad (3.17)$$

where we adopt $|\Im u_i| < \pi$ for $i = 1, \dots, N$ on the principal branch of BC_N Harish-Chandra functions and we sum over elements μ of the $\mathbb{Z}_{\geq 0}$ -cone Q_+ over the positive roots, i.e. the set

$$Q_+ = \left\{ \mu = \sum_{i=1}^{N-1} n_i \alpha_i + \frac{n_N \alpha_N}{2} \mid n_i \in \mathbb{Z}_{\geq 0} \text{ for } i = 1, \dots, N \right\}.$$

For later discussions we note that Q_+ comes equipped with a partial order \preceq where $\mu \preceq \nu$ iff $\nu - \mu \in Q_+$.

It is not difficult to derive recursion relations of the expansion coefficients $\Gamma_\mu(\lambda, k)$ directly from the Calogero-Sutherland eigenvalue problem, see e.g. [76]¹²

$$\langle 2\lambda - \mu, \mu \rangle \Gamma_\mu(\lambda, k) = 2 \sum_{\alpha \in \Sigma^+} k_\alpha \sum_{j \geq 1} \langle \lambda - \rho(k) - \mu + j\alpha, \alpha \rangle \Gamma_{\mu - j\alpha}(\lambda, k). \quad (3.18)$$

This can be solved uniquely, if

$$2\langle \lambda, \mu \rangle - \langle \mu, \mu \rangle \neq 0 \quad \text{for all } \mu \in Q_+. \quad (3.19)$$

¹¹A precise formulation of the condition is given below through the inequalities (3.19) and the subsequent discussion.

¹²The recursion derived in [81] is closely related to this one, specialized for BC_2 . However, our expansion here is in monomials, not in Gegenbauer polynomials, although it is not difficult to go between the two.

The resulting series are known to converge¹³ within the Weyl chamber [76], as is fairly obvious from the physics they describe. Even when one of the conditions (3.19) is violated it is possible to obtain a complete basis of series solutions $\Phi(w\lambda, k; u)$, $w \in W_N$. There is a subset of such cases where things are a bit subtle, namely when λ is chosen such that $\langle \lambda, \alpha^\vee \rangle$ is integer which implies that one of the conditions in eq. (3.19) is violated. For such values of non-generic momenta λ some of the series solutions are logarithmic. This is analogous to usual properties of the Gauss hypergeometric function when the difference of exponents becomes integer [79]. To obtain their expansions, one needs to see which fundamental solutions coincide on the corresponding locus and take the limit of their rescaled difference as a new fundamental solution.

Example. Let us look at the Harish-Chandra functions for $N = 2$ in some detail. Although, we could literally repeat the entire $N = 1$ discussion here, we leave much of it for appendix A.¹⁴ In particular, this appendix contains explicit expressions for the expansion coefficients Γ_μ in the u -expansion (3.17), see eq. (A.11). We do not want to repeat these here and will rather discuss a somewhat intermediate form of a z_i -expansion from which we shall derive many interesting properties of Harish-Chandra functions in the remainder of this and in the next subsection. It is given by

$$\begin{aligned} \Phi(\lambda_1, \lambda_2; k_i; z_1, z_2) &= \frac{1}{4^{2a+1+\epsilon/2-\lambda_1-\lambda_2}} \sum_{n,m=0}^{\infty} \frac{(1/2+a-\lambda_1, 1/2-b-\lambda_1, \frac{\epsilon}{2}-\lambda_1+\lambda_2)_n}{(1-2\lambda_1, 1-\lambda_1+\lambda_2)_n} \\ &\quad \times \frac{(1/2+a-\lambda_2, 1/2-b-\lambda_2, \epsilon/2+\lambda_1-\lambda_2)_m (1-\epsilon/2-\lambda_1+\lambda_2)_{n-m}}{(1-2\lambda_2, 1+\lambda_1-\lambda_2)_m (-\lambda_1+\lambda_2)_{n-m}} \\ &\quad \times {}_4F_3 \left(\begin{matrix} -n, -m, 1-\epsilon/2, 1-\epsilon/2-\lambda_1-\lambda_2 \\ 1-\epsilon/2+\lambda_1-\lambda_2-n, 1-\epsilon/2-\lambda_1+\lambda_2-m, 1-\lambda_1-\lambda_2 \end{matrix}; 1 \right) \\ &\quad \times \frac{1}{n!m!} \left(\frac{z_1}{z_1-1} \right)^{\frac{1+\epsilon}{2}+a-\lambda_1+n} \left(\frac{z_2}{z_2-1} \right)^{\frac{1}{2}+a-\lambda_2+m} {}_2F_1 \left(\begin{matrix} \epsilon/2-\lambda_1+\lambda_2-m+n, \epsilon/2 \\ 1-\lambda_1+\lambda_2-m+n \end{matrix}; \frac{z_1}{z_2} \frac{z_2-1}{z_1-1} \right). \end{aligned} \quad (3.20)$$

As in the discussion for $N = 1$ we choose the principal branch for $(-z_i)^A$, so that $z_i/(z_i - 1) = \cosh^{-2}(u_i/2)$. It is obvious that this function has the correct asymptotic behaviour. Let us stress that, unlike a somewhat similar expansion for conformal blocks that appears in [25], our expansion for Harish-Chandra functions is also valid for non-integer spins l . We can use it to derive a corresponding expansion for conformal blocks once we have explained how to construct blocks from Harish-Chandra functions in the next section. The derivation of eq. (3.20), its features and equivalent expansions are described in appendix A. We note that the z -expansion (3.20) is convergent for arguments in the region

$$\Re z_i < \frac{1}{2}, i = 1, 2, \quad \left| \frac{z_1}{z_1-1} \right| < \left| \frac{z_2}{z_1-1} \right|, \quad (3.21)$$

which includes the entire Weyl chamber, similarly to the BC_1 case. Here we assume that the parameters are generic.

¹³The series converges absolutely and uniformly on compacta in the set of complexified momenta and multiplicities times A^+ , as long as they are chosen to avoid the hyperplanes (3.19).

¹⁴In appendix A, we derive a z -expansion for twisted BC_2 Harish-Chandra function which is related to the present one by formula (3.48).

As an immediate application of the series expansion (3.20) we can evaluate Harish-Chandra functions for some special values of the multiplicities k_i . For $d = \epsilon + 2 = 2$, for example, the multiplicity $k_3 = 0$ vanishes so that one of the upper parameters in the balanced ${}_4F_3(1)$ inside the sum coincides with one of the lower ones. The sum involving the resulting balanced ${}_3F_2(1)$ is summable via Saalschütz identity [82] and we obtain

$$\Phi(\lambda_i; k_1, k_2, k_3 = 0; z_i) = \prod_{i=1,2} \left(\frac{1}{4} \frac{z_i}{z_i - 1} \right)^{1/2+a-\lambda_i} {}_2F_1 \left(\begin{matrix} 1/2+a-\lambda_i, 1/2-b-\lambda_i \\ 1-2\lambda_i \end{matrix}; \frac{z_i}{z_i-1} \right).$$

The result is a product of Harish-Chandra functions for the Pöschl-Teller problem $N = 1$, see eq. (3.3). The case of $d = 4$ is even simpler to evaluate. Indeed, for this value of d , the parameter $k_3 = (d - 2)/2 = 1$ and hence one of the upper parameters in the balanced ${}_4F_3(1)$ tends to zero, so that we obtain

$$\begin{aligned} \Phi(\lambda_1, \lambda_2; k_1, k_2, k_3 = 1; z_1, z_2) &= \\ &= \frac{1}{4} \frac{z_1 z_2}{z_1 - z_2} \prod_{i=1,2} \left(\frac{1}{4} \frac{z_i}{z_i - 1} \right)^{a+1/2-\lambda_i} {}_2F_1 \left(\begin{matrix} 1/2+a-\lambda_i, 1/2-b-\lambda_i \\ 1-2\lambda_i \end{matrix}; \frac{z_i}{z_i-1} \right). \end{aligned}$$

Before we derive further properties of Harish-Chandra functions for $N = 2$ from the series expansion (3.20), we want to review a few more general properties that hold for any N .

The last two formulas for $N = 2$ that we derived from the z -expansion (3.20) possess a nice generalization to arbitrary values of N . For $k_3 = 0, 1$ and generic (non-resonant) values of the eigenvalues λ is known that [83]

$$\Phi(\lambda_i; k_i; z_i) = \Delta_m^{-k_3} \prod_{i=1}^N \left(\frac{1}{4} \frac{z_i}{z_i - 1} \right)^{a+1/2-\lambda_i} {}_2F_1 \left(\begin{matrix} 1/2+a-\lambda_i, 1/2-b-\lambda_i \\ 1-2\lambda_i \end{matrix}; \frac{z_i}{z_i-1} \right),$$

where

$$\Delta_m = \prod_{\alpha \in \Sigma^+, \text{ middle}} \left(e^{\frac{\langle \alpha, u \rangle}{2}} - e^{-\frac{\langle \alpha, u \rangle}{2}} \right)$$

is a Weyl denominator for middle roots. More generally, it is known [83, 84] that Harish-Chandra functions for any positive integer value of the multiplicity k_3 are multilinear combinations of Pöschl-Teller wave functions. The most elegant derivation of such expressions, one that is also completely universal in N , involves $N(N - 1)/2$ differential or difference operators which shift k_3 by one unit. For scalar 4-point blocks similar constructions are known from the work of Dolan-Osborn [25, 26]. We will show how explicit closed formulas for these operators follow from the integrable structure of Calogero-Sutherland models in a forthcoming publication [55].

3.3 Poles and residues of Harish-Chandra functions

Our short excursion to explicit expressions for Harish-Chandra functions which exist for special values of the multiplicities only should not mislead the reader to think that Harish-Chandra functions can only be understood for very special cases. In fact, it is one of the central virtues of Heckman-Opdam theory that one is able to say so much about

Harish-Chandra functions often without knowing their explicit series expansions or integral formulas. As an example for one of the many further properties that is well understood beyond the simple case of $N = 1$ let us mention that $\exp(\langle -\lambda + \rho(k), u \rangle) \Phi(\lambda; k; u)$ are entire functions of the multiplicities k_i and meromorphic function of asymptotic data λ_i , for any choice of u in the fundamental domain. They are known to possess simple poles whenever the set of λ_i satisfies one of the following conditions

$$\langle \lambda_*, \alpha^\vee \rangle = s \quad \text{for } s = 1, 2, \dots, \quad \alpha \in \Sigma^+. \tag{3.22}$$

Let us note that given λ_* satisfying this condition, this λ_* violates the condition (3.19). In fact, the quadratic expression in eq. (3.19) vanishes at least for $\mu = s\alpha^\vee$. The converse is not true, i.e. there exist many values of λ that violate the inequalities (3.19) but do satisfy the condition (3.22). At such values of λ Heckman and Opdam showed that the singularities are only apparent. For the true poles at $\lambda_* = \lambda_{\alpha, n}$, the residues are given by (see e.g. [84])

$$\text{Res}_{(\alpha, s)} \Phi(\lambda, k; z) \sim \Phi(w(\alpha)\lambda_{\alpha, s}, k; z). \tag{3.23}$$

where \sim indicates that the relation with the Harish-Chandra function $\Phi(w(\alpha))$ holds only up to a constant factor. The latter is not known in general, but we shall explain later how it can be determined and provide explicit expressions for $N = 2$ with the help of the series expansion (3.20).

Relation (3.23) is actually a little more subtle than it may appear at first. According to a theorem by Heckman ([76], Cor. 4.2.4), eq. (3.23) holds true as it stands whenever the quadratic equation $\langle 2\lambda_0 - \mu, \mu \rangle = 0$ has exactly one non-zero solution $\mu_0 \in Q_+$. In this case, one has

$$\left\{ \langle 2\lambda - \mu_0, \mu_0 \rangle \Phi(\lambda; k; u) \right\} \Big|_{\lambda_0} = \left\{ (2\lambda - \mu_0, \mu_0) \Gamma_{\mu_0}(\lambda, k) \right\} \Big|_{\lambda_0} \Phi(\lambda_0 - \mu_0; k; u). \tag{3.24}$$

Here, the Harish-Chandra function on the right hand side appears from summing over all terms in the original series with $\mu \succeq \mu_0$ with respect to the partial ordering of Q_+ we introduced above. If the quadratic equation has several solutions in Q_+ , on the other hand, relation (3.23) holds only for a properly defined Harish-Chandra series on the right hand side. One can do this by approaching the desired limit point λ_* along a sequence of irrational values λ for which $\langle 2\lambda - \mu, \mu \rangle = 0$ has a unique solution. Then, provided one knows the series expansion of Harish-Chandra function for such λ , it is possible to calculate its residues by Heckman's theorem.

Example. Let us continue our tradition to describe a few more details in the case of $N = 2$. As we discussed before, the corresponding Weyl group W_2 is eight dimensional. It is generated by the two reflections w_1 and w_2 . All eight elements are listed in table 1 along with their action $\lambda \rightarrow w\lambda$ on the asymptotic data λ of the Harish-Chandra functions. In the case at hand, the shift by $\rho(k)$ reads

$$\rho(k) = 1/2(k_1 + 2k_2 + 2k_3, k_1 + 2k_2) = (a + 1/2 + \epsilon/2, a + 1/2).$$

$w \in W_2$	$w(\lambda_1, \lambda_2)$	$w(\Delta, l)$
e	(λ_1, λ_2)	(Δ, l)
w_1	(λ_2, λ_1)	$(\Delta, 2 - l - d)$
w_2	$(\lambda_1, -\lambda_2)$	$(1 - l, 1 - \Delta)$
$w_1 w_2$	$(-\lambda_2, \lambda_1)$	$(1 - l, \Delta - d + 1)$
$w_2 w_1$	$(\lambda_2, -\lambda_1)$	$(l + d - 1, 1 - \Delta)$
$w_1 w_2 w_1$	$(-\lambda_1, \lambda_2)$	$(l + d - 1, \Delta + 1 - d)$
$w_2 w_1 w_2$	$(-\lambda_2, -\lambda_1)$	$(d - \Delta, l)$
$w_1 w_2 w_1 w_2$	$(-\lambda_1, -\lambda_2)$	$(d - \Delta, 2 - l - d)$

Table 1. Elements of the Weyl group W_2 for the BC_2 root system (first column) along with their action on the asymptotic data $\lambda = (\lambda_1, \lambda_2)$ (second column). The third column is identical to the second, but uses a different parametrization of the asymptotic data that we will discuss in section 4.

The final column of table 1 will be explained in section 4. It describes the asymptotic data in a different parametrization.

If we denote an element of $\mu \in Q_+$ as $\mu = n\alpha_1 + m\alpha_2 = ne_1 + (m - n)e_2 = (n, m - n)$ for $n, m \in \mathbb{Z}_{\geq 0}$, the locally finite set of hyperplanes on which the definition of the Harish-Chandra series requires an appropriate limiting procedure reads

$$2n\lambda_1 + 2(m - n)\lambda_2 = n^2 + (m - n)^2. \tag{3.25}$$

As we stated before, only a small subset of these hyperplanes give actual poles. Namely, according to equation (3.22) the Harish-Chandra functions possess four series of poles in the asymptotic data λ . In fact, while the set Σ^+ of positive roots contains six elements, the solutions of eq. (3.22) for the longest roots $2e_i$ form a subspace of the solutions for the short roots e_i . The relevant four roots are listed table 2 along with the corresponding Weyl reflection written in terms of our fundamental generators w_1 and w_2 . For each of these four cases, the solutions of eq. (3.22) are listed in the third column. They depend on a non-negative integer and one free parameter λ . The residues of our Harish-Chandra functions for these values of λ_* are again given by Harish-Chandra functions with different asymptotic data. The latter is listed in the fourth column of table 2. The information of the third and fourth column is repeated in the last two columns in a different parametrization of the asymptotic data that we explain in section 4.

In order to obtain exact formulas for the residues, including the numerical coefficients, we shall employ the series expansion (3.20). The general idea is simple to state. As we know, the residue should be proportional to a Weyl-reflected Harish-Chandra function, so one just needs to locate the terms of the series expansion that diverge as we send

α	$w(\alpha)$	$(\lambda_1 \lambda_2)_*$	$w(\alpha)(\lambda_1, \lambda_2)_*$	$(\Delta, l)_*$	$w(\alpha)(\Delta, l)_*$
e_1	$w_1 w_2 w_1$	$(s/2, \lambda)$	$(-s/2, \lambda)$	$(l + d - s - 1, l)$	$(l + d - 1, l - s)$
e_2	w_2	$(\lambda, s/2)$	$(\lambda, -s/2)$	$(1 - s - l, l)$	$(1 - l, l + s)$
$e_1 + e_2$	$w_2 w_1 w_2$	$(-\lambda + s, \lambda)$	$(-\lambda, \lambda - s)$	$(d/2 - s, l)$	$(d/2 + s, l)$
$e_1 - e_2$	w_1	$(\lambda + s, \lambda)$	$(\lambda, \lambda + s)$	$(\Delta, 1 - d/2 + s)$	$(\Delta, 1 - s - d/2)$

Table 2. Pole positions (first column) and asymptotic data of their residues (second column) of the Harish-Chandra functions $\Phi(\lambda_1, \lambda_2, k; z)$ for $N = 2$. The third and fourth column contain the same information in a different parametrization of asymptotic data, see section 4.

λ close to a corresponding pole position. Within the set of these divergent terms one needs to identify the one that gives the leading power of $\Phi(w(\alpha)\lambda_{\alpha,s}, k; z)$ and read off the coefficient. Heckman’s theorem quoted above guarantees that we can interchange the summation and the limit. In principle, the described steps can be carried out both for the usual u -expansion (see appendix A) and the z -expansion we spelled out in equation (3.20). Here we shall sketch the derivation from the latter and spell out the explicit residues for all four families of poles.

According to our table, the residue for the series $\lambda_1 = s/2 + \xi$ is proportional to $\Phi(-\frac{s}{2}, \lambda_2; k; z(u))$. It is easy to see that the leading divergent term in the z_i -expansion (3.20) arises from the summands with $n = s, m = 0, p = 0$. Here and in the following, the summation index p refers to the series expansion of the hypergeometric function ${}_2F_1$ in the last line of equation (3.20). Hence, the two hypergeometric functions in the summands of eq. (3.20) can be replaced by $F \sim 1$ in all divergent terms. Keeping in mind that $1/(1 - s - 2\xi)_s \sim (-1)^s/2\xi(s - 1)!$, we obtain

$$\begin{aligned} & \left. \Phi(\lambda, k; z(u)) \right|_{\lambda_1 \sim \frac{s}{2} + \xi} = \\ & = \frac{1}{\xi} \left(\frac{1}{2} \frac{4^s}{s!(s-1)!} \frac{(a + \frac{1-s}{2}, b + \frac{1-s}{2}, \frac{\epsilon}{2} + \lambda_2 - \frac{s}{2}, 1 - \frac{\epsilon}{2} + \lambda_2 - \frac{s}{2})_s}{(\lambda_2 - \frac{s}{2}, 1 + \lambda_2 - \frac{s}{2})_s} x_1^{-a - \frac{1+\epsilon}{2} - \frac{s}{2}} x_2^{-a - \frac{1}{2} + \lambda_2} + \dots \right), \end{aligned}$$

so that the residues are given by

$$\begin{aligned} \text{Res}_{\lambda_1 = s/2} \Phi(\lambda, k; z(u)) &= \frac{1}{2} \frac{4^s}{s!(s-1)!} \frac{(a + \frac{1-s}{2}, b + \frac{1-s}{2}, \frac{\epsilon}{2} + \lambda_2 - \frac{s}{2}, 1 - \frac{\epsilon}{2} + \lambda_2 - \frac{s}{2})_s}{(\lambda_2 - \frac{s}{2}, 1 + \lambda_2 - \frac{s}{2})_s} \quad (3.26) \\ &\times \Phi\left(-\frac{s}{2}, \lambda_2; k; z(u)\right). \end{aligned}$$

The evaluation of the residue for the family $\lambda_2 = s/2 + \xi$ is even simpler. Since it is proportional to $\Phi(\lambda_1, -\frac{s}{2}; k; z(u))$, it must arise from the terms $n = 0, m = s, p = 0$ in eq. (3.20). Once again it is easily seen that the two inner hypergeometric functions that appear in the summands do not contribute so that the residue reads

$$\text{Res}_{\lambda_2 = s/2} \Phi(\lambda, k; z(u)) = \frac{1}{2} \frac{4^s}{s!(s-1)!} \left(a + \frac{1-s}{2}, b + \frac{1-s}{2} \right)_s \Phi\left(\lambda_1, -\frac{s}{2}; k; z(u)\right). \quad (3.27)$$

For the third family of poles at $(\lambda_1, \lambda_2)_* = (-\lambda + s, \lambda)$ the residue is proportional to $\Phi(-\lambda, \lambda - s; k; z(u))$. The relevant terms in the sum arise from $n = m = s, p = 0$. For these values of indices, the inner ${}_2F_1$ still does not contribute, but the ${}_4F_3$ does, as none of the upper parameters is zero anymore. Furthermore, in the present case the Pochhammer symbol that produces the desired pole is among the lower ones in the series expansion of the inner ${}_4F_3$ function. To take it out, we use a Whipple transformation for balanced ${}_4F_3$ [82]

$$\begin{aligned} & {}_4F_3\left(\begin{matrix} -n, -m, 1 - \epsilon/2, 1 - \epsilon/2 - \lambda_1 - \lambda_2 \\ 1 - \epsilon/2 + \lambda_1 - \lambda_2 - n, 1 - \epsilon/2 - \lambda_1 + \lambda_2 - m, 1 - \lambda_1 - \lambda_2 \end{matrix}; 1\right) \\ &= \frac{(1 - \epsilon/2 - \lambda_1 + \lambda_2, 1 - \lambda_1 - \lambda_2 + m)_n}{(1 - \lambda_1 - \lambda_2, 1 - \epsilon/2 - \lambda_1 + \lambda_2 - m)_n} \times \\ & \quad \times {}_4F_3\left(\begin{matrix} -n, -m, \lambda_1 - \lambda_2 - n, 2\lambda_1 - n \\ \epsilon/2 + \lambda_1 - \lambda_2 - n, 1 - \epsilon/2 + \lambda_1 - \lambda_2 - n, \lambda_1 + \lambda_2 - m - n \end{matrix}; 1\right). \end{aligned}$$

Substituting $(\lambda_1, \lambda_2)_*$, taking $n = m = s$ and expanding the balanced ${}_4F_3$ on the right hand side of the last formula in ξ , we see that its leading term is not singular now. Moreover, one upper and one lower parameter in it cancel against each other and we are left with a balanced ${}_3F_2$ function that can be summed by a usual Saalschütz formula [82]:

$${}_4F_3\left(\begin{matrix} -s, -s, -2\lambda + \xi, -2\lambda + s + 2\xi \\ \epsilon/2 - 2\lambda + \xi, 1 - \epsilon/2 - 2\lambda + \xi, -s + \xi \end{matrix}; 1\right) \sim \frac{\left(\frac{\epsilon}{2}, 1 - \frac{\epsilon}{2}\right)_s}{\left(\frac{\epsilon}{2} - 2\lambda, 1 - \frac{\epsilon}{2} - 2\lambda\right)_s} + \dots$$

where the dots denote subleading orders in ξ . All in all, the residue is thus again given just by a product of Pochhammer symbols

$$\begin{aligned} \text{Res}_{\lambda_1 = -\lambda + s, \lambda_2 = \lambda} \Phi(\lambda, k; z(u)) &= \frac{4^{2s}}{s!(s-1)!} \\ &\times \frac{\left(\frac{1}{2} + a + \lambda - s, \frac{1}{2} + b + \lambda - s, \frac{1}{2} + a - \lambda, \frac{1}{2} + b - \lambda, \frac{\epsilon}{2}, 1 - \frac{\epsilon}{2}\right)_s}{(-2\lambda, 1 - 2\lambda)_{2s}} \times \Phi(-\lambda, \lambda - s; k; z(u)). \end{aligned} \tag{3.28}$$

The last sequence of poles at $(\lambda_1, \lambda_2)_* = (\lambda + s, \lambda)$ with residues proportional to $\Phi(\lambda, \lambda + s; k; z(u))$, is again simple. In this case the leading terms arise from $n = m = 0, p = s$, so the inner ${}_4F_3$ does not contribute, whereas the ${}_2F_1$ does. Therefore, we readily obtain

$$\text{Res}_{\lambda_1 = \lambda + s, \lambda_2 = \lambda} \Phi(\lambda, k; z(u)) = \frac{1}{s!(s-1)!} \left(\frac{\epsilon}{2}, 1 - \frac{\epsilon}{2}\right)_s \times \Phi(\lambda, \lambda + s; k; z(u)). \tag{3.29}$$

Notice that all the residues are symmetric under $a, b, \leftrightarrow -a, -b, \epsilon \leftrightarrow 2 - \epsilon$, as they should be in consistency with the symmetries of BC_2 Harish-Chandra function described above. Let us also note that, for generic multiplicities k and eigenvalues λ , all four families contribute an infinite sequences of poles.

3.4 Monodromy group and representation

As in the case of the Pöschl-Teller problem, imposing good features of the wave functions at the walls of A_N^+ requires to consider certain linear combinations of Harish-Chandra

functions. For the BC_N Calogero-Sutherland model, the Weyl chamber A_N^+ possesses N walls $\omega_i, i = 1, \dots, N$, which are in one-to-one correspondence to the generators w_i of the Weyl group. As we have discussed above, there is one more wall ω_0 , but it does not bound the domain A_N^+ and so is not of concern for us, at least for most of this section. Since there are $|W_N|$ Harish-Chandra functions and hence $|W_N|$ coefficients to fix in their linear combinations, one might naively expect that analyticity at the N walls $\omega_i, i = 1, \dots, N$ would leave some coefficients undetermined. It turns out, however, that there exists a unique linear combination (up to a constant factor) that is analytic at all N walls on the boundary of the Weyl chamber.

The prescription to build such analytic wave functions is not difficult to state. Suppose we want the wave function Φ to be analytic at some subset $\omega_{i_1}, \dots, \omega_{i_r}$ consisting of $r \leq N$ of the N walls that bound A_N^+ , i.e. $i_\nu \neq 0$. To each of these walls there is a generator w_{i_ν} of the Weyl group and so our set of r walls is associated with a subgroup $V \subset W_N$ of the Weyl group that is generated by w_{i_1}, \dots, w_{i_r} . Given this subgroup we now define the following superposition of Harish-Chandra functions

$$\Phi^V(\lambda, k; z) = \sum_{w \in V} c(w\lambda, k) \Phi(w\lambda, k; z) \tag{3.30}$$

where the so-called Harish-Chandra c-function¹⁵ reads

$$c(\lambda, k) = \frac{\gamma(\lambda, k)}{\gamma(\rho(k), k)}, \quad \gamma(\lambda, k) = \prod_{\alpha \in \Sigma^+} \gamma_\alpha(\lambda, k), \tag{3.31}$$

$$\gamma_\alpha(\lambda, k) = \frac{\Gamma(\frac{1}{2}k_{\alpha/2} + \langle \lambda, \alpha^\vee \rangle)}{\Gamma(\frac{1}{2}k_{\alpha/2} + k_\alpha + \langle \lambda, \alpha^\vee \rangle)}. \tag{3.32}$$

Any wave function of this form turns out to be regular at the walls $\omega_{i_1}, \dots, \omega_{i_r}$. There are two extreme cases of this construction. If we just demand regularity at a single wall ω_i , then the subgroup V consists of two elements, the identity and the reflection w_i . Hence, from the W_N orbit of $|W_N| = N!2^N$ Harish-Chandra functions $\Phi(w\lambda, k; z)$ we obtain $N!2^{N-1}$ linear combinations that are regular at ω_i . If, on the other hand, we want a wave function that is regular at all the walls $\omega_i, i = 1, \dots, N$ that bound A_N^+ , then the subgroup $V = W_N$ coincides with the Weyl group and we end up with a unique linear combination. Let us note that while this function Φ^W is analytic in a neighborhood of A_N^+ , it fails to be analytic at the wall ω_0 . The function $F^+ = \Phi^W$ is also known as Heckman-Opdam hypergeometric function. More generally, linear combinations of the form (3.30) are referred to Θ -hypergeometric functions, see e.g. [86] and references therein.

One can also build a function $\tilde{\Phi}^W$ that is analytic at $\omega_0, \dots, \omega_{N-1}$, but has monodromy around ω_N . We will give an exact formula in the case of $N = 2$ below. The corresponding wave functions $\Psi^W(\lambda, k; z)$ and $\tilde{\Psi}^W(\lambda, k; z)$ with $\lambda_j = ip_j, p_j \geq 0$, are the physical wave functions of the Calogero-Sutherland problem on A_N^+ and $\tilde{A}_N^+ = A^L$, respectively. Hence, they provide an orthogonal basis within the space of functions on the A_N^+ and A^L [76, 84,

¹⁵This function is an analytic continuation of the so-called Gindikin-Karpelevic c-functions [85] in harmonic analysis.

[87]. In hypergeometric gauge the Hermitian scalar product on A_N^+ reads¹⁶

$$(\phi_1, \phi_2) = \int_{A_N^+} \phi_1(u) \overline{\phi_2(u)} |\Theta(k; u)|^2 du. \quad (3.33)$$

where du is a suitably normalized measure on A_N^+ [87]. The gauge transformed Calogero-Sutherland Hamiltonian $\Theta(k; u)^{-1} H^{\text{CS}} \Theta(k; u)$ is self-adjoint with respect to this inner product.

As in the case of $N = 1$ there exists a nice reformulation in terms of the representation theory of the fundamental group $\mathcal{M}_N = \pi_1(D_N)$. The latter is generated by $N + 1$ elements $g_i, i = 0, \dots, N$ which are associated with closed loops around the $N + 1$ walls $\omega_i, i = 0, \dots, N$ of the Calogero-Sutherland potential [76, 88] (see also [89]). Note that the real codimension of these walls in complex space is two. Since the walls are associated with generators w_i of the affine Weyl group \mathcal{W}_N , so are the generators $g_i \in \mathcal{M}_N$. One may show [88] that the generators $g_i, i = 0, \dots, N$ of the fundamental group $\mathcal{M}_N = \pi(D_N)$ satisfy the following set of relations

$$g_i g_j = g_j g_i \quad \text{for } |i - j| \geq 2, \quad (3.34)$$

$$g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1} \quad \text{for } i = 1, \dots, N - 2, \quad (3.35)$$

$$g_0 g_1 g_0 g_1 = g_1 g_0 g_1 g_0, \quad g_{N-1} g_N g_{N-1} g_N = g_N g_{N-1} g_N g_{N-1}. \quad (3.36)$$

These are very similar to the defining relations of the affine Weyl group, except that in the fundamental group $g_i^2 \neq 1$. So, the relation between the monodromy group of D_N and the affine Weyl group mimics the relation between the braid group and the permutation group. Therefore, the fundamental group \mathcal{M}_N is also referred to as (Artin) affine braid group. Note that the relation in the second row is the usual braid or Yang-Baxter equation. The relations in the third line, which involve the elements g_0 and g_N , on the other hand, resemble forth order reflection equations in which two factors on each side arise from the reflection at a boundary while the other two are associated with scattering in the bulk.

For the affine Weyl group we actually discussed two different presentations, one in terms of $N + 1$ generators $w_i, i = 0, \dots, N$ and a second one that involves $2N$ generators τ_i and w_i with $i = 1, \dots, N$. In complete analogy to the construction of the generators $\tau_i \in W_N$, we can build [76, 90, 91] (see also [92]) a set of commuting generators $l_i \in \mathcal{M}_N$ through

$$l_i = g_i^{-1} \cdots g_{N-1}^{-1} (g_N^{-1} \cdots g_0^{-1}) g_1 \cdots g_{i-1}. \quad (3.37)$$

The reader is invited to verify that $l_i l_j = l_j l_i$. Hence the group elements l_i generate an N -dimensional lattice $\mathbb{Z}^N \cong \Gamma \subset \mathcal{M}_N$. The generators $g_i, i = 1, \dots, N$ act on this lattice as

$$g_i l_i g_i = l_{i+1}, \quad 1 \leq i < N \quad (3.38)$$

$$[g_i, l_j] = 0, \quad |i - j| \geq 2 \text{ or } (i, j) = (N, N - 1). \quad (3.39)$$

Our conventions on directions of the basic loops match [76] (section 4.3) if we identify our generators l_i with the generators l_i used by Heckman as $l_i = \prod_{k=i}^N l_k^{-1}$.

¹⁶This formula defines an inner product only if certain conditions on the multiplicities k_α are satisfied, e.g. if $\Re(k_\alpha) > 0$ for all $\alpha \in \Sigma^+$.

For generic values of λ , the space of Harish-Chandra wave functions $\Phi(w\lambda, k; z)$ forms a $|W_N|$ -dimensional representation of the monodromy group \mathcal{M}_N . This representation is explicitly known due to the work of Heckman and Opdam. With respect to the action of M_i , $i = 1, \dots, N$ the space of Harish-Chandra functions splits into $|W_N|/2$ 2-dimensional subspaces each of which is spanned by a pair of Harish-Chandra functions $\Phi(\lambda, k; z)$ and $\Phi(w_i\lambda, k; z)$. On these subspaces, the action is given by

$$M(g_i) = M_i = C_i^{-1} \begin{pmatrix} 1 & 0 \\ 0 & e^{\pi i(1+\epsilon)} \end{pmatrix} C_i \quad \text{for } i = 1, \dots, N-1, \quad (3.40)$$

$$M(g_N) = M_N = C_N^{-1} \begin{pmatrix} 1 & 0 \\ 0 & e^{2\pi i(a-b)} \end{pmatrix} C_N. \quad (3.41)$$

The matrices C_i , $i = 1, \dots, N$, are defined as

$$C_i(\lambda, k) = \begin{pmatrix} c_i(\lambda, k) & c_i(w_i\lambda, k) \\ c_i(\lambda, k'') & c_i(w_i\lambda, k'') \end{pmatrix},$$

where $k'' = (k_1'', k_2'', k_3'') = (-k_1, 1 - k_2, 1 - k_3)$ denotes an involution of the multiplicities (see also below) and

$$c_i(\lambda, k) = 4^{-\lambda_i + \lambda_{i+1} + \epsilon/2} \frac{\Gamma(\frac{1}{2} + \frac{\epsilon}{2})\Gamma(2\lambda_i - 2\lambda_{i+1})}{\Gamma(\frac{\epsilon}{2} + \lambda_i - \lambda_{i+1})\Gamma(\frac{1}{2} + \lambda_i - \lambda_{i+1})} \quad \text{and} \quad (3.42)$$

$$c_N(\lambda, k) = 4^{-\lambda_N + a + 1/2} \frac{\Gamma(a - b + 1)\Gamma(2\lambda_N)}{\Gamma(1/2 + \lambda_N - b)\Gamma(1/2 + \lambda_N + a)} \quad (3.43)$$

is the same function that appeared in our discussion of the Pöschl-Teller problem. The monodromy representation $M_0 = M(g_0)$ of the element g_0 is once again obtained from M_N by the simple replacement $b \rightarrow -b$ along with a gauge transformation of the form

$$\Omega(\lambda, k) = \text{diag}_{w \in W_N} \left(e^{i\pi \langle w\lambda - \rho, \theta \rangle} \right)$$

where $\theta = \sum_i e_i$ is the sum of short roots.¹⁷ So, in formulas one has

$$M(g_0) = M_0 = \Omega^{-1} \tilde{M}_N \Omega. \quad (3.44)$$

Here \tilde{M}_N denotes the matrix in eq. (3.41) with b replaced by $-b$. Note that the monodromy matrices M_i , $i = 1, \dots, N-1$, for the curves around the walls $\omega_1, \dots, \omega_{N-1}$ do not depend on b and are left invariant by the gauge transformation, i.e. $\Omega^{-1} \tilde{M}_i \Omega = M_i$. One may check that the monodromy matrices M_r , $r = 0, \dots, N$ satisfy all the defining relations of the fundamental group \mathcal{M}_N . In addition they possess the following Hecke property

$$\begin{aligned} (M_r - 1)(M_r - \gamma_r) &= 0, \quad \text{where} \\ \gamma_0 &= e^{2\pi i(a+b)}, \quad \gamma_i = e^{\pi i(1+\epsilon)}, \quad \gamma_N = e^{2\pi i(a-b)} \end{aligned} \quad (3.45)$$

¹⁷The mathematical origin of this θ is that it is twice the minuscule co-weight of root system C_N . In this way, it is related to a symmetry that interchanges the 0-th and N -th affine root, see previous discussion.

for $r = 0, \dots, N$ and $i = 1, \dots, N - 1$. The Hecke property of the monodromy matrix is obvious from the eqs. (3.40), (3.41) and (3.44). The formulas for the monodromy representation we have displayed here imply that the functions Φ^V we defined above indeed have trivial monodromy with respect to g_{i_1}, \dots, g_{i_r} .

One might wonder why these formulas are so similar to the ones in the Pöschl-Teller problem. The reason is however not so difficult to grasp. If we start our curve near asymptotic infinity we can reach each of the walls separately, i.e. we can move near any of the wall ω_i while staying far away from the others $\omega_j, j \neq i$. In this way, the monodromy problem for a given wall can really be solved within the theory of single variable hypergeometric functions. The walls ω_0 and ω_N are in fact entirely equivalent to the walls in the Pöschl-Teller problem while the intermediate ones possess a different dependence on the coupling constants. The latter may be understood through a replacement $b \rightarrow 0$ and $a \rightarrow (\epsilon - 1)/2$ in the expression for M_N .

We have also promised some more comments on the origin of the multiplicities that occur in the second row of the C matrices. These may be traced back to symmetries of the Harish-Chandra functions. Indeed, it is well known that [93]

$$\Phi(\lambda, k''; u) = \Theta_0(-(2k)''; u) \Phi(\lambda, k; u) \quad \text{where} \quad \Theta_0(k; u) = \prod_{\Sigma_0^+} \left(2 \sinh \frac{\langle \alpha, u \rangle}{2} \right)^{k_\alpha} \quad (3.46)$$

and Σ_0^+ denotes the set of middle and long positive roots, i.e. in the product that defines Θ_0 we do not include factors for the short roots. In terms of our parameters (a, b, ϵ) the involution reads $(a'', b'', \epsilon'') = (-a, -b, 2 - \epsilon)$.

Before we return to our discussion of the special case of $N = 2$, we want to conclude the general discussion on the monodromy representation of the affine braid group by stating a general result from [76] (proposition 4.3.10) that provides a criterion for the irreducibility of the $|W_N|$ -dimensional representation in the space of Harish-Chandra functions. According to this theory, the representation we have described is irreducible, provided that the momenta λ and the multiplicities k are sufficiently generic, i.e. that

$$\langle \lambda, \alpha^\vee \rangle \notin \mathbb{Z} \quad \text{and} \quad \langle \lambda, \alpha^\vee \rangle + \frac{k_\alpha/2}{2} + k_\alpha \notin \mathbb{Z} \quad (3.47)$$

for all roots α of the BC_N root system. When one of these conditions is violated, the monodromy representation may contain non-trivial subrepresentations. We will see one such example below.

Example. Let us discuss some more details for the case of $N = 2$ that is most relevant in the context of conformal field theory. We have noted already that the corresponding Weyl group possesses $|W_2| = 8$ elements. The eight Harish-Chandra functions $\Phi(w\lambda, k; z), w \in W_2$ give rise to an eight-dimensional monodromy representation of the fundamental group \mathcal{M}_2 . The latter is generated by three elements g_0, g_1, g_2 subject to the following relations

$$g_0 g_2 = g_2 g_0, \quad g_0 g_1 g_0 g_1 = g_1 g_0 g_1 g_0 \quad g_1 g_2 g_1 g_2 = g_2 g_1 g_2 g_1.$$

These are associated with the three walls in figure 3. Among the linear combinations of Harish-Chandra functions, there are a few that we would like to highlight. The first one is associated with the subgroup $V \subset W_2$ that is generated by w_1 . According to the general theorem we quoted above, the following w_1 symmetric combination of Harish-Chandra functions

$$\Phi^s(\lambda, k; z) = c_1(\lambda_1, \lambda_2, k)\Phi(\lambda_1, \lambda_2, k; z) + c_1(\lambda_2, \lambda_1, k)\Phi(\lambda_2, \lambda_1, k; z).$$

is regular at the wall ω_1 , see figure 3. Note that w_1 acts on momenta of Harish-Chandra functions by exchanging λ_1 and λ_2 . The functions Φ^s coincide with the functions Φ^V with $V = \{e, w_1\} \subset W_2$ we introduced above up to a constant (z -independent) factor. Since Φ^s is analytic at the wall ω_1 , it can be continued from A_2^+ to A_2^E and further into $A_2^L = \tilde{A}_2^+$. On the other hand it fails to be analytic at the walls ω_0 and ω_2 . In order to obtain a regular wave function at the wall ω_2 and hence in a neighborhood of the domain A_2^+ , we must add a whole 8-dimensional orbit of Harish-Chandra functions under the action of the Weyl group on the asymptotic data λ . The coefficients are given by the Harish-Chandra c -function that we defined in eq. (3.31).

If instead we want a wave function that is regular at the walls ω_1 and ω_0 and hence in a neighborhood of \tilde{A}_2^+ we must sum the twisted Harish-Chandra functions¹⁸

$$\begin{aligned} \tilde{\Phi}(\lambda_1, \lambda_2; a, b, \epsilon; u_1(z), u_2(z)) &= e^{i\pi(\epsilon/2 - \lambda_1 + \lambda_2)} \Phi(\lambda_1, \lambda_2; a, b, \epsilon; u_1(z), u_2(z)) \\ &= \Phi(\lambda_1, \lambda_2; a, -b, \epsilon; \tilde{u}_1(z), \tilde{u}_2(z)) \end{aligned} \quad (3.48)$$

over an entire orbit of the Weyl group W_2 with coefficients given by a twisted version \tilde{c} of the Harish-Chandra c -functions (3.31) (i.e. in which b is replaced by $-b$). Since the element w_2 of the Weyl group does not commute with the gauge transformation Ω , the resulting function $\tilde{\Phi}^W$ is not a constant multiple of Φ^W . By construction, the Heckman-Opdam hypergeometric functions $F^+ = \Phi^W$ and $\tilde{F}^+ = F^L = \tilde{\Phi}^W$ trivialize two monodromies each, e.g.

$$M_1 F^+ = F^+, \quad M_2 F^+ = F^+$$

and similarly for $\tilde{F}^+ = F^L$ but with M_0 instead of M_2 . This concludes our discussion of wave functions on A_2^+ and the Lorentzian domain A^L .

The hypergeometric functions we constructed in the previous example provide physical wave functions for the domain A_2^+ and the Lorentzian domain $\tilde{A}_2^+ = A^L$, respectively. Their construction is well known in the mathematical literature. For our purposes below we are also interested in the physical wave functions for the Euclidean domain A^E . As far as we know, there exists no general theory for these functions, but for the specific example of $N = 2$ such wave functions have been discussed in the context of conformal field theory [56, 94]. The former paper also contains a partial characterization in terms of monodromies. Recall that the Euclidean domain A^E is bounded by the wall ω_1 . To be slightly more precise, the strip is bounded by two semi-infinite lines satisfying $u_1 = u_2$ (up to shifts by $2\pi i$) which arise from the wall ω_1 and a compact interval along $u_1 = i\varphi = -u_2$ that

¹⁸We assume $0 < \Im u_1 < 2\pi$, $-2\pi < \Im u_2 < 0$ here.

is associated with the image of the wall ω_1 under the Weyl reflection with w_2 . Consequently, one can characterize the physical wave functions F^E for the Euclidean strip A^E through the following three monodromy conditions

$$\begin{aligned} M_1 F^E(\lambda_1, \lambda_2, k; z_1, z_2) &= F^E(\lambda_1, \lambda_2, k; z_1, z_2) \\ M_2^{-1} M_1 M_2 F^E(\lambda_1, \lambda_2, k; z_1, z_2) &= F^E(\lambda_1, \lambda_2, k; z_1, z_2), \\ \tilde{M}_2^{-1} M_1 \tilde{M}_2 F^E(\lambda_1, \lambda_2, k; z_1, z_2) &= F^E(\lambda_1, \lambda_2, k; z_1, z_2). \end{aligned} \quad (3.49)$$

It is easy to see that these three conditions cannot be solved simultaneously unless the $\lambda_1 - \lambda_2 - k_3$ is a non-negative integer. From now on until the end of this section we will assume that the parameter ϵ is generic, i.e. $\epsilon/2$ is not an integer. In this case, the unique (up to normalization) solution is given by

$$F_{\lambda_1, \lambda_2}^E(z_1, z_2) \sim \tilde{\Phi}(\lambda_1, \lambda_2, k; z_1, z_2) + N(\lambda_1, \lambda_2, k) \tilde{\Phi}(-\lambda_2, -\lambda_1, k; z_1, z_2) \quad (3.50)$$

with

$$\begin{aligned} N(\lambda_1, \lambda_2, k) &= \frac{c_1(-\lambda_2, \lambda_1; k) M_2(\lambda_1, \lambda_2; k)_{12}}{c_1(\lambda_1, -\lambda_2; k) M_2(-\lambda_2, -\lambda_1; k)_{12}} \\ &= 4^{2\lambda_1 + 2\lambda_2} \frac{\Gamma(-\lambda_1 - \lambda_2) \Gamma(\frac{\epsilon}{2} + \lambda_1 + \lambda_2)}{\Gamma(\frac{\epsilon}{2} - \lambda_1 - \lambda_2) \Gamma(\lambda_1 + \lambda_2)} \frac{\Gamma(-2\lambda_2) \Gamma(1 - 2\lambda_2) \Gamma(\frac{1}{2} + \lambda_1 \pm a) \Gamma(\frac{1}{2} + \lambda_1 \pm b)}{\Gamma(2\lambda_1) \Gamma(1 + 2\lambda_1) \Gamma(\frac{1}{2} - \lambda_2 \pm a) \Gamma(\frac{1}{2} - \lambda_2 \pm b)} \end{aligned} \quad (3.51)$$

where $\Gamma(x \pm y)$ is the conventional shorthand for $\Gamma(x \pm y) = \Gamma(x+y)\Gamma(x-y)$ and $M_2(\lambda_1, \lambda_2)_{12}$ denotes a matrix element of the monodromy matrix (3.41) for $N = 2$. While the first two monodromy conditions in the list (3.49) possess four linearly independent solutions, the third condition puts the wave function on the semi-infinite strip and hence imposes strong additional constraints. As in our discussion of the BC_1 theory, regular wave functions can only exist for a discrete set of values $n = \lambda_1 - \lambda_2 - \epsilon/2 = 0, 1, 2, \dots$. Since the strip is infinitely extended in one direction, the sum $\lambda_1 + \lambda_2$ can assume any imaginary value. When $u_1 + u_2$ becomes infinite, the wave functions F^E factorize into a product of Poeschl-Teller wave function of the form (3.6)¹⁹ and a plane wave in the coordinate $u_1 + u_2$. The latter possesses an incoming and an outgoing component.

Let us conclude this discussion of the Euclidean domain with a few general observations concerning the structure of the monodromy representations that occur when we specialize the spin $\lambda_1 - \lambda_2 - \epsilon/2$ to be integer. As we will argue now, the corresponding eight-dimensional representations of the affine braid group are indecomposable but no longer irreducible, i.e. they contain a non-trivial invariant subspace which turns out to be four-dimensional. This is consistent with the criterion (3.47). In fact, for $\alpha = -e_1 + e_2$ it states

$$-\lambda_1 + \lambda_2 + \epsilon/2 \notin \mathbb{Z}$$

which is violated for the representations we are considering. It is indeed easy to see that the four Harish-Chandra functions

$$\tilde{\Phi}(\lambda_1, \lambda_2, k; z_1, z_2), \tilde{\Phi}(-\lambda_2, \lambda_1, k; z_1, z_2), \tilde{\Phi}(\lambda_1, -\lambda_2, k; z_1, z_2), \tilde{\Phi}(-\lambda_2, -\lambda_1, k; z_1, z_2)$$

¹⁹With parameters a, b replaced by $a \rightarrow (\epsilon - 1)/2$ and $b \rightarrow 0$ to account for the strength of the potential at the wall ω_1 .

span a four-dimensional subrepresentation. In fact, while the first two functions are invariant under the action of M_1 , the remaining ones are just mapped onto each other. The monodromy matrix M_2 , on the other hand, mixes the first and the third function and similarly the second and the fourth. All four functions are eigenfunctions of the generators ℓ_i and hence no element within the affine braid group can ever get us out of the space spanned by these four functions. On the other hand, the remaining four Harish-Chandra functions from the eight-dimensional monodromy representation do not span an invariant subspace, i.e. the monodromy representations are indecomposable when $\lambda_1 - \lambda_2 - \epsilon/2$ is integer. Let us also note that the wave function F^E we constructed above, is contained in the four-dimensional invariant subspace.

4 Wave functions and conformal blocks

After this lightning review of Heckman and Opdam’s work on the Calogero-Sutherland problem we want to explain how all this is related to the theory of conformal blocks for four external scalar fields. We shall begin with a brief reminder on the [Casimir equation](#) for conformal blocks before we explain the relation in the case of the $N = 1$ theory that applies to 4-point blocks of bulk fields in 2-dimensional theories as well as to blocks of two scalar bulk fields in the presence of a boundary in any dimension [95]. Then we turn to [conformal 4-point blocks in any dimension](#) and their relation to the BC_2 Calogero-Sutherland model, following [54]. In particular we will construct higher dimensional conformal blocks and their shadows in terms of the BC_2 Harish-Chandra functions.

4.1 Conformal blocks and the Casimir equation

It is time now to embed the theory of scalar conformal blocks into the framework of the Calogero-Sutherland model that we have sketched in the previous [two sections](#). To begin with, let us recall the precise definition of scalar conformal 4-point blocks as solutions of the so-called Casimir equation [25]. We consider the correlation function of four scalar conformal primary fields of weight $\Delta_i, i = 1, \dots, 4$ in a d -dimensional conformal field theory. Using conformal symmetry, this 4-point correlator can be written as

$$\begin{aligned} & \langle \phi_1(x_1)\phi_2(x_2)\phi_3(x_3)\phi_4(x_4) \rangle \\ &= \frac{1}{x_{12}^{\frac{1}{2}(\Delta_1+\Delta_2)} x_{34}^{\frac{1}{2}(\Delta_3+\Delta_4)}} \left(\frac{x_{14}}{x_{24}}\right)^a \left(\frac{x_{14}}{x_{13}}\right)^b G(z, \bar{z}) \end{aligned} \tag{4.1}$$

with $x_{ij} = x_i - x_j$ and $2a = \Delta_2 - \Delta_1, 2b = \Delta_3 - \Delta_4$. The conformal invariants z, \bar{z} are introduced to parameterize the more familiar cross ratios as

$$u = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2} = z\bar{z}, \quad v = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2} = (1-z)(1-\bar{z}). \tag{4.2}$$

For a Euclidean theory, z, \bar{z} are complex conjugates. When we work with Lorentzian signature, on the other hand, the variables z and \bar{z} are real and taken from the interval

$z, \bar{z} \in [0, 1[$. The function G receives contributions from all the primary fields that can appear in the operator product expansion of the fields ϕ_1 and ϕ_2

$$G(z, \bar{z}) = \sum_{\Delta, l} \lambda_l^{12}(\Delta) \lambda_l^{34}(\Delta) G_{\Delta, l}(z, \bar{z}). \tag{4.3}$$

This expansion separates the dynamically determined coefficients λ of the operator product from the kinematic *conformal blocks* $G_{\Delta, l}$. The latter are eigenfunctions of the quadratic conformal Laplacian D_ϵ^2 ,

$$D_\epsilon^2 G(z, \bar{z}) = \frac{1}{2} C_{\Delta, l} G(z, \bar{z}) \tag{4.4}$$

with eigenvalues

$$C_{\Delta, l} = \Delta(\Delta - d) + l(l + d - 2). \tag{4.5}$$

The form of the conformal Laplacian has been worked out in [25],

$$D_\epsilon^2 := D^2 + \bar{D}^2 + \epsilon \left[\frac{z\bar{z}}{\bar{z} - z} (\bar{\partial} - \partial) + (z^2\partial - \bar{z}^2\bar{\partial}) \right] \tag{4.6}$$

where $\epsilon = d - 2$ and

$$D^2 = z^2(1 - z)\partial^2 - (a + b + 1)z^2\partial - abz. \tag{4.7}$$

\bar{D}^2 is defined similarly in terms of \bar{z} . In order to fully determine the conformal blocks, we impose the following boundary condition

$$G_{\Delta, l}(z, \bar{z}) \sim c_l (z\bar{z})^{\frac{1}{2}(\Delta - l)} (z + \bar{z})^l + \dots \tag{4.8}$$

when z, \bar{z} are close to $z, \bar{z} = 0$ and we choose the normalization factor c_l to be

$$c_l = \frac{(\epsilon/2)_l}{(\epsilon)_l}.$$

The condition (4.8) selects a unique solution of the conformal Casimir equation and therefore completes our definition of scalar 4-point blocks.²⁰

4.2 Blocks and the Pöschl-Teller problem

We can gain some first insight into the relation between conformal blocks and Harish-Chandra functions by setting $d = 2$. The conformal Casimir equation then decomposes into two independent equations that determine the dependence of blocks on z and \bar{z} , respectively. Focusing on the z -dependence leads to the following second order differential equation

$$D^2 G_h(z) = h(h - 1) G_h(z).$$

Note that D^2 and hence the eigenfunctions G_h depend on a, b but we do not display this dependence explicitly. The same equation also appears for the blocks of two scalar fields in

²⁰Our normalization conventions match those of [26] (up to a change $z \leftrightarrow \bar{z}$ which is a symmetry of blocks). To match with [56], one should strip off the prefactor c_l .

the presence of a boundary in d -dimensional conformal field theory, see [95]. Following [54] we perform the gauge transformation

$$\Psi(u) := 2^{1-2h} \frac{(1-z)^{\frac{a+b}{2} + \frac{1}{4}}}{\sqrt{z}} G(z) \tag{4.9}$$

where the coordinates z and u are related through eq. (2.28) as before. Note that we consider the region in which $z \in [0, 1]$ which means that $u = \tilde{u} + i\pi$ with

$$z = -\sinh^{-2} \frac{u}{2} = \cosh^{-2} \frac{\tilde{u}}{2}.$$

Inserting these relations it is easy to see that the function Ψ is an eigenfunction of the Pöschl-Teller Hamiltonian with potential $V_{(a,b)}^{\text{PT}}(u)$ where the coupling constants a, b are the same as the parameters a, b in the conformal Casimir equation, i.e. they are related to the holomorphic weights h_i of the external fields through $a = h_2 - h_1$ and $b = h_3 - h_4$. The eigenvalue $\varepsilon = -\lambda^2$, on the other hand, is determined by the holomorphic weight h of the intermediate field $\varepsilon = -(2h - 1)^2$. Hence we read off that $h = 1/2 - \lambda$.

If we compare the relation (4.9) between conformal blocks G and wave functions Ψ with the gauge transformation Θ between wave functions and (twisted) Harish-Chandra functions we find that

$$G(z) = \vartheta(z)\Phi(\lambda, k; z) := 2^{2a+1-2\lambda} z^{-a} \tilde{\Phi}(\lambda, k; z).$$

Note that the gauge transformation $\vartheta(z)$ that relates the twisted Harish-Chandra functions to conformal blocks takes a very simple form. Similarly, the second twisted Harish-Chandra function $\tilde{\Phi}(-\lambda, k; z)$ is related to the so-called shadow block. Finally, there exists a special linear combination of blocks and their shadows that is analytic along the wall that bounds the Lorentzian domain A_1^L , as discussed in section 3.1. These form a complete basis in the space of functions on the half-line.

4.3 Blocks from Harish-Chandra wave functions

As was observed in [54], the Casimir equation for conformal blocks is equivalent to the Calogero-Sutherland model for reflection group BC_2 . The parameters a, b and ϵ that are determined by the conformal weights of the external scalar and the dimension d coincide with the parameters a, b and ϵ in the potential (2.11). To relate the associated Schrödinger problem on the BC_2 Weyl chamber with the eigenvalue equation (4.4) for the conformal Laplacian we employ

$$\Psi(u_1, u_2) := 2^{d-2\Delta} \prod_i \frac{(1-z_i)^{\frac{a+b}{2} + \frac{1}{4}}}{z_i^{\frac{1}{2} + \frac{\epsilon}{2}}} |z_1 - z_2|^{\frac{\epsilon}{2}} G(z_1, z_2), \tag{4.10}$$

where $z_1 = z$ and $z_2 = \bar{z}$. It is not difficult to verify that this gauge transformation, along with the usual relation (2.28) between z_i and u_i , turns the conformal Laplacian into the Calogero-Sutherland Hamiltonian for the potential (2.11), with the eigenvalue

$$\varepsilon = -d(d-2)/4 - (C_{\Delta, d} + 1)/2.$$

In order to identify the conformal block with the precise eigenfunction of the Calogero-Sutherland model, we must also compare the asymptotic behavior (4.8) with the asymptotics (3.16) of twisted Harish-Chandra functions. Note that the limit $u_1, u_2 \rightarrow \infty$ maps to the limit $z_1, z_2 \rightarrow 0$. In Harish-Chandra theory we perform this limit in the Weyl chamber $A_2^L = \tilde{A}_2^+$ where $\tilde{u}_1 > \tilde{u}_2$. This maps to real cross ratios z_1, z_2 with $0 < z_1 < z_2 < 1$. Taking into account the asymptotics of the gauge transformation (4.10) and of the factor Θ , comparison gives

$$\lambda_1 = \frac{1}{2} + \frac{\epsilon}{2} - \frac{1}{2}(\Delta - l), \quad \lambda_2 = \frac{1}{2} - \frac{1}{2}(\Delta + l). \quad (4.11)$$

If we solve these for the conformal weight Δ and the spin l of the exchanged field, we obtain

$$\Delta = \frac{d}{2} - \lambda_1 - \lambda_2, \quad l = \lambda_1 - \lambda_2 - \frac{\epsilon}{2}. \quad (4.12)$$

In order for the twisted Harish-Chandra wave function to possess oscillatory behavior at infinity, the parameters λ_i must be imaginary. The corresponding values of $\Delta = d/2 + ic$ are then associated with the principal continuous series representation of the conformal group [7].

Note that the asymptotic condition (4.8) is symmetric with respect to the exchange of $z = z_1$ and $\bar{z} = z_2$. Hence the block should not be identified with a single twisted Harish-Chandra function $\tilde{\Phi}(\lambda, k; u)$ but rather with the superposition of $\tilde{\Phi}(\lambda, k; u)$ and $\tilde{\Phi}(w_1\lambda, k; u)$. This is the symmetric superposition $\Phi^s = \Phi^V$ with $V = \{e, w_1\}$ we have briefly discussed at the end of the [previous section](#),

$$G_{\Delta,l}(z, \bar{z}) = \frac{2^{4a+d-2\lambda_1-2\lambda_2}}{(z\bar{z})^a} \left[c_{\alpha_1}(\lambda_1, \lambda_2) \tilde{\Phi}(\lambda_1, \lambda_2, k; z_1, z_2) + c_{\alpha_1}(\lambda_2, \lambda_1) \tilde{\Phi}(\lambda_2, \lambda_1, k; z_1, z_2) \right]$$

where $c_{\alpha_1}(\lambda; k) = \frac{\gamma_{\alpha_1}(\lambda, k)}{\gamma_{\alpha_1}(\rho(k), k)} = \frac{\Gamma(\lambda_1 - \lambda_2)\Gamma(\epsilon)}{\Gamma(\lambda_1 - \lambda_2 + \epsilon/2)\Gamma(\epsilon/2)}$. (4.13)

Here the parameters λ on the right hand side are determined from the conformal weight Δ and spin l of the exchanged field through eqs. (4.11), the weights k_α should be fixed from the conformal weights Δ_i of the external scalars and the spacetime dimension d through

$$k_1 = \Delta_4 - \Delta_3, \quad k_2 = \frac{1}{2}(\Delta_2 + \Delta_3 - \Delta_1 - \Delta_4 + 1), \quad k_3 = \frac{d}{2} - 1$$

and the variables u_i are related to the cross ratios by eq. (4.2). By construction, the conformal block trivializes the monodromy M_1 , i.e.

$$M_1 G_{\Delta,l}(z, \bar{z}) = G_{\Delta,l}(z, \bar{z}).$$

This means that it is regular along the wall ω_1 only. On the other hand, the block fails to be regular along the walls ω_0 and ω_2 . We will come back to this issue in the [next section](#). Let us also note that in the case of integer spin, both twisted Harish-Chandra functions contribute to the block if and only if the dimension d is even. Otherwise, the coefficient $c_{\alpha_1}(\lambda_2, \lambda_1)$ in front of the second Harish-Chandra function vanishes due to a divergent

gamma factor in the denominator that does not cancel against a divergence in the numerator. Hence, for integer l and generic dimension d , the conformal block is given by a single Harish-Chandra function.

As discussed in [54], for integer l ²¹ our z -expansion (3.20) for Harish-Chandra functions implies an expansion for conformal block that was first derived by Dolan and Osborn in [25]. Indeed, when $-\epsilon/2 + \lambda_1 - \lambda_2 = l$ is a non-negative integer, we can use Whipple's identity to rewrite the summands in eq. (3.20) through

$$\begin{aligned} & \frac{(\frac{\epsilon}{2} - \lambda_{12})_{n-m}}{(1 - \lambda_{12})_{n-m}} {}_4F_3 \left(\begin{matrix} \epsilon/2 - \lambda_{12} - m + n, 2\lambda_2 - m, \epsilon/2, -m \\ \epsilon/2 - \lambda_{12} - m, \epsilon/2 + \Lambda_{12} - m, 1 - \lambda_{12} - m + n \end{matrix} ; 1 \right) = \\ & = \frac{\Gamma(\epsilon/2, -\epsilon/2 + 2\lambda_1, \epsilon/2 + \lambda_1 \pm \lambda_2)}{\Gamma(\epsilon, 2\lambda_1, \lambda_1 \pm \lambda_2)} \frac{(\epsilon/2 - \lambda_{12})_{n-m}}{(1 - \epsilon/2 - \lambda_{12})_{n-m}} \frac{(1 - 2\lambda_1)_n}{(1 + \epsilon/2 - 2\lambda_1)_n} \frac{(1 - \Lambda_{12})_m}{(1 - \epsilon/2 - \Lambda_{12})_m} \\ & \quad \times {}_4F_3 \left(\begin{matrix} \epsilon/2 - \lambda_{12} - m + n, \epsilon/2 - \lambda_1 \pm \lambda_2, \epsilon/2 \\ \epsilon/2 - \lambda_{12} - m, 1 + \epsilon/2 - 2\lambda_1 + n, \epsilon \end{matrix} ; 1 \right) \end{aligned}$$

where $\lambda_{12} = \lambda_1 - \lambda_2$ and $\Lambda_{12} = \lambda_1 + \lambda_2$. Once this is inserted into eq. (3.20), using the relation (3.48) we recover the expansion from [25]. On the other hand, the expansion formula in [25] fails to work for non-integer spin while our generalized binomial formula for BC_2 provides a valid solution with prescribed asymptotics. In order to correct the expansion of Dolan and Osborn in case of non-integer spin l one should add a second term containing ${}_4F_3$ on the right hand side of the last identity, which then becomes a particular instance of three-term relation for a balanced ${}_4F_3$. Moreover, the two symbols ${}_4F_3$ that appear on the right hand side of this relation are non-terminating, yet still convergent due to balancedness. The analytical continuation of the left hand side, on the other hand, is trivial, since this function stays terminating, due to a negative integer among the upper parameters. Of course, instead of correcting the expansion formula of Dolan and Osborn one can work with our expansion (3.20) which is valid for integer as well as non-integer spin l .

5 Some applications of scattering theory to conformal blocks

Now that we have explained how the conformal blocks are embedded into the Calogero-Sutherland scattering theory, we can begin to apply some of the general results from Heckman-Opdam theory to conformal field theory. The [first subsection](#) aims at understanding poles and residues of conformal blocks and hence their construction through Zamolodchikov-like recurrence relations. We will recover and extend existing results on scalar blocks in the conformal field theory literature, most notably those in [12, 16, 32], from our results on poles and residues of Harish-Chandra functions, see section 3.3. As an application of our discussion of the monodromy representation in section 3.4, we will approach a recent technical observation by Caron-Huot that was instrumental for deriving a [Gribov-Froissart-like formula](#) for conformal correlators in [56].

²¹See appendix B for some details of this limit.

5.1 Poles and residues of blocks

The first application concerns the analysis of poles of conformal blocks and their residues. Such relations play an important role for the explicit evaluation of blocks via Zamolodchikov-type recursions in the numerical bootstrap. In this context they were first discussed in [32],²² see also [35, 38] for subsequent development. The existing results on poles and their residues were derived²³ from representation theory of the conformal algebra. Here we will re-derive the main results for scalar blocks directly from our understanding of the pole structure of Harish-Chandra functions, see eqs. (3.22) and (3.23), but with an important improvement: as we stressed before, our series expansion and all its consequences in sections 3.2, 3.3 are valid for arbitrary complex spin l . Hence we are able to classify poles of scalar conformal blocks and compute their residues for any complex spin l .

Even though the equations (3.26), (3.27), (3.28), (3.29) were obtained for ordinary Harish-Chandra functions, they also hold for twisted Harish-Chandra functions if we replace Φ by $\tilde{\Phi}$, the parameter b by $-b$ and the domain A^+ by A^L , because of the simple relation (3.48) between the two sets of functions. Consequently, according to our general analysis in subsection 3.3, for u in the fundamental domain that contains A^L , complex values of the multiplicities k and complex momenta $\lambda \in \mathbb{C}^2$, the (normalized) twisted Harish-Chandra function $\exp(\langle -\lambda + \rho(k), u \rangle) \tilde{\Phi}(\lambda; k; u)$ is analytic within the (complex) fundamental domain of u variables, entire in k and meromorphic along the hyperplanes defined by $\langle \lambda, \alpha^\vee \rangle = s$ with $s = 1, 2, \dots$. Typically, the twisted Harish-Chandra functions possess simple poles along these hyperplanes, except where two or more of them meet. At such intersection points the pole orders add up.²⁴ We will see some examples in our discussion later on and also explain how to extract the residues in such cases.

Let us recall that for type BC_2 we distinguished four families of poles corresponding to the roots $e_1, e_2, e_1 + e_2, e_1 - e_2$ (see table 3). For compactness we denote the residues of twisted Harish-Chandra function at these simple poles as

$$\begin{aligned}
 r_1 \left(a, b, \epsilon; \frac{s}{2}, \lambda \right) &= \frac{1}{2} \frac{4^s}{s!(s-1)!} \frac{\left(a + \frac{1-s}{2}, -b + \frac{1-s}{2}, \frac{\epsilon}{2} + \lambda - \frac{s}{2}, 1 - \frac{\epsilon}{2} + \lambda - \frac{s}{2} \right)_s}{\left(\lambda - \frac{s}{2}, 1 + \lambda - \frac{s}{2} \right)_s}, \\
 r_2 \left(a, b, \epsilon; \lambda, \frac{s}{2} \right) &= \frac{1}{2} \frac{4^s}{s!(s-1)!} \left(a + \frac{1-s}{2}, -b + \frac{1-s}{2} \right)_s, \\
 r_{12}^+ (a, b, \epsilon; -\lambda + s, \lambda) &= \frac{4^{2s}}{s!(s-1)!} \\
 &\quad \times \frac{\left(\frac{1}{2} + a + \lambda - s, \frac{1}{2} - b + \lambda - s, \frac{1}{2} + a - \lambda, \frac{1}{2} - b - \lambda, \frac{\epsilon}{2}, 1 - \frac{\epsilon}{2} \right)_s}{(-2\lambda, 1 - 2\lambda)_{2s}}, \\
 r_{12}^- (a, b, \epsilon; \lambda + s, \lambda) &= \frac{1}{s!(s-1)!} \left(\frac{\epsilon}{2}, 1 - \frac{\epsilon}{2} \right)_s.
 \end{aligned} \tag{5.1}$$

²²For scalar blocks, the residue formulas were first presented in [12, 16].

²³The derivation of residues for scalar blocks presented in [32] (appendix B.1) is somewhat incomplete for Type III poles, but the results are correct.

²⁴Namely, for a subvariety where n such hyperplanes meet, multiplying our function by n linear forms cancelling those poles, we get an isolated singularity which is removable by Hartogs' theorem [96, 97].

We will often refer to these four series of poles as e_1 , e_2 , $e_1 + e_2$ and $e_1 - e_2$ series, corresponding to the simple roots in the BC_2 root system they are associated with. In our subsequent analysis of poles and residues of conformal blocks we will assume that the dimension $d \geq 2$ is integer in order to reduce the number of cases we have to investigate separately. Let us stress, however, that this restriction is merely of pedagogical origin. On the other hand, we will keep the spin $l = \lambda_1 - \lambda_2 - \epsilon/2$ arbitrary to exploit at least some of the potential our approach has compared to the existing representation theoretic derivations of residue formulas. Of course, our analysis with fixed dimension d will also give us access to integer values of l .²⁵

In order to simplify notations, let us introduce a new symbol \mathcal{G} for the following function

$$\mathcal{G}_{\Delta,l}(\lambda_i; z_i) = G_{\Delta,l}(\lambda_i, z_i) \frac{(z_1 z_2)^a}{4^{2a+\epsilon/2+1-\lambda_1-\lambda_2}}, \tag{5.2}$$

that is obtained by stripping off a simple factor from the block G . According to our formula (4.13) the new functions \mathcal{G} are linear combinations of twisted Harish-Chandra functions of the form

$$\mathcal{G}_{\Delta,l}(\lambda_i; z_i) = c_{\alpha_1}(\lambda_1, \lambda_2) \tilde{\Phi}(\lambda_1, \lambda_2; k; z_1, z_2) + c_{\alpha_1}(\lambda_2, \lambda_1) \tilde{\Phi}(\lambda_2, \lambda_1; k; z_1, z_2).$$

In going from twisted Harish-Chandra functions to these blocks, the pole patterns can change since some singularities of the two summands may cancel each other. Indeed, we now show that all residues related to the root $\alpha_1 = e_1 - e_2$ cancel in the block. Although this follows from general theorems of [86], it is also easy to see using our expressions for residues of Harish-Chandra functions. Let us first check this statement in case ϵ is not a positive even integer. Using eq. (5.1) we then find

$$\begin{aligned} \text{Res}_{\lambda_{12}=s} \mathcal{G}_{\Delta,l} &= \frac{\Gamma(s, \epsilon)}{\Gamma(s + \epsilon/2, \epsilon/2)} \text{Res}_{\lambda_{12}=s} \tilde{\Phi}(\lambda_1, \lambda_2) + \text{Res}_{\lambda_{12}=s} \frac{\Gamma(\lambda_{12}, \epsilon)}{\Gamma(-\lambda_{12} + \epsilon/2, \epsilon/2)} \tilde{\Phi}(\lambda_2, \lambda_1) \\ &= \frac{\Gamma(\epsilon)}{\Gamma(\epsilon/2)^2} \frac{(1 - \epsilon/2)_s}{s!} (1 + (-1)) \tilde{\Phi}(\lambda_2, \lambda_1) = 0. \end{aligned} \tag{5.3}$$

In the first line we set $\lambda_{12} = \lambda_1 - \lambda_2$ and we used the standard notation $\Gamma(A, B) = \Gamma(A)\Gamma(B)$. Indeed, the residues from the first and second term cancel each other. Now let us see what happens for $\epsilon \in 2\mathbb{Z}_{\geq 0}$.²⁶ If $s < \epsilon/2$, our above calculation is still valid. As soon as s becomes equal to $\epsilon/2$ or exceeds this value, the residue of the first summand vanishes due to eq. (5.1) while the residue of the second term vanishes because the ratio as one of the two gamma functions in the second term becomes finite in the limit. The absence of poles in our $e_1 - e_2$ series is of course consistent with [32] where three series of poles were found to occur for a scalar block. More precisely, our first series of poles that is associated with the root e_1 corresponds to Type II in [32] while the series that come with the roots e_2 and $e_1 + e_2$ are Type I and III, respectively.

²⁵As we comment in appendix B, a consistent (usual) rule for obtaining special cases as limits of generic expressions is to first take limits for multiplicity parameters such as the dimension or $\epsilon = d-2$ and then for combinations of the momenta λ_i such as the spin l . Performing the limits in opposite order gives different results.

²⁶For $\epsilon = 0$ there is a singularity in the gamma factors which is obviously removable and just gives the factor of $1/2$.

In all three series, the poles are located along a discrete family of 1-dimensional lines of real momenta $\lambda_i \in \mathbb{R}$. In the first family, $\lambda_1 = s/2$ is constant and $\lambda_2 = \lambda$ is free while in the second family we have $\lambda_2 = s/2$ and $\lambda_1 = \lambda$ is free. The third family that comes with the root $e_1 + e_2$ is described by the equations $\lambda_1 + \lambda_2 = s$. So, obviously, we can have double and even triple intersections of these lines. This motivates to distinguish the following five different cases, see also the discussion below,

- A) $\epsilon \in 2\mathbb{Z}_{\geq 0}$ and $l \in \mathbb{Z}_{\geq 0}$
- B) $\epsilon \in 2\mathbb{Z}_{\geq 0} + 1$ and $l \in \mathbb{Z}_{\geq 0} + 1/2$
- C) $\epsilon \in 2\mathbb{Z}_{\geq 0}$ and $l \in \mathbb{Z}_{\geq 0} + 1/2$
- D) $\epsilon \in 2\mathbb{Z}_{\geq 0} + 1$ and $l \in \mathbb{Z}_{\geq 0}$
- E) $\epsilon \in \mathbb{Z}_{\geq 0}$ and $l \notin \mathbb{Z}_{\geq 0}/2$.

Only two of these cases, namely A and D, involve integer spins and so for these two families our results match the outcome of the analysis in [32]. Case E we will refer to as the case of ‘generic momenta’. It will turn out that cases A and C give rise to double poles while even triple poles appear in case B. Cases D and E, on the other hand possess simple poles only and hence they are the easiest to analyse.

Before we now discuss the poles coming from the Harish-Chandra functions, we want to make one important observation concerning the first e_1 series of poles. From the explicit formula for the residue we listed above one can infer that, when the spin l is integer, i.e. in cases A and D, the residue vanishes for all but the first l values of $s = 1, \dots, l$. Hence in this case we only have a finite number of singular lines of type e_1 . For the values l assumes in case B and C on the other hand, there is an infinite family of e_1 poles with non-vanishing residues. We have depicted the singular lines (light blue) for the cases A-D in figure 5 in order to illustrate the following discussion of these cases.

- A) As we have just discussed, case A involves a finite number of e_1 singularities. After we fix l to be a positive integer (see red line in figure 5), we obtain a finite set of simple poles at $\Delta = l + \epsilon, l + \epsilon - 1, \dots, \epsilon + 1$ from the e_1 series, a finite set of simple poles at $\Delta = \epsilon/2, \epsilon/2 - 1, \dots, 1 - l$ from the $e_1 + e_2$ series and finally, an infinite number of double poles with $\Delta = -l, -l - 1, \dots$ from the collision of the e_2 and the $e_1 + e_2$ series.
- B) This case is similar to the previous, except that now we have infinitely many e_1 singularities. This implies that even triple pole collisions can occur. After we fix l we obtain (see upper right image in figure 5) single poles for $\Delta = l + \epsilon, l + \epsilon - 1, \dots, \epsilon/2 + 1$, (from e_1 series) double poles for $\Delta = \epsilon/2, \epsilon/2 - 1, \dots, 1 - l$ (from the intersection of e_1 and $e_1 + e_2$ series) and an infinite number of triple poles for $\Delta = -l, -l - 1, \dots$ from the triple intersections of all three singularities.
- C) As in the previous case there are infinitely many e_1 singularities, but now the choice of d and l implies that we do not encounter triple poles. In fact, the red line in the lower left image of figure 5 only cuts through intersections of the e_1 and the e_2 series.




Figure 5. Illustration of the poles positions in case A-D. The three sets of blue lines depict the position of singularities for the three different series with e_1 singularities along the vertical lines, e_2 along horizontal and $e_1 + e_2$ along diagonal lines. Note that in cases A and D there is a finite number of e_1 singularities. The dotted red lines depict the choice of spin l , i.e. they run along the curves $\lambda_2 = \lambda_1 - l - \epsilon/2$. For a fixed choice of l we see those as the singularities that lie on the red lines. In several cases these involve collisions of two or even three poles. The corresponding conformal weight at which the singularity occurs is determined as $\Delta = d/2 - \lambda_1 - \lambda_2$.

Consequently, after fixing l we see single poles for $\Delta = l + \epsilon, l + \epsilon - 1, \dots, 1 - l$ and $\Delta = \epsilon/2, \epsilon/2 - 1, \dots$ as well as double poles for $\Delta = -l, -l - 1, \dots$

D, E) All three families of poles are simple.

This completes our analysis of poles of conformal blocks in complex momentum space. Let us note that they all appear for real values of λ_1, λ_2 and outside the cone that is allowed by unitarity bounds.

In our analysis of residues we will first state the complete results for the generic momenta, i.e. case E. For the three different families of poles, the residues are given by

$$\begin{aligned} \text{Res}_{\lambda_1=s/2} \mathcal{G}_{\Delta,l}(z, \bar{z}) = & r_1 \left(a, b, \epsilon; \frac{s}{2}, \lambda_2 \right) \frac{\left(-\frac{s}{2} - \lambda_2 \right)_s}{\left(\frac{\epsilon-s}{2} - \lambda_2 \right)_s} \left[c_{\alpha_1} \left(-\frac{s}{2}, \lambda_2 \right) \tilde{\Phi} \left(-\frac{s}{2}, \lambda_2 \right) \right. \\ & \left. + c_{\alpha_1} \left(\lambda_2, -\frac{s}{2} \right) \tilde{\Phi} \left(\lambda_2, -\frac{s}{2} \right) \frac{\sin \left(\pi \left(\frac{s}{2} - \lambda_2 \right) \right) \sin \left(\pi \left(\frac{s-\epsilon}{2} + \lambda_2 \right) \right)}{\sin \left(\pi \left(\frac{s}{2} + \lambda_2 \right) \right) \sin \left(\pi \left(\frac{s+\epsilon}{2} - \lambda_2 \right) \right)} \right], \end{aligned} \quad (5.4)$$

$$\begin{aligned} \text{Res}_{\lambda_2=s/2} \mathcal{G}_{\Delta,l}(z, \bar{z}) = r_2 \left(a, b, \epsilon; \lambda_1, \frac{s}{2} \right) \frac{\left(\frac{\epsilon-s}{2} + \lambda_1 \right)_s}{\left(-\frac{s}{2} + \lambda_1 \right)_s} \left[c_{\alpha_1} \left(\lambda_1, -\frac{s}{2} \right) \tilde{\Phi} \left(\lambda_1, -\frac{s}{2} \right) \right. \\ \left. + c_{\alpha_1} \left(-\frac{s}{2}, \lambda_1 \right) \tilde{\Phi} \left(-\frac{s}{2}, \lambda_1 \right) \frac{\sin \left(\pi \left(\frac{s}{2} + \lambda_1 \right) \right) \sin \left(\pi \left(\frac{s+\epsilon}{2} - \lambda_1 \right) \right)}{\sin \left(\pi \left(\frac{s}{2} - \lambda_1 \right) \right) \sin \left(\pi \left(\frac{s-\epsilon}{2} + \lambda_1 \right) \right)} \right], \end{aligned} \quad (5.5)$$

$$\begin{aligned} \text{Res}_{\lambda_1=-\lambda+s, \lambda_2=\lambda} \mathcal{G}_{\Delta,l}(z, \bar{z}) = r_{12}^+ (a, b, \epsilon; -\lambda+s, \lambda) \left[c_{\alpha_1} (-\lambda, \lambda-s) \tilde{\Phi} (-\lambda, \lambda-s) \right. \\ \left. + c_{\alpha_1} (\lambda-s, -\lambda) \tilde{\Phi} (\lambda-s, -\lambda) \right] = r_{12}^+ \mathcal{G}_{d/2+s,l}(z, \bar{z}). \end{aligned} \quad (5.6)$$

In the last family, the residue of the block at $(\Delta, l)_* = (d/2 - s, l)$ turns out to be proportional to the block with permuted and reflected momenta, i.e. with $(\Delta, l) = (d/2 + s, l)$. For the first two families, however, the residues of blocks are not proportional to a block anymore, unless the spin l becomes integer in which case

$$\text{Res}_{\lambda_1=s/2} \mathcal{G}_{\Delta,l}(z, \bar{z}) \sim \mathcal{G}_{d-1+l, l-s}(z, \bar{z}), \quad \text{Res}_{\lambda_2=s/2} \mathcal{G}_{\Delta,l}(z, \bar{z}) \sim \mathcal{G}_{1-l, l+s}(z, \bar{z}).$$

These relations for conformal blocks with integer l mimic those of the Harish-Chandra functions, see table 2. Within our list of poles (cases A-D), there are two cases in which the residues of the first two families of poles are proportional to a single block. Actually, this happens in case D since the coefficient in front of the second Harish-Chandra function vanishes, see our discussion in the [previous section](#), so that the block is actually given by a single Harish-Chandra function. In addition, it can also happen in case A for poles of the e_1 series and for those poles of $e_1 + e_2$ series which do not collide to the ones from e_2 series. On the other hand, double poles arising from the colliding series of e_2 and $e_1 + e_2$ singularities need a special treatment. Thus, once again, we are in agreement to the conformal group analysis of [32].

When double or triple pole collision appear in the cases A-C, the residues, of course, will look differently from the expressions we gave above for the generic case. In order to evaluate these such residues for an n -th order pole at some point $\Delta = \Delta^*$ we use the standard prescription

$$\text{Res}_{\Delta=\Delta^*} \mathcal{G}_{\Delta,l} = \lim_{\Delta \rightarrow \Delta^*} \frac{\partial^n}{\partial \Delta^n} \left[(\Delta - \Delta^*)^n \mathcal{G}_{\Delta,l} \right].$$

in our explicit expansion (A.1) (or its u -counterpart). Computing the corresponding residues for case A-C is in principle straightforward now, but we refrain from giving explicit expressions here. Our analysis provides a new view on the ‘irregularities’ that were noticed in [32, 35], when poles of higher orders appear or residues are not proportional to blocks. From our ‘analytically continued’ point of view, such problems are resolved into two separate issues. First, the blocks are linear combinations of twisted Harish-Chandra functions which for generic parameters leads to a linear combination of twisted Harish-Chandra functions rather than a single block. Furthermore, there is an issue of non-generic momenta λ for which care should be taken in defining Harish-Chandra functions as described in sections 3.2, 3.3 and appendix B.

Before we conclude this subsection, we want to discuss in a bit more detail in which respect our analysis advances the one in [32], where the authors analysed conformal blocks as sums over states in certain parabolic Verma modules of the conformal group. While Penedones et al. presented a nice general pattern of pole counting and calculating of residues of conformal blocks labeled by representations of the conformal group, we instead focused on the case of scalar blocks while applying the finer optics of Calogero-Sutherland scattering theory. Our analysis treats blocks as meromorphic functions in the labels of the intermediate field, i.e. the momenta (λ_1, λ_2) for scalar blocks, so that we can really explore a full (complexified) parameter space in momenta λ .²⁷ This also gives some new singularities, in particular the triple poles for odd dimension d and half-integer l . Blocks with non-integer spin l have recently started to play an important role e.g. in the Froissart-Gribov type formula of [56], see also the [next subsection](#). What we have seen here is just a first illustration of a recurrent theme in Calogero-Sutherland theory. In fact, the integrable DAHA [52] setting whose relation to conformal blocks we are developing is tailored to explorations of full complex parameter spaces. We will dwell on algebraic structures and their implications in a forthcoming paper [55]. At the same time, it seems to us that analytical continuation in dimension d and representation labels would be difficult to achieve with the representation theoretic methods of [32], or at least requires to implement significant new technology such as the interpolating Karoubian categories for representation theories of conformal groups [98] to put it on a firm ground.

5.2 On the derivation of a Gribov-Froissart formula for CFT

In a recent paper [56], Caron-Huot derived a Lorentzian inversion formula that allows to obtain the operator product expansion coefficients from the discontinuity of the four-point correlation function $\mathcal{G}(z, \bar{z})$, see also [67] for an alternative derivation and [71] for recent progress in a setup of conformal defects. The formula actually extracts a function that is analytic in the spin l . The mathematical counterpart/origin of formulas of Gribov-Froissart type goes under the name of Ramanujan’s master theorem, see [99] (p.297) and [100] for a more recent review. Related inversion formulas for BC_N Calogero-Sutherland Hamiltonians have been studied in the mathematical literature [101]. To explain the basic setup let us stress that the (twisted) Heckman-Opdam hypergeometric functions on the usual domain A^+ or the Lorentzian domain A^L provide an orthonormal basis of wave functions that in many respects is similar to the basis of exponentials $\exp(iku)$ for functions on the real line. Just as the latter give rise to the usual Fourier transform, Heckman-Opdam functions define an interesting integral transform that has been investigated e.g. in [76, 84, 87, 102]. For the special case of BC_1 , this transform, conventionally known as Jacobi (function) transform [80], has also been considered in the conformal bootstrap literature where it was dubbed “ α space transform”, see [64, 65].

Questions that are natural to study for Fourier transforms are also natural to study for the integral transform that is obtained from (twisted) Heckman-Opdam functions. In

²⁷The restrictions we imposed, namely that the dimension d is integer and that $\Re l$ is positive, are both easy to lift.

particular, one may look at how the behaviour of a function is related to the asymptotic behavior of its transform in the parameters l and Δ for various choices of function spaces [103] on the Weyl chamber or the Lorentzian domain. A large body of such results in mathematics go under the name Tauberian theorems [104]. In the physics literature, similar issues were studied in [70, 105] by application of classical (Hardy-Littlewood and Wiener) Tauberian theorems for the Fourier transform, see also [57, 58, 106] for related physical arguments. In conformal field theory, these provide bounds on the asymptotic behaviour of the spectral density that arise from the short distance behaviour of the correlator. Wiener’s Tauberian theorem tailored²⁸ to the BC_1 case can be found in [107], while Tauberian theorems for special cases of BC_N Heckman-Opdam functions coming from representation theory appear in [108].²⁹

A related line of research concerns Bochner-type positivity statements. In the case of ordinary Fourier(-Stieltjes) transform, the latter states that a function on a real line is a transform of a positive measure iff it is continuous and positive-definite [109]. Generalizations to the context of Heckman-Opdam theory are formulated in terms of positive convolution structures [110]. They were addressed in [111, 112] for BC_1 , in [113] for BC_2 and in [114] for BC_N , but with very specific choices of the multiplicities k . This mathematical framework provides the natural context for Caron-Huot’s Froissart-Gribov formula. In fact, the Lorentzian inversion formula of [56] extracts positive operator product coefficients $c(\Delta, l)$ analytic in spin through an integral transform of the discontinuity of the correlation function $\mathcal{G}(z, \bar{z})$. Instead of the usual Heckman-Opdam hypergeometric function F , Caron-Huot obtains an integral transform defined by the block

$$E(z) = G_{l+d-1, \Delta+1-d}, \tag{5.7}$$

i.e. $E(z)$ is built as in eq. (4.13), but with (λ_1, λ_2) replaced by $(-\lambda_1, \lambda_2)$. Similar integral transforms that are obtained from Θ -hypergeometric functions in the terminology of section 3.4 have also been studied in the mathematical literature [115].

In order to derive the conformal Froissart-Gribov formula, Caron-Huot departs from the Euclidean inversion formula that goes back to the early harmonic analysis on the conformal group [7], see also [94], and performs an analytic continuation to Lorentzian signature. The Euclidean inversion formula involves a transform with the wave functions $F^E = F_{\Delta, l}^E, l \in \mathbb{Z}$ which we discussed at the end of section 3.4. Following [94], Caron-Huot denotes this function by the letter F , a nice coincidence with our interpretation of it as a Euclidean hypergeometric function. In order to pass to Lorentzian signature, one needs a good understanding of the continuation of the functions $F = F^E$. In in context of Heckman-Opdam

²⁸A usual Wiener type Tauberian theorem states that the translations of an integrable function span a dense subspace iff its Fourier transform is everywhere non-zero. For Heckman-Opdam transforms the translation is replaced by a generalized translation that is determined by the positive convolution structure.

²⁹In particular, for the BC_1 case, it would be interesting to see whether the Jacobi transform Tauberian theorem of [107] can improve the conformal field theory analysis of [70] in order to constrain the subleading behaviour of spectral density and, perhaps, to show that the spectrum of the exchanged operators approaches the one of generalized free theory. It would also be nice to see if the BC_2 case can provide a rigorous justification for results in [57, 58] on the lightcone limit, as approached in [70] through the usual Wiener Tauberian theorem.

theory this is provided by the monodromy matrices M_i . Let us note that M_1 and \tilde{M}_2 coincide with the inverse of the monodromies which are encoded in eqs. (A.23) and (A.22) of [56]. From our discussion in section 3.4 we know that they provide an eight-dimensional representation of an affine braid group that satisfies Hecke relations (3.45). This insight provides significant algebraic control of the monodromy properties of Harish-Chandra (or pure, in the terminology of [56]) functions that can be exploited without ever using the explicit representation matrices. In order to demonstrate how constraining these algebraic properties are, we want to rephrase the main derivation of [56] (section 3.3) in our context.

Let us first review the general setup. It will be convenient to introduce the following functions

$$F_{(d/2-\lambda_1-\lambda_2, \lambda_1-\lambda_2-\epsilon/2)}(z, \bar{z}) = 2^{4a+d-2\lambda_1-2\lambda_2} (z\bar{z})^{-a} F_{\lambda_1, \lambda_2}^E(z, \bar{z}), \tag{5.8}$$

$$g_{(d/2-\lambda_1-\lambda_2, \lambda_1-\lambda_2-\epsilon/2)}(z, \bar{z}) = 2^{4a+d-2\lambda_1-2\lambda_2} (z\bar{z})^{-a} \tilde{\Phi}(\lambda_1, \lambda_2, k; z, \bar{z}). \tag{5.9}$$

that agree with the corresponding functions in [56] apart from the normalization. Caron-Huot studies the function F in the complex w -plane. In our u -space, the latter is given by an infinite strip that is perpendicular to the space A_2^E in figure 3 and intersects it along the line $u_1 = \ln \sigma + i\theta$ where $\theta \in [0, 2\pi[$, see the grey strip in figure 6. We can parametrize this strip by a complex coordinate $v \equiv v + 2\pi i$ such that $v \mapsto (u_1 = v + \ln \sigma, u_2 = -v + \ln \sigma)$. The parameter v is related to the variable w used by Caron-Huot through the exponential map, i.e. $w = \exp(v)$. The region in which $\Re v \rightarrow -\infty$ is mapped to $w \sim 0$ while we can reach $w \rightarrow \infty$ by sending $\Re v$ to positive infinity.

The strip we have just described intersects the walls in six different places. In fact, as illustrated in figure 6, there are two intersections with the wall ω_1 in $v = 0, i\pi$. Intersections with the walls ω_2 and ω_0 , on the other hand, are located at $v = \pm \ln \sigma$ and $v = \pm \ln \sigma + i\pi$, respectively. After passing to the w -plane, the intersections with the wall ω_0 become the end-points $w = -\sigma, -1/\sigma$ of the t-channel cuts described by Caron-Huot. The intersections with ω_2 map to the end-points of the u-channel cuts in the w -plane. Caron-Huot now wants to continue the function F to $w = 0$, i.e. $v = -\infty$. Along the way, one has to pass by the intersection with the walls ω_1 and ω_2 . While F is regular at the wall ω_1 , i.e. $M_1 F = M_1$, it picks up some monodromy M_2 as it passes by ω_2 . So, we can take the limit $v = -\infty$ by looking at the function $M_2 F$. According to our general characterization of F , the function $M_2 F$ trivializes the monodromy M_1 , i.e.

$$M_1 M_2 F = M_2 F. \tag{5.10}$$

Let us restrict our discussion to generic dimension d , i.e. we assume that $\epsilon/2$ is not integer. As we explained at the end of the [previous section](#), for such generic d the function F is a linear combination of two Harish-Chandra functions,

$$F(z, \bar{z}) = 2^{4a+d-2\lambda_1-2\lambda_2} (z\bar{z})^{-a} \left(\tilde{\Phi}(\lambda_1, \lambda_2, k; z_1, z_2) + k \tilde{\Phi}(-\lambda_2, -\lambda_1, k; z_1, z_2) \right).$$

When we act with M_2 , two more Harish-Chandra functions appear, namely $\tilde{\Phi}(\lambda_1, -\lambda_2)$ and $\tilde{\Phi}(-\lambda_2, \lambda_1)$. Because of the property (5.10) the corresponding functions g combine into a




Figure 6. Illustration of the upper w -half plane used by Caron-Huot inside the domain of Calogero-Sutherland models.

single block $G_{1-l,1-\Delta}$. This new block that appeared while we continued F through the wall ω_2 to the region of small w does not vanish in the limit $w \sim 0$ and thereby it prevents further evaluation of the integral in [56].

The idea of Caron-Huot is to look for an appropriate combination of Harish-Chandra functions that vanish at $w = \infty$ and can cancel the unwanted divergent terms at $w = 0$. It is easy to see that there are two Harish-Chandra functions that vanish at $w = \infty$, namely the functions $\tilde{\Phi}(\lambda_2, -\lambda_1)$ and $\tilde{\Phi}(-\lambda_1, \lambda_2)$. From the corresponding functions $g_{(l+d-1,1-\Delta)}$ and $g_{(l+d-1,\Delta+1-d)}$, see table 1, we can construct the block $E(z) = G_{l+d-1,\Delta+1-d}$ we introduced before in eq. (5.7). By construction, the function E trivializes the monodromy M_1 , i.e.

$$M_1 E(z) = E(z).$$

In order to evaluate its behavior for small w , one has to continue E from large absolute values of w to very small ones. Along the way, we pass the walls ω_2, ω_1 and ω_2 again. Cancellation of the singular terms then means that there exist constants α and β_i such that

$$M_2 F + \alpha(M_1^{-1} M_2 M_1) M_1 M_2 E = \beta_1 g_{(\Delta, 2-l-d)} + \beta_2 g_{(d-\Delta, 2-l-d)}.$$

This is a precise formulation of Caron Huot's condition (3.13) in [56] in the language we have set up. In order to illustrate how to derive such statements using algebra rather than the precise matrix elements the monodromy matrices, we want to prove, restricting to generic $\epsilon/2 \notin \mathbb{Z}$, a closely related statement, namely that there exist constants α and β such that, the following equation holds

$$M_2 F + \alpha M_2 M_1 M_2 E = \beta E. \tag{5.11}$$

In order to derive this claim we shall consider eq. (5.11) as a formula that allows us to construct the functions F from E as

$$F = \beta M_2^{-1} E - \alpha M_1 M_2 E . \tag{5.12}$$

Through a short computation using braid and Hecke relations of the monodromy matrices, see section 3.4, one may show that the function

$$F' := M_1 M_2 E + \gamma_1 \gamma_2 E$$

which is proportional to the right hand side of the previous equation if $-\beta/\alpha = \gamma_1 \gamma_2$, possesses the following properties

$$M_1 F' = F' \quad \text{and} \quad M_2^{-1} M_1 M_2 F' = F' .$$

On the other hand it is easy to see that F' contains only six of the eight Harish-Chandra functions since we apply at most two monodromy matrices to E . Because F' is an eigenfunction of M_1 with unit eigenvalue we conclude that

$$F' = \alpha_1 g_{(\Delta, l)} + \alpha_2 g_{(d-\Delta, l)} + \beta' E .$$

In addition F' was shown to be an eigenfunction of $M_2^{-1} M_1 M_2$ with unit eigenvalue. This fact imposes strong constraints on the coefficients α_i and β' . In fact, as in our discussion of F^E at the end of section 3.4 we can conclude that α_2/α_1 is given by the expression after eq. (3.50) and $\beta' = 0$. Hence, we have established our formula eq. (5.12) and thereby the equation (5.11) we set out to prove. In the argument we have only used braid and Hecke relations of the monodromy matrices along with the fact that the Harish-Chandra functions $g_{(\Delta, l)}$ and $g_{(\Delta, 2-l-d)}$ are eigenfunctions of the monodromy M_1 with unit eigenvalue, a very particular feature of representations with integer spin l and generic $\epsilon/2 \notin \mathbb{Z}$.

6 Conclusions

In this work we began to elaborate on the very fruitful relation between conformal blocks and wave functions of Calogero-Sutherland models that was uncovered in [54]. In our review of Heckman-Opdam theory we focused on features that may be understood from the quantum mechanical treatment of Calogero-Sutherland scattering states. More advanced properties of the Calogero-Sutherland model that relate to its (super-)integrability were not discussed. These will be briefly outlined below and then presented thoroughly in a forthcoming paper [55].

Even without tools from integrability we were able to recover essentially all important results about scalar 4-point blocks. The basic objects in Heckman-Opdam theory are the Harish-Chandra functions. Special cases of these functions also appeared in the recent work [56] where they were called pure functions. Here we obtained a new series expansion for these functions that is valid for non-integer spin l and we derived a number of important properties from it. In particular, we located all poles in λ -space, which is the space of

conformal weights and spins, and we computed their residues. In addition, we reviewed the classical results of Heckman and Opdam on monodromy properties of Harish-Chandra functions. Some of the monodromy matrices were also computed independently for $N = 2$ in [56]. Here, we learned in addition that they give rise to finite dimensional representation of an affine braid group of reflection type that factors through the corresponding Hecke algebra. This algebraic characterization of the monodromy properties of Harish-Chandra (or pure) functions led to a simple derivation of some central observations in [56]. Once we understood how conformal blocks are built out of Harish-Chandra functions, we were also able to rederive many important results on conformal blocks, see e.g. [25, 26, 32], from our new series expansion and the residue formulas for Harish-Chandra functions. In our discussion we focused on some of the most useful properties of blocks. There exist many other consequences of Heckman-Opdam theory that lead to additional insights, a few of which do not seem to be known in the conformal field theory literature.

We expect that the analytical toolkit we outlined in this paper will prove useful for various approaches to the analytical bootstrap, see in particular [56–71, 116–122] and references therein for some recent work. In particular, our explicit understanding of the properties of conformal blocks should provide better control over large spin expansions, sum rules for operator product coefficients and multi-twist operators.

What makes the connection with Calogero-Sutherland models even more useful is that the techniques we have described actually apply much beyond the theory of scalar 4-point blocks. The most immediate extension concerns the study of defect blocks. In fact, scalar 4-point blocks are a very special case of a larger set of conformal blocks that can be used to expand the correlation function of two conformal defects of dimension $p, q < d$. Scalar 4-point blocks correspond to $p = q = 0$, i.e. a 0-dimensional defect should be considered as a pair of points. Correlations functions of two local fields in the presence of a p -dimensional defect have received some attention for $p = d - 1$ in [95] and more generally in [123]. These involve one and two cross ratios, respectively, and the corresponding blocks turn out to be identical to the ones we have studied here for $N = 1$ and $N = 2$. More recently, Gadde also considered setups with $p \neq 0 \neq q$ which can involve a larger number $N > 2$ of cross ratios [73]. As we will show in a forthcoming paper, the conformal blocks for such configurations of two defects can all be built from the Harish-Chandra functions of BC_N Calogero-Sutherland models. Hence, many of the results and techniques we have described in section 3 and the appendices directly apply to such defect blocks.

Another direction concerns applications to spinning blocks. In [124] we have shown that the Casimir equations for spinning blocks in dimension $d \leq 3$ also take the form of a Calogero-Sutherland eigenvalue problem, though with a matrix valued potential. This observation can be extended to arbitrary dimensions, see [125]. The theory of scattering states for such matrix valued potentials is not really developed, at least for the case of BC_N root systems, but the general strategy we reviewed before should still apply. In the case of A_N root system, a restricted set of matrix valued potentials is well understood [52] and some first steps were taken for the most generic ones in [126]. It would be very interesting to extend the results we have described above to matrix Calogero-Sutherland models for BC_2 roots systems, i.e. to study their symmetries, the fundamental domain, monodromy representations, series expansions etc.

Further extensions to superblocks are also feasible. As long as sufficiently many of the external fields are in short (BPS) multiplets, the associated blocks are the same as for the bosonic conformal symmetry. In general, however, the theory of superblocks with external fields in long multiplets is much more involved. On the other hand, as was demonstrated in [127], crossing symmetry involving long multiplets is significantly more constraining when analysed in terms of superblocks rather than their bosonic decomposition. Hence, it seems worthwhile to develop a systematic theory of superblocks. We believe that the corresponding Casimir equations can still be rewritten as eigenvalue equations for a supersymmetric version of Calogero-Sutherland models so that much of the above could be extended to superblocks. This remains an interesting direction for future research.

Let us stress however, that even in the case of scalar 4-point functions we did not even come close to exploring all the existing features of the model. In fact, the most remarkable property of the Calogero-Sutherland model is its (super-)integrability. It furnishes a wealth of additional and very powerful algebraic structure. So far, the only algebra we have seen above was the Hecke algebra that appeared in the context of the monodromy representation. It acts in the $2^N N!$ -dimensional spaces of Harish-Chandra functions $\Phi(w\lambda; z), w \in W_N$, i.e. in finite dimensional subspaces of functions which all possess the same eigenvalue of the Hamiltonian. This is just the tip of a true iceberg of algebraic structure.

In order to describe the contours of the remaining parts and thereby outline the content of our forthcoming paper [55], it is useful to consider the example of the very simplest quantum mechanical system, i.e. of a system of N freely moving particles on the real line. The standard Hamiltonian $H_{free}^x = -\sum_i \partial_i^2$ of such a system is certainly integrable. Actually, H_{free}^x can be built from a set of N commuting first order operators $\mathcal{Y}_i^{free} = \partial_{x_i} = \partial_i, i = 1, \dots, N$, as $H_{free}^x = -\sum_i \mathcal{Y}_i^2$. Although Calogero-Sutherland particles are not free, they enjoy very similar properties. In fact, it is well known that the Calogero-Sutherland Hamiltonian for BC_N roots systems can be constructed out of a set of N commuting first order operators \mathcal{Y}_i^{CS} that are known as *Dunkl operators*. Of course, these Dunkl operators have to account for the non-trivial potentials in the Calogero-Sutherland model and hence possess a non-trivial dependence of the coordinates. Eigenfunctions of the Dunkl operators belong to a slightly larger class of non-symmetric Harish-Chandra functions. These are similar to the Harish-Chandra functions we discussed above except that they are no longer invariant under the action of the Weyl group W_N on coordinates x_i . Within the space of such non-symmetric Harish-Chandra functions, the Dunkl operators give rise to a systems of Knizhnik-Zamolodchikov-like first order differential equations [128]. Once these are solved, symmetric Harish-Chandra functions and in particular the conformal blocks can be obtained through an appropriate projection. The space of non-symmetric Harish-Chandra functions is acted upon by an infinite dimensional algebraic structure that is generated by elements of the Weyl group, Dunkl operators and multiplication with coordinates: the degenerate double affine Hecke algebra (DAHA) or trigonometric Cherednik algebra [52]. In addition to the monodromy algebra we have seen above, it contains generators that do not commute with the Hamiltonian and hence can relate eigenfunctions with different eigenvalues. In this sense, the degenerate DAHA is a spectrum generating symmetry.

This is not the end of the story. In order to prepare for its next chapter, we return to example of N freely moving particles. The eigenfunctions $\psi_\lambda(x)$ in such a system possess a remarkable property: namely, their dependence on the eigenvalues λ_i is the same as for the variables x_i . This implies that $\psi_\lambda(x)$ are also eigenfunctions of a dual second order operator $H_{free}^y = -\sum_i \partial_{\lambda_i}^2$. We have put a superscript y in this Hamiltonian since it controls dependence of the eigenfunctions ψ on the eigenvalues λ of the first order operators \mathcal{Y}_i . The dual Hamiltonian H_{free}^y has precisely the same form as the original one. In this sense, the theory of a freely moving particles may be considered as self-dual. Some of these facts remain true for our Calogero-Sutherland model. As for free particles, the dependence of Calogero-Sutherland wave functions on the eigenvalue λ is controlled by a dual Hamiltonian H^y . For the hyperbolic Calogero-Sutherland model, the latter is a second order difference operator that describes the so-called *rational Ruijsenaars-Schneider model* [129–131]. This dual Hamiltonian H_{rRS}^y is also integrable, i.e. there is a set of first order difference equations that determines the dependence of eigenfunctions on the eigenvalues λ^i . In the case of the free particle we saw that the dual Hamiltonian H^y had the same form as H^x . This is in sharp contrast to the case of the hyperbolic Calogero-Sutherland system where the dual theory is controlled by difference rather than differential equations. It turns out, however, that the setup can be q -deformed such that the self-duality is restored. In order to do so, the rational Ruijsenaars-Schneider model is deformed into the trigonometric one. After this deformation, the dependence of eigenfunctions on both sets of variables is controlled by the same type of difference equations. Upon deformation the degenerate DAHA we briefly mentioned at the end of the previous paragraph becomes a full fledged DAHA. All this will be explained in detail in our forthcoming paper [55] along with a number of applications.

Acknowledgments

We wish to thank T. Bargheer, S. Caron-Huot, A. Gadde, M. Hogervorst, M. Honda, Z. Komargodski, S. Komatsu, P. Kravchuk, M. Lemos, P. Liendo, G. Mack, D. Mazac, C. Meneghelli, V. Narovlansky, D. Poland, L. Rastelli, B. van Rees, N. Reshetikhin, S. Shatashvili, D. Simmons-Duffin, W. Skiba, Zh. Sobko, G. Torrents and R. Yacoby for useful conversations and/or remarks and especially Ivan Cherednik for illuminating discussions. We are also grateful to the participants and organizers of the ‘Back to bootstrap 2016’, ‘Bootstrap 2017’ and ‘Strings 2017’ conferences, of the 2016 Les Houches Summer School on Integrability and of the 2017 Cargese Summer School for the occasion to discuss our work. MI thanks the DESY Theory Group, the Simons Center for Geometry and Physics, Yale University, the Institute for Advanced Study, Duke University and the Math Department of Chapel Hill University, North Carolina, for a stimulating environment and hospitality during various stages of this work. MI is supported in part by Israel Science Foundation (grant number 1989/14), by the ERC STG grant 335182 and by a Koshland Postdoctoral fellowship, partially financed by the Koshland Foundation.

A z - and x -expansions for BC_2 Harish-Chandra function

As far as we know, the formulas for Harish-Chandra functions in this section are new. Let us recall here that conformal blocks are obtained via formulas (4.13) from twisted Harish-Chandra functions, which are related to the Harish-Chandra function by a simple phase shift as given in our eq. (3.48). According to [86], the resulting expansion is convergent for generic λ on the entire shifted fundamental domain.

We start from the formula (3.20) in the main text. For the twisted Harish-Chandra function $\tilde{\Phi}$ it takes the form

$$\begin{aligned} \tilde{\Phi}(\lambda_1, \lambda_2; k_i; z_1, z_2) &= \frac{1}{4^{2a+1+\epsilon/2-\Lambda_{12}}} \sum_{n,m=0}^{\infty} \frac{(1/2+a-\lambda_1, 1/2+b-\lambda_1, \frac{\epsilon}{2}-\lambda_{12})_n}{(1-2\lambda_1, 1-\lambda_{12})_n} \\ &\times \frac{(1/2+a-\lambda_2, 1/2+b-\lambda_2, \epsilon/2+\lambda_{12})_m (1-\epsilon/2-\lambda_{12})_{n-m}}{(1-2\lambda_2, 1+\lambda_{12})_m (-\lambda_{12})_{n-m}} \\ &\times {}_4F_3 \left(\begin{matrix} -n, -m, 1-\epsilon/2, 1-\epsilon/2-\Lambda_{12} \\ 1-\epsilon/2+\lambda_{12}-n, 1-\epsilon/2-\lambda_{12}-m, 1-\Lambda_{12} \end{matrix}; 1 \right) \quad (\text{A.1}) \\ &\times \frac{1}{n!m!} z_1^{\frac{1+\epsilon}{2}+a-\lambda_1+n} z_2^{\frac{1}{2}+a-\lambda_2+m} {}_2F_1 \left(\begin{matrix} \epsilon/2-\lambda_{12}-m+n, \epsilon/2 \\ 1-\lambda_{12}-m+n \end{matrix}; \frac{z_1}{z_2} \right). \end{aligned}$$

Here and in the following we use the shorthand $\lambda_{12} = \lambda_1 - \lambda_2$ and $\Lambda_{12} = \lambda_1 + \lambda_2$. Before we plunge into details, let us first outline its derivation.

The key is an observation from [54] where we directly related scalar conformal blocks to $q \rightarrow 1^-$ degeneration of virtual Koornwinder polynomials [132]. Due to the fast (and still on-going) progress in the end of 1990's, combinatorial aspects of Koornwinder polynomials are quite well-studied, in particular the so-called binomial expansion of Koornwinder polynomials that were obtained in [133]. Now we will factor this connection through the connection to Harish-Chandra functions of specialized parameters (for BC_2 , it means looking at Harish-Chandra functions with $l = \lambda_1 - \lambda_2 - \epsilon/2$ integer) and then will analytically continue in these parameters. To start with, the (non-symmetric q -) Harish-Chandra functions are related to Macdonald-Koornwinder polynomials in [134]. One may write down a binomial formula for virtual Koornwinder polynomials [132] and then use combinatorial expansions for the interpolating polynomials it contains (combinatorial expansions of BC_N interpolation Macdonald polynomials can be found in [133] and the degeneration to BC_N interpolation Jack polynomials in [135]), see [136] for a nice review. Next one has to perform an analytical continuation of the resulting formula to non-integer differences between the row lengths of Young diagrams. By analysing the asymptotics, one may note that the resulting series expansion is a linear combination of $N!$ asymptotically free solutions (q -Harish-Chandra functions), corresponding to a $S_N \subset W_N$ subgroup of the Weyl group. It is then possible to single out a solution with the correct asymptotics.

This way of obtaining explicit series expansions in z , which solves a rather complicated recurrence relation (3.18) for expansion coefficients (after one passes to x -variable), seems to work, provided one has sufficient control of the analytical continuation of

the constituents of the binomial expansion, for any BC_N Harish-Chandra function. A degeneration $q \rightarrow 1^-$ is possible at any step, depending rather on convenience (purely formally, one can take the limit at any point of the derivation, but sometimes this should be justified with more care). Regardless of that, the complexity of explicit computation increases very fast with N , and we are not aware of any closed form expansion of a generic BC_N Harish-Chandra function valid for any N .

Let us look at $N = 2$ case in more detail. The summand of a generalized binomial formula for virtual Koornwinder polynomials (see [132], formula (7.32)) contains a generalized binomial coefficient (which is proportional to a BC_2 interpolation Macdonald polynomial), a (virtual) BC_2 -interpolation Macdonald polynomial and some remaining ratios of q -Pochhammer symbols. To avoid heavy expressions, we will send $q \rightarrow 1^-$ now. This degeneration has different consequences for the two interpolation polynomials: the one entering the generalized binomial coefficient will go to a BC_2 interpolation Jack polynomial, whereas the virtual BC_2 -interpolation polynomial will go to a virtual Jack polynomial ('virtual' — here just means having a monomial of non-integer powers in front of Gegenbauer polynomial) [135].

Using explicit expressions from [132] and a combinatorial expansion for the BC_2 -interpolation Jack polynomial from [133] combined with [135], one can see that the generalized binomial coefficient (undeformed) is proportional to

$$\begin{aligned} & \frac{(\frac{\epsilon}{2} - \lambda_{12})_{n-m}}{(1 - \lambda_{12})_{n-m}} {}_4F_3 \left(\begin{matrix} \epsilon/2 - \lambda_{12} - m + n, 2\lambda_2 - m, \epsilon/2, -m \\ \epsilon/2 - \lambda_{12} - m, \epsilon/2 + \Lambda_{12} - m, 1 - \lambda_{12} - m + n \end{matrix}; 1 \right) \\ &= \frac{(\frac{\epsilon}{2} - \lambda_{12})_n}{(1 - \lambda_{12})_n} \frac{(\frac{\epsilon}{2} + \lambda_{12}, 1 - \Lambda_{12})_m}{(1 - \frac{\epsilon}{2} \pm \lambda_1 - \lambda_2)_m} \\ & \quad \times {}_4F_3 \left(\begin{matrix} -n, -m, 1 - \epsilon/2, 1 - \epsilon/2 - \Lambda_{12} \\ 1 - \epsilon/2 + \lambda_{12} - n, 1 - \epsilon/2 - \lambda_{12} - m, 1 - \Lambda_{12} \end{matrix}; 1 \right) \end{aligned}$$

which we just related to a balanced ${}_4F_3$ appearing in eq. (3.20) via a Whipple transformation [79], combined with inverting of summation index. Clearly, the analytical continuation in $l = \lambda_1 - \lambda_2 - \epsilon/2$ of the ${}_4F_3$ function on the right-hand side will not change anything (the series will stay terminating), unlike the continuation of the one on the left (there the series is terminating for integer l and becomes non-terminating for non-integer, so another ${}_4F_3$ term should appear, see [82] for a full account on the symmetries of ${}_4F_3$ hypergeometric function).

The 'virtual' Jack polynomial, which as we said for $N = 2$, is just a product of a monomial and a Gegenbauer polynomial, upon analytical continuation in l also becomes a non-terminating series

$$\begin{aligned} J(\lambda_1, \lambda_2; \epsilon; z_1, z_2) &= (z_1 z_2)^{\frac{\epsilon/2 + 1 - \Lambda_{12} + n + m}{2}} \\ & \quad \times {}_2F_1 \left(\begin{matrix} \frac{\epsilon}{2} - \lambda_{12} - m + n, \frac{\epsilon}{2} + \lambda_{12} - n + m \\ \frac{\epsilon + 1}{2} \end{matrix}; \frac{1 - \frac{z_1 + z_2}{2\sqrt{z_1 z_2}}}{2} \right). \end{aligned} \tag{A.2}$$

By using a particular quadratic transformation [79] (chapter 3.1) for ${}_2F_1$, this can also be written as

$$\begin{aligned}
 & c_{\alpha_1}(\lambda_1, \lambda_2; k) \frac{(\lambda_{12})_{m-n}}{(\lambda_{12} + \frac{\epsilon}{2})_{m-n}} \frac{(z_1 z_2)^{n + \frac{1+\epsilon}{2} - \lambda_1}}{(z_1 + z_2)^{n-m + \frac{\epsilon}{2} - \lambda_{12}}} \\
 & \quad \times {}_2F_1\left(\frac{\epsilon/2 - \lambda_{12} + n - m}{2}, \frac{1 + \epsilon/2 - \lambda_{12} + n - m}{2}; \frac{4z_1 z_2}{(z_1 + z_2)^2}\right) \\
 & + c_{\alpha_1}(\lambda_2, \lambda_1; k) \frac{(-\lambda_{12})_{n-m}}{(\frac{\epsilon}{2} - \lambda_{12})_{n-m}} \frac{(z_1 z_2)^{m + \frac{1+\epsilon}{2} - \lambda_2}}{(z_1 + z_2)^{m-n + \frac{\epsilon}{2} + \lambda_{12}}} \\
 & \quad \times {}_2F_1\left(\frac{\epsilon/2 + \lambda_{12} + m - n}{2}, \frac{1 + \epsilon/2 + \lambda_{12} + m - n}{2}; \frac{4z_1 z_2}{(z_1 + z_2)^2}\right). \tag{A.3}
 \end{aligned}$$

Let us dress this back with the above binomial coefficient (and other Pochhammers from the generalized binomial formula that we didn't display), take the first summand and strip off the c_{α_1} prefactor. This then gives us the (twisted)³⁰ Harish-Chandra series decomposition as

$$\begin{aligned}
 \tilde{\Phi}(\lambda_1, \lambda_2; k_i; z_1, z_2) &= \frac{1}{4^{2a+1+\epsilon/2-\Lambda_{12}}} \sum_{n,m=0}^{\infty} \frac{(1/2+a-\lambda_1, 1/2+b-\lambda_1, \frac{\epsilon}{2}-\lambda_{12})_n}{(1-2\lambda_1, 1-\lambda_{12})_n} \\
 & \times \frac{(1/2+a-\lambda_2, 1/2+b-\lambda_2, \epsilon/2+\lambda_{12})_m (1-\epsilon/2-\lambda_{12})_{n-m}}{(1-2\lambda_2, 1+\lambda_{12})_m (-\lambda_{12})_{n-m}} \\
 & \times {}_4F_3\left(\begin{matrix} -n, -m, 1-\epsilon/2, 1-\epsilon/2-\Lambda_{12} \\ 1-\epsilon/2+\lambda_{12}-n, 1-\epsilon/2-\lambda_{12}-m, 1-\Lambda_{12} \end{matrix}; 1\right) \\
 & \times \frac{1}{n!m!} \frac{(z_1 z_2)^{\frac{1+\epsilon}{2}+a-\lambda_1+n}}{(z_1+z_2)^{\frac{\epsilon}{2}-\lambda_{12}+n-m}} {}_2F_1\left(\frac{\epsilon/2-\lambda_{12}+n-m}{2}, \frac{1+\epsilon/2-\lambda_{12}+n-m}{2}; \frac{4z_1 z_2}{(z_1+z_2)^2}\right).
 \end{aligned}$$

Notice that, in accord with the general definition of Harish-Chandra functions, this series development is explicitly W_2 -invariant, the non-trivial symmetry here being exactly $z_1 \leftrightarrow z_2$ since the symmetry with respect to the subgroup $S_2 \subset W_2$ was already taken into account by our definition of z -variables in eq. (2.28). Of course, taking asymptotics at infinity breaks this symmetry explicitly.

To bring the last expansion to the form in eq. (A.1), one should apply another quadratic transformation of the form

$${}_2F_1\left(\frac{A}{2}, \frac{A+1}{2}; \frac{4y}{(y+1)^2}\right) = (1+y)^A {}_2F_1\left(\begin{matrix} A, B \\ A-B+1 \end{matrix}; y\right). \tag{A.4}$$

Here we run into a subtlety (first noticed by Gauss himself) which is sometimes encountered in working with quadratic transformations: the absolute value of y in the above formula should be less than one for it to be valid. Although one might imagine a wider applicability range by analytical continuation, in fact it does not happen. Actually, a second term should appear in the quadratic transformation to correct it in the latter case: compare formula

³⁰As the domain we work on is \tilde{A}^+ .

(3.1.3) in [79] and ex. 6 for chapter 3 *ibid.*³¹ So, we are forced to explicitly choose between two expansion domains related via $z_1 \leftrightarrow z_2$ and to break the symmetry by setting $y = z_1/z_2$ ($|z_1| < |z_2|$ for \tilde{A}^+) along with $A = \epsilon/2 - \lambda_1 + \lambda_2 + n - m$, $B = \epsilon/2$, finally arriving at eq. (A.1).

The expansion (A.1) is absolutely convergent when

$$|z_1| < 1, |z_2| < 1, |z_1| < |z_2|, \quad (\text{A.5})$$

as can be checked by using classical Horn's theorem [137]. For the convergence analysis, it is actually more convenient to pass to a different, resummed form, which makes the convergence region obvious

$$\begin{aligned} \tilde{\Phi}(\lambda_1, \lambda_2; k_i; z_1, z_2) &= \frac{z_1^{\frac{1+\epsilon}{2}+a-\lambda_1} z_2^{\frac{1}{2}+a-\lambda_2}}{4^{2a+1+\epsilon/2-\Lambda_{12}}} \sum_{n,m,p,r=0}^{\infty} \frac{z_1^n z_2^m (z_1 z_2)^p \left(\frac{z_1}{z_2}\right)^r}{n!m!p!r!} \frac{(\frac{\epsilon}{2} - \lambda_{12})_{r+n-m}}{(1 - \lambda_{12})_{r+n-m}} \\ &\times \frac{(1 - \lambda_{12}, 1 - \frac{\epsilon}{2} - \lambda_{12})_{n-m}}{(-\lambda_{12}, \frac{\epsilon}{2} - \lambda_{12})_{n-m}} \frac{(\frac{1}{2} + a - \lambda_1, \frac{1}{2} + b - \lambda_1)_{n+p}}{(1 - 2\lambda_1, 1 - \lambda_{12})_{n+p}} \frac{(\frac{1}{2} + a - \lambda_2, \frac{1}{2} + b - \lambda_2)_{m+p}}{(1 - 2\lambda_2, 1 + \lambda_{12})_{m+p}} \\ &\times \left(\frac{\epsilon}{2} - \lambda_{12}\right)_n \left(\frac{\epsilon}{2} + \lambda_{12}\right)_m \frac{(1 - \frac{\epsilon}{2}, 1 - \frac{\epsilon}{2} - \Lambda_{12})_p}{(1 - \Lambda_{12})_p} \left(\frac{\epsilon}{2}\right)_r. \end{aligned} \quad (\text{A.6})$$

To obtain this formula from eq. (A.1), one can use an identity

$$\sum_{n,m,p=0}^{\infty} F(n, m, p) \frac{y_1^{n+p} y_2^{m-p}}{n!m!p!} = \sum_{n,m,p=0}^{\infty} F(p, m-p, n-p) (-n, -m)_p \frac{y_1^n y_2^{m-n}}{n!m!p!} \quad (\text{A.7})$$

which follows from Bailey lemma [79]. It is also instrumental to keep in mind another particular presentation of the z -expansion which can be obtained by combining Whipple's identity and S_6 symmetry of a terminating well-poised ${}_7F_6$ function [82]

$$\begin{aligned} \tilde{\Phi}(\lambda_1, \lambda_2; k_i; z_1, z_2) &= \frac{1}{4^{2a+1+\epsilon/2-\Lambda_{12}}} \\ &\times \sum_{n,m=0}^{\infty} \frac{(1/2 + a - \lambda_1, 1/2 + b - \lambda_1)_n (1/2 + a - \lambda_2, 1/2 + b - \lambda_2)_m}{(1 - 2\lambda_1)_n (1 - 2\lambda_2)_m} \\ &\times \frac{(\epsilon/2 - \lambda_{12}, 1 - \epsilon/2 - \lambda_{12}, 1 + \epsilon/2)_{n-m} (1 - \epsilon/2 - \Lambda_{12})_m}{(-\lambda_{12}, 1 - \lambda_{12})_{n-m} (1 + \epsilon/2)_n (1 - \Lambda_{12})_m} \\ &\times {}_7F_6 \left(\begin{matrix} \frac{\epsilon}{2} + n - m, 1 + \frac{\epsilon}{4} + \frac{n-m}{2}, 1 - \Lambda_{12} + n, \frac{\epsilon}{2} + \lambda_{12}, \frac{\epsilon}{2} - \lambda_{12} - m + n, \frac{\epsilon}{2}, -m \\ \frac{\epsilon}{4} + \frac{n-m}{2}, \frac{\epsilon}{2} + \Lambda_{12} - m, 1 - \lambda_{12} - m + n, 1 + \lambda_{12}, 1 + n - m, 1 + \epsilon/2 + n \end{matrix}; 1 \right) \\ &\times \frac{(-1)^m (-n)_m}{n!m!} z_1^{\frac{1+\epsilon}{2}+a-\lambda_1+n} z_2^{\frac{1}{2}+a-\lambda_2+m} {}_2F_1 \left(\begin{matrix} \epsilon/2 - \lambda_{12} - m + n, \epsilon/2 \\ 1 - \lambda_{12} - m + n \end{matrix}; \frac{z_1}{z_2} \right). \end{aligned} \quad (\text{A.8})$$

To complete our discussion of the z -expansions, let us finally spell out the one summing over the cone of positive BC_2 roots, which makes the partial ordering of terms in the series

³¹When substituted into the Harish-Chandra expansion, the emergence of a second term can be seen as appropriate to give a continuation of the Harish-Chandra function past the wall ω_1 . In other words, it gives the action of monodromy matrix M_1 , see subsection 3.4.

manifest. To achieve this goal, we again use eq. (A.7) and obtain

$$\begin{aligned}
 \tilde{\Phi}(\lambda_1, \lambda_2; k_i; z_1, z_2) &= \frac{1}{4^{2a+1+\epsilon/2-\Lambda_{12}}} \sum_{n,m=0}^{\infty} \frac{1}{n!m!} z_1^{\frac{1+\epsilon}{2}+a-\lambda_1+n} z_2^{\frac{1}{2}+a-\lambda_2+m-n} \left(\frac{\epsilon}{2}\right)_n \\
 &\times \frac{(1/2+a-\lambda_2, 1/2+b-\lambda_2, 1-\epsilon/2+\lambda_{12})_m (\epsilon/2-\lambda_{12})_{n-m}}{(1-2\lambda_2, \lambda_{12})_m (1-\lambda_{12})_{n-m}} \\
 &\times \sum_{p=0}^{\infty} \frac{(-1)^p}{p!} \frac{\left(-\lambda_{12}-m, 1+\frac{-\lambda_{12}-m}{2}, \frac{1-\frac{\epsilon}{2}-\lambda_{12}-m}{2}, \frac{2-\frac{\epsilon}{2}-\lambda_{12}-m}{2}\right)_p}{\left(\frac{-\lambda_{12}-m}{2}, \frac{1+\frac{\epsilon}{2}-\lambda_{12}-m}{2}, \frac{\frac{\epsilon}{2}-\lambda_{12}-m}{2}\right)_p} \\
 &\times \frac{\left(\frac{\epsilon}{2}-\lambda_{12}+n-m, 1/2+a-\lambda_1, 1/2+b-\lambda_1, -n, -m\right)_p}{\left(1-\frac{\epsilon}{2}-n, 1/2-a+\lambda_2-m, 1/2-b+\lambda_2-m, 1-\lambda_{12}+n-m, 1-\lambda_{12}\right)_p} \\
 &\times \frac{\left(\frac{\epsilon}{2}-\lambda_{12}, 2\lambda_2-m\right)_p}{\left(1-2\lambda_1, 1-\frac{\epsilon}{2}-\lambda_{12}-m\right)_p} {}_4F_3\left(\begin{matrix} -p, p-m, 1-\frac{\epsilon}{2}, 1-\frac{\epsilon}{2}-\Lambda_{12} \\ 1-\frac{\epsilon}{2}+\lambda_{12}-p, 1-\frac{\epsilon}{2}-\lambda_{12}-m+p, 1-\Lambda_{12} \end{matrix}; 1\right).
 \end{aligned} \tag{A.9}$$

Notice that the Pochhammers of index p in the first two lines of the inner double sum are ‘almost well-poised’, i.e. the upper u_a and the lower l_a parameters are related by $l_a = 1 + A - u_a$ with $A = -\lambda_1 + \lambda_2 - m$. This implies that such a double sum should be a close two-variable analogue of a well-poised ${}_7F_6$ function. To see this somewhat more clearly, the last line can be additionally rewritten through Whipple identity as

$$\begin{aligned}
 &\frac{\left(\frac{\epsilon/2-\lambda_{12}-m}{2}, \frac{1+\epsilon/2-\lambda_{12}-m}{2}, 1-2\lambda_1-m, 2\lambda_2-m\right)_p}{\left(\frac{1-2\lambda_1-m}{2}, 1-\frac{2\lambda_1+m}{2}, \epsilon/2-\lambda_{12}-m, 1-\epsilon/2-\lambda_{12}-m\right)_p} \times \\
 &\times {}_7F_6\left(\begin{matrix} -2\lambda_1+p-m, 1+\frac{-2\lambda_1+p-m}{2}, \frac{\epsilon}{2}-\Lambda_{12}, 1-\frac{\epsilon}{2}-\Lambda_{12}, -\lambda_{12}+p-m, p-m, -p \\ \frac{p-m-2\lambda_1}{2}, 1-\frac{\epsilon}{2}-\lambda_{12}+p-m, \frac{\epsilon}{2}-\lambda_{12}+p-m, 1-\Lambda_{12}, 1-2\lambda_1, 1-2\lambda_1+2p-m \end{matrix}; 1\right).
 \end{aligned}$$

In order to obtain expansions in the ‘radial’ coordinates (x_i or u_i), we again start with eq. (A.1). For generic values of parameters, the u -expansion then can be derived in the same way as in the $N = 1$ example from the main text, i.e. through the use of a binomial theorem followed by a change of summation indices

$$\begin{aligned}
 \Phi(\lambda_1, \lambda_2; k_i; u_1, u_2) &= \sum_{i,j,n,m=0}^{\infty} \frac{1}{i!j!n!m!} x_1^{\lambda_1-\frac{1+\epsilon}{2}-a-n-i} x_2^{\lambda_2-\frac{1}{2}-a-m-j} \\
 &\times \frac{(1+\epsilon+2a-2\lambda_1)_{i+2n} (1+2a-2\lambda_2)_{j+2m} \left(\frac{1}{2}+a-\lambda_1, \frac{1}{2}+b-\lambda_1, \frac{\epsilon}{2}-\lambda_{12}\right)_n}{(-\lambda_{12})_{n-m} \left(\frac{\epsilon}{2}+\lambda_{12}\right)_{m-n} (1-2\lambda_1, 1-\lambda_{12}, \frac{1+\epsilon}{2}+a-\lambda_1, 1+\frac{\epsilon}{2}+a-\lambda_1)_n} \\
 &\times \frac{\left(\frac{1}{2}+a-\lambda_2, \frac{1}{2}+b-\lambda_2, \frac{\epsilon}{2}+\lambda_{12}\right)_m}{(1-2\lambda_2, 1+\lambda_{12}, \frac{1}{2}+a-\lambda_2, 1+a-\lambda_2)_m} \\
 &\times {}_4F_3\left(\begin{matrix} -n, -m, 1-\frac{\epsilon}{2}, 1-\frac{\epsilon}{2}-\Lambda_{12} \\ 1-\frac{\epsilon}{2}+\lambda_{12}-n, 1-\frac{\epsilon}{2}-\lambda_{12}-m, 1-\Lambda_{12} \end{matrix}; 1\right) \\
 &\times {}_6F_5\left(\begin{matrix} \frac{\epsilon}{2}-\lambda_{12}-m+n, \frac{\epsilon}{2}, -a+\lambda_2-m, \frac{1}{2}-a+\lambda_2-m, \frac{1+\epsilon}{2}+a-\lambda_1+n+\frac{i}{2}, 1+\frac{\epsilon}{2}+a-\lambda_1+n+\frac{i}{2} \\ 1-\lambda_{12}-m+n, \frac{1+\epsilon}{2}+a-\lambda_1+n, 1+\frac{\epsilon}{2}+a-\lambda_1+n, -a+\lambda_2-m-\frac{j}{2}, \frac{1}{2}-a+\lambda_2-m-\frac{j}{2} \end{matrix}; \frac{x_2}{x_1}\right).
 \end{aligned} \tag{A.10}$$

This form³² seems to be most useful for getting an expansion of the blocks in Jack (Gegenbauer) polynomials of radial variables, which explicitly solves the recursion relations given in [81]. We will not present the result here, as we do not use it. Instead, by changing summation indices and employing again the resummation identity (A.7), we obtain an explicit, manifestly partially ordered (i.e., over the $\mathbb{Z}_{\geq 0}$ -cone of positive roots) expansion of BC_2 Harish-Chandra functions in the coordinates u_1, u_2 as³³

$$\begin{aligned}
 \Phi(\lambda_1, \lambda_2; k_i; u_1, u_2) = & \sum_{i,j=0}^{\infty} e^{u_1(\lambda_1 - \frac{1+\epsilon}{2} - a - i) + u_2(\lambda_2 - \frac{1}{2} - a + i - j)} \frac{1}{i!j!} \\
 & \times \frac{(1+2a-2\lambda_2-2i)_j (\frac{\epsilon}{2}, \frac{\epsilon}{2} - \lambda_{12})_i}{(1-\lambda_{12})_i} \\
 & \times \sum_{n,m,p=0}^{\infty} \frac{(-1)^n}{n!m!p!} {}_4F_3 \left(\begin{matrix} -n, -m, 1 - \frac{\epsilon}{2}, 1 - \frac{\epsilon}{2} - \Lambda_{12} \\ 1 - \frac{\epsilon}{2} + \lambda_{12} - n, 1 - \frac{\epsilon}{2} - \lambda_{12} - m, 1 - \Lambda_{12} \end{matrix}; 1 \right) \\
 & \times \frac{(\lambda_{12} - i)_{m+p-n}}{(1 - \frac{\epsilon}{2} + \lambda_{12} - i)_{m+p-n}} \frac{(-j, 1+2a-2\lambda_2-2i+j)_{m+p}}{(\frac{1}{2} + a - \lambda_2 - i, 1+a-\lambda_2-i)_{m+p}} \\
 & \times \frac{(-\frac{\epsilon}{2} - a + \lambda_1 - i, \frac{1-\epsilon}{2} - a + \lambda_1 - i)_{p-n}}{(-\epsilon - 2a + 2\lambda_1 - 2i)_{p-n}} \frac{(1-\lambda_{12}, 1 - \frac{\epsilon}{2} - \lambda_{12})_{n-m}}{(-\lambda_{12}, \frac{\epsilon}{2} - \lambda_{12})_{n-m}} \\
 & \times \frac{(-p, \frac{1}{2} + a - \lambda_1, \frac{1}{2} + b - \lambda_1, \frac{\epsilon}{2} - \lambda_{12})_n}{(1-2\lambda_1, 1-\lambda_{12})_n} \frac{(\frac{1}{2} + a - \lambda_2, \frac{1}{2} + b - \lambda_2, \frac{\epsilon}{2} + \lambda_{12})_m}{(1-2\lambda_2, 1+\lambda_{12})_m} \frac{(-i)_p}{(1 - \frac{\epsilon}{2} - i)_p}.
 \end{aligned} \tag{A.11}$$

This is the main result of this section. As the coefficient of the expansion term solves the recursion relation (3.18), according to general theorems of Heckman and Opdam [76] for generic values of λ the expansion is convergent on the whole domain A^+ and can be analytically continued to the entire (complex) fundamental domain. The expansion (A.11) combined with Heckman-Opdam results gives a full control over the analytical properties of the BC_2 Harish-Chandra functions.

The two expansions in x coordinates above can be recast in many equivalent forms. All of them can in principle be regarded as instances of sextuple Srivastava-Dooust hypergeometric functions [138], a rather general class of analogues of hypergeometric functions. Since information on the analytical behaviour of a generic member of this family seems to be quite scarce, it is fortunate that we have the well-developed Heckman-Opdam theory at our disposal (and a root symmetry at the root of it). We are not aware of any significant simplification of the above formula for generic values of the parameters, it would be interesting to prove this rigorously.

To conclude this appendix, let us comment on the case of general N . We claim that in fact all the building blocks required for writing down the expansions are already present here, although writing up a general BC_N formula requires extra work along the lines described in the beginning of the appendix. It appears that such general expansions of BC_N

³²See a footnote before the formula (3.4), regarding this and the next expansion.

³³The coefficient of the powers in this expansion, i.e. the inner triple sum of Pochhammer symbols times the balanced ${}_4F_3(1)$ function, is, of course, a finite sum. It can be easily written as a product of Pochhammer quotients and a quadruple hypergeometric polynomial.

Harish-Chandra functions will involve $N(N+1)$ ($x-$) and N^2 ($z-$) summations, which are numbers of roots of the BC_N root system and of its corresponding reduced root subsystem (B_N or C_N), especially when looking at such beautifully symmetric expansions as eq. (A.6). Clearly, for $N = 2$ we get back 6 x -summations and 4 z -summations of the above.

We conjecture that these numbers of summations are minimal possible for the generic values of parameters and, moreover, that the same is true also for Harish-Chandra functions of other root systems, of which the ones associated to exceptional root systems are probably mathematically most interesting, since they cannot be obtained as special cases or limits of the BC case. E.g. for E_8 Harish-Chandra function this would imply a minimum of 120 $x-/z-$ summations. As such explicit general expansions are not yet available, it is an interesting mathematical problem to solve. As explained in [72], the BC_N Harish-Chandra functions describe conformal blocks of a two-point function of defects, N being determined by their dimensions. So, such expansions are of potential interest for the conformal bootstrap programme, provided there is a corresponding analogue of crossing symmetry for such defects.

B Alternative expansions for integer spin

The aim of this appendix is to specialize our general expansion (A.1) to integer spin $l = \lambda_1 - \lambda_2 - \epsilon/2$ and to rewrite it as an expansion in the usual cross ratios z and \bar{z} , or rather variables $u = z\bar{z}$ and $v = (1-z)(1-\bar{z})$. The latter can then be compared with existing expressions from other approaches, such as the shadow or the embedding formalism.

Before we study Harish-Chandra series for such special values of the parameters, we want to make some general comments. Actually, one has to be a bit careful when evaluating formulas that are valid for generic values of the parameters, like the ones for Harish-Chandra functions in the [previous appendix](#), at special values of the parameters such as the momenta λ_i . In order to do so, one should first take a limit for the multiplicities, and only then for the momenta (i.e. the continued spin or conformal dimension). In other words, if we want to study blocks/Harish-Chandra functions for particular spins (or twists) through the general series expansions, our prescription is to fix the dimension d and conformal weights Δ_i of external fields first and only then, after carefully taking into account all implications on the summation range, to send the spin (or twist) to some particular value. This recipe is directly related to the fact that an (appropriately normalized) Harish-Chandra function is an entire function of its multiplicities, but can have simple poles in λ (and further subtleties related to its series expansion for λ along specific affine hyperplanes, see subsections 3.2, 3.3, 5.1). The opposite order of limits, provided it exists at all, also gives some solution of the Calogero-Sutherland problem, but one that may be a non-trivial linear combination of all the $|W_N|$ Harish-Chandra functions. The prescription we propose here is the same as (one of the) usually used when taking limits in formulas for the hypergeometric function ${}_2F_1$ (i.e. for $N = 1$) to special values of its parameters [139]: when e.g. one upper and one lower parameter are negative integers, one takes

$$z^{-\frac{k+m}{2}} {}_2F_1\left(\begin{matrix} -m, b - \frac{k+m}{2} \\ -k - m \end{matrix}; z\right) = \lim_{h \rightarrow -(k+m)/2} \lim_{a \rightarrow -h-m} z^h {}_2F_1\left(\begin{matrix} h + a, h + b \\ 2h \end{matrix}; z\right)$$

rather than

$$\lim_{a \rightarrow (k-m)/2} {}_2F_1 \left(\begin{matrix} a - \frac{k+m}{2}, b - \frac{k+m}{2} \\ -k - m \end{matrix}; z \right),$$

where k, m are positive integers and b is generic. One can show that the second function is a linear combination of the first and its shadow.

After these introductory remarks we now want to start with our general formula (A.1) and set $l = 0$, i.e. $\lambda_2 = \lambda_1 - \frac{\epsilon}{2}$. For the moment we shall assume that the dimension d is not even so that upon specializing λ , the twisted Harish-Chandra function is actually just equal to a block, up to a normalizing prefactor $4^{2a+1+\epsilon-2\lambda_1} (z_1 z_2)^{-a} c_{\alpha_1}(\lambda_1, \lambda_2)$. By applying a Whipple transformation [79] to the inner ${}_4F_3$ function in eq. (A.1), we can perform a sum via Saalschütz formula, yielding³⁴

$$\begin{aligned} \tilde{\Phi}(\lambda_1, \lambda_2; z_1, z_2) \Big|_{l=0} &= \frac{(z_1 z_2)^{\frac{1+\epsilon}{2}+a-\lambda_1}}{4^{2a+1+\epsilon-2\lambda_1}} \sum_{n,m,p=0}^{\infty} \frac{(-z_1)^{n+p} z_2^{m-p}}{n!m!p!} \left(\frac{\epsilon}{2}\right)_{m-n-p} \left(\frac{\epsilon}{2}\right)_p (-m)_{n+p} \\ &\times \frac{\left(\frac{\epsilon}{2}+1\right)_{m-n} \left(\frac{1}{2}+a-\lambda_1, \frac{1}{2}+b-\lambda_1\right)_n \left(\frac{1+\epsilon}{2}+a-\lambda_1, \frac{1+\epsilon}{2}+b-\lambda_1\right)_m}{\left(\frac{\epsilon}{2}\right)_{m-n} (1+\frac{\epsilon}{2}-2\lambda_1)_n (1+\frac{\epsilon}{2}, 1+\epsilon-2\lambda_1)_m}. \end{aligned} \quad (\text{B.1})$$

By changing the order of summation, we can rewrite this expression as

$$\begin{aligned} &= \left(\frac{z_1 z_2}{16}\right)^{\frac{1+\epsilon}{2}+a-\lambda_1} \sum_{n,m=0}^{\infty} \frac{(-z_1)^n z_2^{m-n}}{n!m!} \left(\frac{\epsilon}{2}\right)_{m-n} \left(\frac{\epsilon}{2}\right)_n (-m)_n \frac{\left(\frac{1+\epsilon}{2}+a-\lambda_1, \frac{1+\epsilon}{2}+b-\lambda_1\right)_m}{\left(\frac{\epsilon}{2}, 1+\epsilon-2\lambda_1\right)_m} \times \\ &\times {}_7F_6 \left(\begin{matrix} -\frac{\epsilon}{2}-m, 1+\frac{-\frac{\epsilon}{2}-m}{2}, -n, -m+n, -\epsilon+2\lambda_1-m, \frac{1}{2}+a-\lambda_1, \frac{1}{2}+b-\lambda_1 \\ -\frac{\epsilon}{2}-m, 1-\frac{\epsilon}{2}-m+n, 1-\frac{\epsilon}{2}-n, 1+\frac{\epsilon}{2}-2\lambda_1, \frac{1-\epsilon}{2}-a+\lambda_1-m, \frac{1-\epsilon}{2}-b+\lambda_1-m \end{matrix}; 1 \right). \end{aligned}$$

The inner well-poised ${}_7F_6(1)$ is now to be transformed to another one using the S_6 symmetry of this function. Namely, we can apply a particular identity labeled $W(1; 2) = W(3; 4)$ by Bailey [82], employ the series representation for the resulting inner ${}_7F_6(1)$ function and shift the summation indices. This leads to

$$\begin{aligned} &= \left(\frac{z_1 z_2}{16}\right)^{\frac{1+\epsilon}{2}+a-\lambda_1} \sum_{p=0}^{\infty} \frac{(z_1 z_2)^p}{p!} \frac{\left(\frac{\epsilon}{2}, \epsilon-2\lambda_1, \frac{1+\epsilon}{2} \pm a - \lambda_1, \frac{1+\epsilon}{2} \pm b - \lambda_1\right)_p}{(1+\epsilon-2\lambda_1)_{2p}^2 (1+\frac{\epsilon}{2}-2\lambda_1)_p} \\ &\times {}_2F_1 \left(\begin{matrix} \frac{1+\epsilon}{2}+a-\lambda_1+p, \frac{1+\epsilon}{2}+b-\lambda_1+p \\ 1+\epsilon-2\lambda_1+2p \end{matrix}; z_1 \right) {}_2F_1 \left(\begin{matrix} \frac{1+\epsilon}{2}+a-\lambda_1+p, \frac{1+\epsilon}{2}+b-\lambda_1+p \\ 1+\epsilon-2\lambda_1+2p \end{matrix}; z_2 \right), \end{aligned} \quad (\text{B.2})$$

where n and m summations were absorbed into the two ${}_2F_1$ hypergeometric functions. Using Chu-Vandermonde identity [79] in the form

$$\frac{\left(\frac{\epsilon}{2}\right)_p}{\left(1+\frac{\epsilon}{2}-2\lambda_1\right)_p} = (-1)^p \sum_{r=0}^p \frac{(-p, 1-2\lambda_1)_r}{\left(1+\frac{\epsilon}{2}-2\lambda_1\right)_r} \frac{1}{r!}$$

³⁴The analysis by Horn's theorem [137] shows that the absolute convergence for this series representation of the twisted Harish-Chandra function in fact takes place on a the *extended* domain $\Re u_i > 0$ and $0 < \Im u_i < 2\pi$. This is, of course, consistent with the fact that the single twisted Harish-Chandra function for generic dimension, being proportional to a block, is bound to be regular at the wall $z_1 = z_2$ according to our discussion in subsection 5.1.

and shifting p, r summation indices, we can rewrite our expression further as

$$\begin{aligned}
 &= \left(\frac{z_1 z_2}{16}\right)^{\frac{1+\epsilon}{2}+a-\lambda_1} \sum_{r=0}^{\infty} \frac{(z_1 z_2)^r}{r!} \frac{\left(\frac{1+\epsilon}{2} \pm a - \lambda_1, \frac{1+\epsilon}{2} \pm b - \lambda_1\right)_r}{(1 + \epsilon - 2\lambda_1)_{2r} (1 + \frac{\epsilon}{2} - 2\lambda_1)_r} \\
 &\times \sum_{p=0}^{\infty} \frac{(-z_1 z_2)^p}{p!} \frac{(\epsilon - 2\lambda_1 + 2r, \frac{1+\epsilon}{2} \pm a - \lambda_1 + r, \frac{1+\epsilon}{2} \pm b - \lambda_1 + r)_p}{(\epsilon - 2\lambda_1 + 2r, 1 + \epsilon - 2\lambda_1 + 2r)_{2p}} \\
 &\times {}_2F_1\left(\begin{matrix} \frac{1+\epsilon}{2} + a - \lambda_1 + r + p, \frac{1+\epsilon}{2} + b - \lambda_1 + r + p \\ 1 + \epsilon - 2\lambda_1 + 2r + 2p \end{matrix}; z_1\right) \\
 &\times {}_2F_1\left(\begin{matrix} \frac{1+\epsilon}{2} + a - \lambda_1 + r + p, \frac{1+\epsilon}{2} + b - \lambda_1 + r + p \\ 1 + \epsilon - 2\lambda_1 + 2r + 2p \end{matrix}; z_2\right),
 \end{aligned} \tag{B.3}$$

so that for the inner $(p-)$ sum we can now use a beautiful identity of Burchnell-Chaundy ([140], formula (50))

$$\begin{aligned}
 {}_2F_1\left(\begin{matrix} A, B \\ C \end{matrix}; z_1 + z_2 - z_1 z_2\right) &= \sum_{p=0}^{\infty} \frac{(-z_1 z_2)^p}{p!} \frac{(A, B, C - A, C - B)_p}{(C)_{2p} (C + p - 1)_p} \\
 &\times {}_2F_1\left(\begin{matrix} A + p, B + p \\ C + 2p \end{matrix}; z_1\right) {}_2F_1\left(\begin{matrix} A + p, B + p \\ C + 2p \end{matrix}; z_2\right)
 \end{aligned} \tag{B.4}$$

(provided both sides converge or assuming analytical continuation), and finally arrive at³⁵

$$\begin{aligned}
 \tilde{\Phi}(\lambda_1, \lambda_2; z_1, z_2) \Big|_{l=0} &= \left(\frac{u}{16}\right)^{\frac{1+\epsilon}{2}+a-\lambda_1} \sum_{r,s=0}^{\infty} \frac{u^r (1-v)^s}{r!s!} \frac{\left(\frac{1+\epsilon}{2} + a - \lambda_1, \frac{1+\epsilon}{2} + b - \lambda_1\right)_{r+s}}{(1 + \epsilon - 2\lambda_1)_{2r+s}} \\
 &\times \frac{\left(\frac{1+\epsilon}{2} - a - \lambda_1, \frac{1+\epsilon}{2} - b - \lambda_1\right)_r}{\left(1 + \frac{\epsilon}{2} - 2\lambda_1\right)_r},
 \end{aligned} \tag{B.5}$$

where u, v are the usual cross ratios defined in eq. (4.2). As in section 4, we identify $z_1 = z, z_2 = \bar{z}$. Once we pass to conformal blocks using our relation (4.13), we recognize formula (3.32) in [26] (see also [24]). Dolan and Osborn obtained this formula from the shadow formalism by calculating the corresponding integrals via Mellin transform, ignoring all pole terms from the shadow contributions. As noticed e.g. in [24, 141, 142], this double hypergeometric expression (B.5) can be rewritten as a linear combination of two F_4 Appell functions.³⁶ It is clear that we can lift our restriction on ϵ and admit even values, literally repeating the above calculation for the scalar exchange block which for even $\epsilon \in 2\mathbb{Z}_{\geq 0}$ is composed from two twisted Harish-Chandra functions.

The type of analysis we have presented here and in particular the use of Burchnell-Chaundy type formulas, generalizes much beyond the specific case analysed here and allows to relate expressions from our approach to those obtained through the shadow formalism [26] or results from the embedding formalism [29, 143, 144]. In particular, we can go

³⁵Note that, while previous manipulations just reshuffled summation order in an absolutely convergent series, the application of Burchnell-Chaundy formula changes the domain of (absolute) convergence which now becomes $\{u, v \in \mathbb{C} \mid |1 - v| < 1, \sqrt{|u|} - \sqrt{|1 - v|} < 1\}$.

³⁶Notice that the convergence region after using this connection formula changes to $\{\sqrt{|u|} + \sqrt{|v|} < 1\}$, so a diagonal wall is in fact re-introduced.

through the same steps for any spin $l > 0$ (and ‘push out’ the finite summations associated to the spin being integer) to obtain finite sums of double hypergeometric functions of Srivastava-Daoust (i.e. generalized Kampé de Fériet) type [138]. While the resulting expressions can be used to compare with results from other approaches, we want to stress that the expansion formulas we have derived in appendix A and their direct specializations are typically simpler than those that are obtained by other techniques.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] G. Mack and A. Salam, *Finite component field representations of the conformal group*, *Annals Phys.* **53** (1969) 174 [[INSPIRE](#)].
- [2] A.M. Polyakov, *Conformal symmetry of critical fluctuations*, *JETP Lett.* **12** (1970) 381 [*Pisma Zh. Eksp. Teor. Fiz.* **12** (1970) 538] [[INSPIRE](#)].
- [3] S. Ferrara, A.F. Grillo, G. Parisi and R. Gatto, *Covariant expansion of the conformal four-point function*, *Nucl. Phys.* **B 49** (1972) 77 [Erratum *ibid.* **B 53** (1973) 643] [[INSPIRE](#)].
- [4] S. Ferrara, A.F. Grillo and R. Gatto, *Tensor representations of conformal algebra and conformally covariant operator product expansion*, *Annals Phys.* **76** (1973) 161 [[INSPIRE](#)].
- [5] A.M. Polyakov, *Nonhamiltonian approach to conformal quantum field theory*, *Zh. Eksp. Teor. Fiz.* **66** (1974) 23 [*Sov. Phys. JETP* **39** (1974) 9] [[INSPIRE](#)].
- [6] G. Mack, *Duality in quantum field theory*, *Nucl. Phys.* **B 118** (1977) 445 [[INSPIRE](#)].
- [7] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova and I.T. Todorov, *Harmonic analysis on the n -dimensional Lorentz group and its application to conformal quantum field theory*, *Lect. Notes Phys.* **63** (1977) 1 [[INSPIRE](#)].
- [8] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, *Nucl. Phys.* **B 241** (1984) 333 [[INSPIRE](#)].
- [9] R. Rattazzi, V.S. Rychkov, E. Tonni and A. Vichi, *Bounding scalar operator dimensions in 4D CFT*, *JHEP* **12** (2008) 031 [[arXiv:0807.0004](#)] [[INSPIRE](#)].
- [10] S. El-Showk, M.F. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin and A. Vichi, *Solving the 3d Ising model with the conformal bootstrap*, *Phys. Rev.* **D 86** (2012) 025022 [[arXiv:1203.6064](#)] [[INSPIRE](#)].
- [11] S. El-Showk, M.F. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin and A. Vichi, *Solving the 3d Ising model with the conformal bootstrap II. c -minimization and precise critical exponents*, *J. Stat. Phys.* **157** (2014) 869 [[arXiv:1403.4545](#)] [[INSPIRE](#)].
- [12] F. Kos, D. Poland and D. Simmons-Duffin, *Bootstrapping mixed correlators in the 3d Ising model*, *JHEP* **11** (2014) 109 [[arXiv:1406.4858](#)] [[INSPIRE](#)].
- [13] D. Simmons-Duffin, *A semidefinite program solver for the conformal bootstrap*, *JHEP* **06** (2015) 174 [[arXiv:1502.02033](#)] [[INSPIRE](#)].
- [14] F. Kos, D. Poland, D. Simmons-Duffin and A. Vichi, *Precision islands in the Ising and $O(N)$ models*, *JHEP* **08** (2016) 036 [[arXiv:1603.04436](#)] [[INSPIRE](#)].

- [15] A. Dymarsky, F. Kos, P. Kravchuk, D. Poland and D. Simmons-Duffin, *The 3d stress-tensor bootstrap*, *JHEP* **02** (2018) 164 [[arXiv:1708.05718](#)] [[INSPIRE](#)].
- [16] F. Kos, D. Poland and D. Simmons-Duffin, *Bootstrapping the $O(N)$ vector models*, *JHEP* **06** (2014) 091 [[arXiv:1307.6856](#)] [[INSPIRE](#)].
- [17] F. Kos, D. Poland, D. Simmons-Duffin and A. Vichi, *Bootstrapping the $O(N)$ archipelago*, *JHEP* **11** (2015) 106 [[arXiv:1504.07997](#)] [[INSPIRE](#)].
- [18] L. Iliesiu, F. Kos, D. Poland, S.S. Pufu, D. Simmons-Duffin and R. Yacoby, *Bootstrapping 3d fermions*, *JHEP* **03** (2016) 120 [[arXiv:1508.00012](#)] [[INSPIRE](#)].
- [19] L. Iliesiu, F. Kos, D. Poland, S.S. Pufu and D. Simmons-Duffin, *Bootstrapping 3d fermions with global symmetries*, *JHEP* **01** (2018) 036 [[arXiv:1705.03484](#)] [[INSPIRE](#)].
- [20] R. Rattazzi, S. Rychkov and A. Vichi, *Bounds in 4d conformal field theories with global symmetry*, *J. Phys. A* **44** (2011) 035402 [[arXiv:1009.5985](#)] [[INSPIRE](#)].
- [21] D. Poland, D. Simmons-Duffin and A. Vichi, *Carving out the space of 4d CFTs*, *JHEP* **05** (2012) 110 [[arXiv:1109.5176](#)] [[INSPIRE](#)].
- [22] S.M. Chester and S.S. Pufu, *Towards bootstrapping QED_3* , *JHEP* **08** (2016) 019 [[arXiv:1601.03476](#)] [[INSPIRE](#)].
- [23] S.M. Chester and S.S. Pufu, *Anomalous dimensions of scalar operators in QED_3* , *JHEP* **08** (2016) 069 [[arXiv:1603.05582](#)] [[INSPIRE](#)].
- [24] F.A. Dolan and H. Osborn, *Conformal four point functions and the operator product expansion*, *Nucl. Phys. B* **599** (2001) 459 [[hep-th/0011040](#)] [[INSPIRE](#)].
- [25] F.A. Dolan and H. Osborn, *Conformal partial waves and the operator product expansion*, *Nucl. Phys. B* **678** (2004) 491 [[hep-th/0309180](#)] [[INSPIRE](#)].
- [26] F.A. Dolan and H. Osborn, *Conformal partial waves: further mathematical results*, [arXiv:1108.6194](#) [[INSPIRE](#)].
- [27] M.S. Costa, J. Penedones, D. Poland and S. Rychkov, *Spinning conformal correlators*, *JHEP* **11** (2011) 071 [[arXiv:1107.3554](#)] [[INSPIRE](#)].
- [28] M.S. Costa, J. Penedones, D. Poland and S. Rychkov, *Spinning conformal blocks*, *JHEP* **11** (2011) 154 [[arXiv:1109.6321](#)] [[INSPIRE](#)].
- [29] D. Simmons-Duffin, *Projectors, shadows and conformal blocks*, *JHEP* **04** (2014) 146 [[arXiv:1204.3894](#)] [[INSPIRE](#)].
- [30] F. Rejon-Barrera and D. Robbins, *Scalar-vector bootstrap*, *JHEP* **01** (2016) 139 [[arXiv:1508.02676](#)] [[INSPIRE](#)].
- [31] M.S. Costa and T. Hansen, *Conformal correlators of mixed-symmetry tensors*, *JHEP* **02** (2015) 151 [[arXiv:1411.7351](#)] [[INSPIRE](#)].
- [32] J. Penedones, E. Trevisani and M. Yamazaki, *Recursion relations for conformal blocks*, *JHEP* **09** (2016) 070 [[arXiv:1509.00428](#)] [[INSPIRE](#)].
- [33] A. Castedo Echeverri, E. Elkhidir, D. Karateev and M. Serone, *Deconstructing conformal blocks in 4d CFT*, *JHEP* **08** (2015) 101 [[arXiv:1505.03750](#)] [[INSPIRE](#)].
- [34] L. Iliesiu, F. Kos, D. Poland, S.S. Pufu, D. Simmons-Duffin and R. Yacoby, *Fermion-scalar conformal blocks*, *JHEP* **04** (2016) 074 [[arXiv:1511.01497](#)] [[INSPIRE](#)].

- [35] M.S. Costa, T. Hansen, J. Penedones and E. Trevisani, *Radial expansion for spinning conformal blocks*, *JHEP* **07** (2016) 057 [[arXiv:1603.05552](#)] [[INSPIRE](#)].
- [36] M.S. Costa, T. Hansen, J. Penedones and E. Trevisani, *Projectors and seed conformal blocks for traceless mixed-symmetry tensors*, *JHEP* **07** (2016) 018 [[arXiv:1603.05551](#)] [[INSPIRE](#)].
- [37] A. Castedo Echeverri, E. Elkhidir, D. Karateev and M. Serone, *Seed conformal blocks in 4D CFT*, *JHEP* **02** (2016) 183 [[arXiv:1601.05325](#)] [[INSPIRE](#)].
- [38] D. Karateev, P. Kravchuk and D. Simmons-Duffin, *Weight shifting operators and conformal blocks*, *JHEP* **02** (2018) 081 [[arXiv:1706.07813](#)] [[INSPIRE](#)].
- [39] P. Kravchuk and D. Simmons-Duffin, *Counting conformal correlators*, *JHEP* **02** (2018) 096 [[arXiv:1612.08987](#)] [[INSPIRE](#)].
- [40] P. Kravchuk, *Casimir recursion relations for general conformal blocks*, *JHEP* **02** (2018) 011 [[arXiv:1709.05347](#)] [[INSPIRE](#)].
- [41] F. Calogero, *Solution of the one-dimensional N body problems with quadratic and/or inversely quadratic pair potentials*, *J. Math. Phys.* **12** (1971) 419 [[INSPIRE](#)].
- [42] B. Sutherland, *Exact results for a quantum many body problem in one-dimension. 2*, *Phys. Rev. A* **5** (1972) 1372 [[INSPIRE](#)].
- [43] J. Moser, *Three integrable Hamiltonian systems connected with isospectral deformations*, *Adv. Math.* **16** (1975) 197 [[INSPIRE](#)].
- [44] G.J. Heckman, *Root systems and hypergeometric functions. I*, *Compos. Math.* **64** (1987) 329.
- [45] G.J. Heckman, *Root systems and hypergeometric functions. II*, *Compos. Math.* **64** (1987) 353.
- [46] E.M. Opdam, *Root systems and hypergeometric functions. III*, *Compos. Math.* **67** (1988) 21.
- [47] E.M. Opdam, *Root systems and hypergeometric functions. IV*, *Compos. Math.* **67** (1988) 191.
- [48] T. Koornwinder, *Orthogonal polynomials in two variables which are eigenfunctions of two algebraically independent partial differential operators, I*, *Indag. Math.* **77** (1974) 48.
- [49] T. Koornwinder, *Orthogonal polynomials in two variables which are eigenfunctions of two algebraically independent partial differential operators, II*, *Indag. Math.* **77** (1974) 59.
- [50] T. Koornwinder, *Orthogonal polynomials in two variables which are eigenfunctions of two algebraically independent partial differential operators, III*, *Indag. Math.* **77** (1974) 357.
- [51] T. Koornwinder, *Orthogonal polynomials in two variables which are eigenfunctions of two algebraically independent partial differential operators, IV*, *Indag. Math.* **77** (1974) 370.
- [52] I. Cherednik, *Double affine Hecke algebras*, Cambridge University Press, Cambridge, U.K., (2005).
- [53] I. Macdonald, *Affine Hecke algebras and orthogonal polynomials*, Cambridge tracts in mathematics, Cambridge University Press, Cambridge, U.K., (2003).
- [54] M. Isachenkov and V. Schomerus, *Superintegrability of d -dimensional conformal blocks*, *Phys. Rev. Lett.* **117** (2016) 071602 [[arXiv:1602.01858](#)] [[INSPIRE](#)].
- [55] M. Isachenkov and V. Schomerus, *Integrability of conformal blocks II: algebraic structures*, in preparation.

- [56] S. Caron-Huot, *Analyticity in spin in conformal theories*, *JHEP* **09** (2017) 078 [[arXiv:1703.00278](#)] [[INSPIRE](#)].
- [57] A.L. Fitzpatrick, J. Kaplan, D. Poland and D. Simmons-Duffin, *The analytic bootstrap and AdS superhorizon locality*, *JHEP* **12** (2013) 004 [[arXiv:1212.3616](#)] [[INSPIRE](#)].
- [58] Z. Komargodski and A. Zhiboedov, *Convexity and liberation at large spin*, *JHEP* **11** (2013) 140 [[arXiv:1212.4103](#)] [[INSPIRE](#)].
- [59] L.F. Alday, A. Bissi and T. Lukowski, *Large spin systematics in CFT*, *JHEP* **11** (2015) 101 [[arXiv:1502.07707](#)] [[INSPIRE](#)].
- [60] L.F. Alday and A. Zhiboedov, *An algebraic approach to the analytic bootstrap*, *JHEP* **04** (2017) 157 [[arXiv:1510.08091](#)] [[INSPIRE](#)].
- [61] L.F. Alday, *Solving CFTs with weakly broken higher spin symmetry*, *JHEP* **10** (2017) 161 [[arXiv:1612.00696](#)] [[INSPIRE](#)].
- [62] L.F. Alday, *Large spin perturbation theory for conformal field theories*, *Phys. Rev. Lett.* **119** (2017) 111601 [[arXiv:1611.01500](#)] [[INSPIRE](#)].
- [63] D. Simmons-Duffin, *The lightcone bootstrap and the spectrum of the 3d Ising CFT*, *JHEP* **03** (2017) 086 [[arXiv:1612.08471](#)] [[INSPIRE](#)].
- [64] M. Hogervorst, *Crossing kernels for boundary and crosscap CFTs*, [arXiv:1703.08159](#) [[INSPIRE](#)].
- [65] M. Hogervorst and B.C. van Rees, *Crossing symmetry in alpha space*, *JHEP* **11** (2017) 193 [[arXiv:1702.08471](#)] [[INSPIRE](#)].
- [66] L.F. Alday and S. Caron-Huot, *Gravitational S-matrix from CFT dispersion relations*, [arXiv:1711.02031](#) [[INSPIRE](#)].
- [67] D. Simmons-Duffin, D. Stanford and E. Witten, *A spacetime derivation of the Lorentzian OPE inversion formula*, *JHEP* **07** (2018) 085 [[arXiv:1711.03816](#)] [[INSPIRE](#)].
- [68] D. Li, D. Meltzer and D. Poland, *Conformal bootstrap in the Regge limit*, *JHEP* **12** (2017) 013 [[arXiv:1705.03453](#)] [[INSPIRE](#)].
- [69] M.S. Costa, T. Hansen and J. Penedones, *Bounds for OPE coefficients on the Regge trajectory*, *JHEP* **10** (2017) 197 [[arXiv:1707.07689](#)] [[INSPIRE](#)].
- [70] J. Qiao and S. Rychkov, *A tauberian theorem for the conformal bootstrap*, *JHEP* **12** (2017) 119 [[arXiv:1709.00008](#)] [[INSPIRE](#)].
- [71] M. Lemos, P. Liendo, M. Meineri and S. Sarkar, *Universality at large transverse spin in defect CFT*, [arXiv:1712.08185](#) [[INSPIRE](#)].
- [72] V. Schomerus, *Harmony of defect blocks*, talk given at the workshop *Boundary and defect conformal field theory: open problems and applications*, Chicheley Hall, Buckinghamshire, U.K., September 2017.
- [73] A. Gadde, *Conformal constraints on defects*, [arXiv:1602.06354](#) [[INSPIRE](#)].
- [74] M. Fukuda, N. Kobayashi and T. Nishioka, *Operator product expansion for conformal defects*, *JHEP* **01** (2018) 013 [[arXiv:1710.11165](#)] [[INSPIRE](#)].
- [75] G. Pöschl and E. Teller, *Bemerkungen zur Quantenmechanik des anharmonischen Oszillators* (in German), *Z. Phys.* **83** (1933) 143 [[INSPIRE](#)].

- [76] G. Heckman and H. Schlichtkrull, *Harmonic analysis and special functions on symmetric spaces*, Academic Press, U.S.A., (1994).
- [77] V.I. Danilov, *The geometry of toric varieties*, *Usp. Mat. Nauk* **33** (1978) 85 [*Russ. Math. Surv.* **33** (1978) 97].
- [78] T. Oda, *Convex bodies and algebraic geometry*, Springer-Verlag, Germany, (1988).
- [79] G.E. Andrews, R. Askey and R. Roy, *Special functions*, Cambridge University Press, Cambridge, U.K., (1999).
- [80] T. Koornwinder, *Jacobi functions and analysis on noncompact semisimple Lie groups*, in *Special functions: group theoretical aspects and applications*, Springer, Netherlands, (1984), pg. 1.
- [81] M. Hogervorst and S. Rychkov, *Radial coordinates for conformal blocks*, *Phys. Rev. D* **87** (2013) 106004 [[arXiv:1303.1111](#)] [[INSPIRE](#)].
- [82] W. Bailey, *Generalized hypergeometric series*, Cambridge tracts in mathematics and mathematical physics, Cambridge University Press, Cambridge, U.K., (1935).
- [83] N. Shimeno, *Formula for the hypergeometric functions of type BC_n* , *Pacific J. Math.* **236** (2008) 105.
- [84] E.M. Opdam, *Part I: lectures on Dunkl operators*, *MSJ Mem.* **8** (2000) 2, Mathematical Society of Japan, Tokyo, Japan, (2000).
- [85] S.G. Gindikin and F.I. Karpelevich, *Plancherel measure for symmetric Riemannian spaces of non-positive curvature*, *Sov. Math. Dokl.* **3** (1962) 962.
- [86] A. Pasquale, *Asymptotic analysis of Θ -hypergeometric functions*, *Invent. Math.* **157** (2004) 71.
- [87] E.M. Opdam, *Harmonic analysis for certain representations of graded Hecke algebras*, *Acta Math.* **175** (1995) 75.
- [88] N.V. Dũng, *The fundamental groups of the spaces of regular orbits of the affine Weyl groups*, *Topology* **22** (1983) 425.
- [89] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe*, *Invent. Math.* **12** (1971) 57.
- [90] H. van der Lek, *The homotopy type of complex hyperplane complements*, Ph.D. thesis, Radboud Universiteit, Nijmegen, Netherlands, (1983).
- [91] H. van der Lek, *Extended Artin groups*, *Proc. Symp. Pure Math.* **40** (1981) 117.
- [92] P. Deligne, *Les immeubles des groupes de tresses généralisés* (in French), *Invent. Math.* **17** (1972) 273.
- [93] E.M. Opdam, *An analogue of the Gauss summation formula for hypergeometric functions related to root systems*, *Math. Z.* **212** (1993) 313.
- [94] M.S. Costa, V. Goncalves and J. Penedones, *Conformal Regge theory*, *JHEP* **12** (2012) 091 [[arXiv:1209.4355](#)] [[INSPIRE](#)].
- [95] P. Liendo, L. Rastelli and B.C. van Rees, *The bootstrap program for boundary CFT_d* , *JHEP* **07** (2013) 113 [[arXiv:1210.4258](#)] [[INSPIRE](#)].

- [96] F. Hartogs, *Einige Folgerungen aus der Cauchyschen Integralformel bei Funktionen mehrerer Veränderlichen* (in German), *Sitz. Königl. Bayer. Akad. Wiss. München* **36** (1906) 223.
- [97] W. Osgood, *Topics in the theory of functions of several complex variables*, Dover Publications, New York, U.S.A., (1966).
- [98] P. Deligne, ‘*Catégories tannakiennes*’ in the *Grothendieck Festschrift: a collection of articles written in honor of the 60th birthday of A. Grothendieck*, Birkhäuser, Boston, U.S.A., (1990), pg. 111.
- [99] B. Berndt, *Ramanujan’s notebooks, part I*, Springer-Verlag, New York, U.S.A., (1985).
- [100] T. Amdeberhan, O. Espinosa, I. Gonzalez, M. Harrison, V.H. Moll and A. Straub, *Ramanujan’s master theorem*, *Ramanujan J.* **29** (2012) 103.
- [101] G. Ólafsson and A. Pasquale, *Ramanujan’s master theorem for the hypergeometric Fourier transform associated with root systems*, *J. Fourier Anal. Appl.* **19** (2013) 1150 [[arXiv:1211.0024](#)].
- [102] I. Cherednik, *Inverse Harish-Chandra transform and difference operators*, *Int. Math. Res. Notices* **1997** (1997) 733.
- [103] B. Schapira, *Contributions to the hypergeometric function theory of Heckman and Opdam: sharp estimates, Schwartz space, heat kernel*, *Geom. Funct. Anal.* **18** (2008) 222.
- [104] J. Korevaar, *Tauberian theorems*, Springer-Verlag, Berlin Heidelberg, Germany, (2004).
- [105] D. Pappadopulo, S. Rychkov, J. Espin and R. Rattazzi, *OPE convergence in conformal field theory*, *Phys. Rev. D* **86** (2012) 105043 [[arXiv:1208.6449](#)] [[INSPIRE](#)].
- [106] A.L. Fitzpatrick, J. Kaplan and M.T. Walters, *Universality of long-distance AdS physics from the CFT bootstrap*, *JHEP* **08** (2014) 145 [[arXiv:1403.6829](#)] [[INSPIRE](#)].
- [107] L. Jianming and Z. Weixing, *A Wiener-Tauberian theorem for Fourier-Jacobi transform*, *Approx. Theor. Appl.* **13** (1997) 97.
- [108] E.K. Narayanan and A. Sitaram, *Analogues of the Wiener-Tauberian and Schwartz theorems for radial functions on symmetric spaces*, [arXiv:0905.3018](#).
- [109] S. Bochner, *Vorlesungen über Fouriersche Integrale* (in German), Akademische Verlagsgesellschaft, Leipzig, Germany, (1932).
- [110] C. Berg, J.P.R. Christensen and P. Ressel, *Harmonic analysis on semigroups*, Springer-Verlag, New York, U.S.A., (1984).
- [111] M. Flensted-Jensen and T. Koornwinder, *The convolution structure for Jacobi function expansions*, *Arkiv Matem.* **11** (1973) 245.
- [112] M. Flensted-Jensen and T.H. Koornwinder, *Jacobi functions: the addition formula and the positivity of the dual convolution structure*, *Arkiv Matem.* **17** (1979) 139.
- [113] K. Trimeche, *The positivity of the transmutation operators associated to the Cherednik operators for the root system BC_2* , *Math. J. Okayama Univ.* **58** (2016) 183.
- [114] M. Rösler, *Positive convolution structure for a class of Heckman-Opdam hypergeometric functions of type BC*, *J. Funct. Anal.* **258** (2010) 2779.
- [115] G. Ólafsson and A. Pasquale, *Paley-Wiener theorems for the Θ -spherical transform: an overview*, *Acta Appl. Math.* **81** (2004) 275.

- [116] A. Kaviraj, K. Sen and A. Sinha, *Analytic bootstrap at large spin*, *JHEP* **11** (2015) 083 [[arXiv:1502.01437](#)] [[INSPIRE](#)].
- [117] L.F. Alday and A. Zhiboedov, *Conformal bootstrap with slightly broken higher spin symmetry*, *JHEP* **06** (2016) 091 [[arXiv:1506.04659](#)] [[INSPIRE](#)].
- [118] D. Li, D. Meltzer and D. Poland, *Non-Abelian binding energies from the lightcone bootstrap*, *JHEP* **02** (2016) 149 [[arXiv:1510.07044](#)] [[INSPIRE](#)].
- [119] A. Kaviraj, K. Sen and A. Sinha, *Universal anomalous dimensions at large spin and large twist*, *JHEP* **07** (2015) 026 [[arXiv:1504.00772](#)] [[INSPIRE](#)].
- [120] D. Mazac, *Analytic bounds and emergence of AdS_2 physics from the conformal bootstrap*, *JHEP* **04** (2017) 146 [[arXiv:1611.10060](#)] [[INSPIRE](#)].
- [121] P. Dey, A. Kaviraj and K. Sen, *More on analytic bootstrap for $O(N)$ models*, *JHEP* **06** (2016) 136 [[arXiv:1602.04928](#)] [[INSPIRE](#)].
- [122] D. Jafferis, B. Mukhametzhanov and A. Zhiboedov, *Conformal bootstrap at large charge*, *JHEP* **05** (2018) 043 [[arXiv:1710.11161](#)] [[INSPIRE](#)].
- [123] M. Billò, V. Gonçalves, E. Lauria and M. Meineri, *Defects in conformal field theory*, *JHEP* **04** (2016) 091 [[arXiv:1601.02883](#)] [[INSPIRE](#)].
- [124] V. Schomerus, E. Sobko and M. Isachenkov, *Harmony of spinning conformal blocks*, *JHEP* **03** (2017) 085 [[arXiv:1612.02479](#)] [[INSPIRE](#)].
- [125] V. Schomerus and E. Sobko, *From spinning conformal blocks to matrix Calogero-Sutherland models*, *JHEP* **04** (2018) 052 [[arXiv:1711.02022](#)] [[INSPIRE](#)].
- [126] N. Reshetikhin, *Degenerate integrability of quantum spin Calogero-Moser systems*, *Lett. Math. Phys.* **107** (2017) 187 [[arXiv:1510.00492](#)] [[INSPIRE](#)].
- [127] M. Cornagliotto, M. Lemos and V. Schomerus, *Long multiplet bootstrap*, *JHEP* **10** (2017) 119 [[arXiv:1702.05101](#)] [[INSPIRE](#)].
- [128] I. Cherednik, *Quantum Knizhnik-Zamolodchikov equations and affine root systems*, *Commun. Math. Phys.* **150** (1992) 109 [[INSPIRE](#)].
- [129] S.N.M. Ruijsenaars and H. Schneider, *A new class of integrable systems and its relation to solitons*, *Annals Phys.* **170** (1986) 370 [[INSPIRE](#)].
- [130] J.F. van Diejen, *Integrability of difference Calogero-Moser systems*, *J. Math. Phys.* **35** (1994) 2983.
- [131] J.F. van Diejen, *Deformations of Calogero-Moser systems and finite Toda chains*, *Theor. Math. Phys.* **99** (1994) 549.
- [132] E. Rains, *BC_n -symmetric polynomials*, *Transform. Groups* **10** (2005) 63.
- [133] A. Okounkov, *BC -type interpolation Macdonald polynomials and binomial formula for Koornwinder polynomials*, *Transform. Groups* **3** (1998) 181.
- [134] M. van Meer and J. Stokman, *Double affine Hecke algebras and bispectral quantum Knizhnik-Zamolodchikov equations*, *Int. Math. Res. Notices* **2010** (2010) 969.
- [135] J.V. Stokman and T.H. Koornwinder, *Limit transitions for BC type multivariable orthogonal polynomials*, (2005).
- [136] T.H. Koornwinder, *Okounkov's BC -type interpolation Macdonald polynomials and their $q = 1$ limit*, [arXiv:1408.5993](#).

- [137] J. Horn, *Über die Convergenz der hypergeometrischen Reihen zweier und dreier Veränderlichen* (in German), *Math. Ann.* **34** (1889) 544.
- [138] H. Srivastava and M. Daoust, *Certain generalized Neumann expansions associated with the Kampé de Fériet function*, *Indag. Math.* **175** (1969) 449.
- [139] F.W.J. Olver ed., *NIST handbook of mathematical functions*, Cambridge University Press, Cambridge, U.K., (2010).
- [140] J.L. Burchnall and T.W. Chaundy, *Expansions of Appell's double hypergeometric functions*, *Quart. J. Math.* **11** (1940) 249.
- [141] R.J. Beerends, *Some special values for the BC type hypergeometric function*, *Contemp. Math.* **138** (1992) 27.
- [142] H. Exton, *On the system of partial differential equations associated with Appell's function F_4* , *J. Phys. A* **28** (1995) 631.
- [143] J.-F. Fortin and W. Skiba, *Conformal bootstrap in embedding space*, *Phys. Rev. D* **93** (2016) 105047 [[arXiv:1602.05794](#)] [[INSPIRE](#)].
- [144] J.-F. Fortin and W. Skiba, *Conformal differential operator in embedding space and its applications*, [arXiv:1612.08672](#) [[INSPIRE](#)].