Erratum: Searching for flavored gauge bosons

Eung Jin Chun,^a Arindam Das,^a Jinsu Kim^{b,1} and Jongkuk Kim^c

^aKorea Institute for Advanced Study, Seoul 02455, Korea

^bInstitute for Theoretical Physics, Georg-August University Göttingen,
Friedrich-Hund-Platz 1, Göttingen D-37077, Germany

^cDepartment of Physics, BK21 Physics Research Division, Institute of Basic Science,
Sungkyunkwan University, Suwon 440-746, Korea

E-mail: ejchun@kias.re.kr, arindam@kias.re.kr,
jinsu.kim@theorie.physik.uni-goettingen.de, jongkukkim@skku.edu

ERRATUM TO: JHEP02(2019)093

ARXIV EPRINT: [1811.04320](https://arxiv.org/abs/1811.04320)

We provide a change to the LHC bounds on $B - 3L_i$ ($i = 1, 2, 3$) scenario in ref. [1]. In figure 3 of the original article [1], the LHC bounds on $B - 3L_i$ are depicted as solid and dashed brown curves. They are modified as figure 1 in this erratum.

Accordingly the last paragraph of section 7 of the original article [1] is updated as follows. For all $B - 3L_i$ cases, the LHC bounds become the strongest bounds for large $M_{Z'}$ region; $M_{Z'} \gtrsim 150$ GeV for $i = 1, 2$ cases and $M_{Z'} \gtrsim 500$ GeV for $i = 3$ case. In the $B - 3L_1$ case, the LEP bounds on $e^-e^+ \rightarrow \ell^-\ell^+$ processes give the most stringent bounds on small $M_{Z'}$ region, $10 \text{ GeV} \lesssim M_{Z'} \lesssim 150 \text{ GeV}$. The $B - 3L_2$ case is particularly interesting. Due to the strong bounds from HFAG lepton universality test, only a small region $60 \text{ GeV} \lesssim M_{Z'} \lesssim 150 \text{ GeV}$ is consistent with all the constraints at 2σ .

Acknowledgments

We thank Hongkai Liu for pointing out our mistakes.

¹Corresponding author.

Figure 1. *Upper left panel:* bounds on g' and $M_{Z'}$ for the $B-3L_1$ model. The thin (thick) dashed-red and solid-black curves correspond to 1σ (2σ) bounds from LEP search [2] and SLD/LEP Z-decay lepton universality test [3], respectively. The green region is excluded by BaBar bounds [4]. The brown curve is the LHC bound from ATLAS observed limits. *Upper right panel:* bounds on g' and $M_{Z'}$ for the $B-3L_2$ model. The thin (thick) solid-black curve corresponds to 1σ (2σ) bound from the SLD/LEP Z-decay lepton universality test [3]. The blue (cyan) shaded region between thin (thick) dotted-blue curves is the 1σ (2σ) allowed region by the HFAG lepton universality test [5]. The brown curve is the LHC bound from ATLAS observed limits. The lighter- and darker-grey regions are excluded by neutrino-trident bound [6, 7] and the LHC bound for $Z \rightarrow 4\mu$ [8]. Therefore, only a small region $60 \text{ GeV} \lesssim M_{Z'} \lesssim 150 \text{ GeV}$ is consistent with all the constraints at 2σ . *Lower panel:* bounds on g' and $M_{Z'}$ for the $B-3L_3$ model. The thin (thick) dotted blue and solid black curves correspond to 1σ (2σ) bounds from the HFAG lepton universality test [5] and SLD/LEP Z-decay lepton universality test [3], respectively. The solid and dashed brown curves are the LHC bounds from ATLAS and CMS respectively.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E.J. Chun, A. Das, J. Kim and J. Kim, *Searching for Flavored Gauge Bosons*, *JHEP* **02** (2019) 093 [[arXiv:1811.04320](https://arxiv.org/abs/1811.04320)] [[INSPIRE](#)].
- [2] ALEPH, DELPHI, L3, OPAL collaborations and LEP Electroweak Working Group, *Electroweak Measurements in Electron-Positron Collisions at W-Boson-Pair Energies at LEP*, *Phys. Rept.* **532** (2013) 119 [[arXiv:1302.3415](https://arxiv.org/abs/1302.3415)] [[INSPIRE](#)].

- [3] ALEPH, DELPHI, L3, OPAL, SLD collaborations, LEP Electroweak Working Group, SLD Electroweak Group and SLD Heavy Flavour Group, *Precision electroweak measurements on the Z resonance*, *Phys. Rept.* **427** (2006) 257 [[hep-ex/0509008](#)] [[INSPIRE](#)].
- [4] BABAR collaboration, *Search for a Dark Photon in e^+e^- Collisions at BaBar*, *Phys. Rev. Lett.* **113** (2014) 201801 [[arXiv:1406.2980](#)] [[INSPIRE](#)].
- [5] HFLAV collaboration, *Averages of b-hadron, c-hadron and τ -lepton properties as of summer 2016*, *Eur. Phys. J. C* **77** (2017) 895 [[arXiv:1612.07233](#)] [[INSPIRE](#)].
- [6] CCFR collaboration, *Neutrino tridents and W Z interference*, *Phys. Rev. Lett.* **66** (1991) 3117 [[INSPIRE](#)].
- [7] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, *Neutrino Trident Production: A Powerful Probe of New Physics with Neutrino Beams*, *Phys. Rev. Lett.* **113** (2014) 091801 [[arXiv:1406.2332](#)] [[INSPIRE](#)].
- [8] CMS collaboration, *Search for an $L_\mu - L_\tau$ gauge boson using $Z \rightarrow 4\mu$ events in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **792** (2019) 345 [[arXiv:1808.03684](#)] [[INSPIRE](#)].