Double copy structure of parity-violating CFT correlators

Sachin Jain,^a Renjan Rajan John,^a Abhishek Mehta,^a Amin A. Nizami^b and Adithya Suresh^a

^aIndian Institute of Science Education and Research,
Homi Bhabha Road, Pashan, Pune 411 008, India

^bDepartment of Physics, Ashoka University,
Sonapat, Haryana 131029, India

E-mail: sachin.jain@iiserpune.ac.in,
renjan.john@acads.iiserpune.ac.in,
abhishek.mehta@students.iiserpune.ac.in, aan27cam@gmail.com,
s.adithya@students.iiserpune.ac.in

ABSTRACT: We show that general parity-violating 3d conformal field theories show a double copy structure for momentum space 3-point functions of conserved currents, stress tensor and marginal scalar operators. Splitting up the CFT correlator into two parts — called homogeneous and non-homogeneous — we show that double copy relations exist for each part separately. We arrive at similar conclusions regarding double copy structures using tree-level correlators of massless fields in dS_4 . We also discuss the flat space limit of these correlators. We further extend the double copy analysis to correlators involving higher-spin conserved currents, which suggests that the spin- s current correlator can be thought of as s copies of the spin one current correlator.

KEYWORDS: Conformal Field Theory, Conformal and W Symmetry, Field Theories in Lower Dimensions, Scattering Amplitudes

ARXIV EPRINT: [2104.12803](https://arxiv.org/abs/2104.12803)

Contents

1	Introduction	1
2	Notation and conventions	3
3	CFT correlators	4
3.1	$\langle J_s J_s O_3 \rangle$ for general spin s current	5
3.1.1	$\langle JJO_3 \rangle$	5
3.1.2	$\langle TTO_3 \rangle$	5
3.1.3	$\langle J_s J_s O_3 \rangle$	5
3.2	$\langle J_s J_s J_s \rangle$	6
3.2.1	$\langle JJJ \rangle$	7
3.2.2	$\langle TTT \rangle$	7
3.2.3	$\langle J_s J_s J_s \rangle$	9
4	Double copy structure of CFT correlators	9
4.1	$\langle TTO_3 \rangle$ and $\langle JJO_3 \rangle$	9
4.2	Double copy for higher-spin correlators	10
4.3	Double copy relation for $\langle JJJ \rangle$ and $\langle TTT \rangle$	11
4.4	Double copy structure for higher spin correlators	12
4.5	Spin s current correlator as s copies of the spin one current correlator	13
5	CFT correlators from dS_4 Feynman diagrams and the flat space limit	13
5.1	Amplitudes	13
5.1.1	Gauge amplitudes	14
5.1.2	Gravity amplitudes	15
5.2	Double copy structure of parity-violating amplitudes	17
5.3	CFT correlators from dS_4	18
5.3.1	$\langle JJO_3 \rangle$	19
5.3.2	$\langle TTO_3 \rangle$	20
5.3.3	$\langle JJJ \rangle$	21
5.3.4	$\langle TTT \rangle$	21
6	Discussion	22
A	Expressions in spinor-helicity variables	23
B	$\langle TJJ \rangle$	25
B.1	Mixed gauge-graviton amplitudes	25

C	Proof of some double copy relations	26
C.1	$\mathcal{M}_{W^3} \propto (\mathcal{M}_{F^3})^2$	26
C.2	$(\mathcal{M}_{\phi F^2})^2 \propto (\mathcal{M}_{\phi F \tilde{F}})^2$	27
C.3	$(\mathcal{M}_{F^3})^2 \propto (\mathcal{M}_{F^2 \tilde{F}})^2$	27
D	Momentum space expression of higher spin correlators	28
E	Double copy relations in spinor-helicity notation	28

---

1 Introduction

There has been a remarkable confluence in the study of CFT correlators and scattering amplitudes in recent years. As is well known, scattering amplitudes can be extracted by taking a suitable limit of appropriate CFT correlators - in position, momentum or Mellin space [1–6]. This enables a CFT derivation of various flat space amplitude results [6, 7]. Conversely, amplitude methods have recently been used in the study of CFTs [8, 9].

One of the interesting relationships that exists for flat space scattering amplitudes is the double-copy relation between gauge theory and gravity amplitudes, and the associated color-kinematics duality [10–12]. Here, substitution of color factors by kinematic factors in the numerators generates gravity amplitudes from gauge amplitudes, thereby manifesting a quadratic relationship between these two theories. This means that amplitudes involving gravitons can be built out from those involving gluons. The double copy relation was first observed in Einstein gravity and pure Yang-Mills theory, and later it was extended to a whole host of theories including higher derivative conformal gravity, higher derivative gauge theories and bi-adjoint scalar theories [13–15]. The 3-point structure for the higher-derivative theories is significant because it occurs in the momentum space form of CFT correlators of conserved currents, stress tensors and scalars. Double copy relations also exist for higher point tree amplitudes and loop amplitudes [16, 17]. For a comprehensive review see [18]. The analyses in these works were for the parity-even sector. We will show in this paper that similar relationships between amplitudes continue to hold with the inclusion of possible parity-violating terms in the Lagrangian.

In this work we study 3-point CFT correlators in momentum space. Some recent works where momentum space CFTs have been studied include [19–61]. In particular, the double-copy structures of certain parity-even 3-point functions were inferred in momentum space in [33, 43]. In three dimensions, in addition to the parity-even structures, one also needs to consider parity-odd structures. The most general form of 3-point functions in 3d CFTs is known to be of the form:

$$\langle J_{s_1} J_{s_2} J_{s_3} \rangle = \langle J_{s_1} J_{s_2} J_{s_3} \rangle_{\text{even}} + \langle J_{s_1} J_{s_2} J_{s_3} \rangle_{\text{odd}} \quad (1.1)$$

where J_s is a spin s conserved current. In position space one can show that the parity-even part can be obtained by adding contributions arising from the free-boson and the

free-fermion theories [62–64]. However, interacting 3d CFTs such as Chern-Simons-Matter theories can contain a non-trivial parity-odd sector as well [65–70].

In this paper we will demonstrate double copy relations between general parity-violating CFT_3 3-point correlators involving marginal scalars, spin one and spin two conserved currents. We will also show that a double-copy like structure exists for correlators involving higher spin conserved currents.

To establish our claim, it is convenient to split up CFT correlators into two parts, namely homogeneous and non-homogeneous parts. Their definition will be made clear in the next section. In particular, we show that under double copy relations, the homogeneous part maps to homogeneous part and the non-homogeneous part maps to non-homogeneous part. Let us illustrate this point by considering $\langle TTT \rangle$, the 3-point function of the stress tensor and $\langle JJJ \rangle$, the 3-point function of the conserved spin-1 current. The correlators can be written as:

$$\begin{aligned}\langle JJJ \rangle &= \langle JJJ \rangle_{\text{homogeneous}} + \langle JJJ \rangle_{\text{non-homogeneous}} \\ \langle TTT \rangle &= \langle TTT \rangle_{\text{homogeneous}} + \langle TTT \rangle_{\text{non-homogeneous}}\end{aligned}\tag{1.2}$$

The double copy relation is then given by:

$$\begin{aligned}\langle TTT \rangle_{\text{homogeneous}} &\propto (\langle JJJ \rangle_{\text{homogeneous}})^2 \\ \langle TTT \rangle_{\text{non-homogeneous}} &\propto (\langle JJJ \rangle_{\text{non-homogeneous}})^2\end{aligned}\tag{1.3}$$

where the proportionality factor is momentum dependent and is different for the two cases. It is given explicitly in section 4.3.

We will show that double copy relations hold even with the inclusion of the parity violating contributions. We demonstrate this using the results for parity even CFT correlators from [20, 27, 29, 33] and parity odd correlators from [61, 71], where CFT correlators were obtained by solving conformal Ward identities. There is another interesting way to fix the form of these correlators, initiated in [56]. Here, late-time tree level boundary correlators in Lorentzian dS_4 can be computed by first doing an equivalent calculation in flat Minkowski space. Thereafter, using certain conformal properties of the fields in dS_4 , the corresponding dS_4 correlators are obtained by simply dressing the result with a conformal-time integral factor. This Lorentzian dS_4 correlator also naturally computes a Euclidean CFT_3 correlator. We use this method to independently derive the parity-odd structures for the CFT correlators. This method provides a route to obtaining general parity-violating momentum space CFT_3 correlators without solving conformal Ward identities.

The rest of the paper is organised as follows. In section 2, we introduce the notation used in this paper. In section 3, we give the form of all the relevant CFT correlators. In section 4, we study double copy relations between various pairs of correlators. In section 5, we discuss the flat space limit and write down CFT correlators in terms of tree-level amplitudes without energy conservation for general parity-violating theories of gravitons, gluons and massless scalars in four dimensions. We also discuss here double copy relations for tree level dS_4 correlators and flat space scattering amplitudes. In section 6, we conclude and give some directions for future study. In appendix A we give our results for the scattering amplitudes and the CFT correlators in terms of the spinor helicity variables. We

give the details of the $\langle JJT \rangle$ correlator in appendix B. In appendix C we give the calculational details used in establishing double copy relations. In appendix D we give explicit momentum space results for a few correlators involving higher spin currents. Appendix E contains a few details regarding some constraints on OPE coefficients arising from double copy relations.

2 Notation and conventions

In this paper we denote 4-dimensional Lorentzian momenta and polarisation vectors by k_i^μ and z_i^μ respectively. Here i is a particle index and $\mu = 0, 1, 2, 3$ is the Lorentz index. For massless spin 2 particles the polarisation tensor can be written as an outer product $z_i^{\mu\nu} = z_i^\mu z_i^\nu$. We choose the following gauge to work with null momenta:

$$k_i^\mu = (k_i, \vec{k}_i), \quad z_i^\mu = (0, \vec{z}_i) \quad (2.1)$$

where $k_i = |\vec{k}_i|$ is the magnitude of the 3-momentum.

The 3-dimensional CFT will be Euclidean and current conservation constraints translate to transversality: $k_i \cdot z_i = 0$. We will also take $z_i \cdot z_i = 0$ which in Euclidean signature implies that the components of $\vec{z}_i$ will be complex.

In our computation we will find it useful to introduce the following notation for various combinations of magnitudes of momenta:

$$E = k_1 + k_2 + k_3, \quad b_{ij} = k_i k_j, \quad b_{123} = k_1 k_2 + k_2 k_3 + k_3 k_1, \quad c_{123} = k_1 k_2 k_3 \quad (2.2)$$

We also introduce the following notation:

$$J^2 = (k_1 + k_2 + k_3)(-k_1 + k_2 + k_3)(k_1 - k_2 + k_3)(k_1 + k_2 - k_3) \quad (2.3)$$

We will make use of spinor-helicity notation. The momentum vector p_μ for massless scattering in 4-dimensional flat space-time can be written as $p_\mu \sigma_{\alpha\dot{\alpha}}^\mu = p_{\alpha\dot{\alpha}} = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}}$ where λ denotes a spinor-helicity variable. Since 4d amplitudes are related to 3d CFT correlators it will be useful to have a 3d version of this formalism by utilising the time-like vector $\tau^\mu = (1, 0, 0, 0)$, or $\tau^{\alpha\dot{\beta}} = \epsilon^{\alpha\dot{\beta}}$ which can be used to go from dotted to undotted indices (see appendix B of [33] and [43]). We use this to define $\bar{\lambda}^\alpha \equiv \tau^{\alpha\dot{\beta}} \tilde{\lambda}_{\dot{\beta}}$.

For a correlator comprising conserved currents J_{s_i} the conformal Ward identity in spinor-helicity notation takes the following form:

$$\widetilde{K}^\kappa \left\langle \frac{J_{s_1}}{k_1^{s_1-1}} \frac{J_{s_2}}{k_2^{s_2-1}} \frac{J_{s_3}}{k_3^{s_3-1}} \right\rangle = \text{transverse Ward identity} \quad (2.4)$$

where J_{s_i} are conserved currents with spin s_i and dimension $\Delta = s_i + 1$ and the R.H.S of the above equation is proportional to the transverse Ward identities associated with the correlator. For instance, for the case of $\langle JJJ \rangle$ where J is the spin-1 conserved current, the conformal Ward identity takes the form:

$$\widetilde{K}^\kappa \langle J^- J^- J^- \rangle = 2 \left(z_1^{-\kappa} \frac{k_{1\mu}}{k_1^2} \langle J^\mu J^- J^- \rangle + z_2^{-\kappa} \frac{k_{2\mu}}{k_2^2} \langle J^- J^\mu J^- \rangle + z_3^{-\kappa} \frac{k_{3\mu}}{k_3^2} \langle J^- J^- J^\mu \rangle \right) \quad (2.5)$$

where we see that the r.h.s. of the equation is given by the transverse Ward identities.

The correlator $\langle J_{s_1} J_{s_2} J_{s_3} \rangle$ is given by the sum of two terms that satisfy the homogeneous and non homogeneous equations respectively:

$$\langle J_{s_1} J_{s_2} J_{s_3} \rangle = \langle J_{s_1} J_{s_2} J_{s_3} \rangle_{\text{homogeneous}} + \langle J_{s_1} J_{s_2} J_{s_3} \rangle_{\text{non homogeneous}} \quad (2.6)$$

where $\langle J_{s_1} J_{s_2} J_{s_3} \rangle_{\text{homogeneous}}$ satisfies:

$$\widetilde{K}^\kappa \left\langle \frac{J_{s_1}}{k_1^{s_1-1}} \frac{J_{s_2}}{k_2^{s_2-1}} \frac{J_{s_3}}{k_3^{s_3-1}} \right\rangle_{\text{homogeneous}} = 0 \quad (2.7)$$

and $\langle J_{s_1} J_{s_2} J_{s_3} \rangle_{\text{non homogeneous}}$ satisfies:

$$\widetilde{K}^\kappa \left\langle \frac{J_{s_1}}{k_1^{s_1-1}} \frac{J_{s_2}}{k_2^{s_2-1}} \frac{J_{s_3}}{k_3^{s_3-1}} \right\rangle_{\text{non homogeneous}} = \text{transverse Ward identity} \quad (2.8)$$

The transverse Ward identity is determined in terms of two-point functions. Hence in the momentum space expression for the correlator $\langle J_{s_1} J_{s_2} J_{s_3} \rangle$, we identify the part proportional to the 2-point function coefficient to be the solution to the non-homogeneous Ward identity and the part obtained by setting the coefficient of the 2-point function to zero to be the solution to the homogeneous Ward identity. We will use subscripts **h** and **nh** to denote the solutions to the homogeneous and non-homogeneous equations respectively.

For a correlator with at least one scalar operator O_Δ with conformal dimension Δ the conformal Ward identity has a trivial r.h.s. and is given by:

$$\widetilde{K}^\kappa \left\langle \frac{O_\Delta}{k_1^{\Delta-2}} \frac{J_{s_2}}{k_2^{s_2-1}} \frac{J_{s_3}}{k_3^{s_3-1}} \right\rangle = 0 \quad (2.9)$$

which holds true both when $s_2 = s_3$ and $s_2 \neq s_3$.

We will denote flat space amplitudes by $\mathcal{A}$. The corresponding correlators in dS_4 ,¹ with all insertions at the equal-time spatial-slice $\eta = 0$, will be denoted by $\mathcal{M}$.

3 CFT correlators

In this section we present the momentum space expressions for 3-point CFT_3 correlators comprising spin-1 conserved current J , stress tensor T , higher spin conserved currents J_s with spin $s > 2$ and marginal scalar operators O_3 . The parity-even sector of 3-point CFT correlators has been studied by solving the associated conformal Ward identities in a series of works [20, 25, 27, 29]. In [61, 71] we studied the parity-odd sector of 3-point correlators by solving conformal Ward identities and using the technique of spin-raising and weight-shifting operators in momentum space [58, 59, 72].

We present our results after contracting the momentum space expressions with null transverse polarization vectors.

---

¹Up to overall conformal time integral factors as discussed in section 5.

3.1 $\langle J_s J_s O_3 \rangle$ for general spin s current

In this subsection we write down correlators of the form $\langle J_s J_s O_3 \rangle$ for a general spin s . For $s = 1, 2$ we can write down their explicit form in momentum space easily. A similar momentum space expression for a correlator involving general spin s current is very cumbersome. However, it takes a very simple form when written in terms of spinor-helicity variables. We also note that, as discussed in the previous section, a correlator of the form $\langle J_s J_s O_3 \rangle$ only has a homogeneous part. The homogeneous correlator is given by the contribution from the parity-even and the parity-odd sectors:

$$\langle J_s J_s O_3 \rangle_{\mathbf{h}} = \langle J_s J_s O_3 \rangle_{\text{even}, \mathbf{h}} + \langle J_s J_s O_3 \rangle_{\text{odd}, \mathbf{h}} \quad (3.1)$$

3.1.1 $\langle JJO_3 \rangle$

Let us first consider the 3-point correlator comprising two spin-1 conserved currents and a marginal scalar operator. The momentum space expression for the parity-even part of the correlation function is [20, 29, 33]:

$$\begin{aligned} \langle JJO_3 \rangle_{\text{even}, \mathbf{h}} &= \frac{(E + k_3)}{E^2} \left[2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + E(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2 \right] \\ \langle JJO_3 \rangle_{\text{even}, \mathbf{nh}} &= 0. \end{aligned} \quad (3.2)$$

The momentum space expression for the parity-odd part of the correlator is [61, 71]:

$$\begin{aligned} \langle JJO_3 \rangle_{\text{odd}, \mathbf{h}} &= \frac{(E + k_3)}{E^2} \left[k_2 \epsilon^{k_1 z_1 z_2} - k_1 \epsilon^{k_2 z_1 z_2} \right] \\ \langle JJO_3 \rangle_{\text{odd}, \mathbf{nh}} &= 0. \end{aligned} \quad (3.3)$$

3.1.2 $\langle TTO_3 \rangle$

Let us now consider the 3-point correlator comprising two stress-tensor insertions and a marginal scalar. The parity-even part of the correlator is given by [20, 29, 33]:

$$\begin{aligned} \langle TTO_3 \rangle_{\text{even}, \mathbf{h}} &= k_1 k_2 \frac{E + 3k_3}{E^4} \left[2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + E(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2 \right]^2 \\ \langle TTO_3 \rangle_{\text{even}, \mathbf{nh}} &= 0 \end{aligned} \quad (3.4)$$

whereas the parity-odd part of this correlator is [71]:

$$\begin{aligned} \langle TTO_3 \rangle_{\text{odd}, \mathbf{h}} &= \frac{E + 3k_3}{E^4} (k_2 \epsilon^{k_1 z_1 z_2} - k_1 \epsilon^{k_2 z_1 z_2}) \\ &\quad \times (\vec{z}_1 \cdot \vec{z}_2)(\vec{k}_1 \cdot \vec{k}_2 - k_1 k_2)(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) \\ \langle TTO_3 \rangle_{\text{odd}, \mathbf{nh}} &= 0 \end{aligned} \quad (3.5)$$

3.1.3 $\langle J_s J_s O_3 \rangle$

Correlators of the form $\langle J_s J_s O_3 \rangle$ have unwieldy expressions in momentum space. The easiest way to determine them is to use weight-shifting operators [59]. For this, we first

derive the momentum space expressions for correlators with spin-3 and spin-4 currents using weight-shifting operators [59]:

$$\begin{aligned}\langle J_3 J_3 O_3 \rangle_{\text{even}, \mathbf{h}} &= P_1^{(3)} P_2^{(3)} H_{12}^3 \langle O_2 O_2 O_3 \rangle \\ \langle J_4 J_4 O_3 \rangle_{\text{even}, \mathbf{h}} &= P_1^{(4)} P_2^{(4)} H_{12}^4 \langle O_2 O_2 O_3 \rangle\end{aligned}\quad (3.6)$$

where $P_i^{(s)}$ are spin- s projectors [59] and H_{12} is a bi-local weight shifting operator which raises the spin at points 1 and 2 and lowers the dimensions at points 1 and 2. In momentum space the operator takes the form [59]:

$$H_{12} = 2(\vec{z}_1 \cdot \vec{K}_{12})(\vec{z}_2 \cdot \vec{K}_{12}) - (\vec{z}_1 \cdot \vec{z}_2) K_{12}^2 \quad (3.7)$$

where $K_{12}^i \equiv \frac{\partial}{\partial k_1^i} - \frac{\partial}{\partial k_2^i}$. The explicit form of the correlator in (3.6) is complicated and not reproduced here. A similar expression can be written down for parity-odd contribution.

These correlators when expressed in spinor-helicity variables take a very simple form. For this, let us consider the 3-point correlator of two higher spin conserved currents J_s with spin s and a marginal scalar. The parity-even part of the correlator is given by [71]:

$$\begin{aligned}\langle J_s^- J_s^- O_3 \rangle_{\text{even}, \mathbf{h}} &= \frac{E + (2s - 1)k_3}{E^{2s}} \langle 12 \rangle^{2s} \\ \langle J_s^+ J_s^+ O_3 \rangle_{\text{even}, \mathbf{h}} &= \frac{E + (2s - 1)k_3}{E^{2s}} \langle \overline{12} \rangle^{2s}\end{aligned}\quad (3.8)$$

whereas the parity-odd part of the correlator is [71]:

$$\begin{aligned}\langle J_s^- J_s^- O_3 \rangle_{\text{odd}, \mathbf{h}} &= i \frac{E + (2s - 1)k_3}{E^{2s}} \langle 12 \rangle^{2s} \\ \langle J_s^+ J_s^+ O_3 \rangle_{\text{odd}, \mathbf{h}} &= -i \frac{E + (2s - 1)k_3}{E^{2s}} \langle \overline{12} \rangle^{2s}\end{aligned}\quad (3.9)$$

As stated earlier, the non-homogeneous piece vanishes for both parity-even and parity-odd correlators:

$$\langle J_s J_s O_3 \rangle_{\mathbf{nh}} = 0. \quad (3.10)$$

One can check that for the cases when $s = 1$ and $s = 2$, the momentum space results in (3.2), (3.3), (3.4), and (3.5) when re-expressed in spinor-helicity variables match the above results.

3.2 $\langle J_s J_s J_s \rangle$

In this subsection we write down correlators of the form $\langle J_s J_s J_s \rangle$ for a general s . Unlike $\langle J_s J_s O \rangle$, these correlators have both homogeneous and non-homogeneous pieces:

$$\langle J_s J_s J_s \rangle_{\mathbf{h}} = \langle J_s J_s J_s \rangle_{\text{even}, \mathbf{h}} + \langle J_s J_s J_s \rangle_{\text{odd}, \mathbf{h}}. \quad (3.11)$$

We give the explicit forms for $\langle JJJ \rangle$ and $\langle TTT \rangle$ in momentum space. In both cases, there are exactly two homogeneous pieces, one parity-even and another parity-odd. We also note that there is only one parity-even non-homogeneous contribution. The non-homogeneous part of the parity-odd correlator is always a contact term. For a general spin s , the momentum space expression is very complicated. However, in spinor-helicity variables it becomes simple and we express the homogeneous part in these variables. The analogous expression for non-homogeneous part is not yet known.

3.2.1 $\langle JJJ \rangle$

Let us now consider the 3-point correlator comprising three spin-1 conserved current insertions. The parity-even part of the correlator is given by² [20, 27, 33]:

$$\begin{aligned}\langle JJJ \rangle_{\text{even,h}} &= \frac{c_{JJJ}^{\text{even}}}{E^3} \left[2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_3)(\vec{z}_3 \cdot \vec{k}_1) + E\{k_3(\vec{z}_1 \cdot \vec{z}_2)(\vec{z}_3 \cdot \vec{k}_1) + \text{cyclic}\} \right] \\ \langle JJJ \rangle_{\text{even,nh}} &= -\frac{2c_{JJJ}^{\text{even}}}{E} [(\vec{z}_1 \cdot \vec{z}_2)(\vec{z}_3 \cdot \vec{k}_1) + \text{cyclic}] \end{aligned} \quad (3.12)$$

Note that c_{JJJ}^{even} appears in two point function $\langle J_\mu(k)J_\nu(-k) \rangle_{\text{even}} = c_{JJJ}^{\text{even}}\pi_{\mu\nu}(k)k$ where $\pi_{\mu\nu}(k)$ is the transverse projector:

$$\pi_\alpha^\mu(p) \equiv \delta_\alpha^\mu - \frac{p^\mu p_\alpha}{p^2}. \quad (3.13)$$

The parity-odd part of the correlator is given by [61, 71]

$$\begin{aligned}\langle JJJ \rangle_{\text{odd,h}} &= \frac{c_{JJJ}^{\text{odd}}}{E^3} \left[\left\{ (\vec{k}_1 \cdot \vec{z}_3) \left(\epsilon^{k_3 z_1 z_2} k_1 - \epsilon^{k_1 z_1 z_2} k_3 \right) + (\vec{k}_3 \cdot \vec{z}_2) \left(\epsilon^{k_1 z_1 z_3} k_2 - \epsilon^{k_2 z_1 z_3} k_1 \right) \right. \right. \\ &\quad \left. \left. - (\vec{z}_2 \cdot \vec{z}_3) \epsilon^{k_1 k_2 z_1} E + \frac{k_1}{2} \epsilon^{z_1 z_2 z_3} E(E - 2k_1) \right\} + \text{cyclic perm} \right] \\ \langle JJJ \rangle_{\text{odd,nh}} &= c_{JJJ}^{\text{odd}} \epsilon^{z_1 z_2 z_3} \end{aligned} \quad (3.14)$$

where c_{JJJ}^{odd} arises in parity-odd contribution to the two point function $\langle J_\mu(k)J_\nu(-k) \rangle_{\text{odd}} = c_{JJJ}^{\text{odd}}\epsilon_{\mu\nu k}$. Let us also note that the non-homogeneous contribution to the parity-odd part of $\langle JJJ \rangle$ (term proportional to c_{JJJ}^{odd}) in (3.14) is a contact term.

3.2.2 $\langle TTT \rangle$

Let us now consider the 3-point correlator comprising three stress-tensor insertions. The parity-even contribution to the correlator is given by [20, 27, 33]:

$$\begin{aligned}\langle TTT \rangle_{\text{even,h}} &= \frac{c_{TTT}^{\text{even}} c_{123}^2}{J^2 E^5} \left[(\vec{z}_1 \cdot \vec{k}_2 \vec{z}_2 \cdot \vec{k}_3 \vec{z}_3 \cdot \vec{k}_1)^2 \right. \\ &\quad \left. + \frac{E}{2} \vec{z}_1 \cdot \vec{k}_2 \vec{z}_2 \cdot \vec{k}_3 \vec{z}_3 \cdot \vec{k}_1 (k_3 \vec{z}_1 \cdot \vec{z}_2 \vec{z}_3 \cdot \vec{k}_1 + \text{cyclic}) \right] \\ \langle TTT \rangle_{\text{even,nh}} &= 2c_{TTT}^{\text{even}} \left[\left(\frac{c_{123}}{E^2} + \frac{b_{123}}{E} - E \right) (\vec{z}_1 \cdot \vec{z}_2 \vec{z}_3 \cdot \vec{k}_1 + \text{cyclic})^2 + (k_1^3 + k_2^3 + k_3^3) \mathcal{A}_{ct} \right] \end{aligned} \quad (3.15)$$

where c_{TTT}^{even} appears in the even part of the 2-point function $\langle TT \rangle$ and

$$\mathcal{A}_{ct} = (\vec{z}_1 \cdot \vec{z}_2)(\vec{z}_2 \cdot \vec{z}_3)(\vec{z}_3 \cdot \vec{z}_1). \quad (3.16)$$

We note that the term proportional to $\mathcal{A}_{ct}$ in (3.15) is a contact term and will be ignored below in establishing the double copy relation.

²Throughout this paper we suppress color indices and the structure constants.

We see that apart from the physical pole at $E = 0$, $\langle TTT \rangle_{\text{even}, \mathbf{h}}$ displays an unphysical pole at $J = 0$. However, this is only an artefact of the basis we have chosen to work with and by working in a suitable basis we can get rid of the unphysical pole. For instance, it can be shown that after a clever use of 3d degeneracies, one can express the correlator as follows³

$$\langle TTT \rangle_{\text{even}, \mathbf{h}} = \frac{c_{TTT}^{\text{even}} c_{123}}{E^6} F_2[1, 2, 3] F_2[2, 3, 1] F_2[3, 1, 2] \quad (3.17)$$

where

$$F_2[i, j, l] = \left[(\vec{z}_i \cdot \vec{z}_j) E(E - 2k_l) + 2(\vec{z}_i \cdot \vec{k}_j)(\vec{z}_j \cdot \vec{k}_i) \right]. \quad (3.18)$$

For details of this computation see appendix C.1.

The parity-odd contribution to the correlator is given by [71]

$$\begin{aligned} \langle TTT \rangle_{\text{odd}} = & A_1 \epsilon^{k_3 k_1 z_1} \epsilon^{k_1 k_2 z_2} \epsilon^{k_2 k_3 z_3} (\vec{k}_2 \cdot \vec{z}_1)(\vec{k}_3 \cdot \vec{z}_2)(\vec{k}_1 \cdot \vec{z}_3) \\ & + A_2 \epsilon^{k_2 k_3 z_3} (\vec{k}_1 \cdot \vec{z}_3)(\vec{k}_3 \cdot \vec{z}_2)^2 (\vec{k}_2 \cdot \vec{z}_1)^2 \\ & + A_2 (k_2 \leftrightarrow k_3) \epsilon^{k_1 k_2 z_2} (\vec{k}_1 \cdot \vec{z}_3)^2 (\vec{k}_3 \cdot \vec{z}_2)(\vec{k}_2 \cdot \vec{z}_1)^2 \\ & + A_2 (k_1 \leftrightarrow k_3) \epsilon^{k_3 k_1 z_1} (\vec{k}_1 \cdot \vec{z}_3)^2 (\vec{k}_3 \cdot \vec{z}_2)^2 (\vec{k}_2 \cdot \vec{z}_1) \end{aligned} \quad (3.19)$$

where the homogeneous piece in the form factor is given by:

$$A_{1, \mathbf{h}} = c_{TTT}^{\text{odd}} \frac{c_{123}^2}{2J^4 E^4}, \quad A_{2, \mathbf{h}} = -c_{TTT}^{\text{odd}} \frac{b_{12} c_{123}^2}{2J^4 E^4} \quad (3.20)$$

Just as in the parity-even case, we see that the form factors have an unphysical extra pole at $J = 0$. This can again be gotten rid of by working in a suitable basis where it takes the form⁴

$$\begin{aligned} \langle TTT \rangle_{\text{odd}} = & c_{TTT}^{\text{odd}} \frac{c_{123}}{E^6} \left[2(\vec{z}_1 \cdot \vec{k}_2 \vec{z}_2 \cdot \vec{k}_3 \vec{z}_3 \cdot \vec{k}_1) + E \left((\vec{z}_1 \cdot \vec{z}_2 \vec{z}_3 \cdot \vec{k}_1) k_3 + \text{cyclic} \right) \right] \\ & \left[-(\vec{k}_3 \cdot \vec{z}_2) \left(\epsilon^{k_3 z_1 z_3} k_1 + \epsilon^{k_1 z_1 z_3} (E - k_3) \right) - (\vec{k}_1 \cdot \vec{z}_3) \left(\epsilon^{k_2 z_1 z_2} k_1 + \epsilon^{k_1 z_1 z_2} (E - k_2) \right) \right. \\ & \left. + E(\vec{z}_2 \cdot \vec{z}_3) \epsilon^{k_1 k_2 z_1} - \frac{1}{2} k_1 E(E - 2k_1) \epsilon^{z_1 z_2 z_3} + \text{cyclic perm.} \right] \end{aligned} \quad (3.21)$$

The non-homogeneous part of the parity-odd correlator is determined by the form factors in (3.19):

$$A_{1, \mathbf{nh}} = -c_{TT}^{\text{odd}} \frac{12(k_1^2 + k_2^2 + k_3^2)}{J^4} \quad (3.22)$$

$$A_{2, \mathbf{nh}} = c_{TT}^{\text{odd}} \frac{(k_3^4 + 7k_3^2(k_1^2 + k_2^2) + 4(k_1^4 + 4k_1^2 k_2^2 + k_2^4))}{J^4} \quad (3.23)$$

where c_{TT}^{odd} appears in the parity-odd part of the two point function $\langle TT \rangle$. It can be shown that, the parity-odd non-homogeneous contribution to the correlator is a contact term, see [71] for details. Since the non-homogeneous part is a contact term, this will be ignored while checking double copy relations.

³See section 5.3 for a derivation using gravity.

⁴See section 5.3.

3.2.3 $\langle J_s J_s J_s \rangle$

Let us now consider the 3-point correlator comprising three higher spin conserved currents J_s with spin s . The parity-even contribution to the correlator is given by [71]:

$$\begin{aligned}\langle J_s^- J_s^- J_s^- \rangle_{\text{even}, \mathbf{h}} &= \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle 12 \rangle^s \langle 23 \rangle^s \langle 31 \rangle^s \\ \langle J_s^+ J_s^+ J_s^+ \rangle_{\text{even}, \mathbf{h}} &= \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle \overline{12} \rangle^s \langle \overline{23} \rangle^s \langle \overline{31} \rangle^s\end{aligned}\quad (3.24)$$

The parity-odd contribution to the correlator is given⁵ by [71]

$$\begin{aligned}\langle J_s^- J_s^- J_s^- \rangle_{\text{odd}, \mathbf{h}} &= i \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle 12 \rangle^s \langle 23 \rangle^s \langle 31 \rangle^s \\ \langle J_s^+ J_s^+ J_s^+ \rangle_{\text{odd}, \mathbf{h}} &= -i \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle \overline{12} \rangle^s \langle \overline{23} \rangle^s \langle \overline{31} \rangle^s\end{aligned}\quad (3.25)$$

4 Double copy structure of CFT correlators

In this section we discuss the double copy structure of CFT 3-point correlation functions in momentum space. We will see that the double copy relations are such that homogeneous terms are mapped to homogeneous terms and non-homogeneous terms are mapped to non-homogeneous terms. We will establish our claims in momentum space for $\langle TTO_3 \rangle$ and $\langle JJO_3 \rangle$ and for $\langle TTT \rangle$ and $\langle JJJ \rangle$. We use the spinor-helicity variables to show this for correlators such as $\langle J_s J_s O_3 \rangle$ and $\langle J_s J_s J_s \rangle$.

4.1 $\langle TTO_3 \rangle$ and $\langle JJO_3 \rangle$

The following double copy structure of $\langle TTO_3 \rangle_{\text{even}}$ was established in [33]:

$$\langle TTO_3 \rangle_{\text{even}, \mathbf{h}} = \frac{(E + 3k_3)k_1 k_2}{(E + k_3)^2} \langle JJO_3 \rangle_{\text{even}, \mathbf{h}} \langle JJO_3 \rangle_{\text{even}, \mathbf{h}} \quad (4.1)$$

From the explicit expressions for the correlators in (3.2), (3.3) and (3.5) we notice that the double copy relations extends to the parity-odd sector:

$$\langle TTO_3 \rangle_{\text{odd}, \mathbf{h}} = \frac{(E + 3k_3)k_1 k_2}{(E + k_3)^2} \langle JJO_3 \rangle_{\text{odd}, \mathbf{h}} \langle JJO_3 \rangle_{\text{even}, \mathbf{h}} \quad (4.2)$$

Remarkably, we also notice from (3.3) and (3.4) that $\langle TTO_3 \rangle_{\text{even}}$ is also given by the square of $\langle JJO_3 \rangle_{\text{odd}}$

$$\langle TTO_3 \rangle_{\text{even}, \mathbf{h}} = \frac{(E + 3k_3)k_1 k_2}{(E + k_3)^2} \langle JJO_3 \rangle_{\text{odd}, \mathbf{h}} \langle JJO_3 \rangle_{\text{odd}, \mathbf{h}} \quad (4.3)$$

---

⁵Let us note that the parity-odd correlator, when written in spinor-helicity variables, gets an extra factor of i as compared to parity-even correlator. This arises due to the fact that when converting momentum space parity-odd answer to spinor-helicity variables, the Levi-Civita tensor ϵ^{ijk} is expressed in terms of Pauli matrices and their algebra gives rise to this extra factor. More precisely

$$\text{Tr}(\sigma_i \sigma_j \sigma_k) = 2i \epsilon_{ijk}.$$

The above double copy relations for $\langle TTO_3 \rangle_{\text{even}}$ and $\langle TTO_3 \rangle_{\text{odd}}$ immediately imply the following double copy structure for the complete correlator:

$$\begin{aligned} \langle TTO_3 \rangle_{\text{even},\mathbf{h}} + \langle TTO_3 \rangle_{\text{odd},\mathbf{h}} &= \frac{(E+3k_3)k_1k_2}{(E+k_3)^2} (\langle JJO_3 \rangle_{\text{even},\mathbf{h}} + \langle JJO_3 \rangle_{\text{odd},\mathbf{h}})^2 \\ \implies \langle TTO_3 \rangle_{\mathbf{h}} &= \frac{(E+3k_3)k_1k_2}{(E+k_3)^2} \langle JJO_3 \rangle_{\mathbf{h}}^2 \end{aligned} \quad (4.4)$$

In writing the above double copy relation it is crucial that we have the following relation between $\langle JJO_3 \rangle_{\text{even}}$ and $\langle JJO_3 \rangle_{\text{odd}}$:

$$\langle JJO_3 \rangle_{\text{even},\mathbf{h}}^2 = c \langle JJO_3 \rangle_{\text{odd},\mathbf{h}}^2 \quad (4.5)$$

where c is some constant. As noted in section 2, for correlators such as $\langle TTO_3 \rangle$ and $\langle JJO_3 \rangle$ the conformal Ward identity (in spinor helicity variables) does not have a non-homogeneous term. Hence the double copy structure that we obtained above is purely for correlators that satisfy the homogeneous conformal Ward identity.

4.2 Double copy for higher-spin correlators

We will now extend our analysis of the double copy structure of $\langle TTO_3 \rangle$ to higher spin correlators of the form $\langle J_s J_s O_3 \rangle$. Using (3.6) one can show that the higher spin correlators take the following form:

$$\begin{aligned} \langle J_4 J_4 O_3 \rangle_{\text{even},\mathbf{h}} &= k_1 k_2 \frac{(E+7k_3)}{(E+3k_3)^2} \langle J_2 J_2 O_3 \rangle_{\text{even}}^2 \\ \langle J_3 J_3 O_3 \rangle_{\text{even},\mathbf{h}} &= k_1 k_2 \frac{(E+5k_3)}{(E+k_3)(E+3k_3)} \langle J_2 J_2 O_3 \rangle_{\text{even}} \langle J_1 J_1 O_3 \rangle_{\text{even}}. \end{aligned} \quad (4.6)$$

Calculating the parity-odd contribution to these three point functions is difficult due to the high amount of degeneracy [61, 71]. However, in spinor helicity variables the computation becomes easier. In these variables one has the following remarkable relation between the parity-even and parity-odd contributions [71]:

$$\langle J_s^- J_s^- O_3 \rangle_{\text{even},\mathbf{h}} = i \langle J_s^- J_s^- O_3 \rangle_{\text{odd},\mathbf{h}}, \quad \langle J_s^+ J_s^+ O_3 \rangle_{\text{even},\mathbf{h}} = -i \langle J_s^+ J_s^+ O_3 \rangle_{\text{odd},\mathbf{h}} \quad (4.7)$$

for any s and all the other spinor helicity components are zero. Using this one can generalise (4.6) to include the parity-odd sector. The double copy relation (4.6) then becomes

$$\begin{aligned} \langle J_4 J_4 O_3 \rangle_{\text{even},\mathbf{h}} + \langle J_4 J_4 O_3 \rangle_{\text{odd},\mathbf{h}} &= \frac{k_1 k_2 (E+7k_3)}{(E+3k_3)^2} (\langle J_2 J_2 O_3 \rangle_{\text{even},\mathbf{h}} + \langle J_2 J_2 O_3 \rangle_{\text{odd},\mathbf{h}})^2 \\ \implies \langle J_4 J_4 O_3 \rangle_{\mathbf{h}} &= \frac{k_1 k_2 (E+7k_3)}{(E+3k_3)^2} \langle J_2 J_2 O_3 \rangle_{\mathbf{h}}^2 \\ \langle J_3 J_3 O_3 \rangle_{\text{even},\mathbf{h}} + \langle J_3 J_3 O_3 \rangle_{\text{odd},\mathbf{h}} &= \frac{k_1 k_2 (E+5k_3)}{(E+k_3)(E+3k_3)} (\langle J_2 J_2 O_3 \rangle_{\text{even},\mathbf{h}} + \langle J_2 J_2 O_3 \rangle_{\text{odd},\mathbf{h}}) \\ &\quad \times (\langle J_1 J_1 O_3 \rangle_{\text{even},\mathbf{h}} + \langle J_1 J_1 O_3 \rangle_{\text{odd},\mathbf{h}}) \\ \implies \langle J_3 J_3 O_3 \rangle_{\mathbf{h}} &= \frac{k_1 k_2 (E+5k_3)}{(E+k_3)(E+3k_3)} \langle J_2 J_2 O_3 \rangle_{\mathbf{h}} \langle J_1 J_1 O_3 \rangle_{\mathbf{h}} \end{aligned}$$

To write down the above double copy relations it is crucial that we have the following relation between the parity-odd and parity-even parts of the correlators:

$$\begin{aligned} \langle J_2 J_2 O_3 \rangle_{\text{even}, \mathbf{h}}^2 &= \langle J_2 J_2 O_3 \rangle_{\text{odd}, \mathbf{h}}^2 \\ \langle J_2 J_2 O_3 \rangle_{\text{even}, \mathbf{h}} \langle J_1 J_1 O_3 \rangle_{\text{even}, \mathbf{h}} &= \langle J_2 J_2 O_3 \rangle_{\text{odd}, \mathbf{h}} \langle J_1 J_1 O_3 \rangle_{\text{odd}, \mathbf{h}} \end{aligned} \quad (4.8)$$

As we noted in the case of the double copy structure of $\langle TTO_3 \rangle$ in terms of $\langle JJO_3 \rangle$, the conformal Ward identity for correlators of the form $\langle J_s J_s O_3 \rangle$ does not have a non-homogeneous term. Hence the double copy relations that we arrived at here are purely for the homogeneous terms.

Double copy relation for general spin. One can easily extend the above analysis to correlators of the form $\langle J_s J_s O_3 \rangle$ in the spinor helicity variables, see [71]. The correlation functions are given by:

$$\begin{aligned} \langle J_s^- J_s^- O \rangle_{\text{even}, \mathbf{h}} &= k_1^{s-1} k_2^{s-1} \frac{(E + (2s-1)k_3)}{E^{2s}} \langle 12 \rangle^{2s} \\ \langle J_s^+ J_s^+ O \rangle_{\text{even}, \mathbf{h}} &= k_1^{s-1} k_2^{s-1} \frac{(E + (2s-1)k_3)}{E^{2s}} \langle \bar{1}\bar{2} \rangle^{2s} \\ \langle J_s^+ J_s^- O \rangle_{\text{even}, \mathbf{h}} &= \langle J_s^- J_s^+ O \rangle_{\text{even}, \mathbf{h}} = 0. \end{aligned} \quad (4.9)$$

The spinor helicity components of the odd part of the correlator can be computed using (4.7). One can then derive the following double copy relation for a general correlator of the kind $\langle J_s J_s O_3 \rangle$ that satisfies the homogeneous conformal Ward identity:

$$\begin{aligned} &\langle J_s J_s O \rangle_{\text{even}, \mathbf{h}} + \langle J_s J_s O \rangle_{\text{odd}, \mathbf{h}} \\ &= \frac{k_1 k_2 (E + (2s-1)k_3)}{(E + (2s'-1)k_3)(E + (2s''-1)k_3)} (\langle J_{s'} J_{s'} O \rangle_{\text{even}, \mathbf{h}} + \langle J_{s'} J_{s'} O \rangle_{\text{odd}, \mathbf{h}}) \\ &\quad \times (\langle J_{s''} J_{s''} O \rangle_{\text{even}, \mathbf{h}} + \langle J_{s''} J_{s''} O \rangle_{\text{odd}, \mathbf{h}}) \\ &\implies \langle J_s J_s O \rangle_{\mathbf{h}} = \frac{k_1 k_2 (E + (2s-1)k_3)}{(E + (2s'-1)k_3)(E + (2s''-1)k_3)} \langle J_{s'} J_{s'} O \rangle_{\mathbf{h}} \langle J_{s''} J_{s''} O \rangle_{\mathbf{h}} \end{aligned}$$

where $s' + s'' = s$.

We will now come to more complicated correlators such as $\langle JJJ \rangle$ and $\langle TTT \rangle$ whose conformal Ward identities have a non-homogeneous term and show more interesting double copy relations.

4.3 Double copy relation for $\langle JJJ \rangle$ and $\langle TTT \rangle$

The double copy relation between $\langle JJJ \rangle$ and $\langle TTT \rangle$ is more subtle than those for correlators with a scalar operator insertion. Unlike $\langle TTO_3 \rangle$ or $\langle JJO_3 \rangle$ these correlators have a non-homogeneous term as well and we will see that the double copy structures map homogeneous terms to homogeneous terms and non-homogeneous terms get mapped to non-homogeneous terms.

Homogeneous terms. The following double copy structure was noticed in [33] for the homogeneous term in the even part of $\langle TTT \rangle$:

$$\langle TTT \rangle_{\text{even}, \mathbf{h}} = k_1 k_2 k_3 \langle JJJ \rangle_{\text{even}, \mathbf{h}} \langle JJJ \rangle_{\text{even}, \mathbf{h}} \quad (4.10)$$

From the explicit expressions for the correlators in (3.12), (3.14), and (3.19), we notice:

$$\langle TTT \rangle_{\text{odd}, \mathbf{h}} = k_1 k_2 k_3 \langle JJJ \rangle_{\text{odd}, \mathbf{h}} \langle JJJ \rangle_{\text{even}, \mathbf{h}} \quad (4.11)$$

We also have the remarkable relation that the parity-even part of the homogeneous term is given by the square of the odd part of the homogeneous term in $\langle JJJ \rangle$:

$$\langle TTT \rangle_{\text{even}, \mathbf{h}} = k_1 k_2 k_3 \langle JJJ \rangle_{\text{odd}, \mathbf{h}} \langle JJJ \rangle_{\text{odd}, \mathbf{h}} \quad (4.12)$$

Combining these relations we obtain the following double copy relation for the complete homogeneous term of the $\langle TTT \rangle$ correlator:

$$\begin{aligned} \langle TTT \rangle_{\text{even}, \mathbf{h}} + \langle TTT \rangle_{\text{odd}, \mathbf{h}} &= k_1 k_2 k_3 (\langle JJJ \rangle_{\text{even}, \mathbf{h}} + \langle JJJ \rangle_{\text{odd}, \mathbf{h}})^2 \\ \implies \langle TTT \rangle_{\mathbf{h}} &= k_1 k_2 k_3 \langle JJJ \rangle_{\mathbf{h}}^2 \end{aligned} \quad (4.13)$$

Non-homogeneous terms. From (3.12) and (3.15) we know that $\langle JJJ \rangle_{\text{even}}$ and $\langle TTT \rangle_{\text{even}}$ also have non-trivial non-homogeneous parts between which there exists the following double copy relation:

$$\langle TTT \rangle_{\text{even}, \mathbf{nh}} = (E^3 - E(k_1 k_2 + k_2 k_3 + k_1 k_3) - k_1 k_2 k_3) \langle JJJ \rangle_{\text{even}, \mathbf{nh}}^2 \quad (4.14)$$

This relation is independent of the double copy of the homogeneous part as the pre-factor is different. The non-homogeneous parts of $\langle TTT \rangle_{\text{odd}}$ and $\langle JJJ \rangle_{\text{odd}}$ are trivial as they are contact terms.

4.4 Double copy structure for higher spin correlators

We will now discuss the double copy structures in higher spin correlators. This is most easily done using the spinor-helicity variables. The parity-even part of the homogeneous part of $\langle J_s J_s J_s \rangle$ is given by [71]:

$$\langle J_s J_s J_s \rangle_{\text{even}, \mathbf{h}} = \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle 12 \rangle^s \langle 23 \rangle^s \langle 31 \rangle^s \quad (4.15)$$

As noted in [71], the odd part of the above correlator is given by the same expression up to an overall factor of i :

$$\langle J_s J_s J_s \rangle_{\text{odd}, \mathbf{h}} = i \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle 12 \rangle^s \langle 23 \rangle^s \langle 31 \rangle^s \quad (4.16)$$

From this, we have the following double copy expression for the homogeneous part of the higher spin correlator $\langle J_s J_s J_s \rangle$:

$$\langle J_s J_s J_s \rangle_{\mathbf{h}} = k_1 k_2 k_3 (\langle J^{s'} J^{s'} J^{s'} \rangle \langle J^{s''} J^{s''} J^{s''} \rangle) \quad (4.17)$$

such that $s' + s'' = s$.

4.5 Spin s current correlator as s copies of the spin one current correlator

In this sub-section we note that we can write correlators of the form $\langle J_s J_s O_3 \rangle$ and $\langle J_s J_s J_s \rangle$ as s copies of correlators of the spin-one current. Using the double copy relations in (4.2) recursively we notice that:

$$\langle J_s J_s O \rangle_{\mathbf{h}} = (k_1 k_2)^{s-1} \frac{E + (2s-1)k_3}{(E + k_3)^s} (\langle JJO \rangle_{\mathbf{h}})^s \quad (4.18)$$

Similarly using the double copy relations in (4.17) recursively we notice that:

$$\langle J_s J_s J_s \rangle_{\mathbf{h}} = (k_1 k_2 k_3)^{s-1} (\langle JJJ \rangle_{\mathbf{h}})^s \quad (4.19)$$

5 CFT correlators from dS_4 Feynman diagrams and the flat space limit

In this section, we relate CFT correlators discussed in the previous section to the tree-level amplitude calculated using Feynman diagrams in dS_4 . We also relate CFT correlators to flat space scattering amplitudes.

5.1 Amplitudes

Here we will study 3-point flat-space scattering amplitudes in general parity-violating theories of gravitons, gluons and massless scalars in 4d. These are calculated straightforwardly from a Lagrangian whose cubic vertices will contribute to the 3-point amplitudes. We will use the notation $\mathcal{M}$ to denote the flat space amplitude without energy conservation. This is a useful quantity because $E = k_1 + k_2 + k_3 \neq 0$ in dS_4 and the dS_4 vertex is obtained from $\mathcal{M}$ by multiplying with an overall conformal time integral factor. This also matches the CFT correlators that we computed in section 3. We also take flat space limit of $\mathcal{M}$ and resultant scattering amplitude will be denoted by $\mathcal{A}$. More precisely

$$\mathcal{A} = \lim_{E \rightarrow 0} \mathcal{M}. \quad (5.1)$$

We also express $\mathcal{M}$ in both 4d and 3d notations. See section 2 and the discussion below (2.1) for how to express 4d amplitude in terms of 3d notation. The 3d notation is particularly useful while comparing with the CFT correlators.

We will first describe the gauge theory action, the gravity action and the gravity-gluon interactions that we consider.

Gauge theory action. The gauge theory action that we consider is:

$$S_A = S_{\text{EM}} + S_{\text{even}}^A + S_{\text{odd}}^A \quad (5.2)$$

where S_{EM} is the electromagnetic action and S_{even}^A and S_{odd}^A are gauge invariant parity-preserving and parity-violating actions respectively, given by:

$$S_{\text{even}}^{\text{EM}} = -\frac{1}{4} \int d^4x \sqrt{g} F^2 \quad (5.3)$$

$$S_{\text{even}}^A = \int d^4x \sqrt{g} (\alpha_1^A F^3 + \alpha_2^A \phi F^2) \quad (5.4)$$

$$S_{\text{odd}}^A = \int (\beta_1^A F_{\mu\nu} F_{\rho\sigma} + \beta_2^A F_{\mu\nu} F_{\rho}{}^{\tau} F_{\sigma\tau} + \beta_3^A \phi F_{\mu\nu} F_{\rho\sigma}) dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} \wedge dx^{\sigma} \quad (5.5)$$

where

$$F^3 = F_\alpha{}^\beta F_\beta{}^\gamma F_\gamma{}^\alpha \quad (5.6)$$

Depending on the background, $g_{\mu\nu}$ will be the four dimensional Minkowski or de Sitter metric.

Gravity action. The gravity action that we consider is:

$$S_g = S_{\text{EH}} + S_{\text{even}}^g + S_{\text{odd}}^g \quad (5.7)$$

where S_{EH} is the Einstein-Hilbert action and S_{even}^g and S_{odd}^g are parity-preserving and parity-violating actions respectively, given by:

$$S_{\text{EH}} = \frac{1}{16\pi G} \int d^4x \sqrt{g} (R + \Lambda) \quad (5.8)$$

$$S_{\text{even}}^g = \int d^4x \sqrt{g} (\alpha_1^g W^2 + \alpha_2^g W^3 + \alpha_3^g \phi W^2) \quad (5.9)$$

$$S_{\text{odd}}^g = \int (\beta_1^g W_{\mu\nu\rho\sigma} W_{\alpha\beta}{}^{\rho\sigma} + \beta_2^g W_{\rho\sigma\alpha\beta} W^{\sigma\gamma}{}_{\mu\tau} W_\gamma{}^{\rho\tau}{}_\nu + \beta_3^g \phi W_{\mu\nu\rho\sigma} W_{\alpha\beta}{}^{\rho\sigma}) dx^\mu \wedge dx^\nu \wedge dx^\alpha \wedge dx^\beta \quad (5.10)$$

where

$$W^2 = W_{\mu\nu\rho\sigma} W^{\mu\nu\rho\sigma}, \quad W^3 = W_{\mu\nu\rho\sigma} W^{\rho\sigma\alpha\beta} W_{\alpha\beta\mu\nu}, \quad g = \det(g_{\mu\nu}) \quad (5.11)$$

and $W_{\mu\nu\rho\sigma}$ is the Weyl tensor.

5.1.1 Gauge amplitudes

Gluon-gluon-scalar amplitudes. Let us first compute the contribution to the gluon-gluon-scalar amplitudes due to the interactions corresponding to the coupling α_2^A and β_3^A in (5.4) and (5.5). From the Lagrangian, we first compute the amplitude without energy conservation ($\mathcal{M}$) and express it in both 4d and 3d notation (see (2.1))

$$\begin{aligned} \mathcal{M}_{\phi F^2} &= (k_1 \cdot z_2)(k_2 \cdot z_1) - (k_1 \cdot k_2)(z_1 \cdot z_2) \\ &= (\vec{k}_1 \cdot \vec{z}_2)(\vec{k}_2 \cdot \vec{z}_1) + \frac{E}{2}(E - 2k_3)(\vec{z}_1 \cdot \vec{z}_2) \\ \mathcal{M}_{\phi F\tilde{F}} &= \epsilon^{z_1 k_1 z_2 k_2} = -\epsilon^{z_1 z_2 k_1} k_2 + \epsilon^{z_1 z_2 k_2} k_1 \end{aligned} \quad (5.12)$$

To get actual flat space scattering amplitudes we have to impose energy conservation, i.e. $E \rightarrow 0$. This gives:

$$\begin{aligned} \mathcal{A}_{\phi F^2} &= \lim_{E \rightarrow 0} \mathcal{M}_{\phi F^2} = (\vec{k}_1 \cdot \vec{z}_2)(\vec{k}_2 \cdot \vec{z}_1), \\ \mathcal{A}_{\phi F\tilde{F}} &= \lim_{E \rightarrow 0} \mathcal{M}_{\phi F\tilde{F}} = -\epsilon^{z_1 z_2 k_1} k_2 + \epsilon^{z_1 z_2 k_2} k_1. \end{aligned} \quad (5.13)$$

Gluon-gluon-gluon amplitudes. Let us now compute the contribution to the gluon-gluon-gluon amplitudes due to interactions corresponding to couplings α_1^A and β_2^A in (5.4) and (5.5). The parity-even amplitude is given by:

$$\begin{aligned}\mathcal{M}_{F^3} &= \frac{1}{2} z_{[1\mu} k_{1\nu]} z_{[2\nu} k_{2\rho]} z_{[3\rho} k_{3\mu]} \\ &= (\vec{k}_2 \cdot \vec{z}_1)(\vec{k}_3 \cdot \vec{z}_2)(\vec{k}_1 \cdot \vec{z}_3) + \frac{E}{2} \left(k_1(\vec{k}_2 \cdot \vec{z}_1)(\vec{z}_2 \cdot \vec{z}_3) + \text{cyclic perm.} \right)\end{aligned}\quad (5.14)$$

The parity-odd amplitude is given by:

$$\begin{aligned}\mathcal{M}_{F^2\tilde{F}} &= z_{[2\alpha} k_{2\tau]} z_{[3\tau} k_{3\beta]} \epsilon^{\alpha\beta z_1 k_1} + \text{cyclic perm.} \\ &= \left[-(\vec{k}_1 \cdot \vec{z}_3) \left(\epsilon^{k_3 z_1 z_2} k_1 - \epsilon^{k_1 z_1 z_2} k_3 \right) + (\vec{k}_3 \cdot \vec{z}_2) \left(\epsilon^{k_1 z_1 z_3} k_2 - \epsilon^{k_2 z_1 z_3} k_1 \right) \right. \\ &\quad \left. - (\vec{z}_2 \cdot \vec{z}_3) \epsilon^{k_1 k_2 z_1} E + \frac{k_1}{2} \epsilon^{z_1 z_2 z_3} E(E - 2k_1) \right] + \text{cyclic perm.}\end{aligned}\quad (5.15)$$

The flat space scattering amplitudes is obtained by taking the $E \rightarrow 0$ limit. This gives:

$$\begin{aligned}\mathcal{A}_{F^3} &= \lim_{E \rightarrow 0} \mathcal{M}_{F^3} = (\vec{k}_2 \cdot \vec{z}_1)(\vec{k}_3 \cdot \vec{z}_2)(\vec{k}_1 \cdot \vec{z}_3) \\ \mathcal{A}_{F^2\tilde{F}} &= \left[-(\vec{k}_1 \cdot \vec{z}_3) \left(\epsilon^{k_3 z_1 z_2} k_1 - \epsilon^{k_1 z_1 z_2} k_3 \right) + (\vec{k}_3 \cdot \vec{z}_2) \left(\epsilon^{k_1 z_1 z_3} k_2 - \epsilon^{k_2 z_1 z_3} k_1 \right) \right] + \text{cyclic perm.}\end{aligned}\quad (5.16)$$

We also have contributions to the 3-gluon amplitude from F^2 and $F\tilde{F}$. To obtain these we first calculate:

$$\begin{aligned}\mathcal{M}_{\text{YM}} &= (k_2 \cdot z_1)(z_2 \cdot z_3) + \text{cyclic perm.} = (\vec{k}_2 \cdot \vec{z}_1)(\vec{z}_2 \cdot \vec{z}_3) + \text{cyclic perm.} \\ \mathcal{M}_{F\tilde{F}} &= \epsilon^{k_1 z_1 z_2 z_3} + \text{cyclic perm.} = E \epsilon^{z_1 z_2 z_3}\end{aligned}\quad (5.17)$$

The actual flat space scattering amplitudes are then given by:

$$\begin{aligned}\mathcal{A}_{\text{YM}} &= (\vec{k}_2 \cdot \vec{z}_1)(\vec{z}_2 \cdot \vec{z}_3) + \text{cyclic perm.} \\ \mathcal{A}_{F\tilde{F}} &= 0\end{aligned}\quad (5.18)$$

5.1.2 Gravity amplitudes

Working with a traceless, transverse perturbation around the flat metric, we obtain the following for the Weyl tensor to first order in perturbation:

$$W_{\mu\nu\rho\sigma} = k_{[\mu} z_{\nu]} k_{[\rho} z_{\sigma]}\quad (5.19)$$

where k^μ and z^μ are defined in (2.1).

Graviton-graviton-scalar amplitude. Let us consider now the parity-odd graviton-graviton-scalar amplitude due to the interaction corresponding to the coupling β_3^g in (5.10).

From the Lagrangian we have:

$$\begin{aligned}
 \mathcal{M}_{\phi W \tilde{W}} &= \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} W_{(1)\mu\nu\alpha\beta} W_{(2)}^{\mu\nu}{}_{\gamma\delta} \\
 &= \epsilon(z_1 k_1 z_2 k_2) \left((k_2 \cdot z_1)(k_1 \cdot z_2) + \frac{E}{2}(E - 2k_3)z_1 \cdot z_2 \right) \\
 &= \left((\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + \frac{E}{2}(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2 \right) \left(-k_1 \epsilon^{z_1 z_2 k_2} + k_2 \epsilon^{z_1 z_2 k_1} \right) \quad (5.20)
 \end{aligned}$$

The flat space amplitude is given by:

$$\mathcal{A}_{\phi W \tilde{W}} = (\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) \left(-k_1 \epsilon^{z_1 z_2 k_2} + k_2 \epsilon^{z_1 z_2 k_1} \right) \quad (5.21)$$

Similarly, one can compute the parity-even graviton-graviton-scalar amplitude due to the interaction corresponding to the coupling α_3^g in (5.9) from $\mathcal{M}_{\phi W^2}$:

$$\begin{aligned}
 \mathcal{M}_{\phi W^2} &= \frac{1}{4} W_{(1)\mu\nu\rho\sigma} W_{(2)}^{\mu\nu\rho\sigma} \\
 &= \frac{1}{4} (2(z_1 \cdot k_2)(z_2 \cdot k_1) + E(E - 2k_3)z_1 \cdot z_2)^2 \\
 &= \left((\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + \frac{E}{2}(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2 \right)^2 \quad (5.22)
 \end{aligned}$$

The flat space amplitude is obtained by taking the $E \rightarrow 0$ limit:

$$\mathcal{A}_{\phi W^2} = \left((\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) \right)^2 \quad (5.23)$$

Graviton-graviton-graviton amplitude. Here we calculate graviton-graviton-graviton amplitude from interactions corresponding to couplings α_2^g and β_2^g in (5.9) and (5.10). The parity-even amplitude is given by:

$$\begin{aligned}
 \mathcal{M}_{W^3} &= F_1(1, 2, 3)F_1(2, 3, 1)F_1(3, 1, 2) \\
 &= F_2(1, 2, 3)F_2(2, 3, 1)F_2(3, 1, 2) \quad (5.24)
 \end{aligned}$$

where

$$\begin{aligned}
 F_1(i, j, l) &= [-(z_i \cdot z_j)k_i \cdot k_j + (z_i \cdot k_j)(z_j \cdot k_i)] \\
 F_2(i, j, l) &= \left[(\vec{z}_i \cdot \vec{k}_j)(\vec{z}_j \cdot \vec{k}_i) + \frac{1}{2}(\vec{z}_i \cdot \vec{z}_j)E(E - 2k_l) \right] \quad (5.25)
 \end{aligned}$$

The parity-odd contribution to the amplitude is given by:

$$\begin{aligned}
 \mathcal{M}_{W^2\tilde{W}} &= \frac{1}{2} \left(z_{[1\mu} k_{1\nu]} z_{[2\nu} k_{2\rho]} z_{[3\rho} k_{3\mu]} [z_{[2\alpha} k_{2\tau]} z_{[3\tau} k_{3\beta]} \epsilon^{\alpha\beta z_1 k_1} \right) \\
 &= \frac{1}{2} [2(z_1 \cdot k_2 z_2 \cdot k_3 z_3 \cdot k_1) + E((z_1 \cdot z_2 z_3 \cdot k_1)k_3 + \text{cyclic})] \\
 &\quad [\{(z_3 \cdot k_2)\epsilon(z_2 k_3 z_1 k_1) - (k_2 \cdot k_3)\epsilon(z_1 z_2 z_3 k_1) - (z_2 \cdot z_3)\epsilon(z_1 k_1 k_2 k_3) \\
 &\quad + (z_2 \cdot k_3)\epsilon(k_2 z_3 z_1 k_1)\} + \text{cyclic}] \\
 &= \left[(\vec{z}_1 \cdot \vec{k}_2 \vec{z}_2 \cdot \vec{k}_3 \vec{z}_3 \cdot \vec{k}_1) + \frac{E}{2} ((\vec{z}_1 \cdot \vec{z}_2 \vec{z}_3 \cdot \vec{k}_1)k_3 + \text{cyclic}) \right] \\
 &\quad \left[(\vec{k}_3 \cdot \vec{z}_2) (\epsilon^{k_3 z_1 z_3} k_1 + \epsilon^{k_1 z_1 z_3} (E - k_3)) + (\vec{k}_1 \cdot \vec{z}_3) (\epsilon^{k_2 z_1 z_2} k_1 + \epsilon^{k_1 z_1 z_2} (E - k_2)) \right. \\
 &\quad \left. - E(\vec{z}_2 \cdot \vec{z}_3) \epsilon^{k_1 k_2 z_1} + \frac{1}{2} k_1 E (E - 2k_1) \epsilon^{z_1 z_2 z_3} + \text{cyclic} \right] \quad (5.26)
 \end{aligned}$$

The flat space amplitude, as before, is obtained by taking the $E \rightarrow 0$ limit:

$$\begin{aligned}
 \mathcal{A}_{W^3} &= (\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) \times (\vec{z}_2 \cdot \vec{k}_3)(\vec{z}_3 \cdot \vec{k}_2) \times (\vec{z}_3 \cdot \vec{k}_1)(\vec{z}_1 \cdot \vec{k}_3) \\
 \mathcal{A}_{W^2\tilde{W}} &= \left[(\vec{z}_1 \cdot \vec{k}_2 \vec{z}_2 \cdot \vec{k}_3 \vec{z}_3 \cdot \vec{k}_1) \right. \\
 &\quad \left[(\vec{k}_3 \cdot \vec{z}_2) (\epsilon^{k_3 z_1 z_3} k_1 - \epsilon^{k_1 z_1 z_3} k_3) - (\vec{k}_1 \cdot \vec{z}_3) (\epsilon^{k_1 z_1 z_2} k_2 - \epsilon^{k_2 z_1 z_2} k_1) + \text{cyclic} \right] \quad (5.27)
 \end{aligned}$$

The EH term in (5.8) gives

$$\mathcal{M}_{\text{EH}} = ((k_2 \cdot z_1)(z_2 \cdot z_3) + \text{cyclic perm.})^2 = ((\vec{k}_2 \cdot \vec{z}_1)(\vec{z}_2 \cdot \vec{z}_3) + \text{cyclic})^2. \quad (5.28)$$

In the flat space limit this gives

$$\mathcal{A}_{\text{EH}} = ((\vec{k}_2 \cdot \vec{z}_1)(\vec{z}_2 \cdot \vec{z}_3) + \text{cyclic})^2. \quad (5.29)$$

5.2 Double copy structure of parity-violating amplitudes

In this section, we see how the double copy structure arises directly at the level of 3-point scattering amplitude when parity-violating terms are taken into account. The 3-point amplitudes involved may be written down directly on symmetry grounds, or computed from the action given above. The double copy relations of amplitudes also follows from that for CFT correlators that we saw previously, but here we show that it can be demonstrated directly in 4d flat space.

Scalar-graviton-graviton. From the momentum space expressions for $\mathcal{M}_{\phi F^2}$ and $\mathcal{M}_{\phi F\tilde{F}}$ in (5.12) and the expression for $\mathcal{M}_{\phi W\tilde{W}}$ and $\mathcal{M}_{\phi W^2}$ in (5.20) and in (5.22) we obtain:

$$\begin{aligned}
 \mathcal{M}_{\phi W\tilde{W}} &= \mathcal{M}_{\phi F^2} \mathcal{M}_{\phi F\tilde{F}}, \quad \mathcal{M}_{\phi W^2} = (\mathcal{M}_{\phi F^2})^2 = \mathcal{M}_{\phi F\tilde{F}} \mathcal{M}_{\phi F\tilde{F}} \\
 \mathcal{M}_{\phi W^2} + \mathcal{M}_{\phi W\tilde{W}} &= (\mathcal{M}_{\phi F^2} + \mathcal{M}_{\phi F\tilde{F}})^2 \quad (5.30)
 \end{aligned}$$

In the flat space limit when $E \rightarrow 0$ one has:

$$\begin{aligned}\mathcal{A}_{\phi W \tilde{W}} &= \mathcal{A}_{\phi F^2} \mathcal{A}_{\phi F \tilde{F}}, \quad \mathcal{M}_{\phi W^2} = (\mathcal{A}_{\phi F^2})^2 = \mathcal{A}_{\phi F \tilde{F}} \mathcal{M}_{\phi F \tilde{F}} \\ \mathcal{A}_{\phi W^2} + \mathcal{A}_{\phi W \tilde{W}} &= (\mathcal{A}_{\phi F^2} + \mathcal{A}_{\phi F \tilde{F}})^2\end{aligned}\tag{5.31}$$

Graviton-graviton-graviton. From the momentum space expressions for $\mathcal{M}_{F^3}$ and $\mathcal{M}_{F^2 \tilde{F}}$ in (5.14) and (5.15) and the expression for $\mathcal{M}_{W^2 \tilde{W}}$ in (5.26) one can verify that:

$$\mathcal{M}_{W^2 \tilde{W}} = \mathcal{M}_{F^3} \mathcal{M}_{F^2 \tilde{F}}\tag{5.32}$$

In the $E \rightarrow 0$ limit one has:

$$\mathcal{A}_{W^2 \tilde{W}} = \mathcal{A}_{F^3} \mathcal{A}_{F^2 \tilde{F}}\tag{5.33}$$

We also have the following relation:

$$\mathcal{M}_{W^3} = \mathcal{M}_{F^3}^2\tag{5.34}$$

The details of how one obtains this relation in momentum space are given in appendix (C.1). Thus we notice that the following double copy structure

$$\mathcal{A}_{W^3} = \mathcal{A}_{F^3}^2\tag{5.35}$$

extends beyond the $E \rightarrow 0$ limit. We also notice using the momentum space expressions in (5.15) the following:

$$(\mathcal{M}_{F^3})^2 = (\mathcal{M}_{F^2 \tilde{F}})^2\tag{5.36}$$

Similarly one has the following double copy relation:

$$\begin{aligned}\mathcal{M}_{W^3} + \mathcal{M}_{W^2 \tilde{W}} &= (\mathcal{M}_{F^3} + \mathcal{M}_{F^2 \tilde{F}})^2 \\ \mathcal{A}_{W^3} + \mathcal{A}_{W^2 \tilde{W}} &= (\mathcal{A}_{F^3} + \mathcal{A}_{F^2 \tilde{F}})^2\end{aligned}\tag{5.37}$$

and, also, the well-known relation between Einstein gravity and Yang-Mills amplitudes:

$$\mathcal{M}_{\text{EG}} = (\mathcal{M}_{\text{YM}})^2\tag{5.38}$$

5.3 CFT correlators from dS_4

In this section we will compute tree level dS_4 cosmological correlators using the method of [56] and also including the relevant parity-odd 3-point vertices in the Lagrangian. The calculation with parity-even vertices was done in appendix A of [33].

The idea, due to Maldacena and Pimentel [56], that will be central to our analysis in this section is that certain cosmological correlators in de Sitter can be constructed directly from the corresponding flat-space amplitudes by dressing with conformal time integrals arising from using conformally covariant transformation properties of fields in dS_4 . These

Lorentzian dS_4 correlators also compute boundary (Euclidean) CFT_3 correlators, thereby establishing a relationship between these three quantities.⁶

We will work in (Lorentzian) dS_4 with the metric:

$$ds^2 = \frac{1}{\eta^2}(-d\eta^2 + dx_i^2) \quad (5.39)$$

For calculating dS_4 correlators perturbatively, we first look at the on-shell wavefunctions for (linearised) free fields, which can be massless scalars, gauge or gravitational perturbations. These kind of fields will suffice to calculate the correlators we will be interested in. As is well known, for the scalar we have the Bunch Davies mode function $\phi \sim (1 - ik\eta)\exp(ik\eta)$, the gauge field solution is a plane wave just like in flat space $A_\mu \sim z_\mu \exp(ik\eta)$ and the linearised gravitational perturbation is given by $\gamma_{\mu\nu} \sim z_\mu z_\nu (1 - ik\eta)\exp(ik\eta)$. It was noted in [56], that this results in the Weyl tensors for the linearised gravity perturbations about dS and flat backgrounds being conformally related:

$$W_{(dS)\nu\rho\sigma}^\mu(\exp(ik\eta)(1 - ik\eta)z_\mu z_\nu) = (-ik\eta)W_{(flat)\nu\rho\sigma}^\mu(\exp(ik\eta)z_\mu z_\nu), \quad (5.40)$$

whereas the gauge field strength is the same in both backgrounds. This means that the dS correlators of interest to us are the same as corresponding flat space amplitudes without energy conservation ($\mathcal{M}$) upto conformal time integrals which are easily evaluated. In this section, we will calculate the contribution to 3-point functions from parity-violating interaction terms in the Lagrangian of the form $F\tilde{F}\phi$, $W\tilde{W}\phi$, $F^2\tilde{F}$ and $W^2\tilde{W}$. We will take all parity-odd two-point functions to be vanishing as this is the case for the dS_4 actions considered here.

We will find that our tree-level dS_4 computations will generate the different parts of the corresponding CFT_3 correlator (both parity-even and odd). Therefore, this perturbative approach provides a simple method of fixing the form of CFT correlators without taking recourse to solving conformal Ward identities.

5.3.1 $\langle JJO_3 \rangle$

The term in the action which contributes to the odd part of $\langle JJO_3 \rangle$ is given by:

$$\int \phi F_{\mu\nu} F_{\rho\sigma} dx^\mu \wedge dx^\nu \wedge dx^\rho \wedge dx^\sigma \quad (5.41)$$

The tree-level dS_4 correlator corresponding to this interaction is given by:

$$\langle JJO_3 \rangle_{\text{odd}} = \text{Im} \left[\int_{-\infty}^0 d\eta (1 - ik_3\eta) e^{i\eta E} \right] \mathcal{M}_{\phi F \tilde{F}} = \frac{E + k_3}{E^2} \mathcal{M}_{\phi F \tilde{F}} \quad (5.42)$$

Substituting for $\mathcal{M}_{\phi F \tilde{F}}$ from (5.12) we see that this matches the expression for the homogeneous part of the correlator in (3.3). In the flat space limit we get:

$$\lim_{E \rightarrow 0} \langle JJO_3 \rangle_{\text{odd}} = \frac{k_3}{E^2} \mathcal{A}_{\phi F \tilde{F}} \quad (5.43)$$

where $\mathcal{A}_{\phi F \tilde{F}}$ is given in (5.13).

⁶It is of course true that flat space amplitudes can be generated by an appropriate limit of CFT correlators, the non-trivial part is that we can do the converse at least for certain correlators.

The corresponding parity-even correlator is given by [20, 29, 33]:

$$\langle JJO_3 \rangle_{\text{even}} = \frac{(E + k_3)}{E^2} \mathcal{M}_{\phi F^2} \quad (5.44)$$

This matches the expression for the homogeneous part of the correlator in (3.2) if we use the expression for $\mathcal{M}_{\phi F^2}$ in (5.12). In the flat space limit the correlator takes the form:

$$\lim_{E \rightarrow 0} \langle JJO_3 \rangle_{\text{even}} = \frac{k_3}{E^2} \mathcal{A}_{\phi F^2} \quad (5.45)$$

where $\mathcal{A}_{\phi F^2}$ is given in (5.13).

5.3.2 $\langle TTO_3 \rangle$

Let us now consider the parity-odd part of the correlator $\langle TTO_3 \rangle$. The only contribution to $\langle TTO_3 \rangle_{\text{odd}}$ comes from the term:

$$\int \phi W_{\alpha\beta\mu\nu} W^{\alpha\beta}_{\rho\sigma} dx^\mu \wedge dx^\nu \wedge dx^\rho \wedge dx^\sigma \quad (5.46)$$

The corresponding dS_4 correlator is given by:⁷

$$\begin{aligned} \langle TTO_3 \rangle_{\text{odd}} &= k_1 k_2 \text{Im} \left[\int_{-\infty}^0 d\eta \eta^2 (1 - ik_3 \eta) e^{i\eta E} \right] \mathcal{M}_{\phi W \tilde{W}} \\ &= \frac{k_1 k_2 (E + 3k_3)}{E^4} \mathcal{M}_{\phi W \tilde{W}} \end{aligned} \quad (5.47)$$

We can easily check that with the expression for $\mathcal{M}_{\phi W \tilde{W}}$ in (5.20), this matches the homogeneous part of the correlator in (3.5). In the flat space limit we get:

$$\lim_{E \rightarrow 0} \langle TTO_3 \rangle_{\text{odd}} = \frac{3k_1 k_2 k_3}{E^4} \mathcal{A}_{\phi W \tilde{W}} \quad (5.48)$$

where $\mathcal{A}_{\phi W \tilde{W}}$ is given in (5.21).

The parity-even part of the corresponding correlator is given by [20, 29, 33]:

$$\langle TTO \rangle_{\text{even}} = k_1 k_2 \frac{E + 3k_3}{E^4} \mathcal{M}_{\phi W^2} \quad (5.49)$$

which matches (3.4) if we use the expression for $\mathcal{M}_{\phi W^2}$ given in (5.22).

In the flat space limit we get:

$$\lim_{E \rightarrow 0} \langle TTO \rangle_{\text{even}} = \frac{3k_1 k_2 k_3}{E^4} \mathcal{A}_{\phi W^2} \quad (5.50)$$

where $\mathcal{A}_{\phi W^2}$ is given in (5.23).

---

⁷Since the dS_4 correlators match CFT_3 correlators we will use the notation $\langle JJO_3 \rangle$ instead of $\langle \gamma\gamma\phi \rangle$. We will continue to use similar CFT notations for all dS_4 correlators in this section.

5.3.3 $\langle JJJ \rangle$

Let us now compute the odd part of $\langle JJJ \rangle$. The contribution to $\langle JJJ \rangle_{\text{odd}}$ comes from the following terms:

$$\int F_{\mu\tau} F_{\nu}{}^{\tau} F_{\rho\sigma} dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} \wedge dx^{\sigma}, \quad \int F_{\mu\nu} F_{\rho\sigma} dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} \wedge dx^{\sigma} \quad (5.51)$$

The parts of the tree-level correlator corresponding to these interactions are given by:

$$\begin{aligned} \mathcal{C}_{F^2\tilde{F}} &= \text{Im} \left[\int_{-\infty}^0 d\eta \, \eta^2 e^{i\eta E} \right] \mathcal{M}_{F^2\tilde{F}} \\ \mathcal{C}_{F\tilde{F}} &= \text{Im} \left[\int_{-\infty}^0 d\eta \, e^{i\eta E} \right] \mathcal{M}_{F\tilde{F}} \end{aligned} \quad (5.52)$$

Combining the two, we get the total correlator to be:

$$\begin{aligned} \langle JJJ \rangle_{\text{odd}} &= c_{JJJ}^{\text{odd}} \mathcal{C}_{F^2\tilde{F}} + c_{JJ}^{\text{odd}} \mathcal{C}_{F\tilde{F}} \\ &= c_{JJJ}^{\text{odd}} \frac{\mathcal{M}_{F^2\tilde{F}}}{E^3} + c_{JJ}^{\text{odd}} \frac{\mathcal{M}_{F\tilde{F}}}{E}. \end{aligned} \quad (5.53)$$

One can check using the explicit expression for $\mathcal{M}_{F^2\tilde{F}}$ in (5.15) that the first term in the above equation corresponds to the homogeneous term in (3.14). Similarly, using the explicit expression for $\mathcal{M}_{F\tilde{F}}$ in (5.17) one can see that the second term in the above equation corresponds to the non-homogeneous term in (3.14). As noted earlier, the term proportional to c_{JJ}^{odd} in (5.53) is a contact term. In the flat space limit we obtain:

$$\lim_{E \rightarrow 0} \langle JJJ \rangle_{\text{odd}} = c_{JJJ}^{\text{odd}} \frac{\mathcal{A}_{F^2\tilde{F}}}{E^3} + c_{JJ}^{\text{odd}} \frac{\mathcal{A}_{F\tilde{F}}}{E} = c_{JJJ}^{\text{odd}} \frac{\mathcal{A}_{F^2\tilde{F}}}{E^3} \quad (5.54)$$

where $\mathcal{A}_{F\tilde{F}} = 0$ as shown in (5.18) and the expression for $\mathcal{A}_{F^2\tilde{F}}$ is as in (5.16). The even part of the correlator is given by [20, 27, 33]:

$$\langle JJJ \rangle_{\text{even}} = \frac{c_{JJJ}^{\text{even}}}{E^3} \mathcal{M}_{F^3} - \frac{2c_{JJ}^{\text{even}}}{E} \mathcal{M}_{\text{YM}}. \quad (5.55)$$

From the explicit expression for $\mathcal{M}_{F^3}$ in (5.14) we see that the first term in the above equation corresponds to the homogeneous term in (3.12). Using the expression for $\mathcal{M}_{\text{YM}}$ in (5.17) one can see that the second term in the above equation corresponds to the non-homogeneous term in (3.12). In the flat space limit we obtain:

$$\lim_{E \rightarrow 0} \langle JJJ \rangle_{\text{even}} = \frac{c_{JJJ}^{\text{even}}}{E^3} \mathcal{A}_{F^3} - \frac{2c_{JJ}^{\text{even}}}{E} \mathcal{A}_{\text{YM}} \quad (5.56)$$

where the expression for $\mathcal{A}_{F^3}$ and $\mathcal{A}_{\text{YM}}$ are as in (5.16) and (5.18) respectively.

5.3.4 $\langle TTT \rangle$

The contribution to $\langle TTT \rangle_{\text{odd}}$ comes from the following term in the action:

$$\int W_{\rho\sigma\alpha\beta} W_{\mu\tau}^{\sigma\gamma} W_{\gamma}{}^{\rho\tau}{}_{\nu} dx^{\mu} \wedge dx^{\nu} \wedge dx^{\alpha} \wedge dx^{\beta} \quad (5.57)$$

The tree-level dS_4 correlator corresponding to this term is given by:

$$\begin{aligned}\langle TTT \rangle_{\text{odd}} &= c_{TTT}^{\text{odd}} k_1 k_2 k_3 \text{Im} \left[\int_{-\infty}^0 d\eta \eta^5 e^{i\eta E} \right] \mathcal{M}_{W^2 \tilde{W}} \\ &= c_{TTT}^{\text{odd}} \frac{c_{123}}{E^6} \mathcal{M}_{W^2 \tilde{W}}\end{aligned}\quad (5.58)$$

Using the result for $\mathcal{M}_{W^2 \tilde{W}}$ given in (5.26), one can check that this matches the homogeneous part of the correlator in (3.5). This contribution was also calculated in [56]. In the flat space limit one obtains:

$$\lim_{E \rightarrow 0} \langle TTT \rangle_{\text{odd}} = c_{TTT}^{\text{odd}} \frac{c_{123}}{E^6} \mathcal{A}_{W^2 \tilde{W}} \quad (5.59)$$

where expression for $\mathcal{A}_{W^2 \tilde{W}}$ is given in (5.27). The parity-even contribution to the correlator is given by [20, 27, 33]:

$$\langle TTT \rangle_{\text{even}} = \frac{c_{TTT}^{\text{even}} c_{123}}{E^6} \mathcal{M}_{W^3} + 2c_{TT}^{\text{even}} \left(\frac{c_{123}}{E^2} + \frac{b_{123}}{E} - E \right) \mathcal{M}_{\text{EG}} \quad (5.60)$$

which upon using $\mathcal{M}_{W^3}$ given in (5.24) and $\mathcal{M}_{\text{EH}}$ given in (5.28) produces (3.15) upto contact term.⁸

In the flat space limit we obtain

$$\lim_{E \rightarrow 0} \langle TTT \rangle_{\text{even}} = c_{TTT}^{\text{even}} \frac{c_{123}}{E^6} \mathcal{A}_{W^3} + 2c_{TT}^{\text{even}} \frac{c_{123}}{E^2} \mathcal{A}_{\text{EG}} \quad (5.61)$$

where $\mathcal{A}_{W^3}$ and $\mathcal{A}_{\text{EG}}$ are given in (5.27) and (5.29).

6 Discussion

In this paper we established various double copy relations for parity-violating CFT_3 momentum space 3-point correlators. The double copy structure is a very special property of CFT correlators in momentum space, the analogue of which does not exist in position space. To understand this structure, we divided the momentum space CFT correlation function into two parts, which we called homogeneous and non-homogeneous pieces. It was crucial for our analysis that the homogeneous part consists of two pieces of conformally invariant structures, namely one parity-even and one parity-odd structure whereas the non-homogeneous part has only one parity-even conformally invariant piece - all other contributions are contact terms. Squaring the homogeneous piece could in principle generate three structures. However, interestingly it turns out that squaring the parity-odd and even part produces exactly the same structure, whereas the cross-term which is generated by multiplying the parity-odd and parity-even part gives rise to the needed parity-odd structure.

⁸This expression matches upto a number with the expression for the stress tensor correlator that appears in equation 4.12 of [33] once we identify

$$\mathcal{M}_{W^3} = -\frac{c_{123}E}{16J^2} \mathcal{A}_{F^3} \mathcal{M}_{F^3}.$$

See appendix C.1 for details.

This paper leaves various interesting directions to be followed upon. For example, for a general correlator of conserved currents of the form $\langle J_{s_1} J_{s_2} J_{s_3} \rangle$, we need to understand its structure better to be able make any detailed statement about its double copy. For instance, if we want to understand the parity-odd contribution we need to understand analogue of triangle inequality [64]. We looked only at marginal scalars, but it is of interest to find out if the double copy structure shows up in $\langle J_s J_s O_\Delta \rangle$ if O is a general scalar operator (that is, Δ is arbitrary). Holographic correlators corresponding to gluon and graviton amplitudes in AdS have been computed recently in [73–75]. It would be interesting to explore the double copy structure in this context.

It is also of interest to see whether the double copy relations continue to hold for higher point functions [49]. Establishing double copy relations for 4-point functions is a significantly harder problem. In this case it would be interesting to see if the double copy structure is visible even in some tractable limit or if one can infer the existence of a double copy structure for specific conformal blocks or Polyakov blocks. In this paper we have focused mainly on conformally invariant structures but it would be interesting to see what kind of constraints one needs to impose on OPE coefficients to get double copy relations.

Another interesting direction to study is a momentum space analogue of the analysis of [64]. In [64] conformally invariant parity-even and parity-odd structures were identified in position space. These could then be appropriately composed to get conformally invariant parity-even and parity-odd correlators in position space. The double copy structure of the momentum space CFT correlators that we investigated in this paper already hints towards the existence of such structures in momentum space. See appendix D for some details. We leave such questions for a future investigation.

Acknowledgments

The work of SJ and RRJ is supported by the Ramanujan Fellowship. AM would like to acknowledge the support of CSIR-UGC (JRF) fellowship (09/936(0212)/2019-EMR-I). The work of AS is supported by the KVPY scholarship. We acknowledge our debt to the people of India for their steady support of research in basic sciences.

A Expressions in spinor-helicity variables

In this section we write down various amplitudes in spinor helicity variables. Establishing a double copy at the level of spinor helicity variables is easy and obvious.

Gluon-gluon-scalar amplitudes. In spinor-helicity variables, the non-zero components of $\mathcal{M}$ take the following form:

$$\mathcal{M}_{\phi F^2}^{0--} = 2\langle 12 \rangle^2, \quad \mathcal{M}_{\phi F \tilde{F}}^{0--} = 2i\langle 12 \rangle^2. \quad (\text{A.1})$$

Also by complex conjugation $++$ helicity results also exist. However here as well as below we do not write them explicitly. In spinor-helicity variables flat space amplitudes take

the form

$$\mathcal{A}_{\phi F^2}^{0--} = 2\langle 12 \rangle^2, \quad \mathcal{A}_{\phi F\tilde{F}}^{0--} = 2i\langle 12 \rangle^2 \quad (\text{A.2})$$

where we have written down answer for negative helicity component. Although (A.1) and (A.2) might look similar, later is obtained from former in the $E \rightarrow 0$ limit.

Graviton-graviton-scalar amplitudes. In spinor-helicity variables, the non-zero components of $\mathcal{M}$ take the following form:

$$\mathcal{M}_{\phi W^2}^{0--} = \langle 12 \rangle^4, \quad \mathcal{M}_{\phi W\tilde{W}}^{0--} = 4i\langle 12 \rangle^4. \quad (\text{A.3})$$

In spinor-helicity variables flat space amplitudes take the form

$$\mathcal{A}_{\phi W^2}^{0--} = \langle 12 \rangle^4, \quad \mathcal{A}_{\phi W\tilde{W}}^{0--} = 4i\langle 12 \rangle^4 \quad (\text{A.4})$$

where we have written down answer for negative helicity component. Although (A.3) and (A.4) might look similar, the latter is obtained from the former in the $E \rightarrow 0$ limit.

Double copy. Comparing (A.1) and (A.2) with (A.3) and (A.4) we immediately see the double copy structure between gauge and gravity answers shown in (5.30) and (5.31).

Gluon-gluon-gluon amplitudes. In spinor-helicity variables, the non-zero components of $\mathcal{M}$ are given by:

$$\mathcal{M}_{F^2}^{---} = \frac{E}{k_1 k_2 k_3} \langle 12 \rangle \langle 23 \rangle \langle 31 \rangle, \quad \mathcal{M}_{F^2}^{--+} = \frac{(E - 2k_3)}{k_1 k_2 k_3} \langle 12 \rangle \langle 2\bar{3} \rangle \langle \bar{3}1 \rangle \quad (\text{A.5})$$

$$\mathcal{M}_{F\tilde{F}}^{---} = i \frac{E}{k_1 k_2 k_3} \langle 12 \rangle \langle 23 \rangle \langle 31 \rangle, \quad \mathcal{M}_{F\tilde{F}}^{--+} = i \frac{E}{k_1 k_2 k_3} \langle 12 \rangle \langle 2\bar{3} \rangle \langle \bar{3}1 \rangle \quad (\text{A.6})$$

$$\mathcal{M}_{F^3}^{---} = \langle 12 \rangle \langle 23 \rangle \langle 31 \rangle, \quad \mathcal{M}_{F^2\tilde{F}}^{---} = i\langle 12 \rangle \langle 23 \rangle \langle 31 \rangle \quad (\text{A.7})$$

In the flat space limit we get

$$\mathcal{A}_{F^2}^{---} = 0, \quad \mathcal{A}_{F^2}^{--+} = -2k_3 \langle 12 \rangle \langle 2\bar{3} \rangle \langle \bar{3}1 \rangle \quad (\text{A.8})$$

$$\mathcal{A}_{F\tilde{F}}^{---} = 0, \quad \mathcal{A}_{F\tilde{F}}^{--+} = 0$$

$$\mathcal{A}_{F^3}^{---} = \langle 12 \rangle \langle 23 \rangle \langle 31 \rangle, \quad \mathcal{A}_{F^2\tilde{F}}^{---} = i\langle 12 \rangle \langle 23 \rangle \langle 31 \rangle \quad (\text{A.9})$$

Graviton-graviton-graviton amplitudes. In spinor-helicity variables, the non-zero components of $\mathcal{M}$ are given by:

$$\mathcal{M}_{W^3}^{---} = \langle 12 \rangle^2 \langle 23 \rangle^2 \langle 31 \rangle^2, \quad \mathcal{M}_{W^2\tilde{W}}^{---} = i\langle 12 \rangle^2 \langle 23 \rangle^2 \langle 31 \rangle^2 \quad (\text{A.10})$$

In the flat space limit we get

$$\mathcal{A}_{W^3}^{---} = \langle 12 \rangle^2 \langle 23 \rangle^2 \langle 31 \rangle^2, \quad \mathcal{A}_{W^2\tilde{W}}^{---} = i\langle 12 \rangle^2 \langle 23 \rangle^2 \langle 31 \rangle^2 \quad (\text{A.11})$$

Double copy. Comparing (A.5) and (A.8) with (A.10) and (A.11) we immediately see the double copy structure between gauge and gravity answers shown in (5.37).

B $\langle TJJ \rangle$

We calculate $\langle TJJ \rangle$ using gravity techniques for completeness. This result is not used in the main text.

B.1 Mixed gauge-graviton amplitudes

Gluon graviton interaction. Let us consider interactions between gluons and gravitons which contribute to mixed CFT correlators of the spin-one conserved current J and the stress-tensor T . The interactions we consider are:

$$S^I = \int d^4x \sqrt{g} F^2 + S_{\text{even}}^I + S_{\text{odd}}^I \quad (\text{B.1})$$

where

$$S_{\text{even}}^I = \gamma \int d^4x \sqrt{g} W_{\mu\nu\rho\sigma} F^{\mu\nu} F^{\rho\sigma} \quad (\text{B.2})$$

$$S_{\text{odd}}^I = \tilde{\gamma} \int W_{\mu\nu\rho\sigma} F^{\mu\nu} F_{\alpha\beta} dx^\rho \wedge dx^\sigma \wedge dx^\alpha \wedge dx^\beta \quad (\text{B.3})$$

are the parity-preserving and parity-violating actions, respectively. The tree-level scattering amplitude for parity-even case is given by

$$\begin{aligned} \mathcal{A}_{F^2} &= -(z_1 \cdot z_2)(z_1 \cdot k_3)(z_3 \cdot k_2) - (z_1 \cdot k_2)(z_1 \cdot z_3)(z_2 \cdot k_3) + (z_1 \cdot k_2)(z_2 \cdot z_3)(z_1 \cdot k_3) \\ \mathcal{A}_{WF^2} &= 4(z_1 \cdot k_2)(z_2 \cdot k_1)(z_1 \cdot k_3)(z_3 \cdot k_1) \end{aligned}$$

Let us now compute the odd part of $\langle TJJ \rangle$. The contribution to $\langle TJJ \rangle_{\text{odd}}$ comes from the following term in the action:

$$\int W_{\mu\nu\rho\sigma} F^{\mu\nu} F_{\alpha\beta} dx^\rho \wedge dx^\sigma \wedge dx^\alpha \wedge dx^\beta \quad (\text{B.4})$$

This gives the following parity odd contribution:

$$\begin{aligned} \mathcal{M}_{WF\tilde{F}} &= [2(z_1 \cdot k_2)(z_2 \cdot k_1) + 2(k_1 \cdot k_2)(z_1 \cdot z_2)] \epsilon(z_1 k_1 z_3 k_3) \\ &= [2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + E(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2] \epsilon(z_1 k_1 z_3 k_3) \end{aligned} \quad (\text{B.5})$$

The flat space amplitudes are obtained by taking $E \rightarrow 0$:

$$\mathcal{A}_{WF\tilde{F}} = 2(z_1 \cdot k_2)(z_2 \cdot k_1) \epsilon(z_1 k_1 z_3 k_3) \quad (\text{B.6})$$

The non-zero spinor-helicity components of $\mathcal{M}$ are given by:

$$\mathcal{M}_{F^2}^{--+} = \frac{\langle 12 \rangle^2 \langle \bar{1} \bar{3} \rangle^2}{2k_1^2}, \quad \mathcal{M}_{WF^2}^{---} = 4\langle 12 \rangle^2 \langle 13 \rangle^2, \quad \mathcal{M}_{WF\tilde{F}}^{---} = 4\langle 12 \rangle^2 \langle 13 \rangle^2. \quad (\text{B.7})$$

In spinor-helicity language the non-zero components of the flat space amplitude are given by:

$$\mathcal{A}_{F^2}^{--+} = \frac{\langle 12 \rangle^2 \langle \bar{1} \bar{3} \rangle^2}{2k_1^2}, \quad \mathcal{A}_{WF^2}^{---} = 4\langle 12 \rangle^2 \langle 13 \rangle^2, \quad \mathcal{A}_{WF\tilde{F}}^{---} = 4\langle 12 \rangle^2 \langle 13 \rangle^2. \quad (\text{B.8})$$

Now the parity-odd contribution to correlator $\langle TJJ \rangle$ is given by

$$\begin{aligned}\langle TJJ \rangle_{\text{odd}} &= k_1 \text{Im} \left[\int_{-\infty}^0 d\eta \, \eta^3 e^{i\eta E} \right] \mathcal{M}_{WF\tilde{F}} \\ &= \frac{k_1}{E^4} \mathcal{M}_{WF\tilde{F}}\end{aligned}\quad (\text{B.9})$$

One can check that the correlation function given in (B.9) satisfies the appropriate conformal Ward identity.

C Proof of some double copy relations

C.1 $\mathcal{M}_{W^3} \propto (\mathcal{M}_{F^3})^2$

Here, we briefly show how (5.34) can be derived in momentum space. $\mathcal{M}_{F^3}^2$ is given by

$$\begin{aligned}\mathcal{M}_{F^3}^2 &= (\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{k}_1 \cdot \vec{z}_3)^2 (\vec{k}_3 \cdot \vec{z}_2)^2 + E \left((\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{k}_1 \cdot \vec{z}_3) (\vec{k}_3 \cdot \vec{z}_2) (\vec{z}_2 \cdot \vec{z}_3) + \text{cyclic perm.} \right) \\ &\quad + \frac{E^2}{4} \left(2k_1 k_2 (\vec{k}_2 \cdot \vec{z}_1) (\vec{k}_3 \cdot \vec{z}_2) (\vec{z}_1 \cdot \vec{z}_3) (\vec{z}_2 \cdot \vec{z}_3) + k_1^2 (\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{z}_2 \cdot \vec{z}_3)^2 + \text{cyclic perm.} \right)\end{aligned}\quad (\text{C.1})$$

$\mathcal{M}_{W^3}$ is given by

$$\begin{aligned}\mathcal{M}_{W^3} &= A_1 (k_1, k_2, k_3) (\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{k}_3 \cdot \vec{z}_2)^2 (\vec{k}_1 \cdot \vec{z}_3)^2 \\ &\quad + \left(A_2 (k_1, k_2, k_3) (\vec{z}_1 \cdot \vec{z}_2) (\vec{k}_2 \cdot \vec{z}_1) (\vec{k}_3 \cdot \vec{z}_2) (\vec{k}_1 \cdot \vec{z}_3)^2 + \text{cyclic perm.} \right) \\ &\quad + \left(A_3 (k_1, k_2, k_3) (\vec{z}_1 \cdot \vec{z}_2)^2 (\vec{k}_1 \cdot \vec{z}_3)^2 + \text{cyclic perm.} \right) \\ &\quad + \left(A_4 (k_1, k_2, k_3) (\vec{z}_1 \cdot \vec{z}_3) (\vec{z}_2 \cdot \vec{z}_3) (\vec{k}_2 \cdot \vec{z}_1) (\vec{k}_3 \cdot \vec{z}_2) + \text{cyclic perm.} \right) \\ &\quad + A_5 (k_1, k_2, k_3) (\vec{z}_1 \cdot \vec{z}_2) (\vec{z}_2 \cdot \vec{z}_3) (\vec{z}_1 \cdot \vec{z}_3)\end{aligned}\quad (\text{C.2})$$

where [33]

$$A_1 = 1, \quad A_2 = \frac{E}{2} (2k_3 - E), \quad A_3 = 0, \quad A_4 = \frac{E^2}{4} (E - 2k_1) (E - 2k_1), \quad A_5 = -\frac{J^2 E^2}{8}$$

We can use the following two degeneracies to set A_4 and A_5 to zero [20] in (C.2)

$$(\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{k}_3 \cdot \vec{z}_2)^2 + (k_3^2 - k_1^2 - k_2^2) (\vec{k}_2 \cdot \vec{z}_1) (\vec{k}_3 \cdot \vec{z}_2) (\vec{z}_1 \cdot \vec{z}_2) - \frac{J^2}{4} (\vec{z}_1 \cdot \vec{z}_2)^2 = 0 \quad (\text{C.3})$$

$$\begin{aligned}(\vec{k}_2 \cdot \vec{z}_1) (\vec{k}_3 \cdot \vec{z}_2) (\vec{k}_1 \cdot \vec{z}_3)^2 + k_3^2 (\vec{z}_1 \cdot \vec{z}_2) (\vec{k}_1 \cdot \vec{z}_3)^2 + \frac{1}{2} (k_1^2 - k_2^2 + k_3^2) (\vec{z}_2 \cdot \vec{z}_3) (\vec{k}_2 \cdot \vec{z}_1) (\vec{k}_1 \cdot \vec{z}_3) \\ + \frac{1}{2} (k_2^2 - k_1^2 + k_3^2) (\vec{z}_1 \cdot \vec{z}_3) (\vec{k}_3 \cdot \vec{z}_2) (\vec{k}_1 \cdot \vec{z}_3) + \frac{J^2}{4} (\vec{z}_1 \cdot \vec{z}_3) (\vec{z}_2 \cdot \vec{z}_3) = 0\end{aligned}\quad (\text{C.4})$$

Using this $\mathcal{M}_{W^3}$ can be related to $\mathcal{A}_{F^3} \mathcal{M}_{F^3}$, where

$$\begin{aligned}\mathcal{A}_{F^3} \mathcal{M}_{F^3} &= 2 (\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{k}_1 \cdot \vec{z}_3)^2 (\vec{k}_3 \cdot \vec{z}_2)^2 \\ &\quad + E \left((\vec{k}_2 \cdot \vec{z}_1)^2 (\vec{k}_1 \cdot \vec{z}_3) (\vec{k}_3 \cdot \vec{z}_2) (\vec{z}_2 \cdot \vec{z}_3) k_1 + \text{cyclic perm.} \right)\end{aligned}\quad (\text{C.5})$$

Using the same two degeneracies on (C.1) to remove the term proportional to E^2 , we obtain the following relation.

$$-32 \frac{c_{123}E}{J^2} \mathcal{A}_{F^3} \mathcal{M}_{F^3} = (\mathcal{M}_{F^3})^2 = \mathcal{M}_{W^3} \quad (\text{C.6})$$

Therefore, we see that $(\mathcal{M}_{F^3})^2 = \mathcal{M}_{W^3}$ which implies that the double copy structure holds beyond the flat space limit for the $\langle JJJ \rangle$ and $\langle TTT \rangle$.

C.2 $(\mathcal{M}_{\phi F^2})^2 \propto (\mathcal{M}_{\phi F \tilde{F}})^2$

Some of the equality relation in (5.30) was established in [33]. Here we establish the second part of (5.30). Consider the identity

$$\epsilon_{\mu\nu\rho\sigma} \epsilon^{\alpha\beta\gamma\delta} = \begin{vmatrix} \delta_\mu^\alpha & \delta_\mu^\beta & \delta_\mu^\gamma & \delta_\mu^\delta \\ \delta_\nu^\alpha & \delta_\nu^\beta & \delta_\nu^\gamma & \delta_\nu^\delta \\ \delta_\rho^\alpha & \delta_\rho^\beta & \delta_\rho^\gamma & \delta_\rho^\delta \\ \delta_\sigma^\alpha & \delta_\sigma^\beta & \delta_\sigma^\gamma & \delta_\sigma^\delta \end{vmatrix} \quad (\text{C.7})$$

Since, we have

$$\mathcal{M}_{\phi F \tilde{F}} = \epsilon(z_1 k_1 z_2 k_2) \quad (\text{C.8})$$

Therefore, we may write

$$(\mathcal{M}_{\phi F \tilde{F}})^2 = \begin{vmatrix} 0 & 0 & \vec{z}_1 \cdot \vec{z}_2 & \vec{z}_1 \cdot \vec{k}_2 \\ 0 & 0 & \vec{z}_2 \cdot \vec{k}_1 & k_1 \cdot k_2 \\ \vec{z}_1 \cdot \vec{z}_2 & \vec{z}_2 \cdot \vec{k}_1 & 0 & 0 \\ \vec{z}_1 \cdot \vec{k}_2 & k_1 \cdot k_2 & 0 & 0 \end{vmatrix} = [(\vec{z}_1 \cdot \vec{z}_2) k_1 \cdot k_2 - \vec{z}_1 \cdot \vec{k}_2 \vec{z}_2 \cdot \vec{k}_1]^2 \quad (\text{C.9})$$

Since, in 4D we have $k_1 \cdot k_2 = -\frac{E(E-2k_3)}{2}$, using (5.22)

$$(\mathcal{M}_{\phi F \tilde{F}})^2 \propto (\mathcal{M}_{\phi F^2})^2 \quad (\text{C.10})$$

C.3 $(\mathcal{M}_{F^3})^2 \propto (\mathcal{M}_{F^2 \tilde{F}})^2$

Here we establish the relation (5.36). Consider the identity (C.7). Since, we have

$$\mathcal{M}_{F^2 \tilde{F}} = \epsilon^{\mu\nu\rho\sigma} F_{(1)\mu}{}^\tau F_{(2)\tau\nu} F_{(3)\rho\sigma} \quad (\text{C.11})$$

Therefore, we may write

$$\begin{aligned} (\mathcal{M}_{F^2 \tilde{F}})^2 &= \epsilon^{\mu\nu\rho\sigma} F_{(1)\mu}{}^\tau F_{(2)\tau\nu} F_{(3)\rho\sigma} \epsilon^{\alpha\beta\gamma\delta} F_{(1)\alpha}{}^\lambda F_{(2)\lambda\beta} F_{(3)\gamma\delta} \\ &= \begin{vmatrix} 0 & \delta_{\nu\alpha} & z_{3\alpha} & k_{3\alpha} \\ \delta_{\mu\beta} & 0 & z_{3\beta} & k_{3\beta} \\ z_{3\mu} & z_{3\nu} & 0 & 0 \\ k_{3\mu} & k_{3\nu} & 0 & 0 \end{vmatrix} F_{(1)\mu}{}^\tau F_{(2)\tau\nu} F_{(1)\alpha}{}^\lambda F_{(2)\lambda\beta} \\ &= (F_{(3)})^{\mu\nu} (F_{(3)})^{\alpha\beta} F_{(1)\mu}{}^\tau F_{(2)\tau\nu} F_{(1)\alpha}{}^\lambda F_{(2)\lambda\beta} = (\mathcal{M}_{F^3})^2 \end{aligned} \quad (\text{C.12})$$

D Momentum space expression of higher spin correlators

In this section we give the momentum space expression for the parity-even and parity-odd parts of higher spin correlators $\langle J_s J_s O_3 \rangle$ and $\langle J_s J_s J_s \rangle$ using the relations in sub-section 4.5 and the expressions for $\langle JJO_3 \rangle$ and $\langle JJJ \rangle$ given in section 3:

$$\begin{aligned}
 \langle J_s J_s O_3 \rangle_{\text{even}, \mathbf{h}} &= (k_1 k_2)^{s-1} (E + (2s-1)k_3) \left[\frac{1}{E^2} \left\{ 2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + E(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2 \right\} \right]^s \\
 \langle J_s J_s O_3 \rangle_{\text{odd}, \mathbf{h}} &= (k_1 k_2)^{s-1} \frac{(E + (2s-1)k_3)}{E^{2s}} \left[k_2 \epsilon^{k_1 z_1 z_2} - k_1 \epsilon^{k_2 z_1 z_2} \right] \\
 &\quad \times \left[2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_1) + E(E - 2k_3)\vec{z}_1 \cdot \vec{z}_2 \right]^{s-1} \\
 \langle J_s J_s J_s \rangle_{\text{even}, \mathbf{h}} &= (k_1 k_2 k_3)^{s-1} \left[\frac{1}{E^3} \left\{ 2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_3)(\vec{z}_3 \cdot \vec{k}_1) + E\{k_3(\vec{z}_1 \cdot \vec{z}_2)(\vec{z}_3 \cdot \vec{k}_1) + \text{cyclic}\} \right\} \right]^s \\
 \langle J_s J_s J_s \rangle_{\text{odd}, \mathbf{h}} &= (k_1 k_2 k_3)^{s-1} \frac{1}{E^3} \left[\left\{ (\vec{k}_1 \cdot \vec{z}_3) (\epsilon^{k_3 z_1 z_2} k_1 - \epsilon^{k_1 z_1 z_2} k_3) + (\vec{k}_3 \cdot \vec{z}_2) (\epsilon^{k_1 z_1 z_3} k_2 - \epsilon^{k_2 z_1 z_3} k_1) \right. \right. \\
 &\quad \left. \left. - (\vec{z}_2 \cdot \vec{z}_3) \epsilon^{k_1 k_2 z_1} E + \frac{k_1}{2} \epsilon^{z_1 z_2 z_3} E(E - 2k_1) \right\} + \text{cyclic perm} \right] \\
 &\quad \times \left[\frac{1}{E^3} \left\{ 2(\vec{z}_1 \cdot \vec{k}_2)(\vec{z}_2 \cdot \vec{k}_3)(\vec{z}_3 \cdot \vec{k}_1) + E\{k_3(\vec{z}_1 \cdot \vec{z}_2)(\vec{z}_3 \cdot \vec{k}_1) + \text{cyclic}\} \right\} \right]^{s-1}.
 \end{aligned} \tag{D.1}$$

One can use the Todorov operator [76] to strip off the polarization vectors from the expressions in (D.1). This operator is given by

$$D_z^i = \left(\frac{1}{2} + \vec{z} \cdot \frac{\partial}{\partial \vec{z}} \right) \frac{\partial}{\partial z_i} - \frac{1}{2} z^i \frac{\partial^2}{\partial \vec{z} \cdot \partial \vec{z}} \tag{D.2}$$

Therefore, we have

$$\begin{aligned}
 \langle J^{\mu_1 \dots \mu_s} J^{\nu_1 \dots \nu_s} O_3 \rangle_{\text{even}, \mathbf{h}} &= \prod_{i=1}^s D_{z_1}^{\mu_i} D_{z_2}^{\nu_i} \langle J_s J_s O_3 \rangle_{\text{even}, \mathbf{h}} \\
 \langle J^{\mu_1 \dots \mu_s} J^{\nu_1 \dots \nu_s} O_3 \rangle_{\text{odd}, \mathbf{h}} &= \prod_{i=1}^s D_{z_1}^{\mu_i} D_{z_2}^{\nu_i} \langle J_s J_s O_3 \rangle_{\text{odd}, \mathbf{h}} \\
 \langle J^{\mu_1 \dots \mu_s} J^{\nu_1 \dots \nu_s} J^{\rho_1 \dots \rho_s} \rangle_{\text{even}, \mathbf{h}} &= \prod_{i=1}^s D_{z_1}^{\mu_i} D_{z_2}^{\nu_i} D_{z_3}^{\rho_i} \langle J_s J_s J_s \rangle_{\text{even}, \mathbf{h}} \\
 \langle J^{\mu_1 \dots \mu_s} J^{\nu_1 \dots \nu_s} J^{\rho_1 \dots \rho_s} \rangle_{\text{odd}, \mathbf{h}} &= \prod_{i=1}^s D_{z_1}^{\mu_i} D_{z_2}^{\nu_i} D_{z_3}^{\rho_i} \langle J_s J_s J_s \rangle_{\text{odd}, \mathbf{h}}
 \end{aligned} \tag{D.3}$$

E Double copy relations in spinor-helicity notation

In our discussion of double copy relations we have mostly focussed on the momentum dependent structures of the correlators. However, a strict doubly copy relation would in addition relate the OPE like coefficients in the correlators. In this section we make this clear using the spinor-helicity notation and the double copy structure of $\langle J_{2s} J_{2s} J_{2s} \rangle$.

We have the following non-zero spinor-helicity components of $\langle J_s J_s J_s \rangle$:

$$\begin{aligned}\langle J_s^- J_s^- J_s^- \rangle_{\mathbf{h}} &= (c_{s,\text{even}} + i c_{s,\text{odd}}) \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle 12 \rangle^s \langle 23 \rangle^s \langle 31 \rangle^s \\ \langle J_s^+ J_s^+ J_s^+ \rangle_{\mathbf{h}} &= (c_{s,\text{even}} - i c_{s,\text{odd}}) \frac{(k_1 k_2 k_3)^{s-1}}{E^{3s}} \langle \bar{1}\bar{2} \rangle^s \langle \bar{2}\bar{3} \rangle^s \langle \bar{3}\bar{1} \rangle^s\end{aligned}\quad (\text{E.1})$$

The non-zero spinor-helicity components of $\langle JJJ \rangle$ are:

$$\begin{aligned}\langle J^- J^- J^- \rangle_{\mathbf{h}} &= (c_{1,\text{even}} + i c_{1,\text{odd}}) \frac{1}{E^3} \langle 12 \rangle \langle 23 \rangle \langle 31 \rangle \\ \langle J^+ J^+ J^+ \rangle_{\mathbf{h}} &= (c_{1,\text{even}} - i c_{1,\text{odd}}) \frac{1}{E^3} \langle \bar{1}\bar{2} \rangle \langle \bar{2}\bar{3} \rangle \langle \bar{3}\bar{1} \rangle\end{aligned}\quad (\text{E.2})$$

Demanding the s -copy relations of subsection (4.5) we get the following equations that constrain the coefficients $c_{s,\text{even}}$ and $c_{s,\text{odd}}$ in terms of $c_{1,\text{even}}$ and $c_{1,\text{odd}}$:

$$\begin{aligned}c_{s,\text{even}} + i c_{s,\text{odd}} &= (c_{1,\text{even}} + i c_{1,\text{odd}})^s \\ c_{s,\text{even}} - i c_{s,\text{odd}} &= (c_{1,\text{even}} - i c_{1,\text{odd}})^s\end{aligned}\quad (\text{E.3})$$

Specifically when $s = 2$, for which the conserved current is the stress tensor we obtain:

$$\begin{aligned}c_{TTT,\text{even}} &= c_{1,\text{even}}^2 - c_{1,\text{odd}}^2 \\ c_{TTT,\text{odd}} &= 2c_{1,\text{even}}c_{1,\text{odd}}\end{aligned}\quad (\text{E.4})$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] M. Gary, S.B. Giddings and J. Penedones, *Local bulk S-matrix elements and CFT singularities*, *Phys. Rev. D* **80** (2009) 085005 [[arXiv:0903.4437](https://arxiv.org/abs/0903.4437)] [[INSPIRE](#)].
- [2] M. Gary and S.B. Giddings, *The Flat space S-matrix from the AdS/CFT correspondence?*, *Phys. Rev. D* **80** (2009) 046008 [[arXiv:0904.3544](https://arxiv.org/abs/0904.3544)] [[INSPIRE](#)].
- [3] S. Komatsu, M.F. Paulos, B.C. Van Rees and X. Zhao, *Landau diagrams in AdS and S-matrices from conformal correlators*, *JHEP* **11** (2020) 046 [[arXiv:2007.13745](https://arxiv.org/abs/2007.13745)] [[INSPIRE](#)].
- [4] J. Penedones, *Writing CFT correlation functions as AdS scattering amplitudes*, *JHEP* **03** (2011) 025 [[arXiv:1011.1485](https://arxiv.org/abs/1011.1485)] [[INSPIRE](#)].
- [5] S. Raju, *New Recursion Relations and a Flat Space Limit for AdS/CFT Correlators*, *Phys. Rev. D* **85** (2012) 126009 [[arXiv:1201.6449](https://arxiv.org/abs/1201.6449)] [[INSPIRE](#)].
- [6] A.L. Fitzpatrick and J. Kaplan, *Analyticity and the Holographic S-matrix*, *JHEP* **10** (2012) 127 [[arXiv:1111.6972](https://arxiv.org/abs/1111.6972)] [[INSPIRE](#)].
- [7] A.L. Fitzpatrick and J. Kaplan, *Unitarity and the Holographic S-matrix*, *JHEP* **10** (2012) 032 [[arXiv:1112.4845](https://arxiv.org/abs/1112.4845)] [[INSPIRE](#)].
- [8] S. Caron-Huot, *Analyticity in Spin in Conformal Theories*, *JHEP* **09** (2017) 078 [[arXiv:1703.00278](https://arxiv.org/abs/1703.00278)] [[INSPIRE](#)].

- [9] M. Gillioz, M. Meineri and J. Penedones, *A scattering amplitude in Conformal Field Theory*, *JHEP* **11** (2020) 139 [[arXiv:2003.07361](#)] [[INSPIRE](#)].
- [10] H. Kawai, D.C. Lewellen and S.H.H. Tye, *A Relation Between Tree Amplitudes of Closed and Open Strings*, *Nucl. Phys. B* **269** (1986) 1 [[INSPIRE](#)].
- [11] Z. Bern, J.J.M. Carrasco and H. Johansson, *New Relations for Gauge-Theory Amplitudes*, *Phys. Rev. D* **78** (2008) 085011 [[arXiv:0805.3993](#)] [[INSPIRE](#)].
- [12] Z. Bern, J.J.M. Carrasco and H. Johansson, *Perturbative Quantum Gravity as a Double Copy of Gauge Theory*, *Phys. Rev. Lett.* **105** (2010) 061602 [[arXiv:1004.0476](#)] [[INSPIRE](#)].
- [13] J. Broedel and L.J. Dixon, *Color-kinematics duality and double-copy construction for amplitudes from higher-dimension operators*, *JHEP* **10** (2012) 091 [[arXiv:1208.0876](#)] [[INSPIRE](#)].
- [14] H. Johansson and J. Nohle, *Conformal Gravity from Gauge Theory*, [arXiv:1707.02965](#) [[INSPIRE](#)].
- [15] H. Johansson, G. Mogull and F. Teng, *Unraveling conformal gravity amplitudes*, *JHEP* **09** (2018) 080 [[arXiv:1806.05124](#)] [[INSPIRE](#)].
- [16] Z. Bern, S. Davies, T. Dennen, Y.-t. Huang and J. Nohle, *Color-Kinematics Duality for Pure Yang-Mills and Gravity at One and Two Loops*, *Phys. Rev. D* **92** (2015) 045041 [[arXiv:1303.6605](#)] [[INSPIRE](#)].
- [17] S. He, O. Schlotterer and Y. Zhang, *New BCJ representations for one-loop amplitudes in gauge theories and gravity*, *Nucl. Phys. B* **930** (2018) 328 [[arXiv:1706.00640](#)] [[INSPIRE](#)].
- [18] Z. Bern, J.J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, *The Duality Between Color and Kinematics and its Applications*, [arXiv:1909.01358](#) [[INSPIRE](#)].
- [19] C. Corianò, L. Delle Rose, E. Mottola and M. Serino, *Solving the Conformal Constraints for Scalar Operators in Momentum Space and the Evaluation of Feynman's Master Integrals*, *JHEP* **07** (2013) 011 [[arXiv:1304.6944](#)] [[INSPIRE](#)].
- [20] A. Bzowski, P. McFadden and K. Skenderis, *Implications of conformal invariance in momentum space*, *JHEP* **03** (2014) 111 [[arXiv:1304.7760](#)] [[INSPIRE](#)].
- [21] L. Bonora, A.D. Pereira and B. Lima de Souza, *Regularization of energy-momentum tensor correlators and parity-odd terms*, *JHEP* **06** (2015) 024 [[arXiv:1503.03326](#)] [[INSPIRE](#)].
- [22] L. Bonora and B. Lima de Souza, *Pure contact term correlators in CFT*, *Bled Workshops Phys.* **16** (2015) 22 [[arXiv:1511.06635](#)] [[INSPIRE](#)].
- [23] L. Bonora, M. Cvitan, P. Dominis Prester, B. Lima de Souza and I. Smolić, *Massive fermion model in 3d and higher spin currents*, *JHEP* **05** (2016) 072 [[arXiv:1602.07178](#)] [[INSPIRE](#)].
- [24] L. de Souza, *CFT's, contact terms and anomalies*, Ph.D. Thesis, Scuola Internazionale Superiore di Studi Avanzati, Trieste Italy (2016).
- [25] A. Bzowski, P. McFadden and K. Skenderis, *Scalar 3-point functions in CFT: renormalisation, β -functions and anomalies*, *JHEP* **03** (2016) 066 [[arXiv:1510.08442](#)] [[INSPIRE](#)].
- [26] A. Bzowski, P. McFadden and K. Skenderis, *Evaluation of conformal integrals*, *JHEP* **02** (2016) 068 [[arXiv:1511.02357](#)] [[INSPIRE](#)].
- [27] A. Bzowski, P. McFadden and K. Skenderis, *Renormalised 3-point functions of stress tensors and conserved currents in CFT*, *JHEP* **11** (2018) 153 [[arXiv:1711.09105](#)] [[INSPIRE](#)].

- [28] C. Corianò and M.M. Maglio, *Exact Correlators from Conformal Ward Identities in Momentum Space and the Perturbative TJJ Vertex*, *Nucl. Phys. B* **938** (2019) 440 [[arXiv:1802.07675](#)] [[INSPIRE](#)].
- [29] A. Bzowski, P. McFadden and K. Skenderis, *Renormalised CFT 3-point functions of scalars, currents and stress tensors*, *JHEP* **11** (2018) 159 [[arXiv:1805.12100](#)] [[INSPIRE](#)].
- [30] M. Gillioz, *Momentum-space conformal blocks on the light cone*, *JHEP* **10** (2018) 125 [[arXiv:1807.07003](#)] [[INSPIRE](#)].
- [31] C. Corianò and M.M. Maglio, *Conformal Ward Identities and the Coupling of QED and QCD to Gravity*, *EPJ Web Conf.* **192** (2018) 00047 [[arXiv:1809.05940](#)] [[INSPIRE](#)].
- [32] S. Albayrak and S. Kharel, *Towards the higher point holographic momentum space amplitudes*, *JHEP* **02** (2019) 040 [[arXiv:1810.12459](#)] [[INSPIRE](#)].
- [33] J.A. Farrow, A.E. Lipstein and P. McFadden, *Double copy structure of CFT correlators*, *JHEP* **02** (2019) 130 [[arXiv:1812.11129](#)] [[INSPIRE](#)].
- [34] H. Isono, T. Noumi and G. Shiu, *Momentum space approach to crossing symmetric CFT correlators*, *JHEP* **07** (2018) 136 [[arXiv:1805.11107](#)] [[INSPIRE](#)].
- [35] H. Isono, T. Noumi and G. Shiu, *Momentum space approach to crossing symmetric CFT correlators. Part II. General spacetime dimension*, *JHEP* **10** (2019) 183 [[arXiv:1908.04572](#)] [[INSPIRE](#)].
- [36] H. Isono, T. Noumi and T. Takeuchi, *Momentum space conformal three-point functions of conserved currents and a general spinning operator*, *JHEP* **05** (2019) 057 [[arXiv:1903.01110](#)] [[INSPIRE](#)].
- [37] C. Corianò and M.M. Maglio, *On Some Hypergeometric Solutions of the Conformal Ward Identities of Scalar 4-point Functions in Momentum Space*, *JHEP* **09** (2019) 107 [[arXiv:1903.05047](#)] [[INSPIRE](#)].
- [38] M. Gillioz, *Conformal 3-point functions and the Lorentzian OPE in momentum space*, *Commun. Math. Phys.* **379** (2020) 227 [[arXiv:1909.00878](#)] [[INSPIRE](#)].
- [39] A. Bzowski, P. McFadden and K. Skenderis, *Conformal n-point functions in momentum space*, *Phys. Rev. Lett.* **124** (2020) 131602 [[arXiv:1910.10162](#)] [[INSPIRE](#)].
- [40] M. Gillioz, X. Lu, M.A. Luty and G. Mikaberidze, *Convergent Momentum-Space OPE and Bootstrap Equations in Conformal Field Theory*, *JHEP* **03** (2020) 102 [[arXiv:1912.05550](#)] [[INSPIRE](#)].
- [41] T. Bautista and H. Godazgar, *Lorentzian CFT 3-point functions in momentum space*, *JHEP* **01** (2020) 142 [[arXiv:1908.04733](#)] [[INSPIRE](#)].
- [42] C. Corianò, M.M. Maglio and D. Theofilopoulos, *Four-Point Functions in Momentum Space: Conformal Ward Identities in the Scalar/Tensor case*, *Eur. Phys. J. C* **80** (2020) 540 [[arXiv:1912.01907](#)] [[INSPIRE](#)].
- [43] A.E. Lipstein and P. McFadden, *Double copy structure and the flat space limit of conformal correlators in even dimensions*, *Phys. Rev. D* **101** (2020) 125006 [[arXiv:1912.10046](#)] [[INSPIRE](#)].
- [44] A. Bzowski, P. McFadden and K. Skenderis, *Conformal correlators as simplex integrals in momentum space*, *JHEP* **01** (2021) 192 [[arXiv:2008.07543](#)] [[INSPIRE](#)].

- [45] S. Jain, R.R. John and V. Malvimat, *Momentum space spinning correlators and higher spin equations in three dimensions*, *JHEP* **11** (2020) 049 [[arXiv:2005.07212](#)] [[INSPIRE](#)].
- [46] S. Jain, R.R. John and V. Malvimat, *Constraining momentum space correlators using slightly broken higher spin symmetry*, *JHEP* **04** (2021) 231 [[arXiv:2008.08610](#)] [[INSPIRE](#)].
- [47] C. Corianò and M.M. Maglio, *The Generalized Hypergeometric Structure of the Ward Identities of CFT's in Momentum Space in $d > 2$* , *Axioms* **9** (2020) 54 [[arXiv:2001.09622](#)] [[INSPIRE](#)].
- [48] S. Albayrak, S. Kharel and D. Meltzer, *On duality of color and kinematics in $(A)dS$ momentum space*, *JHEP* **03** (2021) 249 [[arXiv:2012.10460](#)] [[INSPIRE](#)].
- [49] C. Armstrong, A.E. Lipstein and J. Mei, *Color/kinematics duality in AdS_4* , *JHEP* **02** (2021) 194 [[arXiv:2012.02059](#)] [[INSPIRE](#)].
- [50] M. Serino, *The four-point correlation function of the energy-momentum tensor in the free conformal field theory of a scalar field*, *Eur. Phys. J. C* **80** (2020) 686 [[arXiv:2004.08668](#)] [[INSPIRE](#)].
- [51] C. Corianò and M.M. Maglio, *Conformal Field Theory in Momentum Space and Anomaly Actions in Gravity: The Analysis of 3- and 4-Point Functions*, [arXiv:2005.06873](#) [[INSPIRE](#)].
- [52] I. Mata, S. Raju and S. Trivedi, *CMB from CFT*, *JHEP* **07** (2013) 015 [[arXiv:1211.5482](#)] [[INSPIRE](#)].
- [53] A. Ghosh, N. Kundu, S. Raju and S.P. Trivedi, *Conformal Invariance and the Four Point Scalar Correlator in Slow-Roll Inflation*, *JHEP* **07** (2014) 011 [[arXiv:1401.1426](#)] [[INSPIRE](#)].
- [54] N. Kundu, A. Shukla and S.P. Trivedi, *Constraints from Conformal Symmetry on the Three Point Scalar Correlator in Inflation*, *JHEP* **04** (2015) 061 [[arXiv:1410.2606](#)] [[INSPIRE](#)].
- [55] N. Arkani-Hamed and J. Maldacena, *Cosmological Collider Physics*, [arXiv:1503.08043](#) [[INSPIRE](#)].
- [56] J.M. Maldacena and G.L. Pimentel, *On graviton non-Gaussianities during inflation*, *JHEP* **09** (2011) 045 [[arXiv:1104.2846](#)] [[INSPIRE](#)].
- [57] N. Arkani-Hamed, D. Baumann, H. Lee and G.L. Pimentel, *The Cosmological Bootstrap: Inflationary Correlators from Symmetries and Singularities*, *JHEP* **04** (2020) 105 [[arXiv:1811.00024](#)] [[INSPIRE](#)].
- [58] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee and G.L. Pimentel, *The cosmological bootstrap: weight-shifting operators and scalar seeds*, *JHEP* **12** (2020) 204 [[arXiv:1910.14051](#)] [[INSPIRE](#)].
- [59] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee and G.L. Pimentel, *The Cosmological Bootstrap: Spinning Correlators from Symmetries and Factorization*, [arXiv:2005.04234](#) [[INSPIRE](#)].
- [60] E. Skvortsov, *Light-Front Bootstrap for Chern-Simons Matter Theories*, *JHEP* **06** (2019) 058 [[arXiv:1811.12333](#)] [[INSPIRE](#)].
- [61] S. Jain, R.R. John, A. Mehta, A.A. Nizami and A. Suresh, *Momentum space parity-odd CFT 3-point functions*, [arXiv:2101.11635](#) [[INSPIRE](#)].
- [62] J. Maldacena and A. Zhiboedov, *Constraining Conformal Field Theories with A Higher Spin Symmetry*, *J. Phys. A* **46** (2013) 214011 [[arXiv:1112.1016](#)] [[INSPIRE](#)].

- [63] J. Maldacena and A. Zhiboedov, *Constraining conformal field theories with a slightly broken higher spin symmetry*, *Class. Quant. Grav.* **30** (2013) 104003 [[arXiv:1204.3882](#)] [[INSPIRE](#)].
- [64] S. Giombi, S. Prakash and X. Yin, *A Note on CFT Correlators in Three Dimensions*, *JHEP* **07** (2013) 105 [[arXiv:1104.4317](#)] [[INSPIRE](#)].
- [65] O. Aharony, G. Gur-Ari and R. Yacoby, *$d = 3$ Bosonic Vector Models Coupled to Chern-Simons Gauge Theories*, *JHEP* **03** (2012) 037 [[arXiv:1110.4382](#)] [[INSPIRE](#)].
- [66] S. Giombi, S. Minwalla, S. Prakash, S.P. Trivedi, S.R. Wadia and X. Yin, *Chern-Simons Theory with Vector Fermion Matter*, *Eur. Phys. J. C* **72** (2012) 2112 [[arXiv:1110.4386](#)] [[INSPIRE](#)].
- [67] O. Aharony, G. Gur-Ari and R. Yacoby, *Correlation Functions of Large N Chern-Simons-Matter Theories and Bosonization in Three Dimensions*, *JHEP* **12** (2012) 028 [[arXiv:1207.4593](#)] [[INSPIRE](#)].
- [68] S. Giombi, V. Gurucharan, V. Kirilin, S. Prakash and E. Skvortsov, *On the Higher-Spin Spectrum in Large N Chern-Simons Vector Models*, *JHEP* **01** (2017) 058 [[arXiv:1610.08472](#)] [[INSPIRE](#)].
- [69] S.D. Chowdhury, J.R. David and S. Prakash, *Constraints on parity violating conformal field theories in $d = 3$* , *JHEP* **11** (2017) 171 [[arXiv:1707.03007](#)] [[INSPIRE](#)].
- [70] E. Skvortsov, *Light-Front Bootstrap for Chern-Simons Matter Theories*, *JHEP* **06** (2019) 058 [[arXiv:1811.12333](#)] [[INSPIRE](#)].
- [71] S. Jain, R.R. John, A. Mehta, A.A. Nizami and A. Suresh, *Higher spin 3-point functions in 3d CFT using spinor-helicity variables*, [arXiv:2106.00016](#) [[INSPIRE](#)].
- [72] D. Karateev, P. Kravchuk and D. Simmons-Duffin, *Weight Shifting Operators and Conformal Blocks*, *JHEP* **02** (2018) 081 [[arXiv:1706.07813](#)] [[INSPIRE](#)].
- [73] L.F. Alday and X. Zhou, *All Holographic Four-Point Functions in All Maximally Supersymmetric CFTs*, *Phys. Rev. X* **11** (2021) 011056 [[arXiv:2006.12505](#)] [[INSPIRE](#)].
- [74] L.F. Alday and X. Zhou, *All Tree-Level Correlators for M-theory on $AdS_7 \times S^4$* , *Phys. Rev. Lett.* **125** (2020) 131604 [[arXiv:2006.06653](#)] [[INSPIRE](#)].
- [75] L.F. Alday, C. Behan, P. Ferrero and X. Zhou, *Gluon Scattering in AdS from CFT*, *JHEP* **06** (2021) 020 [[arXiv:2103.15830](#)] [[INSPIRE](#)].
- [76] V.K. Dobrev, V.B. Petkova, S.G. Petrova and I.T. Todorov, *Dynamical Derivation of Vacuum Operator Product Expansion in Euclidean Conformal Quantum Field Theory*, *Phys. Rev. D* **13** (1976) 887 [[INSPIRE](#)].