Heterotic Hyper-Kähler flux backgrounds

Nick Halmagyi,^{a,b} Dan Israël,^{a,b} Matthieu Sarkis^{a,b} and Eirik Eik Svanes^c

^a*Sorbonne Universités, UPMC Paris 06,
UMR 7589, LPTHE, 75005, Paris, France*

^b*CNRS, UMR 7589, LPTHE,
75005, Paris, France*

^c*Sorbonne Universités, Institut Lagrange de Paris,
98 bis Bd Arago, 75014, Paris, France*

E-mail: halmagyi@lpthe.jussieu.fr, israel@lpthe.jussieu.fr,
msarkis@lpthe.jussieu.fr, esvanes@lpthe.jussieu.fr

ABSTRACT: We study Heterotic supergravity on Hyper-Kähler manifolds in the presence of non-trivial warping and three form flux with Abelian bundles in the large charge limit. We find exact, regular solutions for multi-centered Gibbons-Hawking spaces and Atiyah-Hitchin manifolds. In the case of Atiyah-Hitchin, regularity requires that the circle at infinity is of the same order as the instanton number, which is taken to be large. Alternatively there may be a non-trivial density of smeared five branes at the bolt.

KEYWORDS: Superstring Vacua, Superstrings and Heterotic Strings, Flux compactifications, Black Holes in String Theory

ARXIV EPRINT: [1706.01725](https://arxiv.org/abs/1706.01725)

Contents

1	Introduction	1
2	Hyper-Kähler heterotic backgrounds	2
2.1	Principal torus bundles and type IIA/IIB solutions	3
3	Gibbons-Hawking: ALE and ALF	3
3.1	Five-brane and magnetic charges	5
3.2	Partial blow-down limits and fivebranes	6
3.3	Double scaling limit	6
4	Atiyah-Hitchin	7
4.1	Bianchi identity	9
4.2	Regularity	9
4.3	Five-brane charge	11
4.4	Gauge field charge	11
5	Fundamental string sources and AdS_3 solutions	11
5.1	Product of Hyper-Kähler manifolds	12
5.2	AdS_3 from Gibbons-Hawking	13
6	Conclusions	14
A	Eguchi-Hanson	15

1 Introduction

Heterotic flux backgrounds are interesting models of string backgrounds. There is no in principle impediment from using the R-NS string worldsheet formalism although compact models with minimal or no supersymmetry remain challenging to construct.

In this work we study local Heterotic flux backgrounds on Hyper-Kähler four manifolds: in particular the Gibbons-Hawking spaces [1] and the Atiyah-Hitchin manifold [2]. Key to our configurations is that the gauge fields are Abelian and we take the large charge limit such that $\text{Tr} F \wedge F$ dominates $\text{Tr} R \wedge R$ in the Bianchi identity. This large-charge limit has been previously studied on the Eguchi-Hanson space [3] and the conifold [4, 5] and this type of limit is familiar from the large-charge supergravity limit crucial to the development of holography [6] in type II and M-theory.

Our strategy is to first compute explicit solutions to the Hermitian-Yang-Mills equations on Hyper-Kähler spaces and then backreact them on the geometry. This backreaction affects only the conformal mode of the metric but generates a non-trivial three-form flux.

According to Aspinwall [7] we are studying *Goldilocks* theories with just the right amount of supersymmetry; perhaps then not surprisingly we solve the supergravity background exactly. For the Atiyah-Hitchin background it is nonetheless somewhat impressive that we can both solve Hermitian-Yang-Mills exactly and integrate analytically the resulting Poisson equation for the backreaction of this instanton.

The instanton we use on the Atiyah-Hitchin manifold is well known from a classic duality paper by Ashoke Sen [8]. Perhaps the central result of our work is that the Heterotic backreaction of this instanton can be made regular. This is somewhat non-trivial since the negative mass of Atiyah-Hitchin induces a negative warp factor thus violating the desired signature of space-time. We circumvent this in two ways: first by allowing the asymptotic circle to be large and secondly by including smeared five-brane sources.

We also study the presence of electric H -flux and fundamental strings. The electric flux modifies the BPS equations in a straightforward way and for each solution with magnetic H -flux, the electric flux can be added through a harmonic function on the Hyper-Kähler manifold. We analyze limits in which we recover AdS_3 geometries but these reduce to the known $\text{AdS}_3 \times S^3 \times HK_4$.

Upon completing this work we were made aware that our BPS equations have turned up in five dimensional supergravity. The local equations we study can essentially be found in [9, 10] but with very different global and regularity requirements. This is not surprising since we can dimensionally reduce our solutions on $\mathbb{R}^5$ to get solutions of ungauged five dimensional supergravity. In addition, these equations also turn up in type II supergravity for T^2 fibrations over Hyper-Kähler spaces and the type II analogue of the Gibbons-Hawking solutions we find have been analyzed in [11]. It is straightforward to convert our solutions on the Atiyah-Hitchin manifold to such type II backgrounds.

2 Hyper-Kähler heterotic backgrounds

The primary backgrounds we consider are of the form $\mathbb{R}^{1,5} \times HK_4$ where HK_4 is a warped Hyper-Kähler-four manifold. We consider a non-trivial three-form flux $\mathcal{H}_{(3)}$, dilaton Φ and Heterotic gauge field F . The background metric ansatz is:

$$ds_{10}^2 = ds_{1,5}^2 + H ds_4^2 \quad (2.1)$$

where ds_4^2 is an Hyper-Kähler metric on a four-manifold HK_4 and H a conformal factor. The BPS equations are fairly standard:

$$e^{2\phi} = H \quad (2.2a)$$

$$\mathcal{H}_{(3)} = - * _4 dH \quad (2.2b)$$

$$J_a \lrcorner F = 0, \quad a = 1, 2, 3 \quad (2.2c)$$

where the $*_4$ is the Hodge dual w.r.t. the Hyper-Kähler metric on HK_4 and J_a are the three Kähler forms.

A major difficulty in finding explicit solutions of Heterotic supergravity with non-trivial three-form flux is to satisfy the Bianchi identity at the appropriate order in α' . Following

earlier works by some of the authors [3–5], our strategy will be to work in a large (fivebrane) charge limit, ensuring that the contribution of the $\text{Tr} R \wedge R$ term is subdominant and can be consistently neglected. The Heterotic Bianchi identity simplifies to

$$d\mathcal{H}_{(3)} = \alpha' \text{Tr} F \wedge F \quad (2.3)$$

implying from the three-form ansatz (2.2b) that:

$$d *_4 dH = -\alpha' \text{Tr} F \wedge F. \quad (2.4)$$

2.1 Principal torus bundles and type IIA/IIB solutions

We consider a more general class of backgrounds, that can be viewed as local models of the principal torus bundles over wrapped K3 surfaces introduced in [12] and discussed in many works including [13–16]. They generalize the solutions based on Eguchi-Hanson space presented in [17]. The general ansatz for such principal two-torus bundle $T^2 \hookrightarrow M_6 \xrightarrow{\pi} HK_4$ over a Hyper-Kähler four-manifold is of the form

$$ds_{10}^2 = ds_{1,3}^2 + H ds_4^2 + \frac{U_2}{T_2} |dx + Tdy + \pi^* \alpha|^2, \quad (2.5)$$

where α is a connection one-form on HK_4 such that $\vartheta = dx + Tdy + \pi^* \alpha$ is a globally defined one-form on M_6 with

$$\frac{1}{2\pi} d\vartheta = \pi^* \varpi, \quad \varpi = \varpi_1 + T\varpi_2, \quad \varpi_i \in H^2(HK_4, \mathbb{Z}), \quad (2.6)$$

and by supersymmetry

$$J^a \wedge \varpi = 0, \quad a = 1, 2, 3. \quad (2.7)$$

The expression for the three-form becomes then

$$\mathcal{H}_{(3)} = - *_4 dH - \frac{\alpha' U_2}{T_2} \text{Re} (*_4 d\vartheta \wedge \bar{\vartheta}). \quad (2.8)$$

By an appropriate choice of $\varpi \in H^2(HK_4, \mathbb{Z})$ one can find solutions with $d\mathcal{H}_{(3)} = 0$, which can also be obtained as supersymmetric solutions of type IIA or type IIB supergravity with NS-NS fluxes, as was discussed in [3] and [18].

3 Gibbons-Hawking: ALE and ALF

We can solve explicitly (2.2) and (2.4) for the multicentered Gibbons-Hawking ALE and ALF spaces, that we denote collectively by M_{GH} . The corresponding Hyper-Kähler metrics are given by:¹

$$ds_4^2 = V(x)^{-1} (d\tau + \omega)^2 + V(x) dx \cdot dx, \quad (3.1a)$$

$$dV = *_3 d\omega, \quad (3.1b)$$

$$V = \epsilon + 2m \sum_{i=1}^k \frac{1}{|\mathbf{x} - \mathbf{x}_i|}, \quad (3.1c)$$

¹As usual $*_3$ is the Hodge dual on $\mathbb{R}^3$.

where $\epsilon = 0$ gives the ALE (multi Eguchi-Hanson) series and $\epsilon = 1$ the ALF (multi Taub-NUT) series. The periodicity of τ is determined by expanding around a pole of $V(\mathbf{x})$ to be:

$$\tau \sim \tau + 8\pi m, \quad (3.2)$$

and the triplet of Kähler forms is given by

$$J_a = (d\tau + \omega) \wedge dx^a - V *_3 dx^a, \quad a = 1, 2, 3. \quad (3.3)$$

We will consider heterotic supergravity solutions for warped ALE or ALF spaces supported by Abelian gauge bundles. To explicitly write the gauge fields we denote

$$V_i = \frac{2m}{|\mathbf{x} - \mathbf{x}_i|}, \quad d\omega_i = *_3 dV_i; \quad (3.4)$$

Then a representative of the topologically non-trivial gauge fields is locally given by

$$A_i = \omega_i - \frac{V_i}{V} (d\tau + \omega). \quad (3.5)$$

We note that

$$\sum_{i=1}^k A_i = \epsilon \frac{d\tau + \omega}{V} - d\tau \quad (3.6)$$

is topologically trivial since $V^{-1}(d\tau + \omega)$ is globally defined. Thus there are $(k-1)$ non-trivial gauge fields, in agreement with the $(k-1)$ non-trivial two-cycles. The corresponding field strengths are

$$F_i = dA_i = V *_3 d \left[\frac{V_i}{V} \right] - d \left[\frac{V_i}{V} \right] \wedge (d\tau + \omega). \quad (3.7)$$

It is straightforward to see that F_i are anti-self dual and solve Hermitian Yang-Mills²

$$*_4 F_i = -F_i, \quad J_a \wedge F_j = 0. \quad (3.8)$$

For the Bianchi identity (2.4) we compute

$$F_i \wedge F_j = -2V *_3 d \left[\frac{V_i}{V} \right] \wedge d \left[\frac{V_j}{V} \right] \wedge (d\tau + \omega) \quad (3.9)$$

$$d *_4 d \left[\frac{V_i V_j}{V} \right] = -F_i \wedge F_j \quad (3.10)$$

so that if we take

$$F = \frac{1}{4m} \sum_{i=1}^k dA_i \mathbf{q}_i \cdot \mathcal{T}, \quad (3.11)$$

²For a one form α on $\mathbb{R}^3$ we have $*_4 \alpha = -(d\tau + \omega) \wedge *_3 \alpha$. In particular this means that a function which is invariant under the $U(1)$ generated by ∂_τ is harmonic on the Gibbons-Hawking space iff it is harmonic on $\mathbb{R}^3$.

where $\mathcal{T} \in \text{U}(1)^{16}$ is in the Cartan subalgebra of $E_8 \times E_8$ or $\text{SO}(32)$ and $\mathbf{q}_i$ the corresponding charge vectors, and solve (2.4), we get the general solution:³

$$H = \delta + h(\mathbf{x}) + \frac{\alpha'}{8m^2V} \sum_{i,j=1}^k V_i V_j \mathbf{q}_i \cdot \mathbf{q}_j. \quad (3.12)$$

Here $\delta = 0, 1$ is an integration constant (not to be confused with ϵ a similar integration constant in the Gibbons-Hawking warp factor V) and $h(\mathbf{x})$ is any harmonic function on M_{GH} . Taking $h(\mathbf{x})$ to be invariant under ∂_τ we have

$$h(\mathbf{x}) = \frac{1}{m} \sum_{\alpha} \frac{q_{\alpha}}{|\mathbf{x} - \mathbf{x}_{\alpha}|} \quad (3.13)$$

corresponding to mobile neutral five-brane sources inserted at $\mathbf{x}_{\alpha}$.

In appendix A we show how the two center solution is related to the Eguchi-Hanson solution that was discussed in particular in [3].

3.1 Five-brane and magnetic charges

At infinity we can compute the five-brane charge using (2.2b) and (3.12). We have

$$\mathcal{H}_{(3)} = (d\tau + \omega) \wedge *_3 dH \quad (3.14)$$

and so⁴

$$dH = -\frac{\alpha'}{4mkr^2} \left(\sum_{i,j=1}^k \mathbf{q}_i \cdot \mathbf{q}_j \right) dr - \frac{1}{m} \sum_{\alpha} \frac{q_{\alpha}}{r^2} + \dots \quad (3.15a)$$

$$\mathcal{H}_{(3)} = \left[-\frac{\alpha'}{4mk} \sum_{i,j=1}^k \mathbf{q}_i \cdot \mathbf{q}_j - \frac{1}{m} \sum_{\alpha} q_{\alpha} \right] (d\tau + \omega) \wedge \Omega_2 + \dots \quad (3.15b)$$

and the Maxwell five-brane charge is

$$\mathcal{Q}_M = \frac{1}{4\pi^2 \alpha'} \int_{S^3/\mathbb{Z}_k} \mathcal{H}_{(3)} = -\frac{2}{k^2} \sum_{i,j=1}^k \mathbf{q}_i \cdot \mathbf{q}_j - \sum_{\alpha} \frac{8q_{\alpha}}{k}. \quad (3.16)$$

One can also define a Page charge, which is quantized, as:

$$\begin{aligned} \mathcal{Q}_P &= \frac{1}{4\pi^2 \alpha'} \int_{S^3/\mathbb{Z}_k} (\mathcal{H}_{(3)} - A \wedge F) \\ &= -\sum_{\alpha} \frac{8q_{\alpha}}{k} \in \mathbb{Z}. \end{aligned} \quad (3.17)$$

³we have chosen to work with Hermitian gauge fields, normalized as $\text{Tr } \mathcal{T}_{\alpha} \mathcal{T}_{\beta} = 2\delta_{\alpha\beta}$.

⁴The volume form of a three-sphere is

$$\begin{aligned} ds_{S^3}^2 &= \frac{1}{4} \left[\frac{1}{4m^2} (d\tau + \omega)^2 + d\Omega_2^2 \right] \\ 2\pi^2 &= \int \text{vol}(S^3) = \frac{1}{8} \int \frac{1}{2m} (d\tau + \omega) \wedge \Omega_2. \end{aligned}$$

Magnetic charges. We take a basis of two cycles to be Δ_i where the poles of the Δ_i are at x_i and x_{i+1} . Then the matrix of magnetic charges associated with the Abelian gauge bundle are:

$$\begin{aligned} q_{i,j} &= \frac{1}{2\pi} \int_{\Delta_j} F_i = \frac{\mathbf{q}_i \cdot \mathcal{T}}{8\pi m} \left(\int d\tau \right) \int_{x_j}^{x_{j+1}} d \left[\frac{V_i}{V} \right] \\ &= \mathbf{q}_i \cdot \mathcal{T} [\delta_{j+1,i} - \delta_{j,i}] . \end{aligned} \quad (3.18)$$

3.2 Partial blow-down limits and fivebranes

The function $V(\mathbf{x})$ has k -poles, now suppose that k' of these poles are co-incident, which correspond to a partial blow-down limit of the ALE or ALF space. In the present situation some of the Abelian instantons (3.7) become point-like as the corresponding two-cycles shrink and we expect heterotic five-brane to appear. We now check that in the region around such a pole we obtain the near horizon of five-brane solution of Callan, Harvey and Strominger [19, 20] where the three-sphere is orbifolded by $\mathbb{Z}_{k'}$.

For simplicity we set $\mathbf{x}_j = 0$ for $j = 1, \dots, k'$, and in the neighborhood of this pole the functions H and V behaves like

$$H \stackrel{r \rightarrow 0^+}{\simeq} \frac{1}{r^2} \frac{\alpha'}{2k'm} Q_5 \quad , \quad V \stackrel{r \rightarrow 0^+}{\simeq} \frac{4mk'}{r^2} \quad (3.19)$$

hence the solution approaches

$$ds_{10}^2 \stackrel{r \rightarrow 0^+}{\simeq} ds_{1,5}^2 + 2\alpha' Q_5 \left[\frac{dr^2}{r^2} + \frac{r^2}{4} \left(\sigma_1^2 + \sigma_3^2 + \left(\frac{\sigma_3}{2k'm} \right)^2 \right) \right] \quad (3.20a)$$

$$\mathcal{H}_{(3)} \stackrel{r \rightarrow 0^+}{\simeq} \frac{\alpha' Q_5}{2} \sigma_1 \wedge \sigma_2 \wedge \frac{\sigma_3}{2k'm} \quad (3.20b)$$

where the five-brane charge is given by:

$$Q_5 = \sum_{i,j=1}^{k'} \mathbf{q}_i \cdot \mathbf{q}_j . \quad (3.21)$$

3.3 Double scaling limit

For the two-center Eguchi-Hanson solution ($k = 2$) there exists an interesting double scaling limit [3], defined as:

$$g_s \rightarrow 0, \quad \lambda := \frac{g_s \sqrt{\alpha'}}{a} \text{ fixed and finite} , \quad (3.22)$$

where a is the distance between the two centers. This limit decouples the asymptotically locally Euclidian region, and λ becomes the effective coupling constant of the interacting string theory.

In the spherical coordinates reviewed in appendix A the metric of the solution becomes

$$ds^2 = ds_{1,5} + \frac{\alpha' Q_5}{2} \left[\frac{dr^2}{r^2(1 - \frac{a^4}{r^2})} + \frac{1}{4} \left(1 - \frac{a^4}{r^2} \right) \sigma_3^2 + d\Omega_2^2 \right] \quad (3.23)$$

and the corresponding heterotic string theory admits an exactly solvable worldsheet CFT. This space has an asymptotic linear dilaton hence admits a holographic description as a little string theory [21] but, unlike the CHS background, is given by a smooth solution of heterotic supergravity.

A double scaling limit can be described in principle for arbitrary k . Let us define $\mathbf{x}_i = a\mathbf{y}_i$ where the coordinates $\mathbf{y}_i$ are dimension-less and a is a common scale factor. The double-scaling limit can be then described exactly as before by eq. (3.22); in practice the double scaling limit amounts to setting $\delta \rightarrow 0$ in (3.12).

It would be interesting to check if one could derive a worldsheet CFT for the double scaled solutions when $k > 2$, in particular whenever the centers are arranged following a simple pattern, for instance a homogeneous distribution on a circle.

4 Atiyah-Hitchin

The Atiyah-Hitchin space M_{AH} is a four-dimensional smooth manifold with an explicit Hyper-Kähler metric which at long distances approximates Taub-NUT with a negative mass parameter. The original work is [2, 22] and an interesting simplification was given in [23]. Our notation will follow a more recent work [10] where M_{AH} was used as a potential Euclidean Hyper-Kähler base manifold for five dimensional supergravity solutions. In [10] regularity required the absence of closed time-like curves and this effectively excluded physical solutions whereas for our computations the non-trivial regularity conditions are essentially just positivity of the warp factor and we will find regular solutions.

The metric is

$$ds_{AH}^2 = \frac{1}{4}a_1^2a_2^2a_3^2d\eta^2 + \frac{1}{4}a_1^2\sigma_1^2 + \frac{1}{4}a_2^2\sigma_2^2 + \frac{1}{4}a_3^2\sigma_3^2 \quad (4.1)$$

with the $SU(2)$ invariant one-forms satisfying $d\sigma_i = \frac{1}{2}\epsilon_{ijk}\sigma_j \wedge \sigma_k$ given by

$$\begin{aligned} \sigma_1 &= \cos\psi d\theta + \sin\psi \sin\theta d\phi \\ \sigma_2 &= \sin\psi d\theta - \cos\psi \sin\theta d\phi \\ \sigma_3 &= d\psi + \cos\theta d\phi \end{aligned}$$

and the a_i are subject to the following system of ODE's:

$$\frac{\dot{a}_1}{a_1} = \frac{1}{2} [(a_2 - a_3)^2 - a_1^2] \quad (4.2)$$

and cyclic permutations (dot is the derivative with respect to η). One defines new functions quadratic in the a_i :

$$w_1 = a_2a_3, \quad w_2 = a_1a_3, \quad w_3 = a_1a_2 \quad (4.3)$$

and then the system of ODE's is then

$$(w_1 + w_2)' = -2 \frac{w_1 w_2}{u^2} \quad (4.4)$$

$$(w_2 + w_3)' = -2 \frac{w_2 w_3}{u^2} \quad (4.5)$$

$$(w_3 + w_1)' = -2 \frac{w_3 w_1}{u^2} \quad (4.6)$$

(prime is derivative with respect to θ) with solution

$$w_1 = -uu' - \frac{1}{2}u^2 \csc \theta \quad (4.7)$$

$$w_2 = -uu' + \frac{1}{2}u^2 \cot \theta \quad (4.8)$$

$$w_3 = -uu' + \frac{1}{2}u^2 \csc \theta, \quad (4.9)$$

where

$$u = \frac{1}{\pi} \sqrt{\sin \theta} K \left(\sin^2 \frac{\theta}{2} \right) \quad (4.10)$$

and η is given in terms of θ through

$$u^2 d\eta = d\theta, \quad \eta = - \int_{\theta}^{\pi} \frac{d\theta}{u^2}. \quad (4.11)$$

For our gauge field ansatz we need an anti-self dual two form on M_{AH} , this is then guaranteed to solve Hermitian Yang-Mills without the need to construct the explicit Hyper-Kähler structure.⁵ In a classic paper on dualities [8], Sen gave an integral expression for exactly such an anti-self dual, harmonic two-form on M_{AH} but the appearance of this two-form dates back to the works [25–27]. Interestingly, from the work [10] we have the closed-form expression of this two-form

$$\Omega = h (a_1^2 dr \wedge \sigma_1 - \sigma_2 \wedge \sigma_3), \quad (4.12)$$

$$h = \frac{u^2}{w_1 \sin \frac{\theta}{2}}. \quad (4.13)$$

In [10] they consider self-dual forms but with a small modification of the frames this is made anti-self dual. More precisely our choice of frames is

$$e_0 = \frac{a_1 a_2 a_3}{2} d\eta, \quad e_i = \frac{a_i}{2} \sigma_i, \quad (4.14)$$

whereas in [10] an additional minus sign in e_0 was used. So we have locally

$$\Omega = -d(h \sigma_1). \quad (4.15)$$

In fact one can construct a triplet of anti-self dual forms Ω_- and a triplet of self-dual forms Ω_+ in a similar manner:

$$\Omega_{i-} = -d(h_i \sigma_i), \quad \Omega_{i+} = d(h_i^{-1} \sigma_i) \quad (4.16)$$

with

$$h_1 = \frac{u^2}{w_1 \sin \frac{\theta}{2}}, \quad h_2 = \frac{u^2}{w_2}, \quad h_3 = \frac{u^2}{w_3 \cos \frac{\theta}{2}}, \quad (4.17)$$

however only $\Omega_{1-} = \Omega$ is normalizable. Given that there is a single non-trivial two-cycle in M_{AH} one might be pleased to know that this normalizable form is dual to this two-cycle but there was no guarantee that the dual two-form would have an $SU(2)$ invariant representative.

⁵One could in principle write down the Hyper-Kähler structure using the results of [24] or by computing the Killing spinors.

4.1 Bianchi identity

We take our gauge field to be

$$F = \Omega \mathbf{q} \cdot \mathcal{H} \quad (4.18)$$

where $\mathcal{H} \in \mathrm{U}(1)^{16}$ is in the Cartan subalgebra of $E_8 \times E_8$ or $\mathrm{SO}(32)$ and $\mathbf{q}$ the corresponding charge vector. The three-form flux is

$$\mathcal{H}_{(3)} = -\frac{H'}{4} \sigma^1 \wedge \sigma^2 \wedge \sigma^3 \quad (4.19)$$

and the Bianchi identity is

$$d *_4 dH = -2q^2 \Omega \wedge \Omega, \quad (4.20)$$

where $q^2 = \mathbf{q} \cdot \mathbf{q}$.

Quite remarkably, one can integrate this Poisson equation explicitly

$$H = h_0 + h_1 \eta + \frac{2q^2}{w_1} \quad (4.21)$$

where $\{h_0, h_1\}$ are constant coefficients of the s-wave harmonic functions on M_{AH} . The last term is manifestly negative definite for the whole region $0 \leq \theta \leq \pi$ but we will see that one can compensate for this by a choice of harmonic function and obtain a positive definite warpfactor.

4.2 Regularity

The regularity of M_{AH} has been previously studied in detail, we repeat it here to help determine regularity of our warp factor.

In the region $\theta \sim \pi$, we define a radial co-ordinate $r = -\log \cos \frac{\theta}{2}$ and using

$$K = r + \log(4) + \dots \quad (4.22)$$

$$u = \frac{\sqrt{2}}{\pi} r e^{-r/2} + \dots \quad (4.23)$$

$$w_1 = -\frac{r}{\pi^2} \quad (4.24)$$

we find that the metric is

$$ds_{AH}^2 = dr^2 + r^2(\sigma_1^2 + \sigma_2^2) + \sigma_3^2 + \dots \quad (4.25)$$

and

$$\frac{1}{w_1} = -\frac{\pi^2}{r} + \mathcal{O}(r^{-2}) \quad (4.26)$$

$$\eta = -\frac{\pi^2}{r} + \frac{\pi \log(4)}{r^2} + \mathcal{O}(r^{-3}) \quad (4.27)$$

so that the asymptotic expansion of the warp factor is

$$H = h_0 - \frac{h_1 + 2q^2}{r} + \dots \quad (4.28)$$

In the region $\theta \sim 0$, we define a new radial variable $\rho = \frac{\theta^2}{64}$ and the metric is

$$ds_{AH}^2 = d\rho^2 + 4\rho^2\sigma_1^2 + \frac{1}{16}(\sigma_2^2 + \sigma_3^2) + \dots \quad (4.29)$$

with

$$\frac{1}{w_1} = -4 + 32\rho^2 + \dots \quad (4.30)$$

$$\eta = \log \rho^2 + \dots \quad (4.31)$$

so that the IR expansion of the warp factor is

$$H = (h_0 - 8q^2) + 64q^2\rho^2 + \dots \quad (4.32)$$

From these expansions we see that with

$$h_0 > 8q^2, \quad h_1 = 0 \quad (4.33)$$

we have a positive warp factor which is regular everywhere. We define a rescaled radial coordinate near $\theta \sim \pi$ to be $\hat{r} = h_0^{1/2}r$ and $\hat{\rho} = h_0^{1/2}\rho$ near $\theta = 0$ so that

$$\theta \sim \pi : \quad ds_{10}^2 = ds_{1,5}^2 + d\hat{r}^2 + \hat{r}^2(\sigma_1^2 + \sigma_2^2) + h_0\sigma_3^2 + \dots \quad (4.34)$$

$$\theta \sim 0 : \quad ds_{10}^2 = ds_{1,5}^2 + d\hat{\rho}^2 + 4\hat{\rho}^2\sigma_1^2 + \frac{h_0}{16}(\sigma_2^2 + \sigma_3^2) + \dots \quad (4.35)$$

and see that the cost of a positive warp factor is that both the circle at infinity and the two-sphere at the bolt are large.

It is also important that the $\text{Tr}R_- \wedge R_-$ term in the Bianchi identity remains small compared to $\text{Tr}F \wedge F$. From explicit computations we find that the only possible divergences in $\text{Tr}R_- \wedge R_-$ appear through the warp factor as⁶ $\{H'/H, H''/H\}$ which by tuning h_0 can be made sufficiently small with respect to $\text{Tr}F \wedge F$. This confirms that our large charge approximation remains valid and these warped Atiyah-Hitchin solutions are good Heterotic backgrounds at leading order.

Alternatively we can obtain a positive warp factor through⁷

$$h_0 = 1, \quad h_1 < -2q^2. \quad (4.36)$$

This corresponds to smearing neutral five-branes on the S^2 at $\theta = 0$. Note that due to this smearing, at the IR ($\theta = 0$) the harmonic function parameterized by h_1 scales like a source in $\mathbb{R}^2$. In the UV ($\theta = \pi$), due to the finite circle, the harmonic function scales like $\frac{1}{r}$ which is that of a source in $\mathbb{R}^3$ not $\mathbb{R}^4$. The solution is of course singular for the usual reason that smeared branes are singular but this is of a good type and is resolved in string theory.

⁶An explicit computation using the Chern connection can be found in [14] and agrees with our conclusion here.

⁷The value of h_0 could be chosen to be another non-zero number $\mathcal{O}(q^0)$.

4.3 Five-brane charge

Computing the five-brane charge requires understanding some global features of M_{AH} . From (4.25) and (4.29) we see that there are two inequivalent, emergent $U(1)$ symmetries in the UV and IR, which are broken in the bulk. From [23] we know that a regular manifold requires the periodicities to be

$$0 \leq \psi \leq 2\pi, \quad 0 \leq \theta \leq \pi, \quad 0 \leq \phi \leq 2\pi \quad (4.37)$$

as well as that the free $\mathbb{Z}_2$ symmetry

$$I_1: \quad \theta \rightarrow \pi - \theta, \quad \phi \rightarrow \pi + \phi, \quad \psi \rightarrow -\psi \quad (4.38)$$

is enforced. The horizontal space in the UV is thus $\mathbb{RP}^3/I_1$.

Using (4.28) we have

$$\mathcal{H}_{(3)} = [(h_1 + 2q^2) + \dots] \wedge \sigma^1 \wedge \sigma^2 \wedge \sigma^3 \quad (4.39)$$

compute the Maxwell five-brane charge to be

$$\mathcal{Q}_M = \frac{1}{4\pi\alpha'} \int_{\mathbb{RP}^3/I_1} \mathcal{H}_{(3)} = h_1 + 2q^2. \quad (4.40)$$

This is not required to be quantized. The Page charge is defined as in (3.17) and we find

$$\mathcal{Q}_P = h_1 \quad (4.41)$$

which must be integral.

4.4 Gauge field charge

The gauge field charge is computed using (4.12) and (4.18) and the IR expansion

$$h = -2 + \dots \quad (4.42)$$

Under the symmetry (4.38), the bolt remains a two sphere⁸ whose volume is 4π . We find

$$\begin{aligned} \frac{1}{2\pi} \int_{S^2} F &= 2\mathbf{q} \cdot \mathcal{H} \frac{1}{2\pi} \int_{S^2} \sigma_2 \wedge \sigma_3 \\ &= 4\mathbf{q} \cdot \mathcal{H} \in \mathbb{Z}. \end{aligned} \quad (4.43)$$

5 Fundamental string sources and AdS_3 solutions

Heterotic backgrounds with an $\mathbb{R}^{1,1}$ factor allow for the inclusion of F1-strings along $\mathbb{R}^{1,1}$ in addition to the magnetic five-branes. The electric source of three form flux induces a non-trivial warp factor and allows for AdS_3 solutions. To include these fundamental strings,

⁸As explained in [23] there is an additional, optional $\mathbb{Z}_2$ symmetry usually denoted I_3 which would convert the bolt into an $\mathbb{RP}^3$.

we first consider internal eight-manifolds X_8 and then specialize the internal manifold to be a product of Hyper-Kähler manifolds.

The metric and three form are

$$ds_{10}^2 = e^{2A} ds_{1,1}^2 + ds_8^2 \quad (5.1)$$

$$H^{(3)} = \text{vol}_2 \wedge h^{(1)} + h^{(3)} \quad (5.2)$$

where $\text{vol}_2 = e^{2A} dx^0 \wedge dx^1$. Then we find that the BPS equations are a slight embellishment of those found in [28]:

$$\Psi \lrcorner d\Psi = d(\phi - A) \quad (5.3)$$

$$H_{(3)} = h_{(1)} \wedge \text{vol}_2 + h_{(3)} \quad (5.4)$$

$$h_{(1)} = -2dA \quad (5.5)$$

$$h_{(3)} = *_8 e^{2(\phi-A)} d \left(e^{2(A-\phi)} \Psi \right) \quad (5.6)$$

where Ψ is the Spin(8) structure on M_8 .⁹

$$\begin{aligned} \Psi = & e^{1234} + e^{1256} + e^{1278} + e^{3456} + e^{3478} + e^{5678} \\ & + e^{1357} - e^{1368} - e^{1458} - e^{1467} - e^{2358} - e^{2367} - e^{2457} + e^{2468} . \end{aligned} \quad (5.8)$$

We must supplement the BPS equations with the Bianchi identity (2.3) and then due to the non-trivial warp-factor A , one must also impose the three form flux equation of motion:

$$0 = d \left(e^{-2\phi} *_8 H_{(3)} \right) \quad \Rightarrow \quad \begin{cases} 0 = d \left(e^{-2\phi} *_8 h_{(1)} \right) \\ 0 = d \left(e^{2(A-\phi)} *_8 h_{(3)} \right) \end{cases} . \quad (5.9)$$

5.1 Product of Hyper-Kähler manifolds

Our solutions with string and five-brane charges have a natural splitting of the internal eight manifold into a product of Hyper-Kähler manifolds¹⁰

$$ds_{10}^2 = e^{2A} ds_{\mathbb{R}^{1,1}}^2 + ds_{M_1}^2 + e^{2B} ds_{M_2}^2 \quad (5.10)$$

where M_i are both HyperKähler four manifolds, whose triplet of Kähler forms we denote

$$\{J_i, \text{Re } \Omega_i, \text{Im } \Omega_i\} . \quad (5.11)$$

The functions A, B depend only on the co-ordinates y_i of M_2 . The Spin(8) structure is given by

$$\Psi = \frac{1}{2} \left(J_1 \wedge J_1 + 2e^{2B} J_1 \wedge J_2 + e^{4B} J_2 \wedge J_2 + e^{2B} (\Omega_1 \wedge \Omega_2 + \bar{\Omega}_1 \wedge \bar{\Omega}_2) \right) . \quad (5.12)$$

⁹We note that with canonical holomorphic frames $E_i = e_{2i-1} + ie_{2i}$ such that $ds_8^2 = E_i \otimes \bar{E}_i$ the SU(4) structure is $J = \frac{1}{2} E_i \wedge \bar{E}_i$, $\Omega = E_1 \wedge E_2 \wedge E_3 \wedge E_4$ and

$$\Psi = \frac{1}{2} (J \wedge J + \Omega + \bar{\Omega}) . \quad (5.7)$$

¹⁰One might consider an additional warp factor in front of $ds_{M_1}^2$ however from [29, 30] we know that this must be constant.

We find the BPS conditions, Bianchi identity and equations of motion give¹¹

$$\phi = A + B \quad (5.13)$$

$$h_{(3)} = - *_{M_2} d e^{2B} \quad (5.14)$$

$$d *_{M_2} d e^{2B} = -\frac{1}{2} \alpha' \text{Tr} F \wedge F \quad (5.15)$$

$$0 = d *_{M_2} d e^{-2A} \quad (5.16)$$

$$0 = J_2 \lrcorner F \quad (5.17)$$

so we see that the only additional pieces of data from the equations in section 2 is that e^{2A} is harmonic on M_2 and the dilaton receives a shift proportional to A . For the Atiyah-Hitchin manifold we can smear F1-strings on the S^2 bolt in much the same way as we have described for smearing 5-branes on the bolt around (4.36), that is by

$$e^{2A} \sim \eta. \quad (5.18)$$

We will now be somewhat more explicit for the Gibbons-Hawking spaces.

5.2 AdS₃ from Gibbons-Hawking

When M_2 is a Gibbons-Hawking space, the U(1) invariant harmonic functions are

$$e^{-2A} = 1 + \sum_r \frac{\hat{q}_r}{|\mathbf{x} - \mathbf{x}_r|} \quad (5.19)$$

corresponding to strings placed along $\mathbb{R}^{1,1}$ and at fixed points of ∂_τ on M_2 .

If in addition we choose to place these strings at poles of V we recover AdS₃ geometries near such a pole. We put k' poles of V as well as the strings at $\mathbf{x}_r = \mathbf{x}_i = 0$ then in the vicinity of x_i we have

$$e^{2A} = \frac{r}{\hat{q}_0}, \dots \quad e^{2B} = \frac{1}{r} \frac{\alpha'}{4m} Q_5 + \dots, \quad V = \frac{2mk'}{r} + \dots \quad (5.20)$$

$$(5.21)$$

so that

$$ds_{10}^2 = \frac{r}{\hat{q}_0} ds_{1,1}^2 + ds_{M_1}^2 + 2\alpha' k'^2 Q_5 \left[\frac{1}{4} \frac{dr^2}{r^2} + ds_{S^3/\mathbb{Z}_k}^2 \right] \quad (5.22)$$

$$= 2\alpha' k'^2 Q_5 \left[ds_{AdS_3}^2 + ds_{S^3/\mathbb{Z}_k}^2 \right] + ds_{M_1}^2 \quad (5.23)$$

$$e^{2\phi} = \frac{\alpha' Q_5}{4m\hat{q}_0} \quad (5.24)$$

where $r = \rho^2$. The F1-charge is given as usual by

$$Q_1 = \frac{4m\hat{q}_0 \text{vol}(M_1)}{\alpha'^3}. \quad (5.25)$$

The gauge field vanishes in this limit and the background is sourced by three-form flux.

¹¹Note that is $E = e^B \tilde{E}$ is an 8d frame $*_8 E \wedge J_1 \wedge J_1 = 2 *_8 E \wedge \text{vol}_{M_1} = 2e^{3B} *_8 \tilde{E} = 2e^{2B} *_8 E$.

6 Conclusions

The key aspect of our solutions with Abelian gauge bundles is that we have taken a large charge limit and consistently suppressed the $\text{Tr } R \wedge R$ term in the Bianchi identity, which is subdominant at leading order in the expansion in $\frac{1}{q^2}$. We have shown how this large charge limit can lead to exact supersymmetric flux backgrounds, and it is particularly interesting the Atiyah-Hitchin manifold can provide a regular background. This configuration requires some ingenuity to counteract the negative mass and result in a background of the correct signature. This Atiyah-Hitchin based solution is distinctly different from those based on Gibbons-Hawking; while the latter can be viewed as marginal deformations of the orbifold of the CHS solutions the finite two-cycle in the Atiyah-Hitchin manifold cannot be blown down. As such we do not have a worldsheet theory from which we can imagine obtaining this as the background geometry.

In these backgrounds, the gauge fields are completely solved for by using the Hermitian Yang-Mills equations which then provide a source for the three-form flux. It is conceivable that non-Abelian bundles could be constructed such that $\text{Tr } F \wedge F$ dominates $\text{Tr } R \wedge R$ everywhere.¹² Since the Kronheimer-Nakajima construction [33] gives a solution of all instantons on ALE Gibbons-Hawking spaces, one could possibly even construct such instantons, however most instantons will provide a source the Bianchi identity whose solution is a general function of four variables and thus unsolvable. A particularly neat class of instantons is based on the 't Hooft ansatz [34]:

$$\begin{aligned} A_0 &= \frac{1}{2} \vec{G} \cdot \vec{\sigma}, & \vec{A} &= \frac{1}{2} \left[\vec{\omega}(\vec{G} \cdot \vec{\sigma}) - V(\vec{G} \times \vec{\sigma}) \right] \\ \vec{G} &= -V^{-1} \vec{\nabla} \log f \end{aligned} \tag{6.1}$$

with f harmonic on $\mathbb{R}^3$. For finite action, the centers of f are constrained to lie at the poles of V . These instantons can have large $\text{Tr } F \wedge F$ in the limit of large number of poles of f but $\text{Tr } R \wedge R$ will not be suppressed.

There are numerous directions for progress on the worldsheet description of these backgrounds. The elliptic genus for type II on ALE spaces has been computed recently [35] based on general developments in this field [36] and we expect to be able to provide a similar solution for these Heterotic models or the type II flux backgrounds of section 2.1. It would also be interesting to provide an exactly solvable worldsheet model of the near-horizon region of the multi-center Gibbons-Hawking backgrounds, generalizing the gauged WZW model of the two-centered solution.

Acknowledgments

N.H. would like to thank Nikolay Bobev and Davide Cassani for discussions. E.S. would like to thank Bobby Acharya for insightful conversations. This work was conducted within the

¹²Interesting five dimensional solutions with non-Abelian gauge fields have appeared recently [31] and the lift to the Heterotic string has been discussed [32]. However it is not clear to us how these solutions will solve the exact Bianchi identity.

ILP LABEX (ANR-10-LABX-63) supported by French state funds managed by the ANR within the *Investissements d’Avenir* program (ANR-11-IDEX-0004-02), by the project QHNS in the program ANR Blanc SIMI5 of Agence National de la Recherche and the CEFIPRA grant 5204-4.

A Eguchi-Hanson

When $k = 2$ and $\epsilon = 0$, the explicit co-ordinate transformation is known [37] from the Gibbons-Hawking space to the Eguchi-Hanson space [38]. In Cartesian co-ordinates the two center ALE Gibbons-Hawking space has

$$\omega = \left[\frac{z - a^2/8}{\sqrt{x^2 + y^2 + (z - a^2/8)^2}} + \frac{z + a^2/8}{\sqrt{x^2 + y^2 + (z + a^2/8)^2}} \right] d\left(\tan^{-1} \frac{y}{x}\right) \quad (\text{A.1})$$

$$V = \frac{1}{\sqrt{x^2 + y^2 + (z - a^2/8)^2}} + \frac{1}{\sqrt{x^2 + y^2 + (z + a^2/8)^2}}. \quad (\text{A.2})$$

Following [37] we have ($a \leq r$):

$$\begin{aligned} x &= \frac{a^2}{8} \sqrt{\frac{r^4}{a^4} - 1} \sin \theta \cos \psi \\ y &= \frac{a^2}{8} \sqrt{\frac{r^4}{a^4} - 1} \sin \theta \sin \psi \\ z &= \frac{1}{8} r^2 \cos \theta \end{aligned}$$

so that

$$\begin{aligned} V &= \frac{16}{a^2} \frac{\frac{r^2}{a^2}}{\frac{r^4}{a^4} - \cos^2 \theta}, \\ \omega &= \frac{2 \cos \theta (\frac{r^2}{a^2} - 1)}{\frac{r^2}{a^2} - \cos \theta} d\psi. \end{aligned} \quad (\text{A.3})$$

As an example, we write explicitly the solution for Heterotic five-branes on Eguchi-Hanson with additional F1-strings.¹³

$$ds_{M_2}^2 = \frac{dr^2}{f^2} + \frac{r^2}{4} [\sigma_1^2 + \sigma_2^2 + f^2 \sigma_3^2] \quad (\text{A.4})$$

$$f^2 = 1 - \frac{a^4}{r^4} \quad (\text{A.5})$$

$$h_{(3)} = 2f^2 r^3 (e^{2B})' \sigma_1 \wedge \sigma_2 \wedge \sigma_3$$

$$F = d\left(\frac{a^2}{r^2} \eta\right) \quad (\text{A.6})$$

$$e^{2B} = 1 + \frac{8\alpha' q^2}{r^2} + \frac{Q_5}{8a^2} \log \left[\frac{r^2/a^2 - 1}{r^2/a^2 + 1} \right] \quad (\text{A.7})$$

¹³One can take M_1 to be T^4 or $K3$ with the Ricci-flat metric.

$$e^{-2A} = 1 + \frac{Q_1}{a^2} \log \left[\frac{r^2/a^2 - 1}{r^2/a^2 + 1} \right] \quad (\text{A.8})$$

$$e^{2\Phi} = e^{2(A+B)} \quad (\text{A.9})$$

In addition to the Heterotic five-branes which resolve the singularity, there are Q_1 mobile F1-strings and Q_5 NS5-branes smeared on the blown-up S^2 . Due the smearing of the strings, the near horizon limit has a log-singularity at $r \sim a$ in the warp factor e^{2A} and thus there is no enhancement to AdS_3 . In the blow-down limit $a \rightarrow 0$ where the Eguchi-Hanson space becomes $\mathbb{C}^2/\mathbb{Z}_2$, the gauge field vanishes and we get the $\mathbb{Z}_2$ orbifold of the usual F1-NS5-solution, the near-horizon limit is $\text{AdS}_3 \times S^3/\mathbb{Z}_2 \times M_1$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] G.W. Gibbons and S.W. Hawking, *Gravitational Multi-Instantons*, *Phys. Lett. B* **78** (1978) 430 [[INSPIRE](#)].
- [2] M.F. Atiyah and N.J. Hitchin, *Low-Energy Scattering of Nonabelian Monopoles*, *Phys. Lett. A* **107** (1985) 21 [[INSPIRE](#)].
- [3] L. Carlevaro, D. Israël and P.M. Petropoulos, *Double-Scaling Limit of Heterotic Bundles and Dynamical Deformation in CFT*, *Nucl. Phys. B* **827** (2010) 503 [[arXiv:0812.3391](#)] [[INSPIRE](#)].
- [4] L. Carlevaro and D. Israël, *Heterotic Resolved Conifolds with Torsion, from Supergravity to CFT*, *JHEP* **01** (2010) 083 [[arXiv:0910.3190](#)] [[INSPIRE](#)].
- [5] N. Halmagyi, D. Israël and E.E. Svanes, *The Abelian Heterotic Conifold*, *JHEP* **07** (2016) 029 [[arXiv:1601.07561](#)] [[INSPIRE](#)].
- [6] J.M. Maldacena, *The large- N limit of superconformal field theories and supergravity*, *Int. J. Theor. Phys.* **38** (1999) 1113 [[hep-th/9711200](#)] [[INSPIRE](#)].
- [7] P.S. Aspinwall, *Compactification, geometry and duality: $N = 2$* , [hep-th/0001001](#) [[INSPIRE](#)].
- [8] A. Sen, *Dyon - monopole bound states, selfdual harmonic forms on the multi - monopole moduli space and $\text{SL}(2, \mathbb{Z})$ invariance in string theory*, *Phys. Lett. B* **329** (1994) 217 [[hep-th/9402032](#)] [[INSPIRE](#)].
- [9] I. Bena and N.P. Warner, *Bubbling supertubes and foaming black holes*, *Phys. Rev. D* **74** (2006) 066001 [[hep-th/0505166](#)] [[INSPIRE](#)].
- [10] I. Bena, N. Bobev and N.P. Warner, *Bubbles on Manifolds with a $U(1)$ Isometry*, *JHEP* **08** (2007) 004 [[arXiv:0705.3641](#)] [[INSPIRE](#)].
- [11] R. Minasian, M. Petrini and A. Zaffaroni, *New families of interpolating type IIB backgrounds*, *JHEP* **04** (2010) 080 [[arXiv:0907.5147](#)] [[INSPIRE](#)].
- [12] K. Dasgupta, G. Rajesh and S. Sethi, *M theory, orientifolds and G-flux*, *JHEP* **08** (1999) 023 [[hep-th/9908088](#)] [[INSPIRE](#)].

- [13] E. Goldstein and S. Prokushkin, *Geometric model for complex nonKähler manifolds with SU(3) structure*, *Commun. Math. Phys.* **251** (2004) 65 [[hep-th/0212307](#)] [[INSPIRE](#)].
- [14] J.-X. Fu and S.-T. Yau, *The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampere equation*, *J. Diff. Geom.* **78** (2008) 369 [[hep-th/0604063](#)] [[INSPIRE](#)].
- [15] K. Becker, M. Becker, J.-X. Fu, L.-S. Tseng and S.-T. Yau, *Anomaly cancellation and smooth non-Kähler solutions in heterotic string theory*, *Nucl. Phys. B* **751** (2006) 108 [[hep-th/0604137](#)] [[INSPIRE](#)].
- [16] K. Becker and S. Sethi, *Torsional Heterotic Geometries*, *Nucl. Phys. B* **820** (2009) 1 [[arXiv:0903.3769](#)] [[INSPIRE](#)].
- [17] J.-X. Fu, L.-S. Tseng and S.-T. Yau, *Local Heterotic Torsional Models*, *Commun. Math. Phys.* **289** (2009) 1151 [[arXiv:0806.2392](#)] [[INSPIRE](#)].
- [18] D. Martelli and J. Sparks, *Non-Kähler heterotic rotations*, *Adv. Theor. Math. Phys.* **15** (2011) 131 [[arXiv:1010.4031](#)] [[INSPIRE](#)].
- [19] C.G. Callan Jr., J.A. Harvey and A. Strominger, *Worldbrane actions for string solitons*, *Nucl. Phys. B* **367** (1991) 60 [[INSPIRE](#)].
- [20] C.G. Callan Jr., J.A. Harvey and A. Strominger, *Supersymmetric string solitons*, [hep-th/9112030](#) [[INSPIRE](#)].
- [21] O. Aharony, M. Berkooz, D. Kutasov and N. Seiberg, *Linear dilatons, NS five-branes and holography*, *JHEP* **10** (1998) 004 [[hep-th/9808149](#)] [[INSPIRE](#)].
- [22] M.F. Atiyah and N.J. Hitchin, *Low-energy scattering of nonAbelian magnetic monopoles*, *Phil. Trans. Roy. Soc. Lond. A* **315** (1985) 459 [[INSPIRE](#)].
- [23] G.W. Gibbons and N.S. Manton, *Classical and Quantum Dynamics of BPS Monopoles*, *Nucl. Phys. B* **274** (1986) 183 [[INSPIRE](#)].
- [24] D. Olivier, *Complex coordinates and Kähler potential for the Atiyah-Hitchin metric*, *Gen. Rel. Grav.* **23** (1991) 1349 [[INSPIRE](#)].
- [25] N.S. Manton, *Monopole and Skyrmion Bound States*, *Phys. Lett. B* **198** (1987) 226 [[INSPIRE](#)].
- [26] G.W. Gibbons and P.J. Ruback, *The Hidden Symmetries of Multicenter Metrics*, *Commun. Math. Phys.* **115** (1988) 267 [[INSPIRE](#)].
- [27] N.S. Manton and B.J. Schroers, *Bundles over moduli spaces and the quantization of BPS monopoles*, *Annals Phys.* **225** (1993) 290 [[INSPIRE](#)].
- [28] J.P. Gauntlett, D. Martelli and D. Waldram, *Superstrings with intrinsic torsion*, *Phys. Rev. D* **69** (2004) 086002 [[hep-th/0302158](#)] [[INSPIRE](#)].
- [29] A. Strominger, *Superstrings with Torsion*, *Nucl. Phys. B* **274** (1986) 253 [[INSPIRE](#)].
- [30] C.M. Hull, *Compactifications of the Heterotic Superstring*, *Phys. Lett. B* **178** (1986) 357 [[INSPIRE](#)].
- [31] P. Meessen, T. Ortín and P.F. Ramirez, *Non-Abelian, supersymmetric black holes and strings in 5 dimensions*, *JHEP* **03** (2016) 112 [[arXiv:1512.07131](#)] [[INSPIRE](#)].
- [32] P.A. Cano, P. Meessen, T. Ortín and P.F. Ramirez, *Non-Abelian black holes in string theory*, [arXiv:1704.01134](#) [[INSPIRE](#)].

- [33] P.B. Kronheimer and H. Nakajima, *Yang-mills instantons on ale gravitational instantons*, *Math. Ann.* **288** (1990) 263.
- [34] H. Boutaleb-Joutei, A. Chakrabarti and A. Comtet, *Gauge Field Configurations in Curved Space-times. 4. Selfdual SU(2) Fields in Multicenter Spaces*, *Phys. Rev. D* **21** (1980) 2280 [[INSPIRE](#)].
- [35] J.A. Harvey, S. Lee and S. Murthy, *Elliptic genera of ALE and ALF manifolds from gauged linear σ -models*, *JHEP* **02** (2015) 110 [[arXiv:1406.6342](#)] [[INSPIRE](#)].
- [36] F. Benini, R. Eager, K. Hori and Y. Tachikawa, *Elliptic Genera of 2d $\mathcal{N} = 2$ Gauge Theories*, *Commun. Math. Phys.* **333** (2015) 1241 [[arXiv:1308.4896](#)] [[INSPIRE](#)].
- [37] M.K. Prasad, *Equivalence of Eguchi-Hanson metric to two-center Gibbons-Hawking metric*, *Phys. Lett. B* **83** (1979) 310 [[INSPIRE](#)].
- [38] T. Eguchi and A.J. Hanson, *Asymptotically Flat Selfdual Solutions to Euclidean Gravity*, *Phys. Lett. B* **74** (1978) 249 [[INSPIRE](#)].